Published in final edited form as:

Sleep Med. 2018 December ; 52: 75–79. doi:10.1016/j.sleep.2018.07.014.

Last Night's Sleep in Relation to Academic Achievement and Neurocognitive Testing Performance in Adolescents with and without ADHD

Caroline N. Cusick, BS¹, Paul A. Isaacson, MA¹, Joshua M. Langberg, PhD², and Stephen P. Becker, PhD^{1,3,4}

¹Division of Behavioral Medicine and Clinical Psychology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

²Department of Psychology, Virginia Commonwealth University, Richmond, VA, USA

³Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

Abstract

Objectives/Background: Objectives were to (1) examine previous night's sleep in relation to next day performance on standardized academic achievement and neurocognitive assessments in adolescents, and (2) explore whether previous night's sleep is differentially associated with testing performance for adolescents with and without attention-deficit/hyperactivity disorder (ADHD).

Participants/Methods: Participants were 300 adolescents (ages 12–14 years; 55% male). Approximately half (53.6%) were diagnosed with ADHD. Adolescents provided ratings of their previous night's sleep quality, sleep duration, and number of night wakings and were administered standardized tests of processing speed and working memory, as well as word reading, numerical operations, and math fluency academic achievement.

Results: In analyses controlling for sex, race, medication use, time of testing, and ADHD group status, more night wakings the previous night were associated with significantly lower numerical operations and math fluency achievement scores and marginally lower working memory scores. Previous night's sleep was not associated with processing speed or reading achievement. ADHD status did not moderate sleep in relation to academic/neurocognitive performance. Participants reporting 2 night wakings the previous night had slightly over half a standard deviation lower scores on average compared to participants reporting 0 night wakings.

Conclusions: This preliminary study suggests that previous night's wakings are associated with poorer mathematics performance the next day, regardless of ADHD status. This may be due to the detrimental effect of interrupted and fragmented sleep on attention and executive control. These

⁴Address correspondence to Stephen P. Becker, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA. Phone: 513-803-2066. stephen.becker@cchmc.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

findings have implications for clinicians and educators when considering contextual factors that may impact academic and neurocognitive testing performance.

Keywords

adolescence; attention-deficit/hyperactivity disorder; math; processing speed; reading; sleep; working memory

Introduction

Sleep problems negatively affect cognitive and academic functioning in youth (Gradisar et al., 2008; Wolfson & Carskadon, 2003). For example, studies examining objective measures of sleep have found that sleep deprivation can also impact psychomotor vigilance and working memory (de Bruin et al., 2017). Perhaps due to rapid developmental changes in the prefrontal cortex, poor sleep is particularly detrimental for academic performance in early adolescence (Dewald et al., 2010). However, most studies have looked broadly at sleep in relation to school performance and relied on self-report ratings of performance (Dewald et al., 2010). There is some indication from nonclinical samples that school performance (Gillen-O'Neel et al., 2013), working memory (Könen et al., 2015), and psychomotor vigilance (Neylan et al., 2010) are impacted by the prior night's sleep. However, several of these studies were conducted in school-aged children (Könen et al., 2015) or adults (Neylan et al., 2010), leaving it largely unclear whether previous night's sleep impacts testing performance the following day for adolescents, a population that frequently obtains insufficient or poor sleep (Crowley et al., 2018), as well as increased intraindividual variability of sleep/wake patterns (i.e., night-to-night variability) (Becker et al., 2017). Examining prior night's sleep in relation to next day's testing performance has important clinical implications for psychoeducational and neuropsychological assessment.

The association between previous night's sleep and testing performance may vary based upon the characteristics of the youth being evaluated. Adolescents with attention-deficit/hyperactivity disorder (ADHD) have poorer sleep than their typically developing peers (Becker, in press; Lunsford-Avery et al., 2016) and are also frequently referred for testing and re-evaluations linked to school services (e.g., Individual Education Plans [IEPs]). Sleep quality, sleep duration, and number of night wakings have been identified in previous research with school-aged children as sleep domains associated with cognitive functioning (Könen et al., 2015; Gruber et al., 2010; Sadeh et al., 2002). However, no study has examined the previous night's sleep in relation to next day test performance in adolescents with ADHD. Accordingly, the present study (1) examined how previous night's sleep (duration, quality, and wakings) is associated with next day performance on academic achievement and neurocognitive assessments in adolescents, and (2) explored whether previous night's sleep was differentially associated with performance for adolescents with and without ADHD.

Methods

Participants

Participants were 300 adolescents (165 male, 135 female) in eighth grade (ages 12–14 years). The original sample consisted of 302 participants, however, two participants that either completed testing at a different location or did not complete the last night's sleep items at the study visit were excluded from analyses. Approximately half of the participants (53.6%; $n=160$) were diagnosed with ADHD. Most participants were White (81.7%), with remaining participants Black (5.3%), Asian (4.7%), American Indian/Alaska Native (0.3%), or Biracial/Multiracial (8.0%); 4.7% of participants identified as Hispanic/Latino. All participants had an estimated IQ ≥ 80 (Range=81–148; $M \pm SD = 107.00 \pm 13.42$) based on the *Wechsler Abbreviated Scale of Intelligence, Second Edition* (WASI-II; Wechsler, 2011).

Procedures

All procedures were approved by an Institutional Review Board. Participants were recruited during the fall semester of their eighth-grade year at two sites in the United States. Parents who contacted the investigators in response to recruitment activities through local schools completed an eligibility screening. Those meeting the screening criteria were scheduled for an evaluation to determine eligibility. We aimed to recruit an approximately equal number of adolescents with and without ADHD. To be eligible for the ADHD group, participants had to meet full DSM-5 diagnostic criteria for either ADHD Predominantly Inattentive Presentation or ADHD Combined Presentation based on the *Parent Children's Interview for Psychiatric Syndromes* (P-ChIPS; Weller, Weller, Fristad, Rooney, & Schecter, 2000). To be eligible for the comparison group, participants were required to have fewer than four symptoms of ADHD in each domain (i.e., inattention and hyperactivity/impulsivity) on the P-ChIPS. To be eligible for either group, participants had to have an estimated IQ ≥ 80 , take core classes in a regular education setting, and could not have a diagnosis of autism, bipolar disorder, a dissociative disorder, a psychotic disorder, or an organic sleep disorder per parent report on a phone screen.

Measures.—Adolescents completed a form created for this study about their sleep the night before the assessment visit. Specifically, adolescents were asked to provide an estimate of their sleep duration as well as ratings of sleep quality and night wakings. The sleep duration item (“*How many hours of actual sleep did you get last night? This number may be different than the number of hours you spent in bed*”) was queried in an open-ended format. Sleep quality (“*How would you rate last night's sleep overall?*”) was rated on a five-point scale (0=Very good; 1=Fairly good; 2=Okay; 3=Fairly bad; 4=Very bad). Night wakings (“*After falling asleep, how many times did you wake up during the night last night?*”) was assessed on a six-point scale (0; 1; 2; 3; 4; 5 or more). Additionally, participants were asked whether (yes/no) they took any medication (e.g., medication for ADHD, another psychiatric disorder, allergies, headache) on the day of testing (“*Did you take any medication today?*”) and this variable was used as a covariate in analyses.

Adolescents were administered the Word Reading, Numerical Operations, and Math Fluency subtests of the *Wechsler Individual Achievement Test–Third Edition* (WIAT-III; Wechsler,

2009) to measure academic achievement. The Processing Speed and Working Memory scales of the *Wechsler Intelligence Scale for Children–Fourth Edition* (WISC-IV; Wechsler, 2003) were also administered. See Supplemental Materials for additional details regarding these tests and evidence for their reliability and validity.

Analyses

Intercorrelations among study variables were conducted. Hierarchical regression analyses were then run to test whether previous night's sleep was associated with test scores above and beyond sex, race, ADHD group status, medication use on the day of testing, and whether testing was conducted during school hours or after 4pm. All covariates were entered on Step 1, followed by the three sleep variables on Step 2. We also explored whether ADHD status moderated any associations between sleep and test performance. Finally, for significant effects, we examined standardized test scores at varying levels of sleep difficulties using one-way analysis of variance (ANOVA).

Results

Descriptive statistics and intercorrelations among study variables are provided in Table S1. Neither sleep duration nor sleep quality were significantly correlated with any of the testing outcomes, though more night wakings was significantly correlated with lower working memory ($p=.02$), numerical operations ($p=.002$), and math fluency ($p<.001$) scores.

Results of the regression analyses are summarized in Table 1. Having an ADHD diagnosis remained significantly associated with all testing outcomes ($ps<.001$). Above and beyond covariates and the variance accounted for by sleep duration and sleep quality, night wakings remained associated with significantly lower numerical operations ($p=.02$) and math fluency ($p<.001$) scores, and marginally associated with lower working memory scores ($p=.07$). None of the last night's sleep domains were associated with processing speed or basic reading performance. Further, ADHD status did not interact with any of the sleep domains in predicting test scores¹.

To further examine the impact of night wakings on mathematics achievement (numerical operations and math fluency), we examined test scores for participants reporting 0, 1, or 2 night wakings the previous night ($ns = 172, 76$, and 52 , respectively). Results are displayed in Figure 1, with the full sample used for analyses since there was no interaction by ADHD group status (though given the main effect of group, separate ADHD and comparison lines are also shown). For numerical operations, participants reporting 0, 1, or 2 night wakings had mean (SD) standard scores of 104.91 (19.65), 101.36 (19.57), and 96.54 (17.45), respectively. The only significant difference was between participants reporting 0 and 2 night wakings ($p=.02$). Participants reporting 0 night wakings had a score on average 8.37 points higher than participants reporting 2 night wakings, a standard score difference of over ½ SD. For math fluency, participants reporting 0, 1, or 2 night wakings had mean

¹Although the objective of our study was to examine whether ADHD moderated the association between previous night's sleep and testing performance, we also explored whether sex moderated this association, as well as the three-way interaction of sleep, sex, and ADHD status. No evidence was found for sex as a moderator or for a three-way interaction in relation to test scores.

(SD) standard scores of 99.24 (16.48), 93.29 (13.17), and 91.10 (12.85), respectively. Participants reporting 0 night wakings had significantly higher math fluency scores than participants reporting either 1 or 2 night wakings (p s=.01 and .002, respectively). Participants reporting 0 night wakings had a score on average 5.95 points higher than participants reporting 1 night waking and 8.14 points higher than participants 2 night wakings, a standard score difference of .40 and .54 SD, respectively.

Discussion

This study examined previous night's sleep in relation to academic and neurocognitive performance the following day in a large sample of adolescents with and without ADHD. We found that more night wakings were significantly associated with lower math achievement scores, with results consistent across both numerical operations and math fluency domains after controlling for a number of relevant variables including sex, ADHD diagnostic status, medication use, and time of the assessment visit. More night wakings were also marginally associated with lower working memory scores. None of these effects were moderated by ADHD diagnostic status, indicating that previous night's sleep impacts performance similarly for adolescents with and without ADHD.

Our findings are in consistent with previous research linking sleep to poorer neurocognitive and academic performance (de Bruin et al., 2017). Particularly in line with our findings, Sadeh and colleagues (2002) found that fragmented sleep "characterized by multiple and/or prolonged night-wakings" (p. 405) may be particularly detrimental for neurobehavioral functioning. Participants in our study who reported more night wakings may have had less opportunity for deeper, more restorative sleep (Sadeh et al., 2002). In contrast, we did not find previous night's sleep duration or sleep quality to be associated with any of the performance domains examined in this study. A recent meta-analysis found that sleep duration was not associated with several aspects of cognitive functioning including processing speed and memory (Short et al., 2018). Further, Gruber et al. (2014) found lower actigraphy-measured sleep efficiency (which is in part calculated based on night wakings) but not sleep duration to be associated with lower grades in healthy school-aged children. Previous studies have linked poorer sleep quality to adolescents' school and achievement performance (Buckhalt et al., 2009; Meijer et al., 2000), making this an important area for further study. Also, none of the sleep domains assessed in this study were associated with processing speed or basic reading. Thus, at least of the sleep domains assessed in this study, more night wakings had the clearest effect on testing performance and this association was clearest for math achievement. Mathematics requires a set of complex cognitive skills, including abstract and logical thinking and the ability to identify patterns in novel situations (see Gruber et al., 2014), and sleep fragmentation related to night wakings may have a particularly detrimental impact on these higher-order cognitive skills.

Strengths of the present study include a large sample, including adolescents with and without ADHD, and multiple measures of academic and neurocognitive performance. However, future research should incorporate objective measures of sleep, such as actigraphy or polysomnography, as well as more nuanced items regarding sleep quality and medication type/dose on the evening preceding testing and the day of testing. It would also be

informative for future studies to formally assess sleep disorders per *International Classification of Sleep Disorders—Second Edition (ICSD-2)*. Additionally, longer periods of sleep assessment may shed light on how cumulative sleep loss, rather than simply previous night's sleep, impacts testing performance. Future studies should also incorporate more comprehensive batteries of academic achievement and neurocognition. Nevertheless, our findings indicate that previous night's sleep, and night wakings specifically, may have important clinical implications. Specifically, contextual factors, such as previous night's sleep, may be important to assess and consider when administering and interpreting testing evaluations. This is especially important in populations that are frequently referred for assessments and re-evaluations that are often used to determine school services.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement:

This research was supported by award number R305A160126 from the Institute of Education Sciences (IES), U.S. Department of Education. Stephen Becker is also supported by award number K23MH108603 from the National Institute of Mental Health (NIMH). The content is solely the responsibility of the authors and does not necessarily represent the official views of the U.S. Department of Education or the National Institutes of Health (NIH).

References

- Becker SP (in press). The triple threat of sleep, adolescence, and ADHD. Hiscock H & Sciberras E (Eds.), *Sleep and ADHD* Amsterdam, Netherlands: Elsevier.
- Becker SP, Sidol CA, Van Dyk TR, Epstein JN, & Beebe DW (2017). Intraindividual variability of sleep/wake patterns in relation to child and adolescent functioning: A systematic review. *Sleep Medicine Reviews*, 34, 94–121. <https://doi.org/10.1016/j.smrv.2016.07.004> [PubMed: 27818086]
- Buckhalt JA, El-Sheikh M, Keller PS, & Kelly RJ (2009). Concurrent and longitudinal relations between children's sleep and cognitive functioning: The moderating role of parent education. *Child Development*, 80, 875–892. <https://doi.org/10.1111/j.1467-8624.2009.01303.x> [PubMed: 19489909]
- Crowley SJ, Wolfson AR, Tarokh L, & Carskadon MA (2018). An update on adolescent sleep: New evidence informing the perfect storm model. *Journal of Adolescence*, 67, 55–65. <https://doi.org/10.1016/j.adolescence.2018.06.001> [PubMed: 29908393]
- de Bruin EJ, van Run C, Staaks J, & Meijer AM (2017). Effects of sleep manipulation on cognitive functioning of adolescents: A systematic review. *Sleep Medicine Reviews*, 32, 45–57 <https://doi.org/10.1016/j.smrv.2016.02.006> [PubMed: 27039223]
- Dewald JF, Meijer AM, Oort FJ, Kerkhof GA, & Bögels SM (2010). The influence of sleep quality, sleep duration and sleepiness on school performance in children and adolescents: A meta-analytic review. *Sleep Medicine Reviews*, 14, 179–189. <https://doi.org/10.1016/j.smrv.2009.10.004> [PubMed: 20093054]
- Gillen-O'Neel C, Huynh VW, & Fuligni AJ (2013). To study or to sleep? The academic costs of extra studying at the expense of sleep. *Child Development*, 84, 133–142. <https://doi.org/10.1111/j.1467-8624.2012.01834.x> [PubMed: 22906052]
- Gradisar M, Terrill G, Johnston A, & Douglas P (2008). Adolescent sleep and working memory performance. *Sleep and Biological Rhythms*, 6, 146–154. <https://doi.org/10.1111/j.1479-8425.2008.00353.x>
- Gruber R, Lavolette R, Deluca P, Monson E, Cornish K, & Carrier J (2010). Short sleep duration is associated with poor performance on IQ measures in healthy school-age children. *Sleep Medicine*, 11, 289–294. <https://doi.org/10.1016/j.sleep.2009.09.007> [PubMed: 20156702]

- Gruber R, Somerville G, Enros P, Paquin S, Kestler M, & Gillies-Poitras E (2014). Sleep efficiency (but not sleep duration) of healthy school-age children is associated with grades in math and languages. *Sleep medicine*, 15, 1517–1525. <https://doi.org/10.1016/j.sleep.2014.08.009> [PubMed: 25441747]
- Köner T, Dirk J, & Schmiedek F (2015). Cognitive benefits of last night's sleep: Daily variations in children's sleep behavior are related to working memory fluctuations. *Journal of Child Psychology and Psychiatry*, 56, 171–182. <https://doi.org/10.1111/jcpp.12296> [PubMed: 25052368]
- Lunsford-Avery JR, Krystal AD, & Kollins SH (2016). Sleep disturbances in adolescents with ADHD: A systematic review and framework for future research. *Clinical Psychology Review*, 50, 159–174. <https://doi.org/10.1016/j.cpr.2016.10.004> [PubMed: 27969004]
- Meijer AM, Habekothé HT, & Van Den Wittenboer GLH (2000). Time in bed, quality of sleep and school functioning of children. *Journal of Sleep Research*, 9, 145–154. 10.1046/J.1365-2869.2000.00198.X [PubMed: 10849241]
- Neylan TC, Metzler TJ, Henn-Haase C, Blank Y, Tarasovsky G, McCaslin SE, ... & Marmar CR. (2010). Prior night sleep duration is associated with psychomotor vigilance in a healthy sample of police academy recruits. *Chronobiology International*, 27, 1493–1508. <https://doi.org/10.3109/07420528.2010.504992> [PubMed: 20795888]
- Sadeh A, Gruber R, & Raviv A (2002). Sleep, neurobehavioral functioning, and behavior problems in school-age children. *Child Development*, 73, 405–417. <https://doi.org/10.1111/1467-8624.00414> [PubMed: 11949899]
- Short MA, Blunden S, Rigney G, Matricciani L, Coussens S, Reynolds CM, & Galland B (2018). Cognition and objectively measured sleep duration in children: A systematic review and meta-analysis. *Sleep Health*, 4, 292–300. <https://doi.org/10.1016/j.sleh.2018.02.004> [PubMed: 29776624]
- Wechsler D (2011). *Wechsler Abbreviated Scale of Intelligence Second Edition (WASI-II)* San Antonio, TX: Pearson.
- Wechsler D (2009). *Wechsler Individual Achievement Test - Third Edition (WIAT-III)* (3rd ed.). San Antonio, TX: Pearson.
- Wechsler D (2003). *Wechsler Intelligence Scale for Children - Fourth Edition (WISC-IV)* (Fourth Edition ed.). San Antonio, TX: Psychological Corporation.
- Weller EB, Weller RA, Fristad MA, Rooney MT, & Schecter J (2000). Children's interview for psychiatric syndromes (ChIPS). *Journal of the American Academy of Child and Adolescent Psychiatry*, 39, 76–84. [doi:10.1097/00004583-200001000-00019](https://doi.org/10.1097/00004583-200001000-00019) [PubMed: 10638070]
- Wolfson AR, & Carskadon MA (2003). Understanding adolescent's sleep patterns and school performance: A critical appraisal. *Sleep Medicine Reviews*, 7, 491–506. [https://doi.org/10.1016/S1087-0792\(03\)90003-7](https://doi.org/10.1016/S1087-0792(03)90003-7) [PubMed: 15018092]

Highlights

- More night wakings the previous night were associated with lower math achievement scores.
- Night wakings were marginally associated with lower working memory scores.
- Previous night's sleep was not associated with processing speed or reading achievement.
- ADHD status did not moderate sleep in relation to academic/neurocognitive performance.
- Assessing and considering previous night's sleep may be important in testing evaluation.

Figure 1. Standardized scores on numerical operations (top panel) and math fluency (bottom panel) between participants reporting zero, one, or two or more night wakings the previous night

Table 1
Hierarchical regression analyses examining previous night's sleep in relation to neurocognitive and academic achievement test scores

	<i>Step 1 Model Summary</i>				<i>Step 2 Model Summary</i>			
	<i>b</i>	<i>SE</i>	β	<i>t</i>	<i>b</i>	<i>SE</i>	<i>B</i>	<i>T</i>
<i>DV: WISC Processing Speed</i>								
Time of testing	.74	1.66	.02	0.44	.82	1.67	.03	0.49
Sex	4.84	1.56	.17	3.11**	4.88	1.56	.17	3.12**
Race	.23	1.95	.01	0.12	-.14	1.97	-.004	-.07
Medication use	3.91	1.86	.13	2.10*	3.89	1.87	.13	2.08*
ADHD status	-11.47	1.73	-.41	-6.62***	-11.12	1.75	-.39	-6.34***
Sleep duration	--	--	--	--	.76	.54	.08	1.40
Sleep quality	--	--	--	--	1.32	1.02	.08	1.29
Night wakings	--	--	--	--	-.82	.72	-.07	-1.14
<i>DV: WISC Working Memory</i>								
Time of testing	3.78	1.76	.12	2.16*	3.91	1.76	.13	2.23*
Sex	-.12	1.64	-.004	-0.07	.04	1.65	.001	0.02
Race	4.86	2.05	.13	2.37*	4.24	2.08	.12	2.04*
Medication use	1.12	1.97	.04	0.57	.99	1.97	.03	0.50
ADHD status	-8.50	1.83	-.30	-4.65***	-7.98	1.85	-.28	-4.32***
Sleep duration	--	--	--	--	.61	.57	.06	1.07
Sleep quality	--	--	--	--	1.19	1.08	.07	1.10
Night wakings	--	--	--	--	-1.36	.76	-.11	-1.79 [†]
<i>DV: WIAT Basic Reading</i>								
Time of testing	3.85	1.48	.15	2.60**	3.94	1.48	.15	2.66**
Sex	-2.61	1.38	-.11	-1.89	-2.48	1.39	-.11	-1.78
Race	.47	1.73	.02	0.27	.03	1.76	.001	0.02
Medication use	1.27	1.66	.05	0.76	1.12	1.66	.04	0.68
ADHD status	-6.29	1.54	-.27	-4.08***	-5.97	1.56	-.25	-3.82***
Sleep duration	--	--	--	--	.13	.48	.02	0.27
Sleep quality	--	--	--	--	.55	.91	.04	0.61
Night wakings	--	--	--	--	-.96	.64	-.09	-1.50
<i>DV: WIAT Numerical Operations</i>								
Time of testing	-4.19	2.36	-.10	-1.78	-4.05	2.35	-.10	-1.73
Sex	-.94	2.20	-.02	-0.43	-.53	2.20	-.01	-0.24
Race	1.12	2.76	.02	0.41	.02	2.78	.000	.01
Medication use	5.30	2.64	.12	2.01*	5.07	2.63	.12	1.92
ADHD status	-15.45	2.45	-.40	-6.30***	-14.48	2.47	-.37	-5.86***

	<i>Step 1 Model Summary</i>				<i>Step 2 Model Summary</i>			
	<i>b</i>	<i>SE</i>	β	<i>t</i>	<i>b</i>	<i>SE</i>	<i>B</i>	<i>T</i>
Sleep duration	--	--	--	--	.73	.76	.05	.95
Sleep quality	--	--	--	--	.80	1.44	.03	0.56
Night wakings	--	--	--	--	-2.34	1.02	-.14	-2.30 [*]
<i>DV: WIAT Math Fluency</i>								
Time of testing	-2.59	1.89	-.08	-1.37	-2.41	1.86	-.07	-1.30
Sex	-4.30	1.76	-.14	-2.44 [*]	-3.79	1.75	-.12	-2.17 [*]
Race	-2.29	2.21	-.06	-1.04	-3.56	2.20	-.09	-1.62
Medication use	1.92	2.11	.06	0.91	1.56	2.09	.05	0.75
ADHD status	-11.21	1.97	-.36	-5.70 ^{***}	-10.18	1.96	-.33	-5.19 ^{***}
Sleep duration	--	--	--	--	.33	.60	.03	0.55
Sleep quality	--	--	--	--	.45	1.14	.02	0.40
Night wakings	--	--	--	--	-2.71	.81	-.20	-3.36 ^{***}

Note. For sex, 0 = male, 1 = female. For race, 0 = non-White, 1 = White. For medication use, 0 = no medication taken on day of testing, 1 = medication taken on day of testing. For time of testing, 0 = testing appointment occurred before 4pm, 1 = testing appointment occurred at 4pm or later. For ADHD status, 0 = comparison, 1 = attention-deficit/hyperactivity disorder (ADHD). WIAT = Wechsler Individual Achievement Test, Third Edition. WISC = Wechsler Intelligence Scale for Children, Fourth Edition.

[†]
 $p = .07$.

^{*}
 $p < .05$.

^{**}
 $p < .01$.

^{***}
 $p < .001$.