Genetic analysis of robustness in meat sheep through body weight and body condition score changes over time¹

Tiphaine Macé,^{†,2} Eliel González-García,^{‡,⊕} Julien Pradel,^{||} Sara Parisot,^{||} Fabien Carrière,^{||} Sebastien Douls,^{||} Didier Foulquié,^{||} and Dominique Hazard[†]

[†]INRA UMR1388 GENPHYSE, 31326 Castanet-Tolosan, France; [‡]SELMET, INRA, CIRAD, Montpellier SupAgro, Univ Montpellier, 34060 Montpellier, France; and ^{||}INRA UE321 La Fage, 12250 Roquefort-sur-soulzon, France

ABSTRACT: Animal robustness may be defined as a complex trait characterizing the ability of an individual to be adapted, productive and healthy under contrasted and fluctuating environmental situations. Such a trait is now considered an essential criterion in order to meet the more ambitious goals of farming sustainability. In ruminants, one of the key mechanisms via which robustness is expressed is the capacity to mobilize or restore body reserves (**BR**) to cope with the challenges of negative energy balances. The objectives of this work were as follows: 1) to estimate the genetic parameters related to **BR** dynamics in ewes over successive production cycles and 2) to investigate **BR** management relationships between different physiological stages. For this, historical individual BW and BCS data from 2,920 phenotyped ewes were used for genetic analysis. The changes in BW (Δ BW) and BCS (Δ BCS) over time were analyzed. Eight physiological stages were considered to investigate these changes over time: mating, early pregnancy, mid-pregnancy, lambing, early suckling, mid-suckling, weaning, and

postweaning. The estimated heritability were low for both Δ BW ($h^2 = 0.13$ to 0.18) and Δ BCS ($h^2 = 0.04$ to 0.16). Moderate to high positive genetic correlations (from 0.48 to 0.91) were obtained between **BR** mobilization phases and between **BR** accretion phases. Similarly, moderate to high negative genetic correlations (from -0.36 to -0.75) were estimated between the **BR** mobilization and accretion periods, suggesting that mechanisms driving **BR** mobilization and accretion processes were genetically correlated. This is the first study in ruminants that demonstrate that the extent and temporal changes in profiles of **BR** mobilization and accretion are heritable and genetically linked, indicating that such traits could be considered in genetic programs aimed at improving robustness. Nevertheless, further research is needed for a more comprehensive understanding of **BR** dynamics, notably by including other physiological parameters (i.e., metabolites and hormones) and additional information on the productive and reproductive life of the ewe.

Key words: body reserves, quantitative genetics, genetic determinism, heritability, Romane sheep

© The Author(s) 2018. Published by Oxford University Press on behalf of the American Society of Animal Science. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

J. Anim. Sci. 2018.96:4501–4511
doi: 10.1093/jas/sky318

¹The authors would like to thank the INRA Divisions of Animal Genetics and Animal Physiology and Livestock Systems for their financial support over the years in providing salaries, experimental costs, and computational facilities. The authors are indebted to all the staff of La Fage experimental

farm for animal management and care as well as for their key role in the experimental data collection.

²Corresponding author: tiphaine.mace@inra.fr

Received May 15, 2018.

Accepted August 1, 2018.

INTRODUCTION

In the future, breeding strategies will include more robust animals in order to improve the sustainability of farming systems (De La Torre et al., 2015; Friggens et al., 2017). It is expected that ruminants will depend more and more on grazing land in harsh environments, which will expose them to more uncertain feed resources due to the unpredictability of the production environment (O'Mara, 2012). Therefore, improving the individual biological capacities of adaptation to such environments is a priority. In that regard, an optimal management of body reserves (BR; i.e., lipid mobilization and accretion processes), an indicator of individual metabolic plasticity (Friggens et al., 2004; Blanc et al., 2006), is considered a relevant trait for these purposes (Phocas et al., 2016). The combined measurement of the BCS and BW has been used in previous studies to analyze the genetic variability of body condition (Edmonson et al., 1989; Arango et al., 2002; Safari et al., 2005; Gizaw et al., 2007; Shackell et al., 2011). The genetic determinism of BCS, however, has only been investigated using single time-point measurements. In the present study, we propose to use variations of BW and BCS over time (i.e., gain or loss of BW and BCS at key physiological stages) to assess BR in a dynamic management perspective while studying its genetic determinism over successive productive cycles with a multitrait approach. Such variations of BW and BCS reflect the biological ability of the ewe to use BR, more or less efficiently, when facing alternating situations (challenges) leading to positive or negative energy balances. To develop this approach, which to the best of our knowledge has not been addressed in previous work, we used historical BW and BCS data (2002 to 2015) recorded longitudinally in Romane meat ewes reared under extensive rangeland conditions. Our objectives were as follows: 1) to analyze, by assessing BW (Δ BW) and BCS (Δ BCS) changes over time, the genetic components of BR dynamics in ewes over successive production cycles and 2) to investigate potential relationships between the BR dynamics and different physiological stages. We hypothesized the existence of the genetic determinism of BR dynamics in sheep as well as related effects between stages and productive cycles.

MATERIALS AND METHODS

Animals and Experimental Farming System

The experiments described here fully comply with applicable legislation on research involving animal subjects in accordance with the European

Union Council directive (2010/63/UE). The researchers carrying out the experiments were certified by the relevant French governmental authority. All experimental procedures were approved and performed under the guidelines for the care and use of experimental animals stated in the ethics policy of the French Ministry of Agriculture.

The experimental animals were *Romane* sheep, a composite line obtained from *Romanov* and *Berrichon du Cher* breeds (Ricordeau et al., 1992). The ewes were reared exclusively outdoor on about 280 ha of rangelands at the INRA Experimental Farm *La Fage* in a flock comprising 250 reproductive females (*Causse du Larzac* 43°54'54.52"N; 3°05'38.11"E; approximately 800 m.a.s.l, Roquefort-sur-Soulzon, Aveyron, south of France). The rangeland is on a limestone plateau and is composed of about 25% shrubs and 75% grass such as *Bromus erectus*, *Stipa pennata*, *Carex humilis*, and *Festuca duriscola*. The rangeland of this farm is divided into "native" (nonfertilized) and fertilized paddocks used during the spring (from lambing to weaning; Molénat et al., 2005). From 2002 to 2015, the average annual temperature was 9.8 °C and the average annual precipitation was 910 mm. During the summer, which is hot and dry (average temperature of 17.8 °C and average rainfall for the season of 142 mm, from June to August), the grass stops growing and regreens in the beginning of autumn. In the winter (average temperature of 2.14 °C and average rainfall of 183 mm, from December to February), the flock is supplemented with hay or silage. Further details regarding the climatic conditions and overall management practices established in this system were reported by Molénat et al. (2005), González-García et al. (2014), and González-García and Hazard (2016).

Before 2010, females were mated at 7 or 19 mo of age depending on their BW. After 2010, all first matings were performed at 19 mo old. Mating occurred in the autumn to obtain a peak of lambing at the beginning of spring (usually mid-April; González-García et al., 2014). *Romane* ewes produced on average 2.2 live lambs per lambing in the period comprised for this study. Lambs were weaned at approximately 2.5 mo (i.e., 75 d). In these experimental conditions, the culling rate considered an annual replacement of 30% of the ewes.

Historical Data and Variables

The BW and BCS measurements used in this study were collected regularly during each female's productive cycle according to a physiological stage schedule. A maximum of 2,920 records per trait were

registered over a 14-yr period (2002 to 2015) with 1,146 females in cycle 1 (i.e., first production cycle, from first to second mating), 1,068 in cycle 2, and 707 in cycle 3 and more. To assess BCS, the original grid described by Russel et al. (1969) was used and subdivided into a 1/10 scale, i.e., a scale from 1 to 5 with 0.1 increments. The BCS measurements were performed by the 2 same operators throughout the entire 14-yr period of this study, with regular training sessions for adjustments and calibration. All measurements were recorded in the Geedoc database (<https://germinal.toulouse.inra.fr/~mcbatut/GEEDOC/>) as well as other additional pedigree information and animal production performance data.

To analyze individual performances related to BR mobilization or accretion, the differences in BW or BCS between 2 physiological stages were calculated and used for interpretation as follows:

$$BW_{i-j} = BW_j - BW_i [I]; BCS_{i-j} = BCS_j - BCS_i [I]$$

where i and j are 2 successive physiological stages.

The intervals were chosen considering key physiological stages which represented periods of BR mobilization and accretion throughout a typical production cycle. Three intervals were defined

in the BR mobilization period: 1) difference between mid-pregnancy (Pb; 80 ± 14 days after mating [DAM], in January) for BW or early pregnancy for BCS (Pa; 40 ± 15 DAM, in December) and lambing (L; 160 ± 16 DAM, in April) (**BW-Pb:L**; **BCS-Pa:L**); 2) difference between lambing and early suckling (Sa; 190 ± 14 DAM, in April) (**BW-L:Sa**; **BCS-L:Sa**); and 3) difference between pregnancy and weaning (W; 250 ± 16 DAM, in June) (**BW-Pb:W**; **BCS-Pa:W**). Three intervals were also defined for the BR accretion period: 1) difference between mating (M; 15 d before mating, in November) and pregnancy (**BW-M:Pb**; **BCS-M:Pa**); 2) difference between weaning and postweaning (Wp; 310 ± 12 DAM, in August) (**BW-W:Wp**; **BCS-W:Wp**); and 3) difference between W and M (**BW-W:M**; **BCS-W:M**) (Figure 1). These intervals were selected in order to illustrate short- and long-term variations of BW and BCS during the BR mobilization or accretion periods throughout each productive cycle.

Descriptive Statistics

Deviations from normality were inspected using the UNIVARIATE procedure of SAS (version 9.4;

Figure 1. Average of body weight (A) and body condition score (B) over a productive cycle. Values are raw means for all ewes and parities involved in the study and for each time point. Each color represents a specific period defined to estimate variation of BW or BCS.

SAS Institute Inc., Cary, NC). None of the variables were transformed given a major deviation from normality was not observed. Analyses of variance taking into account the repeated measures (MIXED procedure of SAS) were used to test the relevant effects and interactions in order to determine factors of variation for BW and BCS. These analyses resulted in the identification of the following significant fixed effects: the Age at first lambing and the Parity of the ewe, the Litter size, and the Year of measurement. The Age effect took into account ewes lambing for the first time at 1 or 2 yr old (classes 1 and 2, respectively). The Parity effect took into account first, second, and third or more lambing (classes 1, 2, and 3, respectively). The Litter effect was classified according to the number of lambs born that remained with the dam during the S stage (i.e., class 1, singletons from L until W; class 2, ewes lambing twins and suckling one; class 3, ewes lambing and suckling twins; and class 4, ewes lambing and suckling more than 2 lambs). The litter effect considered only the litter of the corresponding cycle and not of the previous cycle. Finally, 14 yr were analyzed. The first-order interactions between Age $\times$ Litter and Parity $\times$ Litter were tested. An effect was considered significant if $P < 0.05$.

Genetic Analyses

The variance components for Δ BW and Δ BCS were estimated by restricted maximum-likelihood methodology applied to an animal model using ASREML 1.0 software (Gilmour et al., 2001). Age, Parity, Litter, and Year were considered as fixed effects. Random effects included additive genetic and permanent environmental effects of the ewe. Analyses assumed a repeatability model with measurements across productive cycles considered to be the same traits with constant variances. The following animal mixed model was fitted as follows:

$$y = X\beta + Z_a a + W_c c + e \quad [\text{II}]$$

where y is the vector of observations corresponding to the trait(s) in the analysis; β is the vector of fixed effects; and a and c are the vectors of random ewe additive genetic and permanent environmental effects with incidence matrices X , Z_a , and W_c , respectively, and e is the vector of residual effects. The following (co)variance structure of random effects was assumed as follows:

$$\text{Var} \begin{bmatrix} a \\ c \\ e \end{bmatrix} = \begin{bmatrix} \mathbf{G}_a \otimes \mathbf{A} & 0 & 0 \\ 0 & \mathbf{P}_c \otimes \mathbf{I} & 0 \\ 0 & 0 & \mathbf{R} \otimes \mathbf{I} \end{bmatrix}$$

where $\mathbf{G}_a$ is a (co)variance matrix for direct additive genetic effects; $\mathbf{A}$ is the numerator relationship matrix; $\mathbf{P}_c$ is a (co)variance matrix for the ewe permanent environmental effects; $\mathbf{R}$ is a (co)variance matrix for residual effects; $\mathbf{I}$ are identity matrices of appropriate size; and $\otimes$ is the direct matrix product.

Univariate analyses were performed to estimate variances for each trait. Bivariate analyses were performed to estimate genetic and phenotypic correlations between traits. Variance estimates in the 2-trait analyses were very similar to those from single-trait analyses. From the variance components, 3 parameters were defined as follows: 1) heritability or proportion of total phenotypic variance attributed to the additive genetic effect, $h^2 = \sigma_a^2 / (\sigma_a^2 + \sigma_c^2 + \sigma_e^2)$; 2) proportion of total phenotypic variance attributed to the permanent environmental effect, $c^2 = \sigma_c^2 / (\sigma_a^2 + \sigma_c^2 + \sigma_e^2)$; and 3) proportion of total phenotypic variance attributed to the residual effect, $e^2 = \sigma_e^2 / (\sigma_a^2 + \sigma_c^2 + \sigma_e^2)$. In addition, repeatability (r) was defined as the sum of h^2 and c^2 .

RESULTS

Biological and Year of Measurement Effects

The differences in BW and BCS, Δ BW and Δ BCS, respectively, calculated for the 1,146 ewes over several productive cycles were significantly affected by the Parity and the Age at first lambing of the ewe, the Litter and the Year (Table 1). These fixed factors had significant ($P < 0.01$) effects on Δ BW, except for Age over the BW-W:Wp period. Globally, a significant ($P < 0.01$) increase in the BW loss was observed for BW-Pb:W, BW-Pb:L and a decrease in the BW loss was observed for BW-L:Sa between parities 1 and 2. An increase was also observed in the BW gain for BW-M:Pb and BW-W:M with parity, whereas a decrease was observed for the BW gain during BW-W:Wp (Table 2). Similarly, the BCS loss observed for BCS-Pa:W and BCS-Pa:L decreased with parity, whereas an increase was observed in the BCS gain for BCS-M:Pa between parities 1 and 2. A decrease in the BCS gain was observed with parity for BCS-W:M and for BCS-W:Wp between parities 2 and 3, and parity had no effect for the period BCS-L:Sa. The Litter effect was significant ($P < 0.001$) for all the Δ BCSs evaluated, except for BCS:M-Pa (Table 1). Overall, greater BW gain for the physiological periods BW-M:Pb, BW-W:Wp, and BW-W:M and greater BW loss for BW-Pb:W and BW-Pb:L were observed for larger litter sizes (Table 2). On the contrary, the Litter effect on Δ BW

Table 1. Effects of the ewes' parity and age at first lambing, the litter size, and the year of measurement on the average BW and BCS changes over successive physiological stages

Variable	<i>n</i>	Mean ¹ (SD)	Parity	Litter size	Age at first lambing	Year
BW-M:Pb	2663	8.23 (4.68)	***	***	***	***
BW-Pb:W	2627	-5.74 (6.66)	***	***	***	***
BW-Pb:L	2598	-3.33 (5.45)	***	***	***	***
BW-L:Sa	2717	0.02 (4.08)	**	***	***	***
BW-W:Wp	2078	0.92 (4.05)	***	***	NS	***
BW-W:M	1388	4.13 (4.32)	***	***	***	***
BCS-M:Pa	2706	0.12 (0.21)	**	NS	***	***
BCS-Pa:W	2698	-0.43 (0.27)	***	***	***	***
BCS-Pa:L	2920	-0.32 (0.26)	***	***	***	***
BCS-L:Sa	2811	-0.11 (0.21)	NS	***	***	***
BCS-W:Wp	1889	0.11 (0.21)	**	***	***	***
BCS-W:M	1320	0.25 (0.23)	***	***	***	***

¹BW variation (kg) and BCS variation (points).

n = number of records; SD = standard deviation; NS = nonsignificant; BW = body weight; BCS = body condition score; M = mating; Pa = early pregnancy; Pb = midpregnancy; L = lambing; Sa = early suckling; Sb = midsuckling; W = weaning; Wp = postweaning.

Table 2. Changes in body weight (Δ BW, kg) and body condition score (Δ BCS, points) as a function of the physiological stage, parity, litter size, and age at first lambing of the ewes

Variable	Parity			Litter size				Age at first lambing	
	1	2	3 and +	1	2	3	4 and +	1	2
<i>n</i> (%)	39	36	25	20	44	28	8	46	54
BW-M:Pb	7.32	7.83	8.81	7.00	8.38	7.80	8.85	8.33	7.62
BW-Pb:W	-3.7	-6.20	-8.49	-2.83	-5.31	-7.19	-9.25	-4.64	-7.65
BW-Pb:L	-0.85	-2.70	-4.33	0.01	-3.2	-2.12	-5.31	-1.36	-3.89
BW-L:Sa	-0.39	0.14	0.00	0.49	0.50	-0.32	-0.02	0.46	-0.13
BW-W:Wp	2.05	1.17	1.03	1.10	1.44	1.91	1.95	1.46	1.75
BW-W:M	4.60	4.91	.	3.74	4.11	5.37	5.79	5.28	4.23
BCS-M:Pa	0.09	0.17	0.17	0.14	0.12	0.15	0.15	0.12	0.16
BCS-Pa:W	-0.52	-0.40	-0.40	-0.34	-0.41	-0.49	-0.54 0	-0.42	-0.48
BCS-Pa:L	-0.39	-0.27	-0.26	-0.18	-0.33	-0.29	-0.40	-0.29	-0.33
BCS-L:Sa	-0.09	-0.10	-0.12	-0.08	-0.06	-0.16	-0.12	-0.09	-0.12
BCS-W:Wp	0.09	0.12	0.10	0.07	0.09	0.12	0.11	0.07	0.13
BCS-W:M	0.22	0.18	.	0.14	0.17	0.24	0.25	0.16	0.24

Data (Least squares means) are expressed in kg of BW changes and points of BCS changes (based on the scale of Russel et al. (1969)).

n = number of records in percentage; . = no data; BW = body weight; BCS = body condition score; M = mating; Pa = early pregnancy; Pb = midpregnancy; L = lambing; Sa = early suckling; Sb = midsuckling; W = weaning; Wp = postweaning.

was much lower for BW-L:Sa. The same tendencies were observed for Δ BCS. Regarding the Age effect, the weight gain over the BW-W:M and BW-M:Pb periods decreased with the increase of age at first lambing, whereas the weight loss during BW-Pb:W and BW-Pb:L increased. For BW-L:Sa period, BW gain was observed for ewes that lambed at 1 yr old and weight loss was observed for ewes that lambed at 2 yr old. Regarding Δ BCS, the BCS gain observed over the BCS-M:Pa, BCS-W:Wp, and BCS-W:M periods increased with the increase of age at first lambing. The BCS loss for BCS-Pa:W, BCS-Pa:L, and BCS-L:Sa periods was slightly higher in ewes lambing at 2 yr old than in ewes lambing at 1 yr old.

The Year effect was always highly significant ($P < 0.001$), whatever the physiological stage. The first-order interactions between the fixed effects considered here were not significant.

Heritabilities and Genetic Correlations

Estimates of variance components for Δ BW and Δ BCS are presented in Table 3. Direct heritabilities for Δ BW and Δ BCS ranged between 0.04 ± 0.02 and 0.18 ± 0.04 . The highest heritabilities were found for BW-W:Wp (0.18 ± 0.04), BW-Pb:L (0.17 ± 0.03), and BW-W:M (0.17 ± 0.03), whereas the lowest heritability values were obtained for BCS-L:Sa

(0.04 ± 0.02) and BCS-M:Pa (0.07 ± 0.02). Low heritabilities were found for BCS-Pa:W (0.16 ± 0.03) and BW-M:Pb (0.16 ± 0.03). Estimates for the fraction of permanent environmental variance were null except for BW-M:Pb, BW-Pb:W, and BCS-L:Sa. Repeatability estimates ranged from 0.08 ± 0.02 to 0.17 ± 0.02 for BCS-L:Sa and BCS-Pa:W, respectively. For ΔBW , repeatability estimates varied from 0.17 ± 0.03 to 0.21 ± 0.02 for BW-Pb:L and BW-M:Pb, respectively. Phenotypic variance was high for BW-Pb:W, BW-Pb:L, BW-W:M, BCS-Pa:W, and BCS-Pa:L, and lower for BW-M:Pb, BW-W:Wp, BW-W:M, BCS-M:Pa, BCS-W:Wp, and BCS-W:M periods (Table 3).

Regarding ΔBW , high negative genetic correlations were found between BW-Pb:W and BW-W:Wp, and BW-Pb:L and BW-L:Sa (Tables 4 and 5). Moderate negative genetic correlations were found between BW-M:Pb and BW-W:Wp, BW-M:Pb and BW-W:M, BW-Pb:W and BW-W:M, and BW-L:Sa and BW-W:M, respectively. A moderate positive genetic correlation was found between BW-Pb:W and BW-W:L:Sa, and a high positive genetic correlation was found between BW-W:Wp and BW-W:M. Phenotypic correlations followed the same tendencies as genetic correlations but with lower values. The highest positive phenotypic correlation was found between BW-W:Wp and

Table 3. Estimates ($\pm$ standard errors) of variance components, heritability, and repeatability for BW and BCS changes

Variable	h^2	c^2	e^2	R	σ_p^2
BW-M:Pb	0.16 (0.03)	0.05 (0.03)	0.79 (0.02)	0.21 (0.02)	9.13 (0.27)
BW-Pb:W	0.13 (0.03)	0.06 (0.03)	0.81 (0.03)	0.19 (0.03)	15.57 (0.46)
BW-Pb:L	0.17 (0.03)	0.00 (0.03)	0.83 (0.02)	0.17 (0.03)	17.80 (0.54)
BW-L:Sa	0.13 (0.03)	0.01 (0.02)	0.86 (0.02)	0.14 (0.02)	15.26 (0.44)
BW-W:Wp	0.18 (0.04)	0.03 (0.03)	0.79 (0.03)	0.21 (0.03)	6.95 (0.23)
BW-W:M	0.17 (0.03)	0.00 (0.00)	0.83 (0.04)	0.17 (0.04)	10.28 (0.40)
BCS-M:Pa	0.07 (0.02)	0.00 (0.02)	0.92 (0.02)	0.08 (0.02)	0.041 (0.001)
BCS-Pa:W	0.16 (0.03)	0.02 (0.03)	0.83 (0.02)	0.17 (0.02)	0.056 (0.002)
BCS-Pa:L	0.10 (0.02)	0.02 (0.02)	0.88 (0.02)	0.12 (0.02)	0.055 (0.001)
BCS-L:Sa	0.04 (0.02)	0.04 (0.02)	0.92 (0.02)	0.08 (0.02)	0.030 (0.001)
BCS-W:Wp	0.10 (0.03)	0.01 (0.03)	0.89 (0.03)	0.11 (0.03)	0.032 (0.001)
BCS-W:M	0.10 (0.04)	0.01 (0.05)	0.89 (0.05)	0.11 (0.05)	0.039 (0.002)

h^2 = heritability; c^2 = proportion of total phenotypic variance due to ewe permanent environmental effect; e^2 = proportion of total phenotypic variance due to temporary environmental effects; r = repeatability; σ_p^2 = total phenotypic variance; BW = body weight; BCS = body condition score; M = mating; Pa = early pregnancy; Pb = midpregnancy; L = lambing; Sa = early suckling; Sb = midsuckling; W = weaning; Wp = postweaning.

Table 4. Genetic and phenotypic correlations ($\pm$ standard errors) for BW changes

Variable	BW-M:Pb	BW-Pb:W	BW-Pb:L	BW-L:Sa	BW-W:Wp	BW-W:M
BW-M:Pb	—	−0.17 (0.15)	−0.16 (0.14)	0.18 (0.15)	−0.45 (0.14)	−0.37 (0.16)
BW-Pb:W	−0.47 (0.02)	—	0.03 (0.16)	0.48 (0.16)	−0.68 (0.11)	−0.50 (0.14)
BW-Pb:L	−0.37 (0.02)	0.51 (0.02)	—	−0.63 (0.10)	0.03 (0.15)	−0.03 (0.16)
BW-L:Sa	−0.03 (0.02)	0.16 (0.02)	−0.37 (0.02)	—	−0.30 (0.16)	−0.36 (0.17)
BW-W:Wp	−0.04 (0.02)	−0.45 (0.02)	−0.06 (0.03)	−0.10 (0.02)	—	0.91 (0.06)
BW-W:M	−0.03 (0.03)	−0.43 (0.02)	−0.05 (0.03)	−0.10 (0.03)	0.68 (0.01)	—

Table 5. Genetic and phenotypic correlations ($\pm$ standard errors) for BCS changes

Variable	BCS-M:Pa	BCS-Pa:W	BCS-Pa:L	BCS-L:Sa	BCS-W:Wp	BCS-W:M
BCS-M:Pa	—	−0.52 (0.13)	−0.44 (0.15)	−0.27 (0.26)	−0.02 (0.23)	−0.32 (0.22)
BCS-Pa:W	−0.47 (0.02)	—	0.70 (0.09)	0.52 (0.19)	−0.62 (0.14)	−0.75 (0.14)
BCS-Pa:L	−0.45 (0.02)	0.61 (0.01)	—	0.01 (0.23)	−0.39 (0.19)	−0.56 (0.22)
BCS-L:Sa	−0.00 (0.02)	0.14 (0.02)	−0.34 (0.02)	—	−0.47 (0.25)	−0.58 (0.21)
BCS-W:Wp	−0.00 (0.02)	−0.42 (0.02)	−0.04 (0.02)	−0.03 (0.02)	—	0.57 (0.17)
BCS-W:M	0.01 (0.03)	−0.43 (0.02)	−0.10 (0.03)	−0.03 (0.03)	0.55 (0.02)	—

Phenotypic correlations below the diagonal; genetic correlations above the diagonal; BW: Body Weight; BCS: Body Condition Score; M: Mating; Pa: Early pregnancy; Pb: Mid-pregnancy; L: Lambing; Sa: Early suckling; Sb: Mid-suckling; W: Weaning; Wp: Post-weaning.

BW-W:M, whereas the highest negative phenotypic correlation was observed between BW-M:Pb and BW-Pb:W.

Regarding Δ BCS, high negative genetic correlations were found between BCS-Pa:W and BCS-W:Wp, BCS-Pa:W and BCS-W:M, BCS-Pa:L and BCS-W:M, and BCS-L:Sa and BCS-W:M, respectively (Tables 4 and 5). Moderate negative genetic correlations were found between BCS-M:Pa and BCS-Pa:W, BCS-M:Pa and BCS-Pa:L, BCS-Pa:L and BCS-W:Wp, and BCS-L:Sa and BCS-W:Wp, respectively. A moderate positive genetic correlation was found between BCS-Pa:W and BCS-L:Sa. High positive genetic correlations were found between BCS-Pa:W and BCS-Pa:L, and BCS-W:Wp and BCS-W:M. Phenotypic correlations followed the same tendencies as genetic correlations but with lower values. The highest positive phenotypic correlation was found between BCS-Pa:W and BCS-Pa:L, whereas the highest negative phenotypic correlation was found between BCS-Pa:W and BCS-W:M.

When considering the correlations between Δ BW and Δ BCS, a high negative genetic correlation was observed between BCS-Pa:W and BW-W:Wp (Tables 6 and 7). Moderate negative genetic correlations were observed between BCS-Pa:W and BW-W:M, BCS-Pa:L and BW-W:Wp, BCS-L:Sa and BW-W:Wp, BCS-W:Wp and BW-L:Sa, BCS-W:M and BW-Pb:W, and BCS-W:M and BW-Pb:L, respectively. Moderate positive genetic correlations were observed between BCS-M:Pa and BW-M:Pb, BCS-Pa:W and BW-Pb:W, BCS-Pa:W and BW-L:Sa, BCS-Pa:L and BW-Pb:L, and BCS-W:Wp and

BW-W:Wp, respectively. High positive genetic correlations were observed between BCS-W:Wp and BW-W:M, BCS-W:M and BW-W:Wp, and BCS-W:M and BW-W:M. Phenotypic correlations followed the same tendencies as genetic correlations but with lower values. The highest positive phenotypic correlation was observed between BCS-Pa:W and BW-Pb:W, whereas the highest negative phenotypic correlation was found between BCS-Pa:W and BW-W:M.

DISCUSSION

The BR management over time, named BR dynamics (i.e., biological ability to mobilize and restore BR over time) in the present study, is presently considered as an interesting biological component to include in current and future animal breeding programs to increase robustness in ruminants, in order to improve the sustainability of livestock systems (De La Torre et al., 2015; Phocas et al., 2016; Friggens et al., 2017). This can be applied for systems with a wide range of farming environments and is particularly beneficial to those having to cope for instance with important fluctuations in feed resources.

In this study, we confirmed our global hypothesis regarding the existence of genetic determinism for Δ BW and Δ BCS in sheep.

Overall BR Dynamics

Combined BCS and BW measurements are considered to be typical and relatively easily

Table 6. Genetic correlations ($\pm$ standard errors) between BW and BCS changes

Variables	BW-M:Pb	BW-Pb:W	BW-Pb:L	BW-L:Sa	BW-W:Wp	BW-W:M
BCS-M:Pa	0.46 (0.16)	-0.25 (0.19)	0.02 (0.18)	-0.01 (0.19)	0.30 (0.18)	0.03 (0.20)
BCS-Pa:W	0.01 (0.14)	0.54 (0.12)	0.19 (0.14)	0.29 (0.15)	-0.73 (0.11)	-0.46 (0.15)
BCS-Pa:L	0.19 (0.16)	0.20 (0.16)	0.31 (0.14)	0.12 (0.16)	-0.48 (0.14)	-0.32 (0.17)
BCS-L:Sa	-0.14 (0.23)	0.09 (0.24)	-0.09 (0.23)	0.18 (0.23)	-0.40 (0.21)	-0.11 (0.25)
BCS-W:Wp	-0.18 (0.18)	-0.44 (0.19)	-0.26 (0.17)	-0.40 (0.19)	0.37 (0.18)	0.58 (0.17)
BCS-W:M	-0.38 (0.24)	-0.52 (0.19)	-0.37 (0.18)	-0.32 (0.22)	0.60 (0.18)	0.68 (0.16)

Table 7. Phenotypic correlations ($\pm$ standard errors) between BW and BCS changes

Variables	BW-M:Pb	BW-Pb:W	BW-Pb:L	BW-L:Sa	BW-W:Wp	BW-W:M
BCS-M:Pa	0.12 (0.02)	-0.04 (0.02)	0.02 (0.02)	0.00 (0.02)	0.08 (0.02)	0.05 (0.03)
BCS-Pa:W	-0.00 (0.02)	0.36 (0.02)	0.17 (0.02)	0.08 (0.02)	-0.20 (0.02)	-0.21 (0.03)
BCS-Pa:L	0.02 (0.02)	0.22 (0.02)	0.31 (0.02)	0.03 (0.02)	-0.12 (0.02)	-0.12 (0.03)
BCS-L:Sa	-0.05 (0.02)	0.10 (0.02)	-0.01 (0.02)	0.14 (0.02)	-0.02 (0.02)	-0.03 (0.03)
BCS-W:Wp	-0.03 (0.02)	-0.07 (0.03)	-0.06 (0.03)	-0.05 (0.02)	0.24 (0.02)	0.14 (0.03)
BCS-W:M	-0.04 (0.03)	-0.14 (0.03)	-0.08 (0.03)	-0.04 (0.03)	0.22 (0.03)	0.26 (0.03)

BW = body weight; BCS = body condition score; M = mating; Pa = early pregnancy; Pb = midpregnancy; L = lambing; Sa = early suckling; Sb = midsuckling; W = weaning; Wp = postweaning.

monitored parameters to describe the level of BR in sheep (Thorup et al., 2012; Brown et al., 2015; Morel et al., 2016; Puillet and Martin, 2017). Very few studies, however, are available in the literature regarding BR dynamics over complete and successive production cycles. To investigate the genetic determinism of BR dynamics using a multitrait approach, we defined several time periods based on key physiological stages over a representative production cycle. These periods were also defined in order to assess ewes' ability to mobilize and to restore BR over time and to take into account the maximal amplitude and/or slopes of BR variations.

Overall, in our experimental conditions, BR accretion was observed from weaning to early pregnancy. The increase in ΔBW and ΔBCS during the postweaning period is probably due to a decrease in the ewes' energy requirements. The increase in ΔBW during early pregnancy is most probably related to the anabolism of pregnancy combined with the onset of fetus growth (Bauman and Currie, 1980), but also to the favorable herbage availability and quality on the rangeland at this season of the year (González-García et al., 2014). The BR mobilization was observed from midpregnancy to weaning in our conditions even if a slight, brief increase in BR was observed after early suckling, followed by a decrease in BR at midsuckling. The decrease in BR during pregnancy and suckling is likely due to the known negative energy balance of this physiological stage. The progressive increase in the nutrient requirements of the dam, to support fetus growth and development during pregnancy and then milk production after lambing, contrasts with the linear decrease in the physical intake capacity as pregnancy progresses and the physiological incapacity of the female to meet the peak in nutrient and energy demand that occurs during the first weeks after parturition (Nielsen et al., 2003; Smith et al., 2017).

The biological factors that have an effect on BR changes are similar to those affecting body condition measured at single time point (María and Ascaso, 1999; González-García et al., 2014, 2015). Concerning the litter effect, an overall increase in BR mobilization could be observed via ΔBW and ΔBCS measurements as the litter size increased. This was expected because of the related higher energy requirements for ewes suckling multiple lambs. The ΔBW also increased somewhat as parity increased, especially between mating and midpregnancy and during the postweaning period, which suggests that ewes learn how to better manage BR with age (Clawson et al., 1991).

Concerning the effect of age at first lambing, ewes had higher gain and lower loss of BW when they lambed at 1 yr of age. This might be explained by the breeding system used in our experimental farm where heavier ewes were early mated (i.e., at 7 mo of age), whereas lighter ewes were mated later (i.e., at 19 mo of age). However, the ΔBCS were slightly influenced by the age at first lambing. The year also affected BR variations in our conditions, probably due to variations in biomass availability in the rangeland induced by differences in climatic conditions, thus leading to different nutritional and energy balances among years.

Genetic Parameters

The heritabilities estimated in the present study for ΔBW and ΔBCS measured over time, and based on repeated measurements (i.e., from 0.13 ± 0.03 to 0.18 ± 0.04 and from 0.04 ± 0.02 to 0.16 ± 0.03 , respectively), suggest that the biological capacities determining the nature of BR accretion and mobilization processes are heritable and could be transmitted across generations. To the best of our knowledge, this is the first work reporting heritability estimations considering a dynamic approach of ΔBW and ΔBCS throughout a complete productive cycle (e.g., over several production cycles) in ruminants. Our estimates for ΔBW provide consistent values with those reported previously in sheep for BW gain or loss (Rose et al., 2013). Another study estimated the heritability of a ΔBCS measurement in Holstein cows between the first and tenth weeks after calving and estimated a heritability of 0.09 ± 0.04 (Pryce et al., 2001), which is in agreement with our findings. The heritability estimations calculated for ΔBCS and ΔBW over time were lower than those that we obtained for BCS and BW at a given point, which range from 0.26 to 0.37 for BCS and from 0.44 to 0.58 for BW (Supplementary Table 1). Our single time-point BW and BCS heritability estimations were consistent with previous findings in cows and sheep (Koenen et al., 2001; Mao et al., 2004; Safari et al., 2005; Gizaw et al., 2007; Shackell et al., 2011). Thus, amplitude of variations in BR, i.e., BR mobilization and accretion ability, would be under genetic control but in a lower extent than level of BR. However, although more heritable traits would be better suited to faster genetic progress, BW and BCS levels did not represent the ability of ewes to mobilize or to restore BR as indicated by absence or few moderate genetic correlations between amplitude in BCS or BW variations and levels.

The low proportion of total phenotypic variance for ΔBW and ΔBCS due to the ewe permanent environmental effect suggested that no additional ewe effects apart from the genetic effect affected BR dynamics. Therefore, the repeatability for BR dynamics was close to the heritability. Interestingly, a larger permanent ewe environmental effect was observed, mainly for BW, for measurements at single time points ([Supplementary Table 2](#)). This suggests that no additional ewe effects apart from the genetic effect influenced the BW, and BCS in a lower extent, of a ewe when BW was considered at a given single time point.

High positive genetic correlations for ΔBW and ΔBCS between the early postweaning period and the entire postweaning period confirm that BR accretion (i.e., with the inherent anabolic processes) were predominant during this period. Surprisingly, the increase in ΔBW during the first half of pregnancy was negatively correlated with the increase in ΔBW during the postweaning period and the increase in ΔBCS during early pregnancy was not genetically correlated with the increase in ΔBCS during postweaning. Such negative correlations could be due to different biological mechanisms involved in BW recovery during the postweaning period and the increase in BW related to the growth of the fetus during pregnancy. These results contrasted with the high positive genetic correlations found between single time-point measurements of BW and BCS during the BR accretion period (i.e., during the beginning of postweaning and the entire postweaning period; [Supplementary Table 3](#)). The differences in the high or moderate correlations for single time-point measurements and the nonzero correlations among ΔBCS and ΔBW might be explained by the higher heritability of BW and BCS levels than ΔBW and ΔBCS .

The long BR mobilization period from the second month of pregnancy to weaning was correlated with shorter periods (i.e., from midpregnancy to lambing; from lambing to early suckling) within the same mobilization period, both for ΔBW and ΔBCS . The high positive genetic relationships observed between ΔBCS over the 3 periods (i.e., from midpregnancy to lambing; from lambing to early suckling; and from midpregnancy to weaning) defined to investigate the decrease in BR confirmed that these periods were representative of BR mobilization. In addition, the negative genetic relationships observed between ΔBCS over these 3 periods and the 3 periods characterizing BR increase (i.e., during the beginning of postweaning; during the entire postweaning period; and from mating to

midpregnancy) suggest that the series of mechanisms involved either in BR mobilization and/or accretion processes are probably driven by different physiological factors, but are also genetically related. This result is consistent with common biochemical mechanisms involved in the regulation of lipogenesis and lipolysis in adipose tissue ([Chilliard et al., 1998](#)).

The ΔBCS values were generally independent of the level of BCS (or BW) measured at single time points throughout the productive cycle (e.g., fat ewes did not show higher BR mobilization and vice versa). However, some ΔBCS (and ΔBW) values were genetically correlated with the level of BCS (or BW) ([Supplementary Tables 4–6](#)). The increase in ΔBCS during early pregnancy was positively related to several single time-point BCS measurements (e.g., BCS at early pregnancy, at midpregnancy, at lambing, and at midsuckling). The loss in BCS during pregnancy was negatively correlated with the level of BCS at early pregnancy. These results may be related to the ewes' fitness; indeed, it appeared that the ewes that could not mobilize enough BR during pregnancy tended to activate self-preservation mechanisms. Surprisingly, a similar kind of relationship was not observed during the suckling period.

The present study also highlights high positive genetic correlations between ΔBW and ΔBCS over the same physiological periods. This suggests that variations in both parameters are physiologically concomitant but also genetically linked. Surprisingly, no correlations were found between ΔBW and ΔBCS at early suckling. This could be explained by other factors that greatly affect the manifestation and extent of the expression of these parameters at this stage (e.g., litter size effects on energy balance and year effects). Apart from the correlations at the same physiological stage, almost all the correlations between the BR accretion and mobilization phases were negative. This confirms that ΔBW and ΔBCS are 2 genetically related and somehow mutually exclusive traits, i.e., with well-known successive and complementary physiological cycles.

In conclusion, the present study shows that BR dynamics is a heritable trait and could be a good trait to be considered for improving individual robustness in sheep. Some intervals of BR changes appeared to be more relevant for genetic selection purposes due to higher heritabilities and/or higher genetic correlations between periods. However, to define the best time interval for genetic selection of BR dynamics, further studies are needed

to address hypotheses helping to explain BR changes over time. It would be also interesting to investigate additional direct or indirect relationships between BR dynamics and productive and reproductive traits before considering such traits in selection. Further research efforts are necessary, i.e., including complementary traits (e.g., plasma metabolites and hormone profiles), to determine direct relationships between BR dynamics and the energy metabolism of the animal. This would lead to a better understanding of BR accretion and mobilization mechanisms and help identify individuals or periods with excessive BR mobilization that could have a significant impact on health or survival.

LITERATURE CITED

- Arango, J. A., L. V. Cundiff, and L. D. Van Vleck. 2002. Genetic parameters for weight, weight adjusted for body condition score, height, and body condition score in beef cows. *J. Anim. Sci.* 80:3112–3122.
- Bauman, D. E., and W. B. Currie. 1980. Partitioning of nutrients during pregnancy and lactation: a review of mechanisms involving homeostasis and homeorhesis. *J. Dairy Sci.* 63:1514–1529.
- Blanc, F., F. Bocquier, J. Agabriel, P. D'hour, and Y. Chilliard. 2006. Adaptive abilities of the females and sustainability of ruminant livestock systems. A review. *Anim. Res.* 55:489–510. doi:10.1051/animres:2006040
- Brown, D. J., D. B. Savage, G. N. Hinch, and S. Hatcher. 2015. Monitoring liveweight in sheep is a valuable management strategy: a review of available technologies. *Anim. Prod. Sci.* 55:427–436. doi:10.1071/AN13274
- Chilliard, Y., F. Bocquier, and M. Doreau. 1998. Digestive and metabolic adaptations of ruminants to undernutrition, and consequences on reproduction. *Reprod. Nutr. Dev.* 38:131–152.
- Clawson, A. J., J. D. Garlich, M. T. Coffey, and W. G. Pond. 1991. Nutritional, physiological, genetic, sex, and age effects on fat-free dry matter composition of the body in avian, fish, and mammalian species: a review. *J. Anim. Sci.* 69:3617–3644.
- De La Torre, A., E. Recoules, F. Blanc, I. Ortigues-Marty, P. D'Hour, and J. Agabriel. 2015. Changes in calculated residual energy in variable nutritional environments: an indirect approach to apprehend suckling beef cows' robustness. *Livest. Sci.* 176:75–84. doi:10.1016/j.livsci.2015.03.008
- Edmonson, A. J., I. J. Lean, L. D. Weaver, T. Farver, and G. Webster. 1989. A body condition scoring chart for holstein dairy cows. *J. Dairy Sci.* 72:68–78. doi:10.3168/jds.S0022-0302(89)79081-0
- Friggens, N. C., F. Blanc, D. P. Berry, and L. Puillet. 2017. Review: deciphering animal robustness. A synthesis to facilitate its use in livestock breeding and management. *Animal* 11:2237–2251. doi:10.1017/S175173111700088X
- Friggens, N. C., K. L. Ingvarsen, and G. C. Emmans. 2004. Prediction of body lipid change in pregnancy and lactation. *J. Dairy Sci.* 87:988–1000. doi:10.3168/jds.S0022-0302(04)73244-0
- Gilmour, A. R., B. R. Cullis, S. J. Welham, and R. Thompson. 2001. *ASREML Reference Manual*. 239.
- Gizaw, S., S. Lemma, H. Komen, and J. A. M. Van Arendonk. 2007. Estimates of genetic parameters and genetic trends for live weight and fleece traits in Menz sheep. *Small Rumin. Res.* 70:145–153. doi:10.1016/j.smallrumres.2006.02.007
- González-García, E., V. Gozzo de Figuereido, D. Foulquie, E. Jousserand, P. Autran, S. Camous, A. Tesniere, F. Bocquier, and M. Jouven. 2014. Circannual body reserve dynamics and metabolic profile changes in romane ewes grazing on rangelands. *Domest. Anim. Endocrinol.* 46:37–48. doi:10.1016/j.domaniend.2013.10.002
- González-García, E., and D. Hazard. 2016. Growth rates of Romane ewe lambs and correlated effects of being mated as hoggets or two-tooth ewes on first offspring performance. *Livest. Sci.* 189:63–69. doi:10.1016/j.livsci.2016.04.022
- González-García, E., A. Tesniere, S. Camous, F. Bocquier, F. Barillet, and P. Hassoun. 2015. The effects of parity, litter size, physiological state, and milking frequency on the metabolic profile of lacaune dairy ewes. *Domest. Anim. Endocrinol.* 50:32–44. doi:10.1016/j.domaniend.2014.07.001
- Koenen, E. P., R. F. Veerkamp, P. Dobbelaar, and G. De Jong. 2001. Genetic analysis of body condition score of lactating dutch holstein and red-and-white heifers. *J. Dairy Sci.* 84:1265–1270. doi:10.3168/jds.S0022-0302(01)74588-2
- Mao, I. L., K. Sloniewski, P. Madsen, and J. Jensen. 2004. Changes in body condition score and in its genetic variation during lactation. *Livest. Prod. Sci.* 89:55–65. doi:10.1016/j.livprodsci.2003.12.005
- María, G. A., and M. S. Ascaso. 1999. Litter size, lambing interval and lamb mortality of Salz, Rasa Aragonesa, Romanov and F1 ewes on accelerated lambing management. *Small Rumin. Res.* 32:167–172. doi:10.1016/S0921-4488(98)00176-X
- Molénat, G., D. Foulquie, P. Autran, J. Bouix, D. Hubert, M. Jacquin, and B. Bibe. 2005. Pour un élevage ovin allaitant performant et durable sur parcours: un système expérimental sur le Causse du Larzac. *INRA Prod. Anim.* 18:323–338.
- Morel, P. C. H., N. M. Schreurs, R. A. Corner-Thomas, A. W. Greer, C. M. C. Jenkinson, A. L. Ridler, and P. R. Kenyon. 2016. Live weight and body composition associated with an increase in body condition score of mature ewes and the relationship to dietary energy requirements. *Small Rumin. Res.* 143:8–14. doi:10.1016/j.smallrumres.2016.08.014
- Nielsen, H. M., N. C. Friggens, P. Løvendahl, J. Jensen, and K. L. Ingvarsen. 2003. Influence of breed, parity, and stage of lactation on lactational performance and relationship between body fatness and live weight. *Livest. Prod. Sci.* 79:119–133. doi:10.1016/S0301-6226(02)00146-X
- O'Mara, F. P. 2012. The role of grasslands in food security and climate change. *Ann. Bot.* 110:1263–1270. doi:10.1093/aob/mcs209
- Phocas, F., C. Belloc, J. Bidanel, L. Delaby, J. Y. Dourmad, B. Dumont, P. Ezanno, L. Fortun-Lamothe, G. Foucras, B. Frappat, et al. 2016. Review: towards the agroecological management of ruminants, pigs and poultry through the development of sustainable breeding programmes: I-selection goals and criteria. *Animal* 10:1749–1759. doi:10.1017/S1751731116000926
- Pryce, J. E., M. P. Coffey, and G. Simm. 2001. The relationship between body condition score and reproductive

- performance. *J. Dairy Sci.* 84:1508–1515. doi:10.3168/jds.S0022-0302(01)70184-1
- Puillet, L., and O. Martin. 2017. A dynamic model as a tool to describe the variability of lifetime body weight trajectories in livestock females. *J. Anim. Sci.* 95:4846–4856. doi:10.2527/jas2017.1803
- Ricordeau, G., L. Tchamitchian, J. C. Brunel, T. C. Nguyen, and D. François. 1992. La race ovine INRA 401: un exemple de souche synthétique. *INRA Prod. Anim. (hs)*: 255–262.
- Rose, G., A. Kause, H. A. Mulder, J. H. van der Werf, A. N. Thompson, M. B. Ferguson, and J. A. van Arendonk. 2013. Merino ewes can be bred for body weight change to be more tolerant to uncertain feed supply. *J. Anim. Sci.* 91:2555–2565. doi:10.2527/jas.2012-5539
- Russel, A. J. F., J. M. Doney, and R. G. Gunn. 1969. Subjective assessment of body fat in live sheep. *J. Agric. Sci.* 72:451–454. doi:10.1017/S0021859600024874
- Safari, E., N. M. Fogarty, and A. R. Gilmour. 2005. A review of genetic parameter estimates for wool, growth, meat and reproduction traits in sheep. *Livest. Prod. Sci.* 92:271–289. doi:10.1016/j.livprodsci.2004.09.003
- Shackell, G. H., N. G. Cullen, and G. J. Greer. 2011. Genetic parameters associated with adult ewe liveweight and body condition. In: *Proc. 19th Assoc. Adv. Anim. Breed. Genet.* p. 103–106.
- Smith, G. L., N. C. Friggens, C. J. Ashworth, and M. G. G. Chagunda. 2017. Association between body energy content in the dry period and post-calving production disease status in dairy cattle. *Animal* 11:1590–1598. doi:10.1017/S1751731117000040
- Thorup, V. M., D. Edwards, and N. C. Friggens. 2012. On-farm estimation of energy balance in dairy cows using only frequent body weight measurements and body condition score. *J. Dairy Sci.* 95:1784–1793. doi:10.3168/jds.2011-4631