Published in final edited form as:

Neurosci Biobehav Rev. 2018 July ; 90: 115–129. doi:10.1016/j.neubiorev.2018.04.007.

RDoC-based categorization of amygdala functions and its implications in autism

Thomas Hennessey, PhD^{1,3}, Elissar Andari, PhD², and Donald G. Rainnie, PhD^{1,3}

¹Department of Behavioral Neuroscience and Psychiatric Disorders, Yerkes National Primate Research Center

²Silvio O. Conte Center for Oxytocin and Social Cognition, Department of Psychiatry and Behavioral Sciences, Division of Behavioral Neuroscience and Psychiatric Disorders, Yerkes National Primate Research Center, Emory University

³Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA 30329

Abstract

Confusion endures as to the exact role of the amygdala in relation to autism. To help resolve this we turned to the NIMH's Research Domain Criteria (RDoC) which provides a classification schema that identifies different categories of behaviors that can turn pathologic in mental health disorders, e.g. autism. While RDoC incorporates all the known neurobiological substrates for each domain, this review will focus primarily on the amygdala. We first consider the amygdala from an anatomical, historical, and developmental perspective. Next, we examine the different domains and constructs of RDoC that the amygdala is involved in: Negative Valence Systems, Positive Valence Systems, Cognitive Systems, Social Processes, and Arousal and Regulatory Systems. Then the evidence for a dysfunctional amygdala in autism is presented with a focus on alterations in development, prenatal valproic acid exposure as a model for ASD, and changes in the oxytocin system therein. Finally, a synthesis of RDoC, the amygdala, and autism is offered, emphasizing the task of disambiguation and suggestions for future research.

Keywords

Review; Amygdala; Autism; Research Domain Criteria; Oxytocin; Valproic Acid

1. Introduction

The principal focus of the field of social neuroscience in recent decades has been the elucidation of social brain networks capable of detecting cues, empathizing, interacting with

Corresponding author: Donald Rainnie, 954 Gatewood Rd, Yerkes Primate Center, Emory University, Atlanta, GA 30329, Tel: 404 727-8269, Fax: 404 727-8088, drainni@emory.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

others, and responding adaptively in ambiguous contexts (Dunbar, 2009; Frith, 2007; Stanley and Adolphs, 2013). An imperative of this work has been uncovering how psychiatric diseases impact these networks, particularly in autism spectrum disorder (ASD). One region consistently highlighted by this research as important in social functioning is the amygdala (Bickart et al., 2014).

The amygdala is a prime example of how neuroscience has shifted from a phrenological perspective mapping one function—fear—to one brain region and towards a network paradigm where actions arise from a dynamic affiliation of neural assemblages communicating with one another (Pessoa, 2014; Weiskrantz, 1956). The amygdala is now seen as a hub in several other, distinct networks: establishing valence or salience, cognition, reward, and social learning (Rutishauser et al., 2015). Despite its small size, the amygdala's anatomy and functioning is complex, including disparate nuclei with dense connectivity to cortical and subcortical brain regions.

Increasing knowledge of the amygdala is paralleled by new investigations of the social brain and autism. Since the original descriptions of Kanner (1943) and Asperger (1944), substantial research has been done to expand the understanding of autism leading, along with better awareness, to greatly increased rates of diagnosis (Fombonne, 2009; U.S. Department of Health and Human Services, 2014; Volkmar and Pauls, 2003; Wing, 2005, 1993, 1981; Wing and Potter, 2002). ASD's etiology remains obscure and has been posited to arise from the breakdown of particular genes, networks, or brain regions (Anney et al., 2012; Baron-Cohen et al., 2000; Markram et al., 2007; Poelmans et al., 2013; Voineagu et al., 2011). Presently, it is thought to be the product of altered expression in hundreds of genes, highly heritable, influenced by epigenetic factors, and deficits in connectivity between many different brain regions, including the amygdala (Eapen et al., 2013; Hallmayer et al., 2011; Kosmicki et al., 2017; Mintz, 2017).

Based on the 5th revision of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5, American Psychiatric Association, 2013), ASD is characterized by: 'a) persistent deficits in social interaction and communication as well as b) restricted, repetitive patterns of behaviors, interests, or activities with an early onset of deficits.' The detection of ASD is predicated on clinical measures that rely on parental reports, and on observations of the child's social aptitude during interactive tasks. These measures are crucial for diagnosing autism and categorizing the severity of symptoms. However, the DSM does not account for the vast heterogeneity of ASD and consequently it may not be the best predictor for treatment outcomes; it is thought that only 50% of individuals with ASD who receive treatment achieve significant gains as a result (Stahmer et al., 2010). In addition to improved clinical diagnosis, we need objective tools that acknowledge biological markers to demarcate variation in ASD. Towards this end the NIMH is adopting a new methodology, Research Domain Criteria (RDoC), to better characterize variation from typical behavior and enable precision medicine to improve psychiatric outcomes as it has in oncology and cardiology (Insel et al., 2010; Insel and Cuthbert, 2009). RDoC looks at psychological constructs dimensionally: from healthy to abnormal, examining the underlying genetics, neuropeptides, circuits, and physiology that ultimately gives rise to behavior (Cuthbert and Insel, 2013).

Along with renewed interest in nosology, there are questions regarding the amygdala and its contribution to neuropsychiatric disorders, including autism. Here we propose that by using RDoC to classify amygdala functions into constructs and dimensions, we can better characterize its role in normal functioning and explicate how it is altered in autism. Of the five currently defined RDoC domains—**Negative Valence Systems, Positive Valence Systems, Cognitive Systems, Social Processes, and Arousal and Regulatory Systems**—the amygdala is represented in each, within one or more constructs. The amygdala is not a unitary structure and we will decorticate its role in these domains and constructs by drawing on studies of lesions, animal behavior, in vivo and ex vivo electrophysiology, genetics, functional imaging, and optogenetics.

In this review, we will highlight: how conceptualizations of the amygdala's anatomy, development, and function within the social brain have changed overtime; amygdala functions in RDoC; dysfunction of the amygdala in autism throughout development; linkages to comorbid psychiatric disorders; amygdala alterations seen in the valproic acid (VPA) model of autism; and the effects of oxytocin (OT) on the amygdala and behavior. By understanding the diversity of amygdala functions, and the myriad ways it can be altered in autism, the route to improved therapies and outcomes for people with autism can be illuminated.

2. The Amygdala: anatomy, history, development

2.1 Anatomy

The human amygdala (Fig. 1) comprises 13 nuclei, distinguished by cytoarchitectonics, histochemistry, and connectivity with other brain regions. To greatly simplify the anatomical layout of the amygdala (see Duvarci & Pare, 2014; or Sah, et. al, 2003 for review), there are two main divisions 1) the **basolateral complex (BLA)** encompassing the **lateral (LA)**, **basal (BA)** and **accessory basal (AB)** nuclei with extensive connections to sensory and cortical brain regions and 2) the **central amygdala (CeA)** and its two sub-nuclei, the **central lateral (CeL)**, connected with the BLA, and **central medial (CeM)**, connected with brainstem areas related to the expression of innate behaviors (e.g. fear), along with the **medial amygdala (MeA)** linked to the olfactory system (Janak and Tye, 2015; LeDoux, 2008). The BLA is heavily innervated by sensory systems and the thalamus. The LA and the BA also receives information from higher-order sensory and associative cortices (Adolphs, 2009; Amaral et al., 2008). The BLA has strong efferent projections to reward regions like the nucleus accumbens (NAc), the ventral pallidum in the basal forebrain, and reciprocal connections with memory processing areas, associative and executive cortical sections, and with the CeA (Fig. 2).

2.2 Historical context

The classic studies of Klüver and Bucy in the 1930s first described the involvement of the amygdala in emotional processing by elucidating the behavioral consequences of large, non-selective temporal lesions in monkeys (Klüver and Bucy, 1939, 1937). Lesioned monkeys had 'psychic blindness' stemming from a lack of access to the value of social cues. Subsequent studies with more precise lesions of the amygdala caused more selective

behavioral disruptions: less caution and distrust in approaching novel objects (such as snakes) or human strangers, in line with the known role of the amygdala in resolving unpredictability or ambiguity (Adolphs, 2010; Bauman et al., 2006; Machado et al., 2009).

The most selective lesions of the human amygdala result from vascular damage or from Urbach-Wiethe disease (UWD), a rare autosomal disorder that causes the progressive calcification of the amygdala (Adolphs et al., 1994; Cinaz et al., 1993; Urbach and Wiethe, 1929). SM, the best described case of UWD, and other such patients showed impairments in fear recognition and autonomic conditioned responses in Pavlovian fear conditioning (Adolphs et al., 1994; Bechara et al., 1995; Cristinzio et al., 2010; Siebert et al., 2003). They also have deficits in social judgments from faces, more fixations on the mouth than the eyes during conversations, and general deficits in affective perspective taking (Adolphs et al., 1998; Hillis, 2014; Spezio et al., 2007). Individuals with UWD report experiencing no negative emotions in real life, exhibit no sensitivity to personal space, and fail to tag emotionally charged words as important to remember (Hampton et al., 2007; Hurlemann et al., 2007; Kennedy et al., 2010; Tranel et al., 2006). Research into UWD confirms the role of the amygdala not only in fear processing but also processing salient or unpredictable stimuli, and in social settings.

Historically the amygdala and the rest of the limbic system was thought of as ‘evolutionarily primitive’, and resistant to change (MacLean, 1955, 1952). However, components of the limbic system, principally the BLA, are in fact quite plastic, with evolutionary dynamism that has been attributed to the influence of social factors (Barger et al., 2007). The volume of the monkey’s BLA is about 30-40 times that of the rat, while the CeA and MeA are only four and eight times larger respectively (Chareyron et al., 2011).

Humans in turn have a BLA and, primarily, a lateral division, several times the size of any other ape or monkey (Barger et al., 2007). The LA’s neuronal density is substantially higher in the rat (99,000 neurons/mm³), compared to the monkey (42,000 neurons/mm³), and lowest in the human (9,000 neurons/mm³). This lower density reflects expanded dendritic arbors and glial populations, and greater opportunity for connections with many networks, particularly the social (Pessoa, 2014). Volume, density, and other structural elements of the amygdala are all established during development from fertilization to adulthood.

2.3 Development

In humans the BLA shows synaptogenesis from the first trimester until the seventh month, the CeA is identifiable in the fifth month, and overall neuronal migration is complete by the end of the eighth month of gestation (Schumann et al., 2011; Setzer and Ulfing, 1999; Ulfing et al., 2003). The cellular structure of the amygdala and its neurotransmitter systems are well established at birth, with similar distribution to that seen in adults (Bauman and Amaral, 2005). While the number of neurons in the amygdala does not increase, the volume grows throughout childhood and into adolescence (Payne et al., 2010; Schumann et al., 2004). This is due to significant increases in both the dendritic arbor (Fig. 3) and number of oligodendrocytes that myelinate input to the amygdala, a recapitulation of the evolutionary forces that increased its size and functionality in primates (Chareyron et al., 2012; Ryan et al., 2016).

Simultaneously, more connections are made within the amygdala between the projecting principal and local inhibitory neurons that tune the network, filtering out extraneous information and binding together the input of different sensory modalities (Ryan et al., 2012). The functional role of the amygdala shifts dramatically at these stages, especially in terms of its receptivity to external input and internal modulation. Not until the second week of rat development, (corresponding to a newborn human, Quinn, 2005), are they able to learn associations between painful shocks and odors, and to avoid them (King et al., 2014). During this same period Arruda-Carvalho et al. (2017) demonstrated with viral tracers and optogenetics that connections from the medial prefrontal cortex (mPFC) to the BLA in the mouse go from undetectable in early development (P10) to prominent by early adolescence (P15) and become increasingly functional from then on. This is just one facet of the development of the amygdala, a process that links its different nuclei to the rest of the brain and to each other.

Understanding the underlying neurobiology of the amygdala is essential to seeing how it is involved in both normal and pathological behaviors, and a cornerstone of RDoC (Cuthbert and Insel, 2013). With RDoC as a framework, we will identify the domains and circuitry that include the amygdala and are altered in autism.

3. Amygdala functions and RDoC

Using RDoC, we have here classified amygdala functions in the domains of negative valence, positive valence, cognition, social functions, and arousal. We are interested in both the behaviors and neural circuitry of these different constructs and the relative importance of the amygdala within them.

3.1 Negative Valence Systems (NVS)

Defined as “primarily responsible for responses to aversive situations or contexts such as fear, anxiety and loss,” NVS have a very strong link to the amygdala.

3.1.1 Acute Threat—The role of the amygdala in Acute Threat, i.e. fear learning, may be one of the most studied relationships in all of neuroscience. Weiskrantz (1956) first demonstrated that a lesion of the amygdala prevented monkeys from pairing an aversive unconditioned stimulus, such as a shock, with a neutral cue like a tone, the conditioned stimulus. More detailed studies in rodent models, including seminal papers by LeDoux and others, explicated which sub-regions of the amygdala (Fig. 2) are required: the UA for input, BA for associative learning, and CeM for expression (Clugnet and LeDoux, 1990; Kalin et al., 2004; Keifer et al., 2015; LeDoux et al., 1990, 1988; LeDoux, 1992; Madarasz et al., 2016; Romanski and LeDoux, 1992). The connection from the BLA to the mPFC is particularly important for acute threat; optogenetic depression of this connection both inhibited acquisition, and enhanced extinction, of fear learning (Klavir et al., 2017). Structural and functional imaging studies in humans support a strong amygdala response to fearful faces. Recordings from implanted electrodes in epileptic individuals viewing faces reveal an extremely rapid response to fear, preceding a general response to any expression (Fullana et al., 2015; Méndez-Bértolo et al., 2016). Within the BLA, inhibition and

disinhibition of principal neurons by specific populations of interneurons is required for conditioned and unconditioned stimulus association learning to occur (Wolff et al., 2014).

In terms of neuromodulatory factors (Fig. 2), there are substantial serotonin (5-HT) inputs to the BLA from the dorsal raphe which are sensitive to the presence of stressors (Vertes, 1991). During non-stressful conditions, 5-HT release in the BLA decreases the excitability of neurons, presumably by increasing inhibition and depolarizing GABAergic interneurons (Rainnie, 1999). Conversely, the selective depletion of 5-HT in the LA results in significantly enhanced fear potentiated startle and increased expression of excitatory glutamate receptors (Tran et al., 2013). The role of 5-HT in the amygdala, in the expression of fear and anxiety, as well as the clinical implications of SSRIs are still not fully understood (for review see Burghardt & Bauer, 2013).

Dopamine is also released in the BLA during stressful situations and D1 receptors in the BLA seem to be necessary for the acquisition and expression of Pavlovian fear conditioning (Fadok et al., 2009; Muller et al., 2009). D1 receptors can have effects on long-term potentiation (LTP) induction through the modulation of NMDA receptors and facilitate the transition of salient sensory inputs and fear learning (Li et al., 2011). Dopamine also suppresses certain GABAergic interneurons of the BLA via D2 receptor activation to further boost relevant signals and promote learning (Chu et al., 2012).

3.1.2 Potential Threat and Sustained Threat—Potential Threat (or anxiety) and Sustained Threat (chronic stress) are closely related to each other, and to Acute Threat, differing in the physical or mental proximity to the perceived danger and in the time course (Avery et al., 2015; Davis et al., 2010).

Anxiety can be considered a generalization of fear learning and is an adaptive process from a survival perspective. However, overgeneralization of fear to harmless stimuli can become pathological and result in posttraumatic stress disorder (PTSD), phobias, obsessive compulsive disorder, panic disorder, and generalized anxiety disorder (Lissek et al., 2014). A delicate neuronal balance within the CeA (Fig. 2) regulates fear expression: the CeM is tonically inhibited by the GABAergic output of the CeL, which in turn is innervated by the BLA. If this balance is disrupted, and the CeM disinhibited, fear expression can generalize from conditioned, to non-conditioned stimuli as well (Ciocchi et al., 2010; Keifer et al., 2015). The MeA is also important in the response to stress, and impressive advances in single cell RNA sequencing technology recently identified a subpopulation of cholecystokinin expressing neurons activated by acute restraint stress (Crane et al., 2005; Wu et al., 2017).

The Stoop lab (2008) has shown that two neuropeptides, oxytocin (OT) and vasopressin (VP), have opposite effects on fear expression at the neuronal level in the amygdala. The CeM is excited by VP but fear responses are inhibited in the CeL, which is in turn excited by OT (Huber et al., 2005; Knobloch et al., 2012). Hence, an absence, reduction, or desensitization of oxytocin receptors (OTRs) in the CeL can lead to an imbalance at the neuronal level within the CeA and to generalized fear responses and anxiety. In rodents, chronic treatment with OT has been found to be anxiogenic at high doses (10ng/hr) but at

low doses (1 ng/hr) prevents psychosocial stress induced anxious behaviors (Peters et al., 2014).

There is increasing evidence for OT reducing fear, anxiety, and stress, in part by lowering amygdala activity (Bethlehem et al., 2013; Domes et al., 2007; Labuschagne et al., 2010; Meyer-Lindenberg et al., 2011; Petrovic et al., 2008). Researchers have found that intranasal OT facilitates the extinction of learned fear and reduces anxiety in patients with anxiety disorders (Acheson et al., 2013; Dodhia et al., 2014; Eckstein et al., 2015; Gorka et al., 2015; Labuschagne et al., 2010). It has also been shown that men with PTSD show diminished functional connectivity between the CeM and the ventromedial PFC, while in women with PTSD there is enhanced functional connectivity between the BLA and dorsal anterior cingulate cortex (ACC). OT administration normalized the functional connectivity between the amygdala and frontal areas during resting state in both sexes (Koch et al., 2016). In healthy controls though, Grillon (2013) found that intranasal OT enhanced startle response after unpredictable shock stress, which suggests that care should be taken in translating OT, it may be most efficacious specifically for anxiety related to social behaviors (Neumann and Slattery, 2016).

Considering neuromodulatory factors within the amygdala involved in stress and anxiety, norepinephrine is released into the BLA immediately following a stressful experience and enhances memory of emotionally salient events while 5-HT is released during prolonged, uncontrollable stress (Roosendaal et al., 2009). During stressful conditions there is reduced expression of serotonergic receptors (5-HT_{2A}) leading to a down-regulation in activation of GABAergic BLA interneurons, and increased excitability of BLA neurons (Jiang et al., 2009). Other 5-HT receptors can increase BLA excitability and lead to anxiety related behaviors, but infusion of the 5-HT_{2C} receptor antagonist SB242,084 into the BLA prevents the display of anxiety related behaviors in response to prolonged stressors (Christianson et al., 2010; Pockros-Burgess et al., 2014; Zangrossi and Graeff, 2014).

3.1.3 Frustrative Nonreward—The last construct in NVS linked to the amygdala is Frustrative Nonreward, reactions elicited in response to the withdrawal or prevention of reward (Amsel, 1958; Gallup, 1965; Machado et al., 2009). Frustrative nonreward is distinct from both offensive aggression which, as an expression of dominance or competition for resources is included in the social domain, and defensive aggression, one potential response to acute threat (NVS Working Group, 2011). Tye (2010) and Badrinarayan (2012) demonstrated that there was a distinct population of cells in the amygdala recruited after ‘frustrative events’ which predicted the intensity of response, and that they were unlike those neurons activated in response to freezing inducing shocks (Purgert et al., 2012).

Frustration, harm avoidance, and fear are all powerful motivators of behavior and the amygdala is involved in each construct, processing early autonomic and nonconscious stimuli that signal danger for instance, but it is also vital for the opposite mode of influence in **Positive Valence Systems**.

3.2 Positive Valence Systems (PVS)

In a world of ambiguous rewards and outcomes, the amygdala is vital for updating the value of stimuli based on the current state of the organism. The PVS domain “primarily responsible for responses to positive motivational situations or contexts, such as reward seeking, consummatory behavior, and reward/habit learning” is linked to the amygdala by two related constructs.

3.2.1 Approach Motivation—Approach motivation is the instigation and direction of behavior and generally entails approach when the valence is positive (Elliot and Covington, 2001; Harmon-Jones et al., 2013). Making decisions with incomplete information is a common experience, especially so in ambiguous social encounters. Key for these circumstances is awareness of what the internal state of the individual in that moment is and thus the relative value of incentives; information supplied by the amygdala and areas, like the OFC, that it is connected with (Gottfried et al., 2003; Janak and Tye, 2015). In terms of motivated behavior, the mPFC seems to play an important role in balancing the relative contributions of the NAcc and the amygdala (Ernst et al., 2006). Recent optogenetic studies by Kim et al. (2017, 2016) have also revealed a strong role for a subpopulation of BLA projections to genetically distinct CeL and CeM neurons in driving appetitive behaviors.

3.2.2 Reward Learning—The differential response of the BLA as a result of positive or negative learning (or valence encoding) can be seen in human fMRI studies as well as electrophysiology studies of non-human primates discriminating between cues; when those cues are reversed, some ‘positive’ or ‘negative’ neurons stopped responding, or were inhibited, demonstrating that it was not the sensory features of the cue that were triggering them, but the values they represented (Gottfried et al., 2003; Paton et al., 2006; Shabel and Janak, 2009). In rodents BLA neurons can be distinguished not only based on where they connect to, e.g. the NAc, part of the reward network, but also by what genes they express, as well as their firing patterns, allowing more fine grained dissections of how positive and negative signals are encoded in the brain (Beyeler et al., 2016; Jennings et al., 2013; Namburi et al., 2015).

OTR expression is an important modulator of the reward network. During reward anticipation, healthy individuals carrying the OTR rs2268493 autism risk allele showed decreased activation of the right amygdala and other parts of the reward network compared to non-carriers and heterozygotes (Damiano et al., 2014). Importantly, lesioning the amygdala does not impair all reward based behaviors, only those specifically related to changing reward value (Machado and Bachevalier, 2007). There is evidence of increased BOLD activity in the amygdala during monetary reward-learning and decision-making tasks. For example, the framing effect from economics where a participant’s decisions about which option to select is manipulated by presenting the choice in terms of either potential loss or gain; this effect is correlated with activation of the amygdala (De Martino et al., 2006).

The amygdala’s function in reward processing is part of a network of brain regions (Fig. 2) including the orbital frontal cortex (OFC) and NAc (Montague and Berns, 2002). Lesions that disconnect the amygdala from the OFC results in deficits on reward learning tasks

(Baxter et al., 2000). It appears that the amygdala acquires information about the value of stimuli and the OFC uses this to guide choices (Happé and Frith, 2014). Disrupting the connection between the amygdala and NAc can lead to deficits in instrumental behavior toward rewards (Ambroggi et al., 2008). Following the acquisition of information, the amygdala communicates the sensory aspects of stimuli to the NAc to guide approach behavior and reward learning by modulating dopaminergic neurons (Jackson and Moghaddam, 2001; Stuber et al., 2011).

The PVS domain, and within it the amygdala, has a fundamental role in initiating, and shaping activities, but command of thoughts and actions requires input from **Cognitive Systems**.

3.3 Cognitive Systems

In addition to the primary role of the amygdala in modulating emotional states, it has a crucial function at the perceptual level in integrating sensory information about stimuli. This domain, “responsible for various cognitive processes” covers a very large range of behaviors, one of which is the construct of attention, or salience.

3.3.1 Attention—The process by which the brain assigns attention to various stimuli depends on careful coordination of the top-down, value assigning areas, such as the dorsolateral PFC and inferior parietal cortex, associated respectively with appraisal and reappraisal (Ochsner et al., 2012); and the bottom-up middle frontal gyrus and superior parietal cortex, responsible for directing the visual gaze (Ferri et al., 2016). This coordination takes place within the amygdala in conjunction with the anterior insula and ACC (Fig. 2) as part of the larger saliency network (Jacobs et al., 2012; Rai et al., 2012). Interestingly, there is increased functional connectivity between the amygdala and the precuneus and dorsal ACC when resisting distraction particularly from arousing, negative images (Ferri et al., 2016; Kanske et al., 2011). This balancing of top-down control and bottom-up input requires the precise functioning of the inhibitory circuits within the amygdala that regulate its activity (Rainnie et al., 1991a, 1991b; Truitt et al., 2007).

Perception of fearful stimuli could be considered one highly specialized and conserved task of the amygdala within the larger context of salience assigning (Jacobs et al., 2012). To that end, some inputs from sensory thalami, and nociceptive signals from the pontine parabrachial nucleus (PBN), bypass the LA and project directly to the CeA (Fig. 2), allowing for very rapid and adaptable responses to danger signals (Keifer et al., 2015). The amygdala also has a role in gating connections between the cortex to the hippocampus and striatum tagging experiences with emotional marks, of any valence, which increases their salience and improves recall and learning (Hurlemann et al., 2007; Paz and Pare, 2013; Vuilleumier and Driver, 2007).

Attention control is vitally important for navigating any number of situations, but especially those involving **Social Processes**.

3.4 The Social Processes

These constructs “mediate responses to interpersonal settings of various types, including perception and interpretation of others’ actions.”

3.4.1 Affiliation and Attachment—The first construct in the social domain, Affiliation and Attachment, concerns both the initial engagement in positive interactions as well as the development of bonds that reinforce and motivate the communication constructs. In both non-human primates and humans, amygdala volume is positively correlated with the size of an individual’s social network (Bickart et al., 2011; Zhang et al., 2012). The enlargement of the BLA (Section 2.2) is correlated with, and may be the result of, a concomitant expansion in the cortical regions related to social behavior that it is connected with (Janak and Tye, 2015; Stephan and Andy, 1977). Direct evidence for the role of the amygdala in social motivation can be seen by lesioning the BLA which causes rats to stop preferring prosocial rewards (Hernandez-Lallement et al., 2016). In addition, optogenetic stimulation of BLA projections to the mPFC reduces social interaction and increases anxiety-like behavior, while inhibition of that same pathway increases social interaction in mice (Felix-Ortiz et al., 2016).

Particularly important in the discussion of attachment and the amygdala is the function of OT. This neuropeptide is released in key socio-emotional brain regions, including the amygdala, which is rich in OTRs. OT is known to modulate a wide range of social behaviors, such as maternal attachment, pair bonding, social recognition and empathy-based behaviors (Burkett et al., 2016; Ferguson et al., 2001; Johnson et al., 2016; Rilling and Young, 2014). OT acting on the amygdala also enhances social attachment, possibly by reducing fear and anxiety (section 3.1).

In accordance with the animal literature, there is substantial evidence in humans showing that OT plays a crucial role in maternal care and maternal motivation. Though the circuitry of human parental care is more elaborate than in rodent species, the reward and sub-cortical components involved in the motivational and emotional aspects of parenting are common to both. This circuitry includes (Fig. 2) the hypothalamus, ventral striatum, and amygdala (Feldman, 2015).

3.4.2 Social Communication—The next construct in Social Processes, Social Communication, and the sub-construct Reception of Facial Communication (including either implicit or explicit communication such as affect recognition, facial recognition and characterization) clearly involves the amygdala, particularly in the perception of emotions that signal danger, e.g. fear or anger (Adolphs et al., 1999, 1998; Gamer and Buchel, 2009). The amygdala’s role was formerly thought to be only in automatic and non-conscious rapid processing of such stimuli (LeDoux, 1996; Morris et al., 1999; Öhman et al., 2007). Today, we know that the purview of the amygdala is broad, neuroimaging studies show that the amygdala is activated in response to all faces, and not only to fearful expressions (Fitzgerald et al., 2006). These studies also show that the involvement of the amygdala in processing information from faces is complex, can show inter-individual variability, and is dependent on context (Adams et al., 2003; Kim et al., 2004).

The involvement of the amygdala in controlling attention and ambiguity resolution extends to this construct: SM's fear recognition impairment (Section 2.2) appears to arise from a failure to make spontaneous use of the eye region of faces, a potent source of disambiguating social information (Adolphs et al., 2005). In addition, it has been shown that when interpreting equivocal expressions positive interpretations are associated with decreased amygdala activation, and negative with increased (Kim et al., 2003; M. J. Kim et al., 2011). The amygdala's response to faces is also dependent on the motivational state of the perceiving individual; Radke (2015) discovered that testosterone administration biased individuals towards threat approach and increased amygdala activation, but decreased activity during threat avoidance. Anatomical differences affect facial reception as well; Zhang (2012) found a strong correlation in monkeys between larger amygdala and longer periods of gazing into the eye region of other monkeys. There is also evidence that the function of the amygdala in interpreting gaze changes throughout development, with amygdala size in children positively correlated to *cognitive* mental state inferences and in adults to *emotional* ones (Rice et al., 2014).

Clearly the amygdala has a significant role in social bonding and communication, how important is it for the last domain in RDoC, **Arousal and Regulatory Systems**?

3.5 Arousal and Regulatory Systems

These constructs “are responsible for generating activation of neural systems as appropriate for various contexts, and providing appropriate homeostatic regulation of such systems as energy balance and sleep.”

3.5.1 Arousal—In studies of arousal, which is distinct from both salience and anxiety, activation of the amygdala is linked with increased vigilance (Pessoa, 2011). After electrostimulation of a cat's amygdala Ursin and Kaada (1960) noted an “attention response” and orienting around the environment. Stimulation of the CeA's connection to the basal forebrain results in more activation of cholinergic neurons, including those that innervate the BLA (Fig. 2), increasing the signal-to-noise ratio therein (Davis and Whalen, 2001; Unal et al., 2015). In monkeys that have undergone neonatal removal of the amygdala there is an enhancement of CRF activity and HPA axis activation suggesting that early in development the amygdala acts as a brake on the arousal system but switches to an activator in maturity (Raper et al., 2014).

3.5.2 Circadian Rhythms—Coordinating biological activity to times of day and the light-dark cycle is a function of the suprachiasmatic nucleus (SCN) and the rhythmic expression of ‘clock’ genes that serve as transcription factors (Reppert and Weaver, 2002). These same genes are expressed in other subordinate sites throughout the brain and body, including the amygdala (Harbour et al., 2014).

4. Amygdala dysfunction in Autism

The separation of a heterogeneous clinical population by biomarkers holds great promise in the treatment of many complex psychiatric disorders but particularly in autism (Clementz et al., 2016; Loth et al., 2016). Using biomarkers, such as characteristics of the amygdala, to

parcellate ASD allows the development of targeted therapies and facilitates overall understanding of autism.

4.1 Alterations in Development and Neurobiology

One of the hallmarks of autism is that it is a developmental disorder; though diagnosis typically occurs between 2-7 years, precursor symptoms can be identified even in the first six months of infancy (Hazlett et al., 2017; Jones and Klin, 2013; Mandell et al., 2005; Mazurek et al., 2014). In healthy subjects the amygdala continues to elaborate throughout childhood and adolescence, some connections with forebrain regions are still being strengthened even in adulthood, most notably with the mPFC (Chareyron et al., 2012; Ehrlich et al., 2012; Ernst et al., 2006; Lenroot and Giedd, 2006; Schumann et al., 2011). Indeed, the amygdala has more connections with distinct parts of the brain than almost any other region (Pessoa, 2014, 2008; Petrovich et al., 2001; Young et al., 1994). It is important then to see how exactly prominent characteristics of autism can be linked with altered amygdala development.

While many studies have found enlargement of the amygdala in autism, others have found no change, or a decrease in volume (Aylward et al., 1999; Haar et al., 2016; Nordahl et al., 2012; summary in Allely et al., 2013). We believe this is evidence of subgroups of individuals with autism, and that improvements in care and treatment require a close examination of these seemingly discordant findings, and replication with larger groups (Müller and Amaral, 2017). In addition to altered development in the amygdala, we will look at changes in connectivity with major neural networks, and how these variations compare symptomatically with damage to the amygdala that can come from UWD (section 2.2) or other sources.

4.1.1 Volume differences—Enlarged amygdala were among the first brain anomalies to be identified in individuals with autism (Bauman and Kemper, 1985; Kemper and Bauman, 1993; Schumann et al., 2004; Sparks et al., 2002). This increase is age dependent; Nordahl et al. (2012) found the amygdala was enlarged from 2-4 years and that the rate of increase year over year was higher in children with autism. An enlarged amygdala was also found to correlate with more severe symptoms in toddlers (Schumann et al., 2009). But from 8-14 years and older the increase slowed or reversed, no differences are seen between adolescents with typical development vs. those with autism (Barnea-Goraly et al., 2014).

This early overgrowth in the amygdala in children with autism could be related to later excessive pruning, accounting for volume differences seen between children, but not adults, with autism (Amaral et al., 2008). Dziobek et al. (2010) and Eilam-Stock et al. (2016) found no size differences in adult individuals with autism, although Murphy et al. (2012) found significant increases in amygdala volume in their comparison of individuals with Asperger's vs. typically developing controls. One important proviso is that there is a significant degree of heterogeneity. While Nordahl et al. (2012) found that on average children with autism had enlarged amygdala, there were a considerable number whose amygdala development was on par or decreased relative to typically developing individuals.

Does normalization of amygdala volumes in older adolescents and adult individuals with autism correlate with improvements in social abilities? Although adolescent subjects with autism's amygdala growth does not differ on the whole from typically developing subjects, increases in eye contact in those with autism correlates with increased amygdala size (Barnea-Goraly et al., 2014). In children with autism there is a positive relationship between amygdala volume, eye gaze, and response to joint attention, with smaller amygdala predicting less eye gaze, stronger symptoms, and non-response to bids for attention (Mosconi et al., 2009; Nacewicz et al., 2006). Unaffected siblings also have smaller amygdala and decreased eye gaze but they, like typically developing controls, do not show a relationship between amygdala volume and eye gaze (Dalton et al., 2007). Interestingly, Williams syndrome, a rare autosomal disorder characterized by mild to moderate cognitive impairments and hypersociability, is associated with an *enlarged* amygdala, along with an overall decrease in brain volume (Schumann et al., 2011). Ultimately, the change in size could be a consequence, rather than cause, of social interaction (Zalla and Sperduti, 2013). When the number of individuals in a monkey's social group was altered, the volume of the amygdala also changed (Sallet et al., 2011). If this translates to humans, then amygdala size during development could serve as a biomarker for effective therapeutic intervention.

A stereological comparison of cell types in the amygdala between those with autism and typically developing controls found that before 20 years of age there were no differences, but afterwards there was a significant drop in myelinating oligodendrocytes (Morgan et al., 2014). Thus, it is essential to look beyond gross anatomical differences of the amygdala to alterations in its activity, how it connects with other areas of the social brain, and what significance this has for the pathogenesis of autism.

4.2.1 Abnormal amygdala activity and social cognition—Social deficits are a second hallmark of ASD. The amygdala, along with regions like the superior temporal sulcus (STS) and OFC (Fig. 2), are an essential part of the neurobiological substrate of social cognition. Early neuroimaging of ASD individuals revealed decreased activation in the amygdala and frontal areas during social tasks such as reading the mind in the eyes (Baron-Cohen et al., 1997). Other studies revealed a lack of amygdala activity during the implicit processing of emotional facial stimuli (Critchley et al., 2000). Functional neuroimaging studies have found that abnormal amygdala activation in ASD is, reminiscent of SM, related to decreased eye fixations to faces (Dalton et al., 2005; Kliemann et al., 2012).

In addition to a lack of amygdala activation, individuals with autism show a significant reduction in functional connectivity between the amygdala, principally the BLA, with cortical, visual, and parietal regions involved in perception during resting state (Rausch et al., 2016). This reduction in functional connectivity was also seen between the amygdala and the salience network including the insula (Von Dem Hagen et al., 2014). However, there are a number of attentional and social deficits in ASD, independent of the amygdala, and it's bilateral destruction is insufficient to produce autism, as determined by clinical measures (Paul et al., 2010; Wang et al., 2014).

4.1.3 Damage to the amygdala—In comparative studies between people with autism and focal amygdala lesions (Section 2.2) there are several similarities in their behavioral phenotypes: abnormal social judgment regarding the rating of trustworthiness and approachability in faces, difficulties in identifying emotional features from facial expressions, impairments in judging faces with negative affect such as fear and anger, and abnormalities in attributing social meaning to ambiguous and moving shapes (Adolphs et al., 2001; Klin, 2000; Pelphrey et al., 2002). What's more, individuals whose amygdala are removed due to a tumor or refractory epilepsy also display significant deficits in Theory of Mind tasks, but only if the damage occurred in their childhood (<16 years old, Shaw et al., 2004). Loss of the amygdala from UWD appears to impair the saliency network's (Section 3.3.1) ability to prioritize emotionally charged inputs (Hurlemann et al., 2007). This phenomenon is similar to the weak central coherence model of autism from Happé and Frith (2006): a focus on details rather than the whole.

This suggests autism does not arise exclusively from the functioning of the amygdala, as a complete lesion would be expected to result in the most severe social dysfunction (Paul et al., 2010). It suggests instead a change in *how* the amygdala develops and functions is responsible, and that modeling these developmental disruptions can be a useful way to understand autism.

4.2 Valproic acid and autism

Valproic acid (VPA) is the most widely prescribed anti-epileptic drug in the world, as well as a potent teratogen (Perucca, 2002). VPA exposure during pregnancy is linked with fetal valproate syndrome (FVS) characterized by dysmorphic facial features, spina bifida, lower IQ, delayed language, and autism in approximately 9% of those exposed (Ardinger et al., 1988; Clayton-Smith and Donnai, 1995; Jäger-Roman et al., 1986; Rasalam et al., 2005; Williams et al., 2001; Williams and Hersh, 1997). FVS is thought to result from the effects of VPA on the developing central nervous system in the 1st trimester of pregnancy (Binkerd et al., 1988; Christianson et al., 1994; but see Roullet et al., 2013). VPA is to date the only environmental agent so strongly linked to autism (Bromley et al., 2013; Christensen et al., 2013; Ornoy et al., 2015)

How exactly the epigenetic changes wrought by VPA contribute to FVS and autism is still unknown. By promoting the acetylation of histones, VPA makes DNA more accessible to transcription factors, altering the course of development, and ultimately the neural systems underlying cognitive and social domains (for review see Chomiak et al., 2013). Much of the work in explicating this process has been done in animal models: As originally developed by Rodier (1997) fetal exposure to VPA resulted in rats that were less sociable and more anxious, a finding that has been replicated many times (K. C. Kim et al., 2011; Mabunga et al., 2015; Markram et al., 2008; Roullet et al., 2013 but see Cohen et al., 2013; Štefánik et al., 2015). VPA exposed rats also display hyperserotonemia, one of the oldest known biomarkers of autism (Anderson et al., 2011, 1990; Hranilovic et al., 2007; Narita et al., 2002; Takahashi et al., 1976).

The amygdala is significantly altered along several dimensions following prenatal VPA exposure. In rats, VPA changes the volume of the amygdala and drastically alters dendritic

arbors of principal neurons, following a progression that is comparable to humans, with initial overgrowth of spines at the time of weaning, followed by a retraction in the fully-grown rat (Amaral et al., 2008; Bringas et al., 2013; Sosa-Díaz et al., 2014). This may be related to increases in the excitability of LA neurons, facilitating LTP, and contributing to behavioral alterations in the model (Lin et al., 2013). VPA exposure changes the expression of many amygdala genes including *Homer1*, a notable finding as overexpression of *Homer1* in the amygdala leads to a reduction in social interaction and impaired fear conditioning (Banerjee et al., 2016; Oguchi-Katayama et al., 2013). We have also observed that VPA exposure in the rat significantly increased OTR expression in the female, but not male, CeA (authors, unpublished).

Two caveats: First, though rodent models of autism are crucial for better understanding the neurobiological mechanisms of social functioning, most of the behavioral tests that are used in these experiments do not mirror the complexity of social dysfunction seen in ASD. The autism phenotype is heterogeneous and not restricted to a lack of social approach, the gold standard measure in the majority of animal studies (Moy et al., 2004; Tordjman et al., 2007). Second, the use of a teratogen as the agent for creating autism-like symptoms must be reconciled with autism's high heritability (Hallmayer et al., 2011). But the fact that this model is not the product of a single gene mutation makes it in many ways more representative of the broad swath of individuals whose autism is not related to any known monogenic cause. Autism is associated with higher rates of de novo protein truncating mutations, and more severe autistic symptoms and intellectual disability are associated with the highest rates of mutations (Kosmicki et al., 2017).

4.3 Autism, anxiety, and oxytocin

Besides the core social symptoms in ASD, there are several medical, neurological, and psychiatric comorbidities, including disorders of: anxiety, depression, obsessive-compulsiveness, sleep, irritability, and hyperactivity. Anxiety disorders in particular are much more common in individuals with autism than in the general population, with estimates ranging from 40-55% of young people with autism also exhibiting one or more anxiety disorders (van Steensel et al., 2011; White et al., 2009). Currently, there are two FDA approved drugs (risperidone and aripiprazole) for comorbid symptoms of irritability. There are no approved treatments for the social dysfunctions of ASD or for anxiety disorders in these individuals.

In addition to reducing fear and anxiety (Section 3.1.2) intranasal OT has been shown to promote social functioning, gaze to the eyes, and theory of mind, in both typically developing individuals, and those with autism, as well as activity in, and functional connectivity between, key brain regions involved in social and emotional processing, such as the amygdala (Andari et al., 2016, 2010; Aoki et al., 2014; Auyeung et al., 2015; Gordon, 2014; Guastella et al., 2010; Watanabe et al., 2016, 2015, 2014). A recent meta-analysis reported significant associations between ASD and the OXTR SNPs rs7632287, rs237887, rs2268491 and rs2254298 (LoParo and Waldman, 2015). Importantly, one of these SNPs (rs237887) was found to be strongly associated with recognition memory in individuals with ASD, their parents and their siblings, suggesting a critical role of the OT system in social

recognition (Skuse et al., 2014). We have also demonstrated that the administration of intranasal OT triggered reciprocal cooperation between individuals with ASD and partners in an interactive social game (Andari et al., 2010). Finally, OT differentially altered BOLD activity in the amygdala during an interactive social game, reducing it with an equitable partner but increasing it with an untrustworthy one (Andari et al., 2016).

More research is needed to better understand the mechanism by which OT affects social cognition in these individuals, and whether it can also be a treatment for co-occurring anxiety (Andari, 2016).

5. Conclusion: synthesis of RDoC, the amygdala, and autism

While we have described each of the domains and constructs of RDoC, and their associated neural networks, as independent systems, they exist in dynamic equilibrium. Evaluating competing needs: fear of punishment vs. possibility of reward, or interest in complex systems vs. in conspecifics, is one of the essential tasks of the brain across all species and all levels of development. Disruption of this mechanism, manifested in one construct or many is, in a sense, the definition of mental illness; especially so in autism.

Consider an amygdala dependent ability that cuts across several RDoC domains: disambiguation. It is a difficult task, though one that typically developing individuals accomplish with relative ease: **Attention** must be switched across multiple contextual levels, salient details extracted and judged. The challenge increases if the information is **social** in nature, given the unwritten mores and subtleties of that domain. Imagine how, **stressful**, and **frustrating** that task would be, and how little **motivation** to pursue it, were it not intrinsically **rewarding**. Mental bandwidth is not limitless and, once exhausted, individuals with autism might retreat to spaces where cause and effect are clear, and where they have control over the stimuli they experience i.e. stereotypes.

This line of thinking about autism has several implications. It explains some discordant fMRI findings in the amygdala: hypoactivation when bandwidth demand is low and potent stimuli such as the eyes are unengaged, hyperactivation otherwise, such as when the subject is specifically instructed to look at the eyes (Baron-Cohen et al., 1999; Critchley et al., 2000; Dalton et al., 2005; Kleinhans et al., 2010; Kliemann et al., 2012; Schultz, 2005; Zürcher et al., 2013). The highest degree of amygdala activation, and association with symptom severity, occurs when judging ambiguous expressions (Swartz et al., 2013; Tottenham et al., 2014). Madarasz (2016) directly studied the role of the amygdala in disambiguation by examining changing cue contingencies; neurons in the LA not only store sensory associations, they update dynamically when new information is presented. Kim et al. (2016) found two populations of genetically distinct, positive and negative valence responding neurons in the BLA that not only independently drove or repressed behaviors, but mutually inhibited one another. Meanwhile, Burgos-Robles et al. (2017) demonstrated that when rats were faced with competing fear and reward signals it was a connection from the BLA to the prelimbic mPFC that predicted animal's behavior, and that that outcome could be biased towards or away from freezing by activation or inhibition of that connection. Together these studies suggest a potential neurological substrate involving the amygdala that may account

for why some individuals with autism have difficulty discerning important details within complex inputs and committing to a course of action.

As the classification of the amygdala as a single brain structure was questioned decades ago, there are calls now to abandon ASD as unsuitable for neurobiological research (Davis and Whalen, 2001; Swanson and Petrovich, 1998; Waterhouse et al., 2016). We, like Müller and Amaral (2017), think this is unwarranted. Instead, by embracing RDoC, grasping how variations from neuron to brain are reflected dimensionally across domains of behavior, in typically developing and individuals with autism, we can fulfil the promise of individualized medicine, finding optimal outcomes for everyone with this disorder. We conclude that there are likely dysfunctions in the amygdala of many, but not all, individuals with autism, and that investigation of this relationship remains a very fruitful area of research.

In addition to adoption of RDoC, four specific issues in this field call out for further study: 1) Much of autism research (and biomedical research generally) is done in males, and while a gender disparity in autism does exist, bias in subject selection may serve to exaggerate it (Beery and Zucker, 2011; Giarelli et al., 2010; Mandy et al., 2012; Robinson et al., 2013; Stone et al., 2004; Werling and Geschwind, 2013a, 2013b). Evidence strongly suggests the amygdala is sexually dimorphic, and this may be related to the higher incidence of ASD in males but more studies in females with ASD are needed to explore this possibility (Baron-Cohen et al., 2011, 2005).

2) Most neuroimaging studies of individuals with autism look at only a single time point, while the few longitudinal studies have only 1-4 years between measurements (Courchesne et al., 2011; Hazlett et al., 2017; Mosconi et al., 2009; Nordahl et al., 2012). Since amygdala connections are elaborated well into the second decade of life, imaging studies are clearly needed on this timescale, especially if monitoring the size and growth trajectory of the amygdala could serve as a biomarker for effective therapeutic intervention (Ernst et al., 2006; Lenroot and Giedd, 2006; Ruggeri et al., 2014; Uddin et al., 2011).

3) While fMRI studies are useful for finding large scale changes in activity and connectivity, experiments using implanted electrodes would reveal precisely how information flow between the amygdala and social brain is altered (Rutishauser et al., 2013). Since electrophysiological studies in humans are necessarily in populations with significant comorbidities, such as epilepsy, this is one area that animal models, e.g. VPA, would prove useful (Jeffrey et al., 2013; MacFabe et al., 2007; Meletti et al., 2012; Mormann et al., 2015; Rutishauser et al., 2015; Sato et al., 2011).

4) Promising therapies, such as oxytocin, can be fully realized only through a detailed understanding of how exactly they do, or do not, work in the brains of individuals with autism, especially in critical areas like the amygdala (Bales et al., 2013; Young and Barrett, 2015). To translate research on the effects of acute administration of intranasal OT into clinical therapeutics, it will be necessary to adopt a precision medicine approach to determine OT's targets in autism. More studies and replications of OT treatment's effect on functioning and development, particularly with chronic vs. acute exposure, are needed.

It is our hope that researchers can move to address these and other significant gaps in knowledge, deepening our understanding of a remarkable brain area, and improving the wellbeing of millions of individuals with neuropsychiatric disorders.

Acknowledgments

Funding:

T. H. is supported by NIH grant 5P50MH100023-05

E.A. is supported by NIH grants 1P50MH100023 and P51OD11132 to YNPRC.

D. R. is supported by NIH grants 5P50MH100023-05, 5R01MH072908-12 and 5R01MH069852-12

Bibliography

- Acheson D, Feifel D, De Wilde S, McKinney R, Lohr J, Risbrough V, 2013 The effect of intranasal oxytocin treatment on conditioned fear extinction and recall in a healthy human sample. *Psychopharmacology (Berl)*. 229, 199–208. doi: 10.1007/s00213-013-3099-4 [PubMed: 23644911]
- Adams RB, Gordon HL, Baird AA, Ambady N, Kleck RE, 2003 Effects of Gaze on Amygdala Sensitivity to Anger and Fear Faces. *Science* (80-.). 300, 1536 LP–1536.
- Adolphs R, 2010 What does the amygdala contribute to social cognition? *Ann. N. Y. Acad. Sci* 1191, 42–61. doi: 10.1111/j.1749-6632.2010.05445.x [PubMed: 20392275]
- Adolphs R, 2009 The social brain: neural basis of social knowledge. *Annu. Rev. Psychol* 60, 693–716. doi: 10.1146/annurev.psych.60.110707.163514 [PubMed: 18771388]
- Adolphs R, Gosselin F, Buchanan TW, Tranel D, Schyns P, Damasio AR, 2005 A mechanism for impaired fear recognition after amygdala damage. *Nature* 433, 68–72. [PubMed: 15635411]
- Adolphs R, Sears L, Piven J, 2001 Abnormal Processing of Social Information from Faces in Autism. *J. Cogn. Neurosci* 13, 232–240. doi: 10.1162/089892901564289 [PubMed: 11244548]
- Adolphs R, Tranel D, Damasio AR, 1998 The human amygdala in social judgment. *Nature* 393, 470–474. doi: 10.1038/30982 [PubMed: 9624002]
- Adolphs R, Tranel D, Damasio H, Damasio A, 1994 Impaired recognition of emotion in facial expressions following bilateral damage to the human amygdala. *Nature*. doi:10.1038/372669a0
- Adolphs R, Tranel D, Hamann S, Young ACAA, Phelps E, Lee GP, Damasio AR, 1999 Recognition of Facial Emotion in Nine Subjects with Bilateral Amygdala Damage. *Neuropsychologia* 37, 1111–1117. [PubMed: 10509833]
- Allely CS, Gillberg C, Wilson P, 2013 Neurobiological abnormalities in the first few years of life in individuals later diagnosed with Autistic Spectrum Disorder: A review of recent data. *Behav.Neurol* 2014.
- Amaral DG, Schumann CM, Nordahl CW, 2008 Neuroanatomy of autism. *Trends Neurosci* 31, 137–45. doi: 10.1016/j.tins.2007.12.005 [PubMed: 18258309]
- Ambroggi F, Ishikawa A, Fields HL, Nicola SM, 2008 Basolateral amygdala neurons facilitate reward-seeking behavior by exciting nucleus accumbens neurons. *Neuron* 59, 648–61. doi: 10.1016/j.neuron.2008.07.004 [PubMed: 18760700]
- American Psychiatric Association, 2013 Diagnostic and statistical manual of mental disorders (5th ed.), 5th ed. Washington, DC.
- Amsel A, 1958 The role of frustrative nonreward in noncontinuous reward situations. *Psychol. Bull* 55, 102–119. doi: 10.1037/h0043125 [PubMed: 13527595]
- Andari E, 2016 The Need for a Theoretical Framework of Social Functioning to Optimize Targeted Therapies in Psychiatric Disorders. *Biol. Psychiatry* 79, e5–e7. doi:10.1016/j.biopsych.2015.11.014 [PubMed: 26723109]
- Andari E, Duhamel J-R, Zalla T, Herbrecht E, Leboyer M, Sirigua A, 2010 Promoting social behavior with oxytocin in high-functioning autism spectrum disorders. *Proc. Natl. Acad. Sci* 107, 4389–4394. doi: 10.1073/pnas.0910249107 [PubMed: 20160081]

- Andari E, Richard N, Leboyer M, Sirigu A, 2016 Adaptive coding of the value of social cues with oxytocin, an fMRI study in autism spectrum disorder. *Cortex* 76, 79–88. doi:10.1016/j.cortex.2015.12.010 [PubMed: 26872344]
- Anderson GM, Hertzog ME, McBride P. a, 2011 Brief Report: Platelet-Poor Plasma Serotonin in Autism. *J. Autism Dev. Disord*, doi: 10.1007/s10803-011-1371-1
- Anderson GM, Home WC, Chattegee D, Cohen DJ, 1990 The Hyperserotonemia of Autism. *Ann. N. Y. Acad. Sci* 600, 331–340. [PubMed: 2252319]
- Anney R, Klei L, Pinto D, Almeida J, Bacchelli E, Baird G, Bolshakova N, B?lte S, Bolton PF, Bourgeron T, Brennan S, Brian J, Casey J, Conroy J, Correia C, Corsello C, Crawford EL, De jonge M, Delorme R, Duketis E, Duque F, Estes A, Farrar P, Fernandez BA, Folstein SE, Fombonne E, Gilbert J, Gillberg C, Glessner JT, Green A, Green J, Guter SJ, Heron EA, Holt R, Howe JL, Hughes G, Hus V, Iglizzi R, Jacob S, Kenny GP, Kim C, Kolevzon A, Kustanovich V, Lajonchere CM, Lamb JA, Law-Smith M, Leboyer M, Le couteur A, Leventhal BL, Liu XQ, Lombard F, Lord C, Lotspeich L, Lund SC, Magalhaes TR, Mantoulan C, McDougle CJ, Melhem NM, Merikangas A, Minshew NJ, Mirza GK, Munson J, Noakes C, Nygren G, Papanikolaou K, Pagnamenta AT, Parrini B, Paton T, Pickles A, Posey DJ, Poustka F, Ragoussis J, Regan R, Roberts W, Roeder K, Roge B, Rutter ML, Schlitt S, Shah N, Sheffield VC, Soorya L, Sousa I, Stoppioni V, Sykes N, Tancredi R, Thompson AP, Thomson S, Tryfon A, Tsiantis I, Van Engeland H, Vincent JB, Volkmar F, Vorstman JAS, Wallace S, Wing K, Wittemeyer K, Wood S, Zurawiecki D, Zwaigenbaum L, Bailey AJ, Battaglia A, Cantor RM, Coon H, Cuccaro ML, Dawson G, Ennis S, Freitag CM, Geschwind DH, Haines JL, Klauck SM, McMahon WM, Maestrini E, Miller J, Monaco AP, Nelson SF, Numberger JJ, Oliveira G, Parr JR, Pericak-Vance MA, Piven J, Schellenberg GD, Scherer SW, Vicente AM, Wassink TH, Wijsman EM, Betancur C, Buxbaum JD, Cook EH, Gallagher L, Gill M, Hallmayer J, Paterson AD, Sutcliffe JS, Szatmari P, Veland VJ, Hakonarson H, Devlin B, 2012 Individual common variants exert weak effects on the risk for autism spectrum disorders. *Hum. Mol. Genet* 21, 4781–4792. doi:10.1093/hmg/dds301 [PubMed: 22843504]
- Aoki Y, Yahata N, Watanabe T, Takano Y, Kawakubo Y, Kuwabara H, Iwashiro N, Natsubori T, Inoue H, Suga M, Takao H, Sasaki H, Gono W, Kunimatsu A, Kasai K, Yamasue H, 2014 Oxytocin improves behavioural and neural deficits in inferring others' social emotions in autism. *Brain* 137, 3073–3086. doi:10.1093/brain/awu231 [PubMed: 25149412]
- Ardinger HH, Atkin JF, Blackston RD, Elsas LJ, Clarren SK, Livingstone S, Flannery DB, Pellock JM, Harrod MJ, Lammer EJ, 1988 Verification of the fetal valproate syndrome phenotype. *Am. J. Med. Genet* 29, 171–85. doi:10.1002/ajmg.1320290123 [PubMed: 3125743]
- Arruda-Carvalho M, Wu W-C, Cummings KA, Clem RL, 2017 Optogenetic examination of prefrontal-amygdala synaptic development. *J. Neurosci* 37, 3097–16. doi: 10.1523/JNEUROSCI.3097-16.2017
- Asperger H, 1944 Die “Autistische Psychopathen” im Kindesalter. *Arch. Psychiatr. Nervenkr* 117, 76–136. doi: 10.1007/BF01837709
- Auyeung B, Lombardo MV, Heinrichs M, Chakrabarti B, Sule A, Deakin JB, Bethlehem RAI, Dickens L, Mooney N, Sippel JAN, Thiemann P, Baron-Cohen S, 2015 Oxytocin increases eye contact during a real-time, naturalistic social interaction in males with and without autism. *Transl. Psychiatry* 5, e507. doi: 10.1038/tp.2014.146 [PubMed: 25668435]
- Avery SN, Clauss JA, Blackford JU, 2015 The Human BNST: Functional Role in Anxiety and Addiction. *Neuropsychopharmacology* 41, 126–141. doi:10.1038/npp.2015.185 [PubMed: 26105138]
- Aylward EH, Minshew NJ, Goldstein G, Honeycutt N. a, Augustine a M., Yates KO, Barta PE, Pearlson GD, 1999 MRI volumes of amygdala and hippocampus in non-mentally retarded autistic adolescents and adults. *Neurology* 53, 2145–50. doi: 10.1212/WNL.53.9.2145 [PubMed: 10599796]
- Badrinarayan a., Prater KE, Orsini C. a., 2012 The Role of the Central Amygdala in Selecting Circuits and Responses. *J. Neurosci* 32, 8431–8433. doi: 10.1523/JNEUROSCI.1820-12.2012 [PubMed: 22723682]
- Bales KL, Perkeybile AM, Conley OG, Lee MH, Guynes CD, Downing GM, Yun CR, Solomon M, Jacob S, Mendoza SP, 2013 Chronic intranasal oxytocin causes long-term impairments in partner

- preference formation in male prairie voles. *Biol. Psychiatry* 74, 180–188. doi: 10.1016/j.biopsych.2012.08.025 [PubMed: 23079235]
- Banerjee A, Luong JA, Ho A, Saib AO, Ploski JE, 2016 Overexpression of Homer1a in the basal and lateral amygdala impairs fear conditioning and induces an autism-like social impairment. *Mol. Autism* 7, 16. doi:10.1186/s13229-016-0077-9 [PubMed: 26929812]
- Barger N, Stefanacci L, Semendeferi K, 2007 A comparative volumetric analysis of the amygdaloid complex and basolateral division in the human and ape brain. *Am. J. Phys. Anthropol* 134, 392–403. doi: 10.1002/ajpa.20684 [PubMed: 17661399]
- Barnea-Goraly N, Frazier TW, Piacenza L, Minshew NJ, Keshavan MS, Reiss AL, Hardan AY, 2014 A preliminary longitudinal volumetric MRI study of amygdala and hippocampal volumes in autism. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* 48, 124–128. doi:10.1016/j.pnpbp.2013.09.010
- Baron-Cohen S, Baron-Cohen S, Ring HA, Ring HA, Wheelright S, Wheelright S, Bullmore ET, Bullmore ET, Brammer MJ, Brammer MJ, Simmons AM, Simmons AM, Williams SC, Williams SC, 1999 Social intelligence in the normal and autistic brain: an fMRI study. *Eur. J. Neurosci* 11, 1891–1898. doi: 10.1046/j.1460-9568.1999.00621.x [PubMed: 10336657]
- Baron-Cohen S, Jolliffe T, Mortimore C, Robertson M, 1997 Another Advanced Test of Theory of Mind: Evidence from Very High Functioning Adults with Autism or Asperger Syndrome. *J. Child Psychol. Psychiatry* 38, 813–822. doi: 10.1111/j.1469-7610.1997.tb01599.x [PubMed: 9363580]
- Baron-Cohen S, Knickmeyer RC, Belmonte MK, 2005 Sex differences in the brain: implications for explaining autism. *Science* 310, 819–23. doi: 10.1126/science.1115455 [PubMed: 16272115]
- Baron-Cohen S, Lombardo MV, Auyeung B, Ashwin E, Chakrabarti B, Knickmeyer R, 2011 Why are Autism Spectrum conditions more prevalent in Males? *PLoS Biol* 9. doi:10.1371/journal.pbio.1001081
- Baron-Cohen S, Ring HA, Bullmore ET, Wheelwright S, Ashwin C, Williams SC, 2000 The amygdala theory of autism. *Neurosci. Biobehav. Rev* 24, 355–64. [PubMed: 10781695]
- Bauman M, Kemper TL, 1985 Histoanatomic observations of the brain in early infantile autism. *Neurology* 35, 866–74. doi: 10.1212/WNL.35.6.866 [PubMed: 4000488]
- Bauman MD, Amaral DG, 2005 The distribution of serotonergic fibers in the macaque monkey amygdala: An immunohistochemical study using antisera to 5-hydroxytryptamine. *Neuroscience* 136, 193–203. doi:10.1016/j.neuroscience.2005.07.040 [PubMed: 16182456]
- Bauman MD, Toscano JE, Mason WA, Lavenex P, Amaral DG, 2006 The expression of social dominance following neonatal lesions of the amygdala or hippocampus in rhesus monkeys (*Macaca mulatta*). *Behav. Neurosci* 120, 749–60. doi: 10.1037/0735-7044.120.4.749 [PubMed: 16893283]
- Baxter MG, Parker A, Lindner CCC, Izquierdo AD, Murray E. a, 2000 Control of response selection by reinforcer value requires interaction of amygdala and orbital prefrontal cortex. *J. Neurosci* 20, 4311–9. doi:10.1523/JNEUROSCI.4311-00.2000 [PubMed: 10818166]
- Bechara A, Tranel D, Damasio H, Adolphs R, others, 1995 Double dissociation of conditioning and declarative knowledge relative to the amygdala and hippocampus in humans. *Science* (80-.). 269, 1115.
- Beery AK, Zucker I, 2011 Sex bias in neuroscience and biomedical research. *Neurosci. Biobehav. Rev* 35, 565–572. doi:10.1016/j.neubiorev.2010.07.002 [PubMed: 20620164]
- Bethlehem RAI, van Honk J, Auyeung B, Baron-Cohen S, 2013 Oxytocin, brain physiology, and functional connectivity: A review of intranasal oxytocin fMRI studies. *Psychoneuroendocrinology* 38, 962–974. doi:10.1016/j.psyneuen.2012.10.011 [PubMed: 23159011]
- Beyeler A, Namburi P, Glover GF, Simonnet C, Calhoun GG, Conyers GF, Luck R, Wildes CP, Tye KM, 2016 Divergent Routing of Positive and Negative Information from the Amygdala during Memory Retrieval. *Neuron* 90, 348–361. doi: 10.1016/j.neuron.2016.03.004 [PubMed: 27041499]
- Bickart KC, Dickerson BC, Barrett LF, 2014 The amygdala as a hub in brain networks that support social life. *Neuropsychologia* 63, 235–48. doi:10.1016/j.neuropsychologia.2014.08.013 [PubMed: 25152530]
- Bickart KC, Wright CI, Dautoff RJ, Dickerson BC, Barrett LF, 2011 Amygdala volume and social network size in humans. *Nat. Neurosci* 14, 163–164. doi:10.1038/nn.2724 [PubMed: 21186358]

- Binkerd PE, Rowland JM, Nau H, Hendrickx AG, 1988 Evaluation of valproic acid (VPA) developmental toxicity and pharmacokinetics in Sprague-Dawley rats. *Fundam. Appl. Toxicol* 11, 485–93. doi: 10.1093/toxsci/11.1.485 [PubMed: 3146521]
- Bringas ME, Carvajal-Flores FN, López-Ramírez T. a, Atzori M, Flores G, 2013 Rearrangement of the dendritic morphology in limbic regions and altered exploratory behavior in a rat model of autism spectrum disorder. *Neuroscience* 241, 170–187. doi: 10.1016/j.neuroscience.2013.03.030 [PubMed: 23535253]
- Bromley RL, Mawer GE, Briggs M, Cheyne C, Clayton-Smith J, García-Fiñana M, Kneen R, Lucas SB, Shallcross R, Baker GA, 2013 The prevalence of neurodevelopmental disorders in children prenatally exposed to antiepileptic drugs. *J. Neurol. Neurosurg. Psychiatry* 84, 637–43. doi: 10.1136/jnnp-2012-304270 [PubMed: 23370617]
- Burghardt NS, Bauer EP, 2013 Acute and chronic effects of selective serotonin reuptake inhibitor treatment on fear conditioning: Implications for underlying fear circuits. *Neuroscience* 247, 253–272. doi: 10.1016/j.neuroscience.2013.05.050 [PubMed: 23732229]
- Burgos-Robles A, Kimchi EY, Izadmehr EM, Porzenheim MJ, Ramos-Guasp WA, Nieh EH, Felix-Ortiz AC, Namburi P, Leppla CA, Presbrey KN, Anandalingam KK, Pagan-Rivera PA, Anahtar M, Beyeler A, Tye KM, 2017 Amygdala inputs to prefrontal cortex guide behavior amid conflicting cues of reward and punishment. *Nat. Neurosci* 20, 824–835. doi: 10.1038/nn.4553 [PubMed: 28436980]
- Burkett JP, Andari E, Johnson ZV, Curry DC, de Waal FBM, Young LJ, 2016 Oxytocin-dependent consolation behavior in rodents. *Science* 351, 375–8. doi: 10.1126/science.aac4785 [PubMed: 26798013]
- Chareyron LJ, Banta Lavenex P, Amaral DG, Lavenex P, 2011 Stereological analysis of the rat and monkey amygdala. *J. Comp. Neurol* 519, 3218–3239. doi: 10.1002/cne.22677 [PubMed: 21618234]
- Chareyron LJ, Lavenex PB, Amaral DG, Lavenex P, 2012 Postnatal development of the amygdala: A stereological study in macaque monkeys. *J. Comp. Neurol* 520, 1965–1984. doi: 10.1002/cne.23023 [PubMed: 22173686]
- Chomiak T, Turner N, Hu B, 2013 What We Have Learned about Autism Spectrum Disorder from Valproic Acid. *Pathol. Res Int* 2013, 712758. doi: 10.1155/2013/712758
- Christensen J, Grønborg TK, Sørensen MJ, Schendel D, Parner ET, Pedersen LH, Vestergaard M, 2013 Prenatal valproate exposure and risk of autism spectrum disorders and childhood autism. *JAMA* 309, 1696–703. doi:10.1001/jama.2013.2270 [PubMed: 23613074]
- Christianson AL, Chesler N, Kromberg JG, 1994 Fetal valproate syndrome: clinical and neurodevelopmental features in two sibling pairs. *Dev. Med. Child Neurol* 36, 361–9. [PubMed: 7512516]
- Christianson JP, Ragole T, Amat J, Greenwood BN, Strong PV, Paul ED, Fleshner M, Watkins LR, Maier SF, 2010 5-Hydroxytryptamine 2C Receptors in the Basolateral Amygdala Are Involved in the Expression of Anxiety After Uncontrollable Traumatic Stress. *Biol. Psychiatry* 67, 339–345. doi:10.1016/j.biopsych.2009.09.011 [PubMed: 19914601]
- Chu H-Y, Ito W, Li J, Morozov a., 2012 Target-Specific Suppression of GABA Release from Parvalbumin Interneurons in the Basolateral Amygdala by Dopamine. *J. Neurosci* 32, 14815–14820. doi: 10.1523/JNEUROSCI.2997-12.2012 [PubMed: 23077066]
- Cinaz P, Güvenir T, Gönlü en G, 1993 Lipoid proteinosis: Urbach-Wiethe disease. *Acta Paediatr* 82, 892–3. doi: 10.1159/000027461 [PubMed: 8241657]
- Ciocchi S, Herry C, Grenier F, Wolff SBE, Letzkus JJ, Vlachos I, Ehrlich I, Sprengel R, Deisseroth K, Stadler MB, Müller C, Lüthi A, 2010 Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature* 468, 277–282. doi:10.1038/nature09559 [PubMed: 21068837]
- Clayton-Smith J, Donnai D, 1995 Fetal valproate syndrome. *J. Med. Genet* 32, 724–7. [PubMed: 8544193]
- Clementz BA, Sweeney JA, Hamm JP, Ivleva EI, Ethridge LE, Pearlson GD, Keshavan MS, Tamminga CA, 2016 Identification of Distinct Psychosis Biotypes Using Brain-Based Biomarkers. *Am. J. Psychiatry* 173, 373–84. doi: 10.1176/appi.ajp.2015.14091200 [PubMed: 26651391]

- Clugnet MC, LeDoux JE, 1990 Synaptic plasticity in fear conditioning circuits: Induction of LTP in the lateral nucleus of the amygdala by stimulation of the medial geniculate body. *J. Neurosci* 10, 2818–2824. [PubMed: 2388089]
- Cohen OS, Varlinskaya EI, Wilson CA, Glatt SJ, Mooney SM, 2013 Acute prenatal exposure to a moderate dose of valproic acid increases social behavior and alters gene expression in rats. *Int. J. Dev. Neurosci* 31, 740–750. doi:10.1016/j.ijdevneu.2013.09.002 [PubMed: 24055786]
- Courchesne E, Campbell K, Solso S, 2011 Brain growth across the life span in autism: Age-specific changes in anatomical pathology. *Brain Res* 1380, 138–145. doi:10.1016/j.brainres.2010.09.101 [PubMed: 20920490]
- Crane JW, French KR, Buller KM, 2005 Patterns of neuronal activation in the rat brain and spinal cord in response to increasing durations of restraint stress. *Stress* 8, 199–211. doi: 10.1080/10253890500333817 [PubMed: 16236624]
- Cristinzio C, N'Diaye K, Seeck M, Vuilleumier P, Sander D, 2010 Integration of gaze direction and facial expression in patients with unilateral amygdala damage. *Brain* 133, 248–261. doi: 10.1093/brain/awp255 [PubMed: 19828596]
- Critchley HD, Daly EM, Bullmore ET, Williams SC, Van Amelsvoort T, Robertson DM, Rowe A, Phillips M, McAlonan G, Howlin P, Murphy DG, 2000 The functional neuroanatomy of social behaviour: changes in cerebral blood flow when people with autistic disorder process facial expressions. *Brain* 123, 2203–2212. doi: 10.1093/brain/123.11.2203 [PubMed: 11050021]
- Cuthbert BN, Insel TR, 2013 Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC Med* 11, 126. doi: 10.1186/1741-7015-11-126 [PubMed: 23672542]
- Dalton KM, Nacewicz BM, Alexander AL, Davidson RJ, 2007 Gaze-Fixation, Brain Activation, and Amygdala Volume in Unaffected Siblings of Individuals with Autism. *Biol. Psychiatry* 61, 512–520. doi:10.1016/j.biopsych.2006.05.019 [PubMed: 17069771]
- Dalton KM, Nacewicz BM, Johnstone T, Schaefer HS, Gernsbacher MA, Goldsmith HH, Alexander AL, Davidson RJ, 2005 Gaze fixation and the neural circuitry of face processing in autism. *Nat. Neurosci* 8, 519–526. doi:10.1038/nn1421 [PubMed: 15750588]
- Damiano CR, Aloï J, Dunlap K, Burrus CJ, Mosner MG, Kozink RV, McLaurin RE, Mullette-Gillman O. a, Carter RM, Huettel S. a, McClernon FJ, Ashley-Koch A, Dichter GS, 2014 Association between the oxytocin receptor (OXTR) gene and mesolimbic responses to rewards. *Mol. Autism* 5, 7. doi: 10.1186/2040-2392-5-7 [PubMed: 24485285]
- Davis M, Walker DL, Miles L, Grillon C, 2010 Phasic vs sustained fear in rats and humans: role of the extended amygdala in fear vs anxiety. *Neuropsychopharmacology* 35, 105–35. doi:10.1038/npp.2009.109 [PubMed: 19693004]
- Davis M, Whalen PJ, 2001 The amygdala: vigilance and emotion. *Mol. Psychiatry* 6, 13–34. doi: 10.1038/sj.mp.4000812 [PubMed: 11244481]
- De Martino B, Kumaran D, Seymour B, Dolan RJ, 2006 Frames, biases, and rational decision-making in the human brain. *Science* 313, 684–7. doi: 10.1126/science.1128356 [PubMed: 16888142]
- Dodhia S, Hosanagar A, Fitzgerald DA, Labuschagne I, Wood AG, Nathan PJ, Phan KL, 2014 Modulation of Resting-State Amygdala-Frontal Functional Connectivity by Oxytocin in Generalized Social Anxiety Disorder. *Neuropsychopharmacology* 39, 2061–2069. doi:10.1038/npp.2014.53 [PubMed: 24594871]
- Domes G, Heinrichs M, Gläscher J, Büchel C, Braus DF, Herpertz SC, 2007 Oxytocin Attenuates Amygdala Responses to Emotional Faces Regardless of Valence. *Biol. Psychiatry* 62, 1187–1190. doi:10.1016/j.biopsych.2007.03.025 [PubMed: 17617382]
- Dunbar RIM, 2009 The social brain hypothesis and its implications for social evolution. *Ann. Hum. Biol* 36, 562–72. doi: 10.1080/03014460902960289 [PubMed: 19575315]
- Duvarci S, Pare D, 2014 Amygdala microcircuits controlling learned fear. *Neuron* 82, 966–980. doi: 10.1016/j.neuron.2014.04.042 [PubMed: 24908482]
- Dziobek I, Bahnemann M, Convit A, Heekeren HR, 2010 The role of the fusiform-amygdala system in the pathophysiology of autism. *Arch. Gen. Psychiatry* 67, 397–405. doi: 10.1001/archgenpsychiatry.2010.31 [PubMed: 20368515]

- Eapen V, Crnec R, Walter A, 2013 Exploring Links between Genotypes, Phenotypes, and Clinical Predictors of Response to Early Intensive Behavioral Intervention in Autism Spectrum Disorder. *Front. Hum. Neurosci* 7, 567 doi: 10.3389/fnhum.2013.00567 [PubMed: 24062668]
- Eckstein M, Becker B, Scheele D, Scholz C, Preckel K, Schlaepfer TE, Grinevich V, Kendrick KM, Maier W, Hurlmann R, 2015 Oxytocin facilitates the extinction of conditioned fear in humans. *Biol. Psychiatry* 78, 194–202. doi:10.1016/j.biopsych.2014.10.015 [PubMed: 25542304]
- Ehrlich DE, Ryan SJ, Rainnie DG, 2012 Postnatal development of electrophysiological properties of principal neurons in the rat basolateral amygdala. *J. Physiol* 590, 4819–38. doi: 10.1113/jphysiol.2012.237453 [PubMed: 22848043]
- Eilam-Stock T, Wu T, Spagna A, Egan LJ, Fan J, 2016 Neuroanatomical alterations in high-functioning adults with autism spectrum disorder. *Front. Neurosci* 10, 1–10. doi: 10.3389/fnins.2016.00237 [PubMed: 26858586]
- Elliot AJ, Covington MV, 2001 Approach and Avoidance Motivation. *Educ. Psychol. Rev* 13, 73–92. doi:10.1023/A:1009009018235
- Ernst M, Pine DS, Hardin M, 2006 Triadic model of the neurobiology of motivated behavior in adolescence. *Psychological Med* 36, 299–312. doi:10.1017/S0033291705005891
- Fadok JP, Dickerson TM, Palmiter RD, 2009 Dopamine is necessary for cue-dependent fear conditioning. *J Neurosci* 29, 11089–11097. doi: 10.1523/JNEUROSCI.1616-09.2009 [PubMed: 19741115]
- Feldman R, 2015 The adaptive human parental brain: implications for children's social development. *Trends Neurosci* 38, 387–99. doi: 10.1016/j.tins.2015.04.004 [PubMed: 25956962]
- Felix-Ortiz AC, Burgos-Robles A, Bhagat ND, Leppla CA, Tye KM, 2016 Bidirectional modulation of anxiety-related and social behaviors by amygdala projections to the medial prefrontal cortex. *Neuroscience* 321, 197–209. doi: 10.1016/j.neuroscience.2015.07.041 [PubMed: 26204817]
- Ferguson JN, Aldag JM, Insel TR, Young LJ, 2001 Oxytocin in the Medial Amygdala is Essential for Social Recognition in the Mouse. *J. Neurosci* 21, 8278–8285. doi:10.1523/JNEUROSCI.1158-01.2001 [PubMed: 11588199]
- Ferri J, Schmidt J, Hajcak G, Canli T, 2016 Emotion regulation and amygdala-precuneus connectivity: Focusing on attentional deployment. *Cogn. Affect. Behav. Neurosci* 1–12. doi: 10.3758/s13415-016-0447-y [PubMed: 26785840]
- Fitzgerald DA, Angstadt M, Jelsone LM, Nathan PJ, Phan KL, 2006 Beyond threat: Amygdala reactivity across multiple expressions of facial affect. *Neuroimage* 30, 1441–1448. doi: 10.1016/j.neuroimage.2005.11.003 [PubMed: 16368249]
- Fombonne E, 2009 Epidemiology of pervasive developmental disorders. *Pediatr. Res* 65, 591–8. doi: 10.1203/PDR.0b013e31819e7203 [PubMed: 19218885]
- Frith CD, 2007 The social brain? *Philos. Trans. R. Soc. Lond. B. Biol. Sci* 362, 671–8. doi: 10.1098/rstb.2006.2003 [PubMed: 17255010]
- Fullana M, Harrison B, Soriano-Mas C, Vervliet B, Cardoner N, Àvila-Parcet A, Radua J, 2015 Neural signatures of human fear conditioning: an updated and extended meta-analysis of fMRI studies. *Mol. Psychiatry* 21, 500–508. doi: 10.1038/mp.2015.88 [PubMed: 26122585]
- Gallup GG, 1965 Aggression in rats as a function of frustrative nonreward in a straight alley 3, 99–100.
- Gamer M, Buchel C, 2009 Amygdala activation predicts gaze toward fearful eyes. *J Neurosci* 29, 9123–9126. doi:10.1523/jneurosci.1883-09.2009 [PubMed: 19605649]
- Giarelli E, Wiggins LD, Rice CE, Levy SE, Kirby RS, Pinto-Martin J, Mandell D, 2010 Sex differences in the evaluation and diagnosis of autism spectrum disorders among children. *Disabil. Health J* 3, 107–116. doi:10.1016/j.dhjo.2009.07.001 [PubMed: 21122776]
- Gordon I, 2014 Can Oxytocin Improve Core Brain and Behavioral Features of Autism Spectrum Disorders in Children?, in: Decety J, Christen Y (Eds.), *New Frontiers in Social Neuroscience*. Springer International Publishing, Cham, pp. 189–199. doi: 10.1007/978-3-319-02904-7_11
- Gorka SM, Fitzgerald DA, Labuschagne I, Hosanagar A, Wood AG, Nathan PJ, Phan KL, 2015 Oxytocin Modulation of Amygdala Functional Connectivity to Fearful Faces in Generalized Social Anxiety Disorder. *Neuropsychopharmacology* 40, 278–286. doi:10.1038/npp.2014.168 [PubMed: 24998619]

- Gottfried JA, O'Doherty J, Dolan RJ, 2003 Value in Human Amygdala and Orbitofrontal Cortex. *Science* (80-.). 301, 1104–1108. doi: 10.1126/science.1087919
- Grillon C, Krinsky M, Charney DR, Vytal K, Ernst M, Cornwell B, 2013 Oxytocin increases anxiety to unpredictable threat. *Mol. Psychiatry* 18, 958–960. doi:10.1038/mp.2012.156 [PubMed: 23147382]
- Guastella AJ, Einfeld SL, Gray KM, Rinehart NJ, Tonge BJ, Lambert TJ, Hickie IB, 2010 Intranasal Oxytocin Improves Emotion Recognition for Youth with Autism Spectrum Disorders. *Biol. Psychiatry* 67, 692–694. doi:10.1016/j.biopsych.2009.09.020 [PubMed: 19897177]
- Haar S, Berman S, Behrmann M, Dinstei I, 2016 Anatomical Abnormalities in Autism? *Cereb. Cortex* 26, 1440–1452. doi:10.1093/cercor/bhu242 [PubMed: 25316335]
- Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, Torigoe T, Miller J, Fedele A, Collins J, Smith K, Lotspeich L, Croen L. a., Ozonoff S, Lajonchere C, Grether JK, Risch N, 2011 Genetic Heritability and Shared Environmental Factors Among Twin Pairs With Autism. *Arch. Gen. Psychiatry* 68, 1095–1102. doi:10.1001/archgenpsychiatry.2011.76 [PubMed: 21727249]
- Hampton AN, Adolphs R, Tyszka MJ, O'Doherty JP, 2007 Contributions of the Amygdala to Reward Expectancy and Choice Signals in Human Prefrontal Cortex. *Neuron* 55, 545–555. doi: 10.1016/j.neuron.2007.07.022 [PubMed: 17698008]
- Happé F, Frith U, 2014 Annual research review: Towards a developmental neuroscience of atypical social cognition. *J. Child Psychol. Psychiatry*. 55, 553–7. doi: 10.1111/jcpp.12162 [PubMed: 24963529]
- Happé F, Frith U, 2006 The weak coherence account: Detail-focused cognitive style in autism spectrum disorders. *J. Autism Dev. Disord* 36, 5–25. doi: 10.1007/s10803-005-0039-0 [PubMed: 16450045]
- Harbour VL, Weigl Y, Robinson B, Amir S, 2014 Phase differences in expression of circadian clock genes in the central nucleus of the amygdala, dentate gyrus, and suprachiasmatic nucleus in the rat. *PLoS One* 9, 1–9. doi: 10.1371/journal.pone.0103309
- Harmon-Jones E, Harmon-Jones C, Price TF, 2013 What is Approach Motivation? *Emot. Rev* 5, 291–295. doi: 10.1177/1754073913477509
- Hazlett HC, Gu H, Munsell BC, Kim SH, Styner M, Wolff JJ, Elison JT, Swanson MR, Zhu H, Botteron KN, Collins DL, Constantino JN, Dager SR, Estes AM, Evans AC, Fonov VS, Gerig G, Kostopoulos P, Mckinstry RC, Pandey J, Paterson S, Pruett JR, Jr, Schultz RT, Shaw DW, Zwaigenbaum L, Piven J, 2017 Early brain development in infants at high risk for autism spectrum disorder. *Nat. Publ. Gr* 542, 348–351. doi: 10.1038/nature21369
- Hernandez-Lallement J, van Wingerden M, Schäble S, Kalenscher T, 2016 Basolateral amygdala lesions abolish mutual reward preferences in rats. *Neurobiol. Learn. Mem* 127, 1–9. doi: 10.1016/j.nlm.2015.11.004 [PubMed: 26596916]
- Hillis AE, 2014 Inability to empathize: brain lesions that disrupt sharing and understanding another's emotions. *Brain* 137, 981–97. doi:10.1093/brain/awt317 [PubMed: 24293265]
- Hranilovic D, Bujas-Petkovic Z, Vragovic R, Vuk T, Hock K, Jernej B, 2007 Hyperserotonemia in adults with autistic disorder. *J. Autism Dev. Disord* 37, 1934–1940. doi:10.1007/s10803-006-0324-6 [PubMed: 17165147]
- Huber D, Veinante P, Stoop R, 2005 Vasopressin and oxytocin excite distinct neuronal populations in the central amygdala. *Science* 308, 245–8. doi: 10.1126/science.1105636 [PubMed: 15821089]
- Hurlemann R, Wagner M, Hawellek B, Reich H, Pieperhoff P, Amunts K, Oros-Peusquens AM, Shah NJ, Maier W, Dolan RJ, 2007 Amygdala control of emotion-induced forgetting and remembering: Evidence from Urbach-Wiethe disease. *Neuropsychologia* 45, 877–884. doi: 10.1016/j.neuropsychologia.2006.08.027 [PubMed: 17027866]
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Sanislow C, Wang P, 2010 Research Domain Criteria (RDoC): Toward a new classification framework for research on mental disorders. *Am. J. Psychiatry* 167, 748–751. doi: 10.1176/appi.ajp.2010.09091379 [PubMed: 20595427]
- Insel TR, Cuthbert BN, 2009 Endophenotypes: Bridging Genomic Complexity and Disorder Heterogeneity. *Biol. Psychiatry* 66, 988–989. doi: 10.1016/j.biopsych.2009.10.008 [PubMed: 19900610]

- Jackson ME, Moghaddam B, 2001 Amygdala regulation of nucleus accumbens dopamine output is governed by the prefrontal cortex. *J. Neurosci* 21, 676–681. doi:10.1523/JNEUROSCI.1116-01.2001 [PubMed: 11160446]
- Jacobs RHAH, Renken R, Aleman A, Cornelissen FW, 2012 The amygdala, top-down effects, and selective attention to features. *Neurosci. Biobehav. Rev* 36, 2069–2084. doi:10.1016/j.neubiorev.2012.05.011 [PubMed: 22728112]
- Jäger-Roman E, Deichl A, Jakob S, Hartmann AM, Koch S, Rating D, Steldinger R, Nau H, Helge H, 1986 Fetal growth, major malformations, and minor anomalies in infants born to women receiving valproic acid. *J. Pediatr* 108, 997–1004. doi: 10.1016/S0022-3476(86)80949-0 [PubMed: 3086531]
- Janak PH, Tye KM, 2015 From circuits to behaviour in the amygdala. *Nature* 517, 284–292. doi: 10.1038/nature14188 [PubMed: 25592533]
- Jeffrey M, Lang M, Gane J, Wu C, Burnham WM, Zhang L, 2013 A reliable method for intracranial electrode implantation and chronic electrical stimulation in the mouse brain. *BMC Neurosci* 14, 82. doi: 10.1186/1471-2202-14-82 [PubMed: 23914984]
- Jennings JH, Sparta DR, Stamatakis AM, Ung RL, Pleil KE, Kash TL, Stuber GD, 2013 Distinct extended amygdala circuits for divergent motivational states. *Nature* 496, 224–228. doi: 10.1038/nature12041 [PubMed: 23515155]
- Jiang X, Xing G, Yang C, Verma A, Zhang L, Li H, 2009 Stress impairs 5-HT_{2A} receptor-mediated serotonergic facilitation of GABA release in juvenile rat basolateral amygdala. *Neuropsychopharmacology* 34, 410–423. doi:10.1038/npp.2008.71 [PubMed: 18536707]
- Johnson ZV, Walum H, Jamal YA, Xiao Y, Keebaugh AC, Inoue K, Young LJ, 2016 Central oxytocin receptors mediate mating-induced partner preferences and enhance correlated activation across forebrain nuclei in male prairie voles. *Horm. Behav* 79, 8–17. doi: 10.1016/j.yhbeh.2015.11.011 [PubMed: 26643557]
- Jones W, Klin A, 2013 Attention to eyes is present but in decline in 2-6-month-old infants later diagnosed with autism. *Nature* 504, 427–31. doi: 10.1038/nature12715 [PubMed: 24196715]
- Kalin NH, Shelton SE, Davidson RJ, 2004 The Role of the Central Nucleus of the Amygdala in Mediating Fear and Anxiety in the Primate. *J. Neurosci* 24, 5506–5515. doi: 10.1523/JNEUROSCI.0292-04.2004 [PubMed: 15201323]
- Kanner L, 1943 Autistic disturbances of affective contact. *Nerv. Child*
- Kanske P, Heissler J, Schönfelder S, Bongers A, Wessa M, 2011 How to regulate emotion? Neural networks for reappraisal and distraction. *Cereb. Cortex* 21, 1379–1388. doi: 10.1093/cercor/bhq216 [PubMed: 21041200]
- Keifer OP, Hurt RC, Ressler KJ, Marvar PJ, 2015 The Physiology of Fear: Reconceptualizing the Role of the Central Amygdala in Fear Learning. *Physiology (Bethesda)*. 30, 389–401. doi: 10.1152/physiol.00058.2014 [PubMed: 26328883]
- Kemper TL, Bauman ML, 1993 The contribution of neuropathologic studies to the understanding of autism. *Neurol. Clin* 11, 175–87. [PubMed: 8441369]
- Kennedy DP, Gläscher J, Tyszka JM, Adolphs R, 2010 Personal Space Regulation by the Human Amygdala. *Nat. Neurosci* 12, 1226–1227. doi: 10.1038/nn.2381
- Kim H, Somerville LH, Johnstone T, Alexander AL, Whalen PJ, 2003 Inverse amygdala and medial prefrontal cortex responses to surprised faces. *Neuroreport* 14, 2317–2322. doi: 10.1097/01.wnr.0000101520.44335.20 [PubMed: 14663183]
- Kim H, Somerville LH, Johnstone T, Polis S, Alexander AL, Shin LM, Whalen PJ, 2004 Contextual modulation of amygdala responsivity to surprised faces. *J. Cogn. Neurosci* 16, 1730–1745. doi: 10.1162/0898929042947865 [PubMed: 15701225]
- Kim J, Pignatelli M, Xu S, Itohara S, Tonegawa S, 2016 Antagonistic negative and positive neurons of the basolateral amygdala. *Nat. Neurosci* 19, 1636–1646. doi:10.1038/nn.4414 [PubMed: 27749826]
- Kim J, Zhang X, Muralidhar S, LeBlanc SA, Tonegawa S, 2017 Basolateral to Central Amygdala Neural Circuits for Appetitive Behaviors. *Neuron* 93, 1464–1479.e5. doi: 10.1016/j.neuron.2017.02.034 [PubMed: 28334609]

- Kim KC, Kim P, Go HS, Choi CS, Yang S, Il, Cheong JH, Shin CY, Ko KH, 2011 The critical period of valproate exposure to induce autistic symptoms in Sprague-Dawley rats. *Toxicol. Lett* 201, 137–142. doi:10.1016/j.toxlet.2010.12.018 [PubMed: 21195144]
- Kim MJ, Loucks RA, Palmer AL, Brown AC, Solomon KM, Marchante AN, Whalen PJ, 2011 The structural and functional connectivity of the amygdala: from normal emotion to pathological anxiety. *Behav. Brain Res* 223, 403–10. doi:10.1016/j.bbr.2011.04.025 [PubMed: 21536077]
- King EC, Pattwell SS, Glatt CE, Lee FS, 2014 Sensitive periods in fear learning and memory. *Stress* 17, 13–21. doi: 10.3109/10253890.2013.796355 [PubMed: 23611461]
- Klavir O, Prigge M, Sarel A, Paz R, Yizhar O, 2017 Manipulating fear associations via optogenetic modulation of amygdala inputs to prefrontal cortex. *Nat. Neurosci* 20, 836–844. doi:10.1038/nn.4523 [PubMed: 28288126]
- Kleinmans NM, Richards T, Weaver K, Johnson LC, Greenson J, Dawson G, Aylward E, 2010 Association between amygdala response to emotional faces and social anxiety in autism spectrum disorders. *Neuropsychologia* 48, 3665–70. doi: 10.1016/j.neuropsychologia.2010.07.022 [PubMed: 20655320]
- Kliemann D, Dziobek I, Hatri A, Baudewig J, Heekeren HR, 2012 The role of the amygdala in atypical gaze on emotional faces in autism spectrum disorders. *J. Neurosci* 32, 9469–76. doi: 10.1523/JNEUROSCI.5294-11.2012 [PubMed: 22787032]
- Klin A, 2000 Attributing social meaning to ambiguous visual stimuli in higher-functioning autism and Asperger syndrome: The Social Attribution Task. *J. Child Psychol. Psychiatry*. 41, 831–46. doi: 10.1111/1469-7610.00671 [PubMed: 11079426]
- Klüver H, Bucy PC, 1939 PRELIMINARY ANALYSIS OF FUNCTIONS OF THE TEMPORAL LOBES IN MONKEYS. *Arch. Neurol. Psychiatry* 42, 979–1000.
- Klüver H, Bucy PC, 1937 “Psychic blindness” and other symptoms following bilateral temporal lobectomy in Rhesus monkeys. *Am. J. Physiol* 119, 352–353.
- Knobloch HS, Charlet A, Hoffmann L, Eliava M, Khrulev S, Cetin A, Osten P, Schwarz M, Seeburg P, Stoop R, Grinevich V, 2012 Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73, 553–566. doi:10.1016/j.neuron.2011.11.030 [PubMed: 22325206]
- Koch SB, van Zuiden M, Nawijn L, Frijling JL, Veltman DJ, Olff M, 2016 Intranasal Oxytocin Administration Dampens Amygdala Reactivity towards Emotional Faces in Male and Female PTSD Patients. *Neuropsychopharmacology* 41, 1495–504. doi:10.1038/npp.2015.299 [PubMed: 26404844]
- Kosmicki JA, Samocha KE, Howrigan DP, Sanders SJ, Slowikowski K, Lek M, Karczewski KJ, Cutler DJ, Devlin B, Roeder K, Buxbaum JD, Neale BM, MacArthur DG, Wall DP, Robinson EB, Daly MJ, 2017 Refining the role of de novo protein-truncating variants in neurodevelopmental disorders by using population reference samples. *Nat. Genet* 49, 504–510. doi: 10.1038/ng.3789 [PubMed: 28191890]
- Labuschagne I, Phan KL, Wood A, Angstadt M, Chua P, Heinrichs M, Stout JC, Nathan PJ, 2010 Oxytocin Attenuates Amygdala Reactivity to Fear in Generalized Social Anxiety Disorder. *Neuropsychopharmacology* 35, 2403–2413. doi:10.1038/npp.2010.123 [PubMed: 20720535]
- LeDoux J, Cicchetti P, Xagoraris A, Romanski L, 1990 The lateral amygdaloid nucleus: sensory interface of the amygdala in fear conditioning. *J. Neurosci* 10(4), 1062–1069. doi:2329367 [PubMed: 2329367]
- LeDoux JE, 2008 Amygdala. *Scholarpedia* 3, 2698.
- LeDoux JE, 1996 *The Emotional Brain: The mysterious underpinnings of emotional life*. Simon and Schuster, New York.
- LeDoux JE, 1992 Brain mechanisms of emotion and emotional learning. *Curr. Opin. Neurobiol* 2, 191–197. doi: 10.1016/0959-4388(92)90011-9 [PubMed: 1638153]
- LeDoux JE, Iwata J, Cicchetti P, Reis DJ, 1988 Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J. Neurosci* 8, 2517–2529. [PubMed: 2854842]
- Lenroot RK, Giedd JN, 2006 Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neurosci. Biobehav. Rev* 30, 718–729. doi: 10.1016/j.neubiorev.2006.06.001 [PubMed: 16887188]

- Li C, Dabrowska J, Hazra R, Rainnie DG, 2011 Synergistic activation of dopamine D1 and TrkB receptors mediate gain control of synaptic plasticity in the basolateral amygdala. *PLoS One* 6, e26065. doi: 10.1371/journal.pone.0026065
- Lin H-C, Gean P-W, Wang C-C, Chan Y-H, Chen PS, 2013 The amygdala excitatory/inhibitory balance in a valproate-induced rat autism model. *PLoS One*. doi:10.1371/journal.pone.0055248
- Lissek S, Kaczurkin AN, Rabin S, Geraci M, Pine DS, Grillon C, 2014 Generalized anxiety disorder is associated with overgeneralization of classically conditioned fear. *Biol. Psychiatry* 75, 909–915. doi:10.1016/j.biopsych.2013.07.025 [PubMed: 24001473]
- LoParo D, Waldman ID, 2015 The oxytocin receptor gene (OXTR) is associated with autism spectrum disorder: a meta-analysis. *Mol. Psychiatry* 20, 640–646. doi:10.1038/mp.2014.77 [PubMed: 25092245]
- Loth E, Spooren W, Ham LM, Isaac MB, Auriche-Benichou C, Banaschewski T, Baron-Cohen S, Broich K, Bölte S, Bourgeron T, Charman T, Collier D, de Andres-Trelles F, Durston S, Ecker C, Elferink A, Haberkamp M, Hemmings R, Johnson MH, Jones EJH, Khwaja OS, Lenton S, Mason L, Mantua V, Meyer-Lindenberg A, Lombardo MV, O'Dwyer L, Okamoto K, Pandina GJ, Pani L, Persico AM, Simonoff E, Tauscher-Wisniewski S, Llinares-Garcia J, Vamvakas S, Williams S, Buitelaar JK, Murphy DGM, 2016 Identification and validation of biomarkers for autism spectrum disorders. *Nat. Rev. Drug Discov* 15, 70–3. doi:10.1038/nrd.2015.7 [PubMed: 26718285]
- Mabunga DFN, Gonzales ELT, Kim J-W, Kim KC, Shin CY, 2015 Exploring the Validity of Valproic Acid Animal Model of Autism. *Exp. Neurobiol* 24, 285–300. doi: 10.5607/en.2015.24.4.285 [PubMed: 26713077]
- MacFabe DF, Cain DP, Rodriguez-Capote K, Franklin AE, Hoffman JE, Boon F, Taylor AR, Kavaliers M, Ossenkopp KP, 2007 Neurobiological effects of intraventricular propionic acid in rats: Possible role of short chain fatty acids on the pathogenesis and characteristics of autism spectrum disorders. *Behav. Brain Res* 176, 149–169. doi:10.1016/j.bbr.2006.07.025 [PubMed: 16950524]
- Machado CJ, Bachevalier J, 2007 The effects of selective amygdala, orbital frontal cortex or hippocampal formation lesions on reward assessment in nonhuman primates. *Eur. J. Neurosci* 25, 2885–2904. doi: 10.1111/j.1460-9568.2007.05525.x [PubMed: 17561849]
- Machado CJ, Kazama AM, Bachevalier J, 2009 Impact of amygdala, orbital frontal, or hippocampal lesions on threat avoidance and emotional reactivity in nonhuman primates. *Emotion* 9, 147–163. doi: 10.1037/a0014539 [PubMed: 19348528]
- MacLean PD, 1955 The limbic system (“visceral brain”) and emotional behavior. *AMA. Arch. Neurol. Psychiatry* 73, 130–4. [PubMed: 13227663]
- MacLean PD, 1952 Some psychiatric implications of physiological studies on frontotemporal portion of limbic system (visceral brain). *Electroencephalogr. Clin. Neurophysiol* 4, 407–418. doi: 10.1016/0013-4694(52)90073-4 [PubMed: 12998590]
- Madarasz TJ, Diaz-Mataix L, Akhand O, Ycu EA, LeDoux JE, Johansen JP, 2016 Evaluation of ambiguous associations in the amygdala by learning the structure of the environment. *Nat. Neurosci* 19, 965–972. doi: 10.1038/nn.4308 [PubMed: 27214568]
- Mandell DS, Novak MM, Zubrisky CD, 2005 Factors associated with age of diagnosis among children with autism spectrum disorders. *Pediatrics* 116, 1480–6. doi:10.1542/peds.2005-0185 [PubMed: 16322174]
- Mandy W, Chilvers R, Chowdhury U, Salter G, Seigal A, Skuse D, 2012 Sex differences in autism spectrum disorder: Evidence from a large sample of children and adolescents. *J. Autism Dev. Disord* 42, 1304–1313. doi: 10.1007/s10803-011-1356-0 [PubMed: 21947663]
- Markram H, Rinaldi T, Markram K, 2007 The intense world syndrome--an alternative hypothesis for autism. *Front. Neurosci* 1, 77–96. doi: 10.3389/neuro.01.1.1.006.2007 [PubMed: 18982120]
- Markram K, Rinaldi T, La Mendola D, Sandi C, Markram H, 2008 Abnormal fear conditioning and amygdala processing in an animal model of autism. *Neuropsychopharmacology* 33, 901–912. doi:DOI 10.1038/sj.npp.1301453 [PubMed: 17507914]
- Mazurek MO, Handen BL, Wodka EL, Nowinski L, Butter E, Engelhardt CR, 2014 Age at first autism spectrum disorder diagnosis: the role of birth cohort, demographic factors, and clinical features. *J. Dev. Behav. Pediatr* 35, 561–569. doi: 10.1097/DBP.0000000000000097 [PubMed: 25211371]

- Meletti S, Cantalupo G, Benuzzi F, Mai R, Tassi L, Gasparini E, Tassinari CA, Nichelli P, 2012 Fear and happiness in the eyes: An intra-cerebral event-related potential study from the human amygdala. *Neuropsychologia* 50, 44–54. doi:10.1016/j.neuropsychologia.2011.10.020 [PubMed: 22056505]
- Méndez-Bértolo C, Moratti S, Toledano R, Lopez-Sosa F, Martínez-Alvarez R, Mah YH, Vuilleumier P, Gil-Nagel A, Strange BA, 2016 A fast pathway for fear in human amygdala. *Nat. Neurosci* 19, 1041–1049. doi: 10.1038/nn.4324 [PubMed: 27294508]
- Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M, 2011 Oxytocin and vasopressin in the human brain: social neuropeptides for translational medicine. *Nat. Rev. Neurosci* 12, 524–538. doi: 10.1038/nrn3044 [PubMed: 21852800]
- Mintz M, 2017 Evolution in the Understanding of Autism Spectrum Disorder: Historical Perspective. *Indian J. Pediatr* 84, 44–52. doi:10.1007/s12098-016-2080-8 [PubMed: 27053182]
- Montague PR, Berns GS, 2002 Neural economics and the biological substrates of valuation. *Neuron* 36, 265–284. doi:10.1016/S0896-6273(02)00974-1 [PubMed: 12383781]
- Morgan JT, Barger N, Amaral DG, Schumann CM, 2014 Stereological study of amygdala glial populations in adolescents and adults with autism spectrum disorder. *PLoS One* 9, 29–30. doi: 10.1371/journal.pone.0110356
- Mormann F, Niediek J, Tudusciuc O, Quesada CM, Coenen VA, Elger CE, Adolphs R, 2015 Neurons in the human amygdala encode face identity, but not gaze direction. *Nat Neurosci* 18, 1568–1570. [PubMed: 26479589]
- Morris JS, Ohman A, Dolan RJ, 1999 A subcortical pathway to the right amygdala mediating “unseen” fear. *Proc. Natl. Acad. Sci* 96, 1680–1685. doi: 10.1073/pnas.96.4.1680 [PubMed: 9990084]
- Mosconi MW, Cody-Hazlett H, Poe MD, Gerig G, Gimpel-Smith R, Piven J, 2009 Longitudinal Study of Amygdala Volume and Joint Attention in 2- to 4-Year-Old Children With Autism. *Arch. Gen. Psychiatry* 66, 509. doi:10.1001/archgenpsychiatry.2009.19 [PubMed: 19414710]
- Moy SS, Nadler JJ, Perez A, Barbaro RP, Johns JM, Magnuson TR, Piven J, Crawley JN, 2004 Sociability and preference for social novelty in five inbred strains: an approach to assess autistic-like behavior in mice. *Genes Brain Behav* 3, 287–302. doi: 10.1111/j.1601-1848.2004.00076.xGBB076 [pii] [PubMed: 15344922]
- Muller JF, Mascagni F, McDonald AJ, 2009 Dopaminergic innervation of pyramidal cells in the rat basolateral amygdala. *Brain Struct. Funct* 213, 275–288. doi:10.1007/s00429-008-0196-y [PubMed: 18839210]
- Müller R-A, Amaral DG, 2017 Editorial: Time to give up on Autism Spectrum Disorder? *Autism Res* 10, 10–14. doi: 10.1002/aur.1746 [PubMed: 28130875]
- Murphy CM, Deeley Q, Daly EM, Ecker C, O’Brien FM, Hallahan B, Loth E, Toal F, Reed S, Hales S, Robertson DM, Craig MC, Mullins D, Barker GJ, Lavender T, Johnston P, Murphy KC, Murphy DG, 2012 Anatomy and aging of the amygdala and hippocampus in autism spectrum disorder: An in vivo magnetic resonance imaging study of Asperger syndrome. *Autism Res* 5, 3–12. doi: 10.1002/aur.227 [PubMed: 21948742]
- Nacewicz BM, Dalton KM, Johnstone T, Long MT, McAuliff EM, Oakes TR, Alexander AL, Davidson RJ, 2006 Amygdala volume and nonverbal social impairment in adolescent and adult males with autism. *Arch. Gen. Psychiatry* 63, 1417–1428. doi:10.1001/archpsyc.63.12.1417 [PubMed: 17146016]
- Namburi P, Beyeler A, Yorozu S, Calhoon GG, Halbert S. a., Wichmann R, Holden SS, Mertens KL, Anahtar M, Felix-Ortiz AC, Wickersham IR, Gray JM, Tye KM, 2015 A circuit mechanism for differentiating positive and negative associations. *Nature* 520, 675–678. doi: 10.1038/nature14366 [PubMed: 25925480]
- Narita N, Kato M, Tazoe M, Miyazaki K, Narita M, Okado N, 2002 Increased monoamine concentration in the brain and blood of fetal thalidomide- and valproic acid-exposed rat: putative animal models for autism. *Pediatr. Res* 52, 576–579. [PubMed: 12357053]
- Neumann ID, Slattery DA, 2016 Oxytocin in General Anxiety and Social Fear: A Translational Approach. *Biol. Psychiatry* 79, 213–221. doi: 10.1016/j.biopsych.2015.06.004 [PubMed: 26208744]

- Nordahl CW, Scholz R, Yang X, Buonocore MH, Simon T, Rogers S, Amaral DG, 2012 Increased rate of amygdala growth in children aged 2 to 4 years with autism spectrum disorders: a longitudinal study. *Arch. Gen. Psychiatry* 69, 53–61. doi:10.1001/archgenpsychiatry.2011.145 [PubMed: 22213789]
- NVS Working Group, 2011 NVS Workshop Proceedings [WWW Document]. URL <https://www.nimh.nih.gov/research-priorities/rdoc/negative-valence-systems-workshop-proceedings.shtml> (accessed 1.1.16).
- Ochsner KN, Silvers JA, Buhle JT, 2012 Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Ann. N. Y. Acad. Sci* 1251, E1–24. doi: 10.1111/j.1749-6632.2012.06751.x [PubMed: 23025352]
- Oguchi-Katayama A, Monma A, Sekino Y, Moriguchi T, Sato K, 2013 Comparative gene expression analysis of the amygdala in autistic rat models produced by pre- and post-natal exposures to valproic acid. *J. Toxicol. Sci* 38, 391–402. doi: 10.2131/jts.38.391 [PubMed: 23665938]
- Öhman A, Carlsson K, Lundqvist D, Ingvar M, 2007 On the unconscious subcortical origin of human fear. *Physiol. Behav* 92, 180–185. doi: 10.1016/j.physbeh.2007.05.057 [PubMed: 17599366]
- Ornøy A, Weinstein-Fudim L, Ergaz Z, 2015 Prenatal factors associated with autism spectrum disorder (ASD). *Reprod. Toxicol* 56, 155–169. doi:10.1016/j.reprotox.2015.05.007 [PubMed: 26021712]
- Paton JJ, Belova MA, Morrison SE, Salzman CD, 2006 The primate amygdala represents the positive and negative value of visual stimuli during learning. *Nature* 439, 865–870. doi:10.1038/nature04490 [PubMed: 16482160]
- Paul LK, Corsello C, Tranel D, Adolphs R, 2010 Does bilateral damage to the human amygdala produce autistic symptoms? *J. Neurodev. Disord* 2, 165–173. doi: 10.1007/s11689-010-9056-1 [PubMed: 20700516]
- Payne C, Machado CJ, Bliwise NG, Bachevalier J, 2010 Maturation of the hippocampal formation and amygdala in *Macaca mulatta*: A volumetric magnetic resonance imaging study. *Hippocampus* 20, 922–935. doi: 10.1002/hipo.20688 [PubMed: 19739247]
- Paz R, Pare D, 2013 Physiological basis for emotional modulation of memory circuits by the amygdala. *Curr. Opin. Neurobiol* 23, 381–386. doi: 10.1016/j.conb.2013.01.008 [PubMed: 23394774]
- Pelphrey KA, Sasson NJ, Reznick JS, Paul G, Goldman BD, Piven J, 2002 Visual Scanning of Faces in Autism. *J. Autism Dev. Disord* 32, 249–261. doi:10.1023/A:1016374617369 [PubMed: 12199131]
- Perucca E, 2002 Pharmacological and therapeutic properties of valproate: a summary after 35 years of clinical experience. *CNS Drugs* 16, 695–714. doi: 10.2165/00023210-200216100-00004 [PubMed: 12269862]
- Pessoa L, 2014 Understanding brain networks and brain organization. *Phys. Life Rev* 11, 400–35. doi: 10.1016/j.plrev.2014.03.005 [PubMed: 24819881]
- Pessoa L, 2011 Reprint of: Emotion and cognition and the amygdala: from “what is it?” to “what’s to be done?”. *Neuropsychologia* 49, 681–94. doi:10.1016/j.neuropsychologia.2011.02.030 [PubMed: 21414465]
- Pessoa L, 2008 On the relationship between emotion and cognition. *Nat. Rev. Neurosci* 9, 148–58. doi: 10.1038/nrn2317 [PubMed: 18209732]
- Peters S, Slattery DA, Uschold-Schmidt N, Reber SO, Neumann ID, 2014 Dose-dependent effects of chronic central infusion of oxytocin on anxiety, oxytocin receptor binding and stress-related parameters in mice. *Psychoneuroendocrinology* 42, 225–236. doi: 10.1016/j.psyneuen.2014.01.021 [PubMed: 24636519]
- Petrovic P, Kalisch R, Singer T, Dolan RJ, 2008 Oxytocin Attenuates Affective Evaluations of Conditioned Faces and Amygdala Activity. *J. Neurosci* 28, 6607–6615. doi: 10.1523/JNEUROSCI.4572-07.2008 [PubMed: 18579733]
- Petrovich GD, Canteras NS, Swanson LW, 2001 Combinatorial amygdalar inputs to hippocampal domains and hypothalamic behavior systems. *Brain Res. Rev* 38, 247–289. doi: 10.1016/S0165-0173(01)00080-7 [PubMed: 11750934]
- Pockros-Burgess L. a, Pentkowski NS, Der-Ghazarian T, Neisewander JL, 2014 Effects of the 5-HT_{2C} receptor agonist CP809101 in the amygdala on reinstatement of cocaine-seeking behavior and

anxiety-like behavior. *Int. J. Neuropsychopharmacol* 1–12. doi: 10.1017/S1461145714000856 [PubMed: 23953038]

- Poelmans G, Franke B, Pauls DL, Glennon JC, Buitelaar JK, 2013 AKAPs integrate genetic findings for autism spectrum disorders. *Transl. Psychiatry* 3, e270. doi: 10.1038/tp.2013.48 [PubMed: 23756379]
- Purgert RJ, Wheeler DS, McDannald M. a., Holland PC, 2012 Role of Amygdala Central Nucleus in Aversive Learning Produced by Shock or by Unexpected Omission of Food. *J. Neurosci* 32, 2461–2472. doi: 10.1523/JNEUROSCI.5090-11.2012 [PubMed: 22396420]
- Quinn R, 2005 Comparing rat's to human's age: how old is my rat in people years? *Nutrition* 21, 775–7. doi: 10.1016/j.nut.2005.04.002 [PubMed: 15925305]
- Radke S, Volman I, Mehta P, van Son V, Enter D, Sanfey a., Toni I, de Bruijn E.R. a., Roelofs K, 2015 Testosterone biases the amygdala toward social threat approach. *Sci. Adv* 1, e1400074–e1400074. doi: 10.1126/sciadv.1400074 [PubMed: 26601187]
- Rai R, Ranjan M, Pradhan BB, Chatterjee S, 2012 Atypical Regulation of Virulence-Associated Functions by a Diffusible Signal Factor in *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant-Microbe Interact* 25, 789–801. doi:10.1094/MPMI-11-11-0285-R [PubMed: 22352717]
- Rainnie D, 1999 Serotonergic modulation of neurotransmission in the rat basolateral amygdala. *J. Neurophysiol* 69–85. [PubMed: 10400936]
- Rainnie DG, Asprodini EK, Shinnick-Gallagher P, 1991a Excitatory transmission in the basolateral amygdala. *J. Neurophysiol* 66, 986–98. [PubMed: 1684383]
- Rainnie DG, Asprodini EK, Shinnick-Gallagher P, 1991b Inhibitory transmission in the basolateral amygdala. *J. Neurophysiol* 66, 999–1009. [PubMed: 1684384]
- Raper J, Stephens SBZ, Henry A, Villarreal T, Bachevalier J, Wallen K, Sanchez MM, 2014 Neonatal Amygdala Lesions Lead to Increased Activity of Brain CRF Systems and Hypothalamic-Pituitary-Adrenal Axis of Juvenile Rhesus Monkeys. *J. Neurosci* 34, 11452–11460. doi: 10.1523/JNEUROSCI.0269-14.2014 [PubMed: 25143624]
- Rasalam AD, Hailey H, Williams JHG, Moore SJ, Turnpenny PD, Lloyd DJ, Dean JCS, 2005 Characteristics of fetal anticonvulsant syndrome associated autistic disorder. *Dev. Med. Child Neurol* 47, 551–5. doi:doi:10.1038/sj.eye.6700371 [PubMed: 16108456]
- Rausch A, Zhang W, Haak KV, Mennes M, Hermans EJ, van Oort E, van Wingen G, Beckmann CF, Buitelaar JK, Groen WB, 2016 Altered functional connectivity of the amygdaloid input nuclei in adolescents and young adults with autism spectrum disorder: a resting state fMRI study. *Mol. Autism* 7, 13. doi:10.1186/s13229-015-0060-x [PubMed: 26823966]
- Reppert SM, Weaver DR, 2002 Coordination of circadian timing in mammals. *Nature* 418, 935–41. doi:10.1038/nature00965 [PubMed: 12198538]
- Rice K, Viscomi B, Riggins T, Redcay E, 2014 Amygdala volume linked to individual differences in mental state inference in early childhood and adulthood. *Dev. Cogn. Neurosci* 8, 153–163. doi: 10.1016/j.dcn.2013.09.003 [PubMed: 24139023]
- Rilling JK, Young LJ, 2014 The biology of mammalian parenting and its effect on offspring social development. *Science* (80-.). 345, 771 LP–776.
- Robinson EB, Lichtenstein P, Anckarsäter H, Happé F, Ronald A, 2013 Examining and interpreting the female protective effect against autistic behavior. *Proc. Natl. Acad. Sci. U. S. A* 110, 5258–62. doi:10.1073/pnas.1211070110 [PubMed: 23431162]
- Rodier PM, Ingram JL, Tisdale B, Croog VJ, 1997 Linking etiologies in humans and animal models: Studies of autism, in: *Reproductive Toxicology*, pp. 417–422. doi:10.1016/S0890-6238(97)80001-U [PubMed: 9100317]
- Romanski LM, LeDoux JE, 1992 Equipotentiality of thalamo-amygdala and thalamo-cortico-amygdala circuits in auditory fear conditioning. *J. Neurosci* 12, 4501–4509. [PubMed: 1331362]
- Roosendaal B, McEwen BS, Chattarji S, 2009 Stress, memory and the amygdala. *Nat. Rev. Neurosci* 10, 423–433. doi:10.1038/nrn2651 [PubMed: 19469026]
- Roulet FI, Lai JKY, Foster JA, 2013 In utero exposure to valproic acid and autism--a current review of clinical and animal studies. *Neurotoxicol. Teratol* 36, 47–56. doi:10.1016/j.ntt.2013.01.004 [PubMed: 23395807]

- Ruggeri B, Sarkans U, Schumann G, Persico AM, 2014 Biomarkers in autism spectrum disorder: The old and the new. *Psychopharmacology (Berl)*. 231, 1201–1216. doi:10.1007/s00213-013-3290-7 [PubMed: 24096533]
- Rutishauser U, Mamelak AN, Adolphs R, 2015 The primate amygdala in social perception - insights from electrophysiological recordings and stimulation. *Trends Neurosci* 38, 295–306. doi: 10.1016/j.tins.2015.03.001 [PubMed: 25847686]
- Rutishauser U, Tudusciuc O, Wang S, Mamelak AN, Ross IB, Adolphs R, 2013 Single-Neuron Correlates of Atypical Face Processing in Autism. *Neuron* 80, 887–899. doi: 10.1016/j.neuron.2013.08.029 [PubMed: 24267649]
- Ryan SJ, Ehrlich DE, Jasnow AM, Daftary S, Madsen TE, Rainnie DG, 2012 Spike-timing precision and neuronal synchrony are enhanced by an interaction between synaptic inhibition and membrane oscillations in the amygdala. *PLoS One* 7. doi: 10.1371/journal.pone.0035320
- Ryan SJ, Ehrlich DE, Rainnie DG, 2016 Morphology and dendritic maturation of developing principal neurons in the rat basolateral amygdala. *Brain Struct. Funct* 221, 839–854. doi: 10.1007/s00429-014-0939-x [PubMed: 25381464]
- Sah P, Faber ESL, Lopez De Armentia M, Power J, 2003 The amygdaloid complex: anatomy and physiology. *Physiol. Rev* 83, 803–844. doi: 10.1152/physrev.00002.2003 [PubMed: 12843409]
- Sallet J, Mars RB, Noonan MP, Andersson JL, O'Reilly JX, Jbabdi S, Croxson PL, Jenkinson M, Miller KL, Rushworth MFS, 2011 Social network size affects neural circuits in macaques. *Science* 334, 697–700. doi: 10.1126/science.1210027 [PubMed: 22053054]
- Sato W, Kochiyama T, Uono S, Matsuda K, Usui K, Inoue Y, Toichi M, 2011 Rapid Amygdala Gamma Oscillations in Response to Eye Gaze. *PLoS One* 6, e28188. doi: 10.1371/journal.pone.0028188 [PubMed: 22140541]
- Schultz RT, 2005 Developmental deficits in social perception in autism: the role of the amygdala and fusiform face area. *Int. J. Dev. Neurosci* 23, 125–41. doi:10.1016/j.ijdevneu.2004.12.012 [PubMed: 15749240]
- Schumann CM, Barnes CC, Lord C, Courchesne E, 2009 Amygdala Enlargement in Toddlers with Autism Related to Severity of Social and Communication Impairments. *Biol. Psychiatry* 66, 942–949. doi: 10.1016/j.biopsych.2009.07.007 [PubMed: 19726029]
- Schumann CM, Bauman MD, Amaral DG, 2011 Abnormal structure or function of the amygdala is a common component of neurodevelopmental disorders. *Neuropsychologia* 49, 745–759. doi: 10.1016/j.neuropsychologia.2010.09.028 [PubMed: 20950634]
- Schumann CM, Hamstra J, Goodlin-Jones BL, Lotspeich LJ, Kwon H, Buonocore MH, Lammers CR, Reiss AL, Amaral DG, 2004 The amygdala is enlarged in children but not adolescents with autism; the hippocampus is enlarged at all ages. *J. Neurosci* 24, 6392–401. doi: 10.1523/JNEUROSCI.1297-04.2004 [PubMed: 15254095]
- Setzer M, Ulfing N, 1999 Differential expression of calbindin and calretinin in the human fetal amygdala. *Microsc. Res. Tech* 46, 1–17. doi:10.1002/(SICI)1097-0029(19990701)46:1<1::AID-JEMT1>3.0.CO;2-P [PubMed: 10402268]
- Shabel SJ, Janak PH, 2009 Substantial similarity in amygdala neuronal activity during conditioned appetitive and aversive emotional arousal. *Proc. Natl. Acad. Sci. U. S. A* 106, 15031–15036. doi: 10.1073/pnas.0905580106 [PubMed: 19706473]
- Shaw P, Lawrence EJ, Radbourne C, Bramham J, Polkey CE, David AS, 2004 The impact of early and late damage to the human amygdala on “theory of mind” reasoning. *Brain* 127, 1535–1548. doi: 10.1093/brain/awh168 [PubMed: 15155523]
- Siebert M, Markowitsch HJ, Bartel P, 2003 Amygdala, affect and cognition: Evidence from 10 patients with Urbach-Wiethe disease. *Brain* 126, 2627–2637. doi:10.1093/brain/awg271 [PubMed: 12937075]
- Skuse DH, Lori A, Cubells JF, Lee I, Conneely KN, Puura K, Lehtimäki T, Binder EB, Young LJ, 2014 Common polymorphism in the oxytocin receptor gene (OXTR) is associated with human social recognition skills. *Proc. Natl. Acad. Sci. U. S. A* 111, 1987–92. doi: 10.1073/pnas.1302985111 [PubMed: 24367110]

- Sosa-Díaz N, Bringas ME, Atzori M, Flores G, 2014 Prefrontal cortex, hippocampus, and basolateral amygdala plasticity in a rat model of autism spectrum. *Synapse* 68, 468–473. doi:10.1002/syn.21759 [PubMed: 24985713]
- Sparks BF, Friedman SD, Shaw DW, Aylward EH, Echelard D, Artru AA, Maravilla KR, Giedd JN, Munson J, Dawson G, Dager SR, 2002 Brain structural abnormalities in young children with autism spectrum disorder. *Neurology* 59□ □, 184–192. doi:10.1212/WNL.59.2.184 [PubMed: 12136055]
- Spezio ML, Huang P-YS, Castelli F, Adolphs R, 2007 Amygdala Damage Impairs Eye Contact During Conversations with Real People. *J. Neurosci* 27, 3994–3997. doi: 10.1523/JNEUROSCI.3789-06.2007 [PubMed: 17428974]
- Stahmer AC, Schreibman L, Cunningham AB, 2010 Toward a technology of treatment individualization for young children with autism spectrum disorders. *Brain Res* 1380, 229–239. doi: 10.1016/j.brainres.2010.09.043 [PubMed: 20858466]
- Stanley DA, Adolphs R, 2013 Toward a neural basis for social behavior. *Neuron* 80, 816–826. doi: 10.1016/j.neuron.2013.10.038 [PubMed: 24183030]
- Štefánik P, Olexová L, Kršková L, 2015 Increased sociability and gene expression of oxytocin and its receptor in the brains of rats affected prenatally by valproic acid. *Pharmacol. Biochem. Behav* 131, 42–50. doi: 10.1016/j.pbb.2015.01.021 [PubMed: 25662821]
- Stephan H, Andy OJ, 1977 Quantitative comparison of the amygdala in insectivores and primates. *Cells Tissues Organs* 98, 130–153.
- Stone JL, Merriman B, Cantor RM, Yonan AL, Gilliam TC, Geschwind DH, Nelson SF, 2004 Evidence for sex-specific risk alleles in autism spectrum disorder. *Am. J. Hum. Genet* 75, 1117–23. doi: 10.1086/426034 [PubMed: 15467983]
- Stuber GD, Sparta DR, Stamatakis AM, van Leeuwen WA, Hardjoprajitno JE, Cho S, Tye KM, Kempadoo KA, Zhang F, Deisseroth K, Bonci A, 2011 Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. *Nature* 475, 377–80. doi: 10.1038/nature10194 [PubMed: 21716290]
- Swanson LW, Petrovich GD, 1998 What is the amygdala? *Trends Neurosci* 21, 323–331. doi:10.1016/S0166-2236(98)01265-X [PubMed: 9720596]
- Swartz JR, Wiggins JL, Carrasco M, Lord C, Monk CS, 2013 Amygdala habituation and prefrontal functional connectivity in youth with autism spectrum disorders. *J. Am. Acad. Child Adolesc. Psychiatry* 52, 84–93. doi:10.1016/j.jaac.2012.10.012 [PubMed: 23265636]
- Takahashi S, Kanai H, Miyamoto Y, 1976 Reassessment of elevated serotonin levels in blood platelets in early infantile autism. *J. Autism Child. Schizophr* 6, 317–26. [PubMed: 1036738]
- Tordjman S, Drapier D, Bonnot O, Gaignic R, Fortes S, Cohen D, Millet B, Laurent C, Roubertoux PL, 2007 Animal models relevant to schizophrenia and autism: Validity and limitations. *Behav. Genet* 37, 61–78. doi:10.1007/s10519-006-9120-5 [PubMed: 17160702]
- Tottenham N, Hertzog ME, Gillespie-Fynch K, Gilhooly T, Millner AJ, Casey BJ, 2014 Elevated amygdala response to faces and gaze aversion in autism spectrum disorder. *Soc. Cogn. Affect. Neurosci* 9, 106–17. doi:10.1093/scan/nst050 [PubMed: 23596190]
- Tran L, Lasher BK, Young K. a, Keele NB, 2013 Depletion of serotonin in the basolateral amygdala elevates glutamate receptors and facilitates fear-potentiated startle. *Transl. Psychiatry* 3, e298–8. doi:10.1038/tp.2013.66 [PubMed: 24002084]
- Tranel D, Gullickson G, Koch M, Adolphs R, 2006 Altered experience of emotion following bilateral amygdala damage. *Cogn. Neuropsychiatry* 11, 219–232. doi: 10.1080/13546800444000281 [PubMed: 17354069]
- Truitt WA, Sajdyk TJ, Dietrich AD, Oberlin B, McDougale CJ, Shekhar A, 2007 From anxiety to autism: Spectrum of abnormal social behaviors modeled by progressive disruption of inhibitory neuronal function in the basolateral amygdala in Wistar rats. *Psychopharmacology (Berl)*. 191, 107–118. doi: 10.1007/s00213-006-0674-y [PubMed: 17277936]
- Tye KM, Cone JJ, Schairer WW, Janak PH, 2010 Amygdala neural encoding of the absence of reward during extinction. *J. Neurosci* 30, 116–125. doi: 10.1523/JNEUROSCI.4240-09.2010 [PubMed: 20053894]

- U.S. Department of Health and Human Services, 2014 Prevalence of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 sites, United States, 2010. MMWR. Surveill. Summ 63, 1–21. doi:24670961
- Uddin LQ, Menon V, Young CB, Ryali S, Chen T, Khouzam A, Minshew NJ, Hardan AY, 2011 Multivariate searchlight classification of structural magnetic resonance imaging in children and adolescents with autism. *Biol. Psychiatry* 70, 833–841. doi: 10.1016/j.biopsych.2011.07.014 [PubMed: 21890111]
- Ulfing N, Setzer M, Bohl J, 2003 Ontogeny of the human amygdala. *Ann. New York Acad ...* 985, 22–33. doi: 10.1111/j.1749-6632.2003.tb07068.x
- Unal CT, Pare D, Zaborszky L, 2015 Impact of basal forebrain cholinergic inputs on basolateral amygdala neurons. *J. Neurosci* 35, 853–63. doi:10.1523/JNEUROSCI.2706-14.2015 [PubMed: 25589777]
- Urbach E, Wiethe C, 1929 Lipoidosis cutis et mucosae. *Virchows Arch. f{ü}r Pathol. Anat. und Physiol. und f{ü}r Klin. Medizin* 273, 285–319. doi:10.1007/BF02158983
- Ursin H, Kaada BR, 1960 Functional localization within the amygdaloid complex in the cat. *Electroencephalogr. Clin. Neurophysiol* 12, 1–20. doi: 10.1016/0013-4694(60)90058-4 [PubMed: 13840488]
- van Steensel FJA, Bögels SM, Perrin S, 2011 Anxiety Disorders in Children and Adolescents with Autistic Spectrum Disorders: A Meta-Analysis. *Clin. Child Fam. Psychol. Rev* 14, 302–317. doi: 10.1007/s10567-011-0097-0 [PubMed: 21735077]
- Vertes RP, 1991 A PHA-L analysis of ascending projections of the dorsal raphe nucleus in the rat. *J. Comp. Neurol* 313, 643–68. doi: 10.1002/cne.903130409 [PubMed: 1783685]
- Viviani D, Stoop R, 2008 Opposite effects of oxytocin and vasopressin on the emotional expression of the fear response. *Prog. Brain Res* 170, 207–218. doi: 10.1016/S0079-6123(08)00418-4 [PubMed: 18655884]
- Voineagu I, Wang X, Johnston P, Lowe JK, Tian Y, Horvath S, Mill J, Cantor RM, Blencowe BJ, Geschwind DH, 2011 Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature* 474, 380–384. doi:10.1038/nature10110 [PubMed: 21614001]
- Volkmar FR, Pauls D, 2003 Autism. *Lancet* 362, 1133–1141. doi: 10.1016/S0140-6736(03)14471-6 [PubMed: 14550703]
- Von Dem Hagen EAH, Stoyanova RS, Rowe JB, Baron-Cohen S, Calder AJ, 2014 Direct gaze elicits atypical activation of the theory-of-mind network in Autism spectrum conditions. *Cereb. Cortex* 24, 1485–1492. doi:10.1093/cercor/bht003 [PubMed: 23324559]
- Vuilleumier P, Driver J, 2007 Modulation of visual processing by attention and emotion: windows on causal interactions between human brain regions. *Philos. Trans. R. Soc. B Biol. Sci* 362, 837–855. doi: 10.1098/rstb.2007.2092
- Wang SSH, Kloth AD, Badura A, 2014 The Cerebellum, Sensitive Periods, and Autism. *Neuron* 83, 518–532. doi:10.1016/j.neuron.2014.07.016 [PubMed: 25102558]
- Watanabe T, Abe O, Kuwabara H, Yahata N, Takano Y, Iwashiro N, Natsubori T, Aoki Y, Takao H, Kawakubo Y, Kamio Y, Kato N, Miyashita Y, Kasai K, Yamasue H, 2014 Mitigation of sociocommunicational deficits of autism through oxytocin-induced recovery of medial prefrontal activity: a randomized trial. *JAMA psychiatry* 71, 166–75. doi: 10.1001/jamapsychiatry.2013.3181 [PubMed: 24352377]
- Watanabe T, Kuroda M, Kuwabara H, Aoki Y, Iwashiro N, Tatsunobu N, Takao H, Nippashi Y, Kawakubo Y, Kunimatsu A, Kasai K, Yamasue H, 2015 Clinical and neural effects of six-week administration of oxytocin on core symptoms of autism. *Brain* 138, 3400–3412. doi: 10.1093/brain/awv249 [PubMed: 26336909]
- Watanabe T, Otowa T, Abe O, Kuwabara H, Aoki Y, Natsubori T, Takao H, Kakiuchi C, Kondo K, Ikeda M, Iwata N, Kasai K, Sasaki T, Yamasue H, 2016 Oxytocin receptor gene variations predict neural and behavioral response to oxytocin in autism. *Soc. Cogn. Affect. Neurosci.* doi:10.1093/scan/nsw150
- Waterhouse L, London E, Gillberg C, 2016 ASD Validity. *Rev. J. Autism Dev. Disord* 3, 302–329. doi: 10.1007/s40489-016-0085-x

- Weiskrantz L, 1956 Behavioral changes associated with ablation of the amygdaloid complex in monkeys. *J. Comp. Physiol. Psychol* 49, 381–391. doi: 10.1037/h0088009 [PubMed: 13345917]
- Werling DM, Geschwind DH, 2013a Sex differences in autism spectrum disorders. *Curr. Opin. Neurol* 26, 146–53. doi:10.1097/WCO.0b013e32835ee548 [PubMed: 23406909]
- Werling DM, Geschwind DH, 2013b Understanding sex bias in autism spectrum disorder. *Proc. Natl. Acad. Sci. U. S. A* 110, 4868–9. doi:10.1073/pnas.1301602110 [PubMed: 23476067]
- White S, Oswald D, Ollendick T, Scahill L, 2009 Anxiety in children and adolescents with autism spectrum disorders. *Clin. Psychol. Rev* 29, 216–229. doi:10.1016/j.cpr.2009.01.003 [PubMed: 19223098]
- Williams G, King J, Cunningham M, Stephan M, Kerr B, Hersh JH, 2001 Fetal valproate syndrome and autism: additional evidence of an association. *Dev. Med. Child Neurol* 43, 202–206. doi: 10.1111/j.1469-8749.2001.tb00188.x [PubMed: 11263692]
- Williams PG, Hersh JH, 1997 A male with fetal valproate syndrome and autism. *Dev. Med. Child Neurol* doi: 10.1111/j.1469-8749.1997.tb07500.x
- Wing L, 2005 Reflections on opening Pandora's box. *J. Autism Dev. Disord* 35, 197–203. doi: 10.1007/s10803-004-1998-2 [PubMed: 15909406]
- Wing L, 1993 The definition and prevalence of autism: A review. *Eur. Child Adolesc. Psychiatry* 2, 61–74. doi: 10.1007/BF02098832 [PubMed: 21590530]
- Wing L, 1981 Asperger's syndrome: a clinical account. *Psychol. Med* 11, 115–29. [PubMed: 7208735]
- Wing L, Potter D, 2002 The epidemiology of autistic spectrum disorders: Is the prevalence rising? *Ment. Retard. Dev. Disabil. Res. Rev* 8, 151–161. doi: 10.1002/mrdd.10029 [PubMed: 12216059]
- Wolff SBE, Gründemann J, Tovote P, Krabbe S, Jacobson GA, Müller C, Herry C, Ehrlich I, Friedrich RW, Letzkus JJ, Lüthi A, 2014 Amygdala interneuron subtypes control fear learning through disinhibition. *Nature* 509, 453–8. doi:10.1038/nature13258 [PubMed: 24814341]
- Wu YE, Pan L, Zuo Y, Li X, Hong W, 2017 Detecting Activated Cell Populations Using Single-Cell RNA-Seq. *Neuron* 96, 313–329.e6. doi: 10.1016/j.neuron.2017.09.026 [PubMed: 29024657]
- Young LJ, Barrett CE, 2015 Can oxytocin treat autism? *Science* (80-.). 347, 825–826. doi: 10.1126/science.aaa8120
- Young MP, Scannell JW, Burns GA, Blakemore C, 1994 Analysis of connectivity: neural systems in the cerebral cortex. *Rev. Neurosci* 5, 227–50. [PubMed: 7889215]
- Zalla T, Sperduti M, 2013 The amygdala and the relevance detection theory of autism: an evolutionary perspective. *Front. Hum. Neurosci* doi: 10.3389/fnhum.2013.00894
- Zangrossi H, Graeff FG, 2014 Serotonin in anxiety and panic: Contributions of the elevated T-maze. *Neurosci. Biobehav. Rev* 46, 397–406. doi: 10.1016/j.neubiorev.2014.03.007 [PubMed: 24657635]
- Zhang B, Noble PL, Winslow JT, Pine DS, Nelson EE, 2012 Amygdala volume predicts patterns of eye fixation in rhesus monkeys. *Behav. Brain Res* 229, 433–437. doi:10.1016/j.bbr.2012.01.009 [PubMed: 22285417]
- Zürcher NR, Donnelly N, Rogier O, Russo B, Hippolyte L, Hadwin J, Lemonnier E, Hadjikhani N, 2013 It's All in the Eyes: Subcortical and Cortical Activation during Grotesqueness Perception in Autism. *PLoS One* 8, 1–14. doi: 10.1371/journal.pone.0054313

Highlights

- A novel approach for categorizing numerous amygdala functions with RDoC criteria
- Advances in understanding of the amygdala, autism, and their interrelation.
- Oxytocin and VPA: two tools for discerning dysfunction in autism and the amygdala

Figure 1:
The human amygdala, *from* Schumann 2011.

Figure 2: Components, inputs, and outputs of the amygdala.

The BLA has bidirectional connections to sensory and cortical areas including the Insula, Fusiform gyrus, Prefrontal Cortex (PFC), Orbital Frontal Cortex (OFC), Anterior Cingulate Cortex (ACC), and Superior Temporal Sulcus (STS), as well as temporal memory areas including CA1 and 3 of the Hippocampus, and Entorhinal cortex. The BLA innervates the Ventral Striatum, including the Nucleus Accumbens core (NAcc) and shell (NAcs), and the Bed Nucleus of the Stria Terminalis (BNST) which is also connected with the CeA. Monoaminergic modulation of the BLA comes from the Dorsal Raphe, Ventral Tegmental Area (VTA), Substantia Nigra (SN), and Locus Coeruleus (LC). The CeA is connected to the Basal Forebrain including the Medial Septal Nucleus (mSN) and the Substantia Inominata (SI), which contains the Nucleus Basalis and together with the Ventral Pallidum (innervated by the BLA) provides cholinergic input to the BA. The CeA also connects to regions in the brainstem, midbrain, and pons, such as the Parabrachial nucleus (PBN), Caudal Pontine Reticular Nucleus (PnC), Periaqueductal Gray (PAG), and Nucleus of the Solitary Tract (NTS). The CeA projects to regions of the hypothalamus including the Dorsomedial Hypothalamic Nucleus (DMH), Posterior Paraventricular Thalamus (pPVT), Lateral Hypothalamus (LH), and Suprachiasmatic Nucleus (SCN). Dashed line from Thalamus to CeA indicates a small number of direct projections there vs. substantial innervation of the LA.

Figure 3:
Developmental expansion of BLA dendritic arbors, *from* Ryan 2016.