Published in final edited form as:

Neurobiol Aging. 2019 January ; 73: 190–199. doi:10.1016/j.neurobiolaging.2018.09.031.

UNC13A polymorphism contributes to frontotemporal disease in sporadic amyotrophic lateral sclerosis

Katerina Placek^a, G. Michael Baer^b, Lauren Elman^c, Leo McCluskey^c, Laura Hennessy^a, Pilar M. Ferraro^a, Edward B. Lee^b, Virginia M.-Y. Lee^b, John Q. Trojanowski^b, Viviana M. Van Deerlin^b, Murray Grossman^a, David J Irwin^{a,b}, and Corey T. McMillan^{a,*}

^aUniversity of Pennsylvania, Department of Neurology, Penn Frontotemporal Degeneration Center, Philadelphia, PA

^bUniversity of Pennsylvania, Department of Pathology & Laboratory Medicine, Center for Neurodegenerative Disease Research, Philadelphia, PA

^cUniversity of Pennsylvania, Penn Comprehensive ALS Center, Philadelphia, PA

Abstract

The majority (90-95%) of amyotrophic lateral sclerosis (ALS) is sporadic, and ~50% of patients develop symptoms of FTD associated with shorter survival. The genetic polymorphism rs12608932 in *UNC13A* confers increased risk of sporadic ALS and sporadic frontotemporal degeneration (FTD), and modifies survival in ALS. Here, we evaluate whether rs12608932 is also associated with frontotemporal disease in sporadic ALS. We identified reduced cortical thickness in sporadic ALS with T1-MRI (N=109) relative to controls (N=113), and observed that minor allele (C) carriers exhibited greater reduction of cortical thickness in dorsal prefrontal, ventromedial prefrontal, anterior temporal, and middle temporal cortices and worse performance on a frontal-lobe mediated cognitive test (reverse digit span). In sporadic ALS with autopsy data (N=102), minor allele homozygotes exhibited greater burden of pTDP-43 pathology in the middle frontal, middle temporal, and motor cortices. Our findings demonstrate converging evidence that rs12608932 may modify frontotemporal disease in sporadic ALS, and suggest that rs12608932 may function as a prognostic indicator and could be used to define patient endophenotypes in clinical trials.

*Corresponding Author: University of Pennsylvania, Department of Neurology, Penn Frontotemporal Degeneration Center, 3400 Spruce Street, 3 West Gates, Philadelphia, PA, 19104, mcmillan@pennmedicine.upenn.edu; (215) 349-5863; (f) (215) 349-8464.
Author Contributions: KP, CTM drafting/revising manuscript for content, KP, GMB, DJI, CTM, study concept/design, KP, GMB, LE, LM, LH, PMF, VM-YL, JQT, VMVD, MG, DJI, CM performed analysis/interpretation of the data, KP, CM performed statistical analysis, and JQT, VM-YL, MG, CM obtained funding.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure Statement

All authors report no conflicts of interest.

1 Introduction

Amyotrophic lateral sclerosis (ALS) is a multi-system disorder primarily characterized by progressive degeneration of the upper and lower motor neurons (UMN, LMN) and affected individuals typically survive 2-5 years from symptom onset (Chiò et al., 2009). An estimated ~50% of individuals with ALS also develop impairments in at least one cognitive domain including executive function, social behavior, or language indicative of extramotor neurodegeneration in the frontal and temporal lobes, and about 10% of these individuals develop multi-domain impairments consistent with frank frontotemporal degeneration (FTD) (Montuschi et al., 2015; Ringholz et al., 2005). While ALS and FTD commonly feature tau DNA-binding protein-43kDa (TDP-43) pathology (Neumann et al., 2006) and can both be caused by pathogenic *C9orf72* repeat expansions (DeJesus-Hernandez et al., 2011; Renton et al., 2011), the mechanisms influencing the risk of progression from ALS to develop cognitive impairment and FTD have been under-evaluated. The presence of FTD is consistently associated with shorter survival in ALS patients (Elamin et al., 2011; Govaarts et al., 2016), and therefore it is critical to identify risk factors for frontotemporal disease in ALS.

The vast majority of ALS is considered “sporadic”, with only a small proportion of approximately 5-10% of ALS patients featuring a familial history or autosomal dominant source of disease (e.g. *C9orf72* expansions) (Taylor et al., 2016). Therefore, it is important to consider sources of common genetic variation that may influence risk of disease – including FTD – in ALS. Case-control genome-wide association studies (GWAS) have identified several single nucleotide polymorphisms (SNPs) associated with increased odds of having ALS (van Rheenen et al., 2016) or FTD (Van Deerlin et al., 2010). More recently, two loci, including rs3849942 in the *C9orf72* gene and rs4239633 in the *UNC13A* gene demonstrate shared genetic overlap between ALS and FTD (Karch et al., 2018). The observed rs3849942 SNP is a marker of the *C9orf72* expansion haplotype (Jones et al., 2013) with no additional genetic influence reported. However, loci in *UNC13A* including rs12608932, that is in high linkage disequilibrium (LD) with rs4239633 ($D' = 0.83$), are associated with increased risk of sporadic ALS (van Es et al., 2009) and sporadic FTD with underlying TDP-43 pathology (Diekstra et al., 2014). Clinical studies have further suggested that the minor allele (C) of rs12608932 (MAF=0.43) is associated with shorter survival in sporadic ALS, which has been demonstrated in multiple populations and under both additive and recessive minor allele models (Chiò et al., 2013; Diekstra et al., 2012; Vidal-Taboada et al., 2015).

In this report we perform a multimodal evaluation of rs12608932 to further investigate evidence for shared risk between ALS and FTD. Specifically, we test the hypothesis that the disease-associated allele (C) in rs12608932 is associated with frontotemporal cortical disease in sporadic ALS, including reduced cortical thickness, impaired cognitive performance, and increased vulnerability to TDP-43 pathology.

2 Materials and Methods

We evaluated 190 sporadic ALS cases distributed across a neuroimaging, neuropathological, and/or neuropsychological cohort, described in detail below and summarized in Supplementary Figure 1.

2.1 Neuroimaging cohort

We retrospectively evaluated 109 sporadic (see Genetic Screening) ALS patients recruited for research between 2004 and 2017 from the Penn Comprehensive ALS Clinic and Penn Frontotemporal Degeneration Center at the University of Pennsylvania Perelman School of Medicine. All patients were diagnosed with ALS by a board-certified neurologist (L.E., L.M., M.G., D.I) using the revised El Escorial criteria (Brooks et al., 2000), including possible, probable, and definite ALS. All patients were also assessed for frontotemporal dysfunction using established criteria (Strong et al., 2009) and those patients enrolled in research prior to 2009 were retrospectively evaluated through a chart review; in total 26 patients were diagnosed with ALS-FTD and 11 patients were diagnosed with ALS-cognitive impairment (ALSci).

To identify regions of significant cortical thinning in ALS, we recruited 113 demographically-comparable healthy controls who self-reported a negative history for neurologic or psychiatric disease and scored >27 on the MMSE (Table 1). There were no statistically significant differences in age, education, or sex between controls and ALS (all p values > .05). We assessed participant race and ethnicity via self-report. While population diversity is known to influence allele frequency across individuals, rs12608932 has a relatively equal minor allele frequency across populations (e.g., European 0.35, Africans 0.33, American 0.30) and post-hoc analyses limited to the majority race and ethnicity of our patient population (e.g. white non-Latino) remain significant or approach significance.

Demographic features of ALS and controls are summarized in Table 1.

All participants participated in an informed consent procedure approved by an Institutional Review Board convened at the University of Pennsylvania.

2.2 Autopsy cohort

We evaluated neuropathological data from 102 sporadic ALS autopsy cases (Table 2) who were diagnosed by a board-certified neuropathologist (JQT, EBL) with ALS due to TDP-43 pathology using immunohistochemistry (Lippa et al., 2009; Neumann et al., 2006) and published criteria (Mackenzie et al., 2011). This cohort included 21 patients from the ALS neuroimaging cohort (repeated analyses excluding these individuals summarized in Supplementary Table 1 revealed similar findings summarized in Supplementary Results). Autopsy cases were identified from the Integrated Neurodegenerative Disease Database (Toledo et al., 2014) after excluding cases with a family history of neurodegenerative disease or a known genetic mutation associated with ALS (see Genetic Screening). Within this autopsy cohort we rated the extent of phosphorylated TDP-43 (pTDP-43) intraneuronal inclusions (dots, wisps, skeins) for each sampled region on a semi-quantitative ordinal scale: 0 = none/rare, 1 = mild, 2 = moderate, 3 = severe/

numerous(Toledo et al., 2014). All neuropathological ratings were performed by an expert neuropathologist (JQT, EBL) blinded to genotype.

2.3 Alzheimer's Disease Neuroimaging Initiative (ADNI) cohort

To evaluate the disease specificity of any observed neuroanatomic and genetic associations we additionally performed cortical thickness analyses in 84 amyloid-negative (florbetapir SUVR <1.11) (Landau et al., 2012), cognitively-normal healthy controls from the publicly-available Alzheimer's Disease Neuroimaging Initiative (ADNI) database (see Supplementary Methods & Materials for details).

2.4 Genetic Screening

DNA was extracted from peripheral blood or frozen brain tissue following the manufacturer's protocols (Flexigene (Qiagen) or QuickGene DNA whole blood kit (Autogen) for blood, and QIAasympy DNA Mini Kit (Qiagen) for brain tissue). All patients were screened for *C9orf72* hexanucleotide repeat expansions using a modified repeat-primed polymerase-chain reaction (PCR) as previously described(Suh et al., 2015), and we excluded any patient with ≥ 30 hexanucleotide repeats. Of the remaining individuals, we evaluated family history using a three-generation pedigree history as previously reported(Wood et al., 2013). For cases that had a family history of the same disease we sequenced 45 genes previously associated with neurodegenerative disease(Toledo et al., 2014), including four genes known to be associated with ALS (e.g. *SOD1*(Rosen, 1993), *TARDBP*(Kabashi et al., 2008), *FUS*(Kwiatkowski et al., 2009) (Vance et al., 2009), and *VCP*(Johnson et al., 2010)). Sequencing was performed using a custom-targeted next-generation sequencing panel (MiND-Seq)(Toledo et al., 2014), and analyzed using Mutation Surveyor software (Soft Genetics, State College, PA). We excluded any individuals identified as having a pathogenic mutation.

2.5 SNP Genotyping

All neuroimaging and neuropathology cases were genotyped using peripheral or brain DNA, extracted as above, for rs12608932 using a custom-designed Pan Neurodegenerative Disease-oriented Risk Allele (PANDoRA) panel designed to genotype 92 common and risk allele variants identified in association and other studies as modifying risk of disease or a phenotype for several neurodegenerative diseases, including ALS as well as FTD, Parkinson's disease, and Alzheimer's disease (Toledo et al., 2014). While our analyses focus on a single genotype, rs12608932, this approach provides a cost-effective manner to genotyping that can be used for future comparative studies as previously reported by our center (McMillan et al., 2018) (McMillan et al., 2014). Briefly, the 92 SNP Type assays were designed by D3 Assay Design tool (Fluidigm). Allele-specific PCR was performed using the 96.96 Dynamic Array integrated fluidic circuits (Fluidigm) and genotyping was carried out using the BioMark HD system (Fluidigm) according to the manufacturer's protocol. The genotype call data was collected and analyzed using BioMark Genotyping Analysis software.

2.6 Clinical Evaluations

Detailed clinical evaluations were available for a subset of 88 (79%) sporadic ALS patients in the neuroimaging cohort. These patients were clinically assessed within approximately two months of neuroimaging acquisition date ($M = 1.62$, $SD = 2.63$) for motor function using the Revised ALS Functional Rating Scale (ALSFRS-R)(Cedarbaum et al., 1999) and cognitive function using the Forward and Reverse Digit Span, the Visual-Verbal Test (VVT), letter fluency, and the Mini-Mental State Exam (MMSE). Forward digit span, a measure of auditory-verbal short-term memory, and reverse digit span, a measure of auditory-verbal working memory, were assessed in an untimed manner with repetition of increasingly longer sequences until the patient erred; the maximum number of digits on a correct trial was recorded (Wechsler, 1945). Our group has previously demonstrated that the VVT is a brief, untimed measure of cognitive flexibility with minimal motor demands appropriate for use in an ALS patient population(Evans et al., 2015). Participants first identify a feature shared by three of four simple geometric designs, and are next challenged to identify a different feature shared by another combination of three of the four same geometric designs; a discrepancy between the number correct on the first and second identifications of 10 trials is considered a sign of reduced mental flexibility. Letter fluency is a measure of executive control and verbal ability(Abrahams et al., 2000), and was assessed by the number of unique words beginning with “F” a patient was able to generate in one minute (excluding proper nouns and numbers); patients did not complete letter fluency if bulbar upper or lower motor symptoms were present upon exam. The MMSE is a 30-point questionnaire that evaluates global dementia severity(Crum et al., 1993); we calculated the percentage correct on MMSE as some patients were not able to complete the entire test due to motor disability.

In addition to reporting raw patient performance on neuropsychological assessments, we report the percent of patients impaired on each task relative to published normative data of healthy controls matched to the mean age and mean education of our patients where available (Tombaugh et al., 1999; Weintraub et al., 2009) (see Table 1). We report patient performance on the Visual Verbal Test relative to normative data based on healthy controls recruited by the Penn Frontotemporal Degeneration Center who were matched to the mean age and mean education of our patients ($N=31$ (17 Female); Age, years: $M=60.58$, $SD=12.80$; Education, years: $M=15.29$, $SD=2.52$). We defined patient impairment on each task as performance at or greater than 1.96 standard deviations, equivalent to $p<0.05$, below normative data from healthy controls.

2.7 Neuroimaging Acquisition & Processing

High-resolution T1-weighted MPRAGE structural scans were acquired for neuroimaging participants using a 3T Siemens Tim Trio scanner with an 8-channel head coil, with $T=1620\text{ms}$, $T=3.09\text{ms}$, flip angle= 15° , 192×256 matrix, and 1mm^3 voxels. T1-weighted MRI images were then preprocessed using ANTs Cortical Thickness software(Tustison et al., 2014). Each individual dataset was deformed into a standard local template space in a canonical stereotactic coordinate system. Advanced Normalization Tools (ANTs) provides a highly accurate registration routine using symmetric and topology-preserving diffeomorphic deformations to minimize bias toward the reference space and to capture the deformation necessary to aggregate images in a common space. Then, we used N4 bias correction to

minimize heterogeneity (Tustison et al., 2010) and the ANTs Atropos tool to segment images into six tissue classes (cortex, white matter, CSF, subcortical grey structures, brainstem, and cerebellum) using template-based priors and to generate probability maps of each tissue. Voxel-wise cortical thickness was measured in millimeters (mm) from the pial surface and then transformed into Montreal Neurological Institute (MNI) space, smoothed using a 2 sigma full-width half-maximum Gaussian kernel, and downsampled to 2mm isotropic voxels.

2.8 Neuroimaging analysis

We used *randomise* software from FSL to perform nonparametric, permutation-based statistical analyses of cortical thickness images from our ALS and control neuroimaging cohorts and the ADNI control cohort. Permutation-based statistical testing is robust to concerns regarding multiple comparisons since, rather than a traditional assessment of two sample distributions, this method assesses a true assignment of factors (e.g., genotype) to cortical thickness relative to many (e.g., 10,000) random assignments (Winkler et al., 2014).

First, we identified regions of reduced cortical thickness in ALS relative to healthy controls, constraining analysis using an explicit mask restricted to high probability cortex ($>.4$). We report clusters that survive $p < .005$ threshold-free cluster enhancement (TFCE) (Smith and Nichols, 2009) corrected for family-wise error (FWE) relative to 10,000 random permutations.

Next, we evaluated whether rs12608932 genotype relates to magnitude of reduced cortical thickness in ALS, covarying for disease duration, age, and sex in an effort to control for factors associated with reduced cortical thickness but not specifically associated with SNP genotype. We restricted this analysis to a mask defined by regions of reduced cortical thickness in ALS relative to controls so that we could focus our interpretation in the context of cortical degeneration affected by ALS. We report clusters that survive $p < .01$ (uncorrected) threshold and cluster extent threshold of >20 adjacent voxels relative to 10,000 random permutations; we employ an uncorrected threshold to minimize the chance of Type II error (not observing a true result).

We performed comparable analyses to evaluate SNP genotype relative to cortical thickness in the ADNI control cohort and adopted the same statistical thresholds as described above.

2.9 Statistical Analyses

All additional statistical analysis were performed using R (www.r-project.org). For assessment of ordinal neuropathology data we performed ordinal logistic regression using the *MASS* package in R to investigate whether burden of TDP-43 pathology differed according to genotype at rs12608932, covarying for age, sex, and disease duration at death. For clinical comparisons, we used multiple linear regression to evaluate the association between genotype at rs12608932 and performance on forward and reverse digit span, the MMSE, the VVT, and letter fluency; we included age, sex, disease duration, ALSFRS-R total score, and cognitive diagnosis (i.e. diagnosis of ALS-FTD, ALSci) as covariates in each analysis.

3 Results

3.1 Reduced cortical thickness in ALS associated with the rs12608932 minor allele

A group comparison of ALS patients relative to healthy controls revealed reduced cortical thickness in the bilateral frontal and temporal lobes, consistent with the pattern of cortical degeneration associated with ALS-FTD spectrum disorders (see Figure 1, Table 3).

To evaluate disease-specific genotype and neuroanatomic associations we restricted our subsequent analyses to regions of reduced cortical thickness observed in ALS. Evaluation of rs12608932 under a minor allele additive model revealed reduced cortical thickness associated with the minor allele (C) in the right anterior temporal lobe, bilateral ventromedial prefrontal cortex, left middle temporal gyrus, and left dorsal prefrontal cortex (see Figure 1, Table 3); we observed similar findings under a minor allele recessive model (see Supplementary Results including Supplementary Table 2 and Supplementary Figure 2). Investigation of the inverse association between rs12608932 genotype (i.e. number of major non-risk allele “A”) and cortical thickness yielded no statistically significant findings for either additive or recessive models (not shown), suggesting that the rs12608932 minor allele C is specifically associated with reduced cortical thickness in these regions.

To examine the disease specificity of the relationship between rs12608932 genotype and reduced cortical thickness in ALS, we performed comparable analyses in the ADNI control cohort restricted to the same regions of reduced cortical thickness observed in ALS. We observed no relationship between cortical thickness and genotype at rs12608932 under minor allele additive or recessive models (not shown).

3.2 Working memory performance associated with rs12608932

Given the observed association between rs12608932 and frontal and temporal cortices, we evaluated each clinical assessment to investigate potential clinical consequences of the observed neuroanatomic and genetic associations.

We first investigated whether ALSFRS-R relates to rs12608932 genotype. Under a minor allele additive model, minor allele heterozygotes (AC) ($\beta=3.41$, $p=.04$) but not minor allele homozygotes at rs12608932 (CC) ($\beta=1.97$, $p=.35$) demonstrated higher ALSFRS-R total score compared to major allele homozygotes (AA); disease duration ($\beta=-.037$, $p=.1$), age ($\beta=.059$, $p=.38$), sex ($\beta=1.53$, $p=.33$), and cognitive diagnosis (ALSci: $\beta=-2.34$, $p=.37$; ALS-FTD: $\beta=-0.88$, $p=.62$) were not found to significantly affect ALSFRS-R Total in this model. However, under a minor allele recessive model, we did not observe a significant difference in ALSFRS-R Total score between minor allele heterozygotes (AC) and minor allele homozygotes (CC) at rs12608932 ($\beta=0.34$, $p=.87$) compared to major allele homozygotes (AA). To account for the observed difference in ALSFRS-R under a minor allele additive model, we covaried for ALSFRS-R Total score in all subsequent regressions.

Under a minor allele additive model, we observed a trend whereby minor allele homozygotes at rs12608932 (CC) ($\beta=-.81$, $p=.086$) performed worse than major allele homozygotes (AA) on reverse digit span after accounting for cognitive diagnosis (ALSci: $\beta=-1.73$, $p=.0014$; ALS-FTD: $\beta=-2.09$, $p<.00001$); heterozygous genotype (AC) ($\beta=.32$, $p=$

38), disease duration ($\beta = -.0022$, $p = .64$), ALSFRS-R total score ($\beta = -.012$, $p = .60$), and age ($\beta = -.0021$, $p = .89$) did not relate to reverse digit span performance in this model.

Under a minor allele recessive model, minor allele homozygotes at rs12608932 (CC) ($\beta = -.96$, $p = .029$) performed significantly worse than major allele homozygotes (AA) and heterozygotes (AC) on reverse digit span after accounting for cognitive diagnosis (ALSci: $\beta = -1.69$, $p = .0017$; ALS-FTD: $\beta = -2.08$, $p < .00001$); disease duration ($\beta = -.0018$, $p = .69$), ALSFRS-R total score ($\beta = -.0013$, $p = .95$), age ($\beta = -.0013$, $p = .76$), and sex ($\beta = .55$, $p = .11$), did not significantly relate to reverse digit span performance in this model.

Performance on other neuropsychological tests was not found to associate with rs12608932 genotype, including forward digit span (CC: $\beta = -.03$, $p = .93$; CA: $\beta = .40$, $p = .20$), letter fluency (CC: $\beta = .26$, $p = .85$; CA: $\beta = 1.61$, $p = .15$), MMSE (CC: $\beta = .036$, $p = .28$; CA: $\beta = .006$, $p = .82$), and the VVT (CC: $\beta = -.22$, $p = .67$; CA: $\beta = -.40$, $p = .32$). We observed similar results under minor allele recessive models: forward digit span (CC: $\beta = -.22$, $p = .57$), letter fluency (CC: $\beta = -.39$, $p = .76$), MMSE (CC: $\beta = .034$, $p = .26$), and the VVT (CC: $\beta = .12$, $p = .80$).

3.3 pTDP-43 pathologic burden associated with rs12608932 in middle frontal and temporal cortices

To evaluate converging evidence for our observed genetic and neuroanatomic associations, we assessed ordinal neuropathologist ratings of pTDP-43 pathologic burden in the middle frontal, temporal, and motor cortices (see Figure 2), which were associated with rs12608932 in the above neuroimaging analyses.

Minor allele homozygotes (CC) were 8.26 times (95% CI: 2.01, 38.64; $p = .0043$) more likely and heterozygotes (AC) were 5.53 times (95% CI: 1.78, 21.21; $p = .0057$) more likely to have higher TDP-43 burden in the middle frontal cortex relative to major allele homozygotes (AA). Minor allele homozygotes (CC) were 4.40 times (95% CI: 1.25, 16.23; $p = .022$) more likely to have higher TDP-43 burden in the temporal cortex and – reaching marginal significance – 3.04 times (95% CI: 0.98, 9.72; $p = .056$) more likely to have higher TDP-43 burden in the motor cortex relative to major allele homozygotes (AA); minor allele heterozygotes (AC) were not more likely to have higher TDP-43 burden relative to major allele homozygotes in either region (both p values $> .1$).

4 Discussion

In this study, we evaluated whether rs12608932 genotype in *UNC13A* was associated with frontotemporal disease in sporadic ALS using a novel multimodal approach integrating genetic, neuroimaging, clinical, and neuropathology data. Our results indicate that sporadic ALS patients who are carriers of the rs12608932 minor allele (C) show reduced cortical thickness in regions including the dorsal and ventromedial prefrontal cortex, anterior and middle temporal cortex, and premotor cortex, and that minor allele homozygotes (CC) demonstrate worse performance on reverse digit span, a frontal-lobe mediated cognitive test. We did not observe a relationship between rs12608932 genotype and cortical thickness in the amyloid-negative ADNI healthy control cohort, suggesting that the association between rs12608932 and reduced cortical thickness is specific to ALS. Furthermore, in our sporadic

ALS autopsy cohort, we demonstrate that carriers of the rs12608932 minor allele have increased odds of pTDP-43 pathologic burden in the middle frontal cortex, middle temporal cortex, and motor cortex, consistent with our neuroimaging findings. To our knowledge, our study provides novel evidence that the minor allele of rs12608932 in *UNC13A* is associated with *in vivo* frontotemporal cortical atrophy, impaired cognitive performance, and greater burden of pTDP-43 pathological inclusions in sporadic ALS.

rs12608932 was first identified through a two-stage GWAS as a susceptibility locus for sporadic ALS with a combined $P = 2.53 \times 10^{-14}$ (van Es et al., 2009). rs12608932 maps to a haplotype block within the boundaries of gene *UNC13A*, which regulates presynaptic vesicle priming and glutamate release at neuromuscular synapses, and mice lacking the *UNC13A* homolog have arrested synaptic vesicle maturation and disrupted glutamatergic transmission (Augustin et al., 1999). Subsequent population-based study indicated the minor allele (C) at rs12608932 as a risk factor for shorter survival in sporadic ALS under both additive and recessive models (Chiò et al., 2013; Diekstra et al., 2012; Vidal-Taboada et al., 2015), and as a modifier of physical symptom progression on the ALSFRS-R (Vidal-Taboada et al., 2015). In addition to risk and progression of sporadic ALS, rs12608932 was also identified to serve as a risk locus for sporadic FTLTDP (Diekstra et al., 2014), suggesting rs12608932 as a potential link between ALS and FTLTDP. More recently, an additional SNP in *UNC13A* in high LD with rs12608932 (rs4239633; $D' = 0.83$) was identified as demonstrating selective genetic overlap between ALS and FTD in GWAS meta-analysis (Karch et al., 2018).

Our findings corroborate rs12608932 as a genetic link between ALS and FTD with TDP-43 pathology, and specifically demonstrate that the minor allele is associated with reduced cortical thickness, worse working memory performance, and greater burden of TDP-43 pathology in the frontal and temporal lobes. Importantly, we use continuous disease traits from multiple modalities (i.e. structural imaging, cognitive testing, neuropathology data) to present converging evidence that rs12608932 confers increased risk of frontotemporal disease in sporadic ALS and relates to patient cognitive performance. This approach offers an advance over discovery GWAS which compare only categorical clinical designations (e.g. ALS vs. healthy controls), and allows detailed phenotypic characterization associated with rs12608932 genotype.

Furthermore, the observed relationship between rs12608932 and both frontotemporal cortical thinning and burden of TDP-43 pathology are consistent with the disease anatomy of ALS-FTD spectrum disorders. Structural MRI studies have previously demonstrated progressive frontotemporal gray matter degeneration over disease course (Keil et al., 2012; Kwan et al., 2013; Menke et al., 2014; Müller et al., 2016; Senda et al., 2011; Verstraete et al., 2012; 2014), and that degree of frontotemporal cortical thinning relates to cognitive-behavioral phenotype (Agosta et al., 2016; Schuster et al., 2014). Additionally, staging of neuropathological burden suggests stereotyped propagation of TDP-43 from motor regions (brainstem, spinal cord) to frontal and temporal neocortex over the course of disease (Braak et al., 2013; Brettschneider et al., 2013). Our finding of greater TDP-43 pathologic burden associated with the rs12608932 in frontal and temporal neocortex may thus be interpreted to reflect more progressive disease propagation in sporadic ALS patients who are minor allele

carriers. This finding could also be related to TDP-43 pathologic subtype(Mackenzie et al., 2011), and future work is necessary to 1) evaluate the influence of TDP-43 pathologic subtype on the anatomic distribution and phenotypic presentation (e.g. comorbid symptoms of frontotemporal disease) of disease, and 2) investigate how rs12608932 genotype relates to TDP-43 pathologic subtype. With this in mind, rs12608932 genotype may potentially be used prognostically to evaluate risk of frontotemporal cortical disease in ALS, which has been previously associated with reduced survival(Elamin et al., 2011; Govaarts et al., 2016).

The observed relationships between rs12608932, reduced cortical thickness in the dorsal and ventromedial prefrontal cortices, and worse performance on reverse digit span are congruent with the neuroanatomy of working memory. Indeed, functional activation of the dorsal and ventromedial prefrontal cortices relates to memory load function and information retrieval on tasks of working memory, respectively(Rypma and D'Esposito, 1999).

We also observed a relationship between rs12608932 and reduced cortical thickness in the right orbital frontal cortex, right anterior temporal lobe, and left middle temporal gyrus. Our group has previously shown that reduced cortical thickness in the orbital frontal cortex is associated with behavioral disinhibition and apathy in patients with FTD(Massimo et al., 2009), while others have shown that cortical thinning in the right anterior temporal lobe and left middle temporal gyrus are associated with impaired semantic knowledge and theory of mind in patients with FTD(Irish et al., 2014; Rohrer et al., 2009). Impairments in behavior and language are common to both ALS and FTD(Beeldman et al., 2018), and prospective studies are necessary to evaluate potential rs12608932 associations relative to language and behavioral function using more comprehensive neuropsychological batteries like the Edinburgh Cognitive Assessment Scale (ECAS) (Abrahams et al., 2014), which were not available in this retrospective study.

Our findings add to an increasing body of evidence in support of a clinicopathologic continuum between ALS and FTD with underlying TDP-43 pathology, and specifically suggest that genetic polymorphisms may relate to the phenotypic presentation of frontotemporal disease in sporadic ALS. This is consistent with prior work from our group demonstrating that TDP-43 pathology-associated SNPs relate to selective neuroanatomic distribution of cortical atrophy and white matter degeneration in patients with sporadic forms of FTD(McMillan et al., 2014), modify disease onset and survival in FTD with *C9orf72* repeat expansions (Chen-Plotkin et al., 2014) (Gallagher et al., 2014), and confer risk for impaired executive function in ALS(Vass et al., 2010).

The identification of genetic polymorphisms associated with disease phenotypes holds important implications for both basic science research and translational application. Our observed association between rs12608932 and frontotemporal disease in sporadic ALS motivates further investigation into potential mechanisms of disease vulnerability associated with genetic polymorphisms. While the function of gene *UNC13A* has previously been characterized(Augustin et al., 1999), future work is necessary to determine the extent to which rs12608932 genotype provides an actual disease modifier or is an association related to matching downstream transcription sites or other long-range interactions(Li et al., 2016). In regards to translational application, our findings may contribute to patient stratification

for clinical trials. Prior research has demonstrated baseline differences in patients stratified for SNP genotype and indicate a contribution of genetic polymorphisms in dose-response to treatment (Wang et al., 2011), leading to the incorporation of genetic polymorphisms in the design and analysis of clinical trials (Zhang et al., 2017).

Several caveats should be considered in the present study. Our data were limited according to strict inclusion criteria to investigate the clinical, pathologic, and regional anatomical differences in sporadic ALS patients relative to rs12608932 genotype. While we establish multiple sources of converging evidence for rs12608932 genotype relating to frontal disease, replication of the present findings in a large independent cohort using a prospective design is necessary. Furthermore, additional research is required to determine the extent to which pTDP-43 pathological burden directly or indirectly relates to reductions in cortical thickness. In this study, our evaluation of cognitive performance was retrospective and limited to measures broadly assessing executive function and global cognition. Future studies using revised diagnostic criteria for frontotemporal dysfunction in ALS (Strong et al., 2017) and specialized assessment of cognitive function in patients with ALS, such as the Edinburgh Cognitive and Behavioral ALS Screen (Abrahams et al., 2014), are necessary to also assess impairments in other domains including language and behavior. Here, we focus on genetic contributions to frontotemporal disease in sporadic ALS. However, environmental factors such as those associated with cognitive reserve have been demonstrated to influence frontotemporal disease neuroanatomy in FTD (Massimo et al., 2018) (Massimo et al., 2015) (Placek et al., 2016), also when considered in addition to genetic polymorphisms (Premi et al., 2017). Additional study is necessary to examine frontotemporal disease in sporadic ALS relative to both genetic polymorphisms and environmental factors associated with cognitive reserve.

With these caveats in mind, our research demonstrates converging clinical, neuroimaging, and pathologic evidence supporting the hypothesis that the common genetic polymorphism rs12608932 contributes to frontotemporal disease phenotype in sporadic ALS. These findings stimulate investigation into additional genetic contributors to the nature of disease in sporadic ALS, and suggest their importance in prognostic consideration and treatment trials in patients with ALS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported in part by the National Institutes of Health (NS106754, AG043503, AG017586, NS092091), the ALS Association, and the Dana Foundation.

References

Abrahams S, Leigh PN, Harvey A, Vythelingum GN, Gris D, Goldstein LH, 2000 Verbal fluency and executive dysfunction in amyotrophic lateral sclerosis (ALS). *Neuropsychologia* 38, 734–747. [PubMed: 10689049]

- Abrahams S, Newton J, Niven E, Foley J, Bak TH, 2014 Screening for cognition and behaviour changes in ALS. *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration* 15, 9–14. doi: 10.3109/21678421.2013.805784 [PubMed: 23781974]
- Agosta F, Ferraro PM, Riva N, Spinelli EG, Chiò A, Canu E, Valsasina P, Lunetta C, Iannaccone S, Copetti M, Prudente E, Comi G, Falini A, Filippi M, 2016 Structural brain correlates of cognitive and behavioral impairment in MND. *Hum Brain Mapp* n/a–n/a. doi:10.1002/hbm.23124
- Augustin I, Rosenmund C, Südhof TC, Brose N, 1999 Munc13-1 is essential for fusion competence of glutamatergic synaptic vesicles. *Nature* 400, 457–461. doi:10.1038/22768 [PubMed: 10440375]
- Beeldman E, Raaphorst J, Klein Twennaar M, Govaarts R, Pijnenburg YAL, de Haan RJ, de Visser M, Schmand BA, 2018 The cognitive profile of behavioural variant FTD and its similarities with ALS: a systematic review and meta-analysis. *Journal of Neurology, Neurosurgery & Psychiatry* jnnp–2017–317459–9. doi:10.1136/jnnp-2017-317459
- Braak H, Brettschneider J, Ludolph AC, Lee VM, Trojanowski JQ, Del Tredici K, 2013 Amyotrophic lateral sclerosis--a model of corticofugal axonal spread. *Nature Reviews Neurology* 9, 708–714. doi: 10.1038/nrneurol.2013.221 [PubMed: 24217521]
- Brettschneider J, Del Tredici K, Toledo JB, Robinson JL, Irwin DJ, Grossman M, Suh E, Deerlin VM, Wood EM, Baek Y, Kwong L, Lee EB, Elman L, McCluskey L, Fang L, Feldengut S, Ludolph AC, Lee VMY, Braak H, Trojanowski JQ, 2013 Stages of pTDP-43 pathology in amyotrophic lateral sclerosis. *Ann Neurol* 74, 20–38. doi:10.1002/ana.23937 [PubMed: 23686809]
- Brooks BR, Miller RG, Swash M, Munsat TL, World Federation of Neurology Research Group on Motor Neuron Diseases, 2000 El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis., in: Presented at the Amyotrophic lateral sclerosis and other motor neuron disorders : official publication of the World Federation of Neurology, Research Group on Motor Neuron Diseases, pp. 293–299.
- Cedarbaum JM, Stambler N, Malta E, Fuller C, Hilt D, Thurmond B, Nakanishi A, 1999 The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. BDNF ALS Study Group (Phase III). *Journal of the Neurological Sciences* 169, 13–21. [PubMed: 10540002]
- Chen-Plotkin A, Gallagher M, Suh E, Grossman M, Elman L, McCluskey L, Trojanowski J, Lee V, Van Deerlin V, 2014 TMEM106B is a Genetic Modifier of Frontotemporal Lobar Degeneration with C9orf72 Hexanucleotide Repeat Expansions (I5–1.008). *Neurology* 82, I5–1.008.
- Chiò A, Logroscino G, Hardiman O, Swingler R, Mitchell D, Beghi E, Traynor BG, Consortium, O.B.O.T.E., 2009 Prognostic factors in ALS: A critical review. *Amyotrophic Lateral Sclerosis* 10, 310–323. doi:10.3109/17482960802566824 [PubMed: 19922118]
- Chiò A, Mora G, Restagno G, Brunetti M, Ossola I, Barberis M, Ferrucci L, Canosa A, Manera U, Moglia C, Fuda G, Traynor BJ, Calvo A, 2013 UNC13A influences survival in Italian amyotrophic lateral sclerosis patients: a population-based study. *Neurobiology of Aging* 34, 357.e1–5. doi: 10.1016/j.neurobiolaging.2012.07.016
- Crum RM, Anthony JC, Bassett SS, Folstein MF, 1993 Population-Based Norms for the Mini-Mental State Examination by Age and Educational Level. *JAMA* 269, 2386–2391. doi:10.1001/jama.1993.03500180078038 [PubMed: 8479064]
- DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson AM, Finch NA, Flynn H, Adamson J, Kouri N, Wojtas A, Sengdy P, Hsiung G-YR, Karydas A, Seeley WW, Josephs KA, Coppola G, Geschwind DH, Wszolek ZK, Feldman H, Knopman DS, Petersen RC, Miller BL, Dickson DW, Boylan KB, Graff-Radford NR, Rademakers R, 2011 Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72, 245–256. doi:10.1016/j.neuron.2011.09.011 [PubMed: 21944778]
- Diekstra FP, Van Deerlin VM, Van Swieten JC, Al-Chalabi A, Ludolph AC, Weishaupt JH, Hardiman O, Landers JE, Brown RH, van Es MA, Pasterkamp RJ, Koppers M, Andersen PM, Estrada K, Rivadeneira F, Hofman A, Uitterlinden AG, Van Damme P, Melki J, Meininger V, Shatunov A, Shaw CE, Leigh PN, Shaw PJ, Morrison KE, Fogh I, Chiò A, Traynor BJ, Czell D, Weber M, Heutink P, de Bakker PIW, Silani V, Robberecht W, van den Berg LH, Veldink JH, 2014 C9orf72 and *UNC13A* are shared risk loci for amyotrophic lateral sclerosis and frontotemporal dementia: a genome-wide meta-analysis. *Ann Neurol* 76, 120–133. doi:10.1002/ana.24198 [PubMed: 24931836]

- Diekstra FP, van Vught PWJ, van Rheenen W, Koppers M, Pasterkamp RJ, van Es MA, Schelhaas HJ, de Visser M, Robberecht W, Van Damme P, Andersen PM, van den Berg LH, Veldink JH, 2012 UNC13A is a modifier of survival in amyotrophic lateral sclerosis. *Neurobiology of Aging* 33, 630.e3–8. doi:10.1016/j.neurobiolaging.2011.10.029
- Elamin M, Phukan J, Bede P, Jordan N, Byrne S, Pender N, Hardiman O, 2011 Executive dysfunction is a negative prognostic indicator in patients with ALS without dementia. *Neurology* 76, 1263–1269. doi:10.1212/WNL.0b013e318214359f [PubMed: 21464431]
- Evans J, Olm C, McCluskey L, Elman L, Boller A, Moran E, Rascovsky K, Bisbing T, McMillan CT, Grossman M, 2015 Impaired cognitive flexibility in amyotrophic lateral sclerosis. *Cogn Behav Neurol* 28, 17–26. doi:10.1097/WNN.000000000000049 [PubMed: 25812127]
- Gallagher MD, Suh E, Grossman M, Elman L, McCluskey L, Van Swieten JC, Al-Sarraj S, Neumann M, Gelpi E, Ghetti B, Rohrer JD, Halliday G, Van Broeckhoven C, Seilhean D, Shaw PJ, Frosch MP, Alafuzoff I, Antonell A, Bogdanovic N, Brooks W, Cairns NJ, Cooper-Knock J, Cotman C, Cras P, Cruts M, De Deyn PP, DeCarli C, Dobson-Stone C, Engelborghs S, Fox N, Galasko D, Gearing M, Gijssels I, Grafman J, Hartikainen P, Hatanpää KJ, Highley JR, Hodges J, Hulette C, Ince PG, Jin L-W, Kirby J, Kofler J, Kril J, Kwok JBJ, Levey A, Lieberman A, Llado A, Martin J-J, Masliah E, McDermott CJ, McKee A, McLean C, Mead S, Miller CA, Miller J, Munoz DG, Murrell J, Paulson H, Piguet O, Rossor M, Sanchez-Valle R, Sano M, Schneider J, Silbert LC, Spina S, van der Zee J, Van Langenhove T, Warren J, Wharton SB, White CL, Woltjer RL, Trojanowski JQ, Lee VMY, Van Deerlin V, Chen-Plotkin AS, 2014 TMEM106B is a genetic modifier of frontotemporal lobar degeneration with C9orf72 hexanucleotide repeat expansions. *Acta Neuropathologica* 127, 407–418. doi:10.1007/s00401-013-1239-x [PubMed: 24442578]
- Govaarts R, Beeldman E, Kampelmacher MJ, van Tol M-J, van den Berg LH, vander Kooi AJ, Wijkstra PJ, Zijnen-Suyker M, Cobben NAM, Schmand BA, de Haan RJ, de Visser M, Raaphorst J, 2016 The frontotemporal syndrome of ALS is associated with poor survival. *Journal of Neurology* 1–8. doi:10.1007/s00415-016-8290-1
- Irish M, Hodges JR, Piguet O, 2014 Right anterior temporal lobe dysfunction underlies theory of mind impairments in semantic dementia. *Brain* 137, 1241–1253. doi:10.1093/brain/awu003 [PubMed: 24523434]
- Johnson JO, Mandrioli J, Benatar M, Abramzon Y, Van Deerlin VM, Trojanowski JQ, Gibbs JR, Brunetti M, Gronka S, Wu J, Ding J, McCluskey L, Martinez-Lage M, Falcone D, Hernandez DG, Arepalli S, Chong S, Schymick JC, Rothstein J, Landi F, Wang Y-D, Calvo A, Mora G, Sabatelli M, Monsurro MR, Battistini S, Salvi F, Spataro R, Sola P, Borghero G, ITALSGEN Consortium, Galassi G., Scholz SW, Taylor JP, Restagno G, Chiò A, Traynor BJ, 2010 Exome sequencing reveals VCP mutations as a cause of familial ALS. *Neuron* 68, 857–864. doi:10.1016/j.neuron.2010.11.036 [PubMed: 21145000]
- Jones AR, Woollacott I, Shatunov A, Cooper-Knock J, Buchman V, Sproviero W, Smith B, Scott KM, Balendra R, Abel O, McGuffin P, Ellis CM, Shaw PJ, Morrison KE, Farmer A, Lewis CM, Leigh PN, Shaw CE, Powell JF, Al-Chalabi A, 2013 Residual association at C9orf72 suggests an alternative amyotrophic lateral sclerosis-causing hexanucleotide repeat. *Neurobiology of Aging* 34, 2234.e1–7. doi:10.1016/j.neurobiolaging.2013.03.003
- Kabashi E, Valdmanis PN, Dion P, Spiegelman D, McConkey BJ, Velde CV, Bouchard J-P, Lacomblez L, Pochigaeva K, Salachas F, Pradat P-F, Camu W, Meininger V, Dupre N, Rouleau GA, 2008 TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet* 40, 572–574. doi:10.1038/ng.132 [PubMed: 18372902]
- Karch CM, Wen N, Fan CC, Yokoyama JS, Kouri N, Ross OA, Höglinger G, Müller U, Ferrari R, Hardy J, Schellenberg GD, Sleiman PM, Momeni P, Hess CP, Miller BL, Sharma M, Van Deerlin V, Smeland OB, Andreassen OA, Dale AM, Desikan RS, International Frontotemporal Dementia (FTD)–Genomics Consortium, International Collaboration for Frontotemporal Dementia, Progressive Supranuclear Palsy (PSP) Genetics Consortium, and International Parkinson's Disease Genomics Consortium, 2018 Selective Genetic Overlap Between Amyotrophic Lateral Sclerosis and Diseases of the Frontotemporal Dementia Spectrum. *JAMA Neurol* doi:10.1001/jamaneurol.2018.0372
- Keil C, Prell T, Peschel T, Hartung V, Dengler R, Grosskreutz J, 2012 Longitudinal diffusion tensor imaging in amyotrophic lateral sclerosis. *BMC Neuroscience* 2012 13:1 13, 1. doi: 10.1186/1471-2202-13-141

- Kwan JY, Meoded A, Danielian LE, Wu T, Floeter MK, 2013 Structural imaging differences and longitudinal changes in primary lateral sclerosis and amyotrophic lateral sclerosis. *NeuroImage: Clinical* 2, 151–160. doi:10.1016/j.nicl.2012.12.003
- Kwiatkowski TJ, Bosco DA, Leclerc AL, Tamrazian E, Vanderburg CR, Russ C, Davis A, Gilchrist J, Kasarskis EJ, Munsat T, Valdmanis P, Rouleau GA, Hosler BA, Cortelli P, de Jong PJ, Yoshinaga Y, Haines JL, Pericak-Vance MA, Yan J, Ticozzi N, Siddique T, McKenna-Yasek D, Sapp PC, Horvitz HR, Landers JE, Brown RH, 2009 Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323, 1205–1208. doi:10.1126/science.1166066 [PubMed: 19251627]
- Landau SM, Mintun MA, Joshi AD, Koeppe RA, Petersen RC, Aisen PS, Weiner MW, Jagust WJ, 2012 Amyloid deposition, hypometabolism, and longitudinal cognitive decline. *Ann Neurol* 72, 578–586. doi:10.1002/ana.23650 [PubMed: 23109153]
- Li H, Achour I, Bastarache L, Berghout J, Gardeux V, Li J, Lee Y, Pesce L, Yang X, Ramos KS, Foster I, Denny JC, Moore JH, Lussier YA, 2016 Integrative genomics analyses unveil downstream biological effectors of disease-specific polymorphisms buried in intergenic regions. *NPJ Genom Med* 1, 10. doi:10.1038/npjgenmed.2016.6
- Lippa CF, Rosso AL, Stutzbach LD, Neumann M, Lee VMY, Trojanowski JQ, 2009 Transactive response DNA-binding protein 43 burden in familial Alzheimer disease and Down syndrome. *Arch Neurol* 66, 1483–1488. doi:10.1001/archneurol.2009.277 [PubMed: 20008652]
- Mackenzie IRA, Neumann M, Baborie A, Sampathu DM, Plessis Du D, Jaros E, Perry RH, Trojanowski JQ, Mann DMA, Lee VMY, 2011 A harmonized classification system for FTLTD-TDP pathology. *Acta Neuropathol* 122, 111–113. doi:10.1007/s00401-011-0845-8 [PubMed: 21644037]
- Massimo L, Powers C, Moore P, Vesely L, Avants B, Gee J, Libon DJ, Grossman M, 2009 Neuroanatomy of apathy and disinhibition in frontotemporal lobar degeneration. *Dement Geriatr Cogn Disord* 27, 96–104. doi:10.1159/000194658 [PubMed: 19158440]
- Massimo L, Xie SX, Rennert L, Fick DM, Halpin A, Placek K, Williams A, Rascovsky K, Irwin DJ, Grossman M, McMillan CT, 2018 Occupational attainment influences longitudinal decline in behavioral variant frontotemporal degeneration. *Brain Imaging Behav* 16, 1–9. doi:10.1007/s11682-018-9852-x
- Massimo L, Zee J, Xie SX, McMillan CT, Rascovsky K, Irwin DJ, Kolanowski A, Grossman M, 2015 Occupational attainment influences survival in autopsy-confirmed frontotemporal degeneration. *Neurology* 84, 2070–2075. doi:10.1212/WNL.0000000000001595 [PubMed: 25904687]
- McMillan CT, Lee EB, Jefferson-George K, Naj A, Van Deerlin VM, Trojanowski JQ, Wolk DA, 2018 Alzheimer's genetic risk is reduced in primary age-related tauopathy: a potential model of resistance? *Annals of Clinical and Translational Neurology* 5, 927–934. doi:10.1002/acn3.581 [PubMed: 30128317]
- McMillan CT, Toledo JB, Avants BB, Cook PA, Wood EM, Suh E, Irwin DJ, Powers J, Olm C, Elman L, McCluskey L, Schellenberg GD, Lee VMY, Trojanowski JQ, Van Deerlin VM, Grossman M, 2014 Genetic and neuroanatomic associations in sporadic frontotemporal lobar degeneration. *Neurobiology of Aging* 35, 1473–1482. doi:10.1016/j.neurobiolaging.2013.11.029 [PubMed: 24373676]
- Menke RAL, Körner S, Filippini N, Douaud G, Knight S, Talbot K, Turner MR, 2014 Widespread grey matter pathology dominates the longitudinal cerebral MRI and clinical landscape of amyotrophic lateral sclerosis. *Brain* 137, 2546–2555. doi:10.1093/brain/awu162 [PubMed: 24951638]
- Montuschi A, Iazzolino B, Calvo A, Moglia C, Lopiano L, Restagno G, Brunetti M, Ossola I, Presti Lo, A., Cammarosano S, Canosa A, Chio A, 2015 Cognitive correlates in amyotrophic lateral sclerosis: a population-based study in Italy. *Journal of Neurology, Neurosurgery & Psychiatry* 86, 168–173. doi:10.1136/jnnp-2013-307223
- Müller H-P, Turner MR, Grosskreutz J, Abrahams S, Bede P, Govind V, Prudlo J, Ludolph AC, Filippi M, Kassubek J, Neuroimaging Society in ALS (NiSALS) DTI Study Group, 2016 A large-scale multicentre cerebral diffusion tensor imaging study in amyotrophic lateral sclerosis. *Journal of Neurology, Neurosurgery & Psychiatry* jnnp-2015-311952. doi:10.1136/jnnp-2015-311952
- Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, Bruce J, Schuck T, Grossman M, Clark CM, McCluskey LF, Miller BL, Masliah E, Mackenzie IR, Feldman H, Feiden W, Kretschmar HA, Trojanowski JQ, Lee VMY, 2006 Ubiquitinated TDP-43 in frontotemporal

lobar degeneration and amyotrophic lateral sclerosis. *Science* 314, 130–133. doi:10.1126/science.1134108 [PubMed: 17023659]

- Placek K, Massimo L, Olm C, Ternes K, Firn K, Van Deerlin V, Lee EB, Trojanowski JQ, Lee VMY, Irwin D, Grossman M, McMillan CT, 2016 Cognitive reserve in frontotemporal degeneration. *Neurology* 87, 1813–1819. doi:10.1212/WNL.0000000000003250 [PubMed: 27683847]
- Premi E, Grassi M, van Swieten J, Galimberti D, Graff C, Masellis M, Tartaglia C, Tagliavini F, Rowe JB, Laforce R, Finger E, Frisoni GB, de Mendonça A, Sorbi S, Gazzina S, Cosseddu M, Archetti S, Gasparotti R, Manes M, Alberici A, Cardoso MJ, Bocchetta M, Cash DM, Ourselin S, Padovani A, Rohrer JD, Borroni B, Genetic FTD Initiative (GENFI), 2017 Cognitive reserve and TMEM106B genotype modulate brain damage in presymptomatic frontotemporal dementia: a GENFI study. *Brain*. doi:10.1093/brain/awx103
- Renton AE, Majounie E, Waite A, Simón-Sánchez J, Rollinson S, Gibbs JR, Schymick JC, Laaksovirta H, Van Swieten JC, Myllykangas L, Kalimo H, Paetau A, Abramzon Y, Remes AM, Kaganovich A, Scholz SW, Duckworth J, Ding J, Harmer DW, Hernandez DG, Johnson JO, Mok K, Ryten M, Trabzuni D, Guerreiro RJ, Orrell RW, Neal J, Murray A, Pearson J, Jansen IE, Sondervan D, Seelaar H, Blake D, Young K, Halliwell N, Callister JB, Toulson G, Richardson A, Gerhard A, Snowden J, Mann D, Neary D, Nalls MA, Peuralinna T, Jansson L, Isovita V-M, Kaivorinne A-L, Hölttä-Vuori M, Ikonen E, Sulkava R, Benatar M, Wu J, Chiò A, Restagno G, Borghero G, Sabatelli M, ITALSGEN Consortium, Heckerman D, Rogaeva E, Zinman L, Rothstein JD, Sendtner M, Drepper C, Eichler EE, Alkan C, Abdullaev Z, Pack SD, Dutra A, Pak E, Hardy J, Singleton A, Williams NM, Heutink P, Pickering-Brown S, Morris HR, Tienari PJ, Traynor BJ, 2011 A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72, 257–268. doi:10.1016/j.neuron.2011.09.010 [PubMed: 21944779]
- Ringholz GM, Appel SH, Bradshaw M, Cooke NA, Mosnik DM, Schulz PE, 2005 Prevalence and patterns of cognitive impairment in sporadic ALS. *Neurology* 65, 586–590. doi:10.1212/01.wnl.0000172911.39167.b6 [PubMed: 16116120]
- Rohrer JD, Warren JD, Modat M, Ridgway GR, Douiri A, Rossor MN, Ourselin S, Fox NC, 2009 Patterns of cortical thinning in the language variants of frontotemporal lobar degeneration. *Neurology* 72, 1562–1569. doi:10.1212/WNL.0b013e3181a4124e [PubMed: 19414722]
- Rosen DR, 1993 Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 364, 362. doi:10.1038/364362c0 [PubMed: 8332197]
- Rypma B, D'Esposito M, 1999 The roles of prefrontal brain regions in components of working memory: Effects of memory load and individual differences. *PNAS* 96, 6558–6563. doi:10.1073/pnas.96.11.6558 [PubMed: 10339627]
- Schuster C, Kasper E, Dyrba M, Machts J, Bittner D, Kaufmann J, Mitchell AJ, Benecke R, Teipel S, Vielhaber S, Prudlo J, 2014 Cortical thinning and its relation to cognition in amyotrophic lateral sclerosis. *Neurobiology of Aging* 35, 240–246. doi:10.1016/j.neurobiolaging.2013.07.020 [PubMed: 23992619]
- Senda J, Kato S, Kaga T, Ito M, Atsuta N, Nakamura T, Watanabe H, Tanaka F, Naganawa S, Sobue G, 2011 Progressive and widespread brain damage in ALS: MRI voxel-based morphometry and diffusion tensor imaging study 12, 59–69. doi:10.3109/17482968.2010.517850
- Smith SM, Nichols TE, 2009 Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *NeuroImage* 44, 83–98. doi:10.1016/j.neuroimage.2008.03.061 [PubMed: 18501637]
- Strong MJ, Abrahams S, Goldstein LH, Woolley S, McLaughlin P, Snowden J, Mioshi E, Roberts-South A, Benatar M, Hortobágyi T, Rosenfeld J, Silani V, Ince PG, Turner MR, 2017 Amyotrophic lateral sclerosis - frontotemporal spectrum disorder (ALS-FTSD): Revised diagnostic criteria. *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration* 1–22. doi:10.1080/21678421.2016.1267768
- Strong MJ, Grace GM, Freedman M, Lomen-Hoerth C, Woolley S, Goldstein LH, Murphy J, Shoesmith C, Rosenfeld J, Leigh PN, Bruijn L, Ince P, Figlewicz D, 2009 Consensus criteria for the diagnosis of frontotemporal cognitive and behavioural syndromes in amyotrophic lateral sclerosis 10, 131–146. doi:10.1080/17482960802654364
- Suh E, Lee EB, Neal D, Wood EM, Toledo JB, Rennert L, Irwin DJ, McMillan CT, Krock B, Elman LB, McCluskey LF, Grossman M, Xie SX, Trojanowski JQ, Deerlin VM, 2015 Semi-automated

quantification of C9orf72 expansion size reveals inverse correlation between hexanucleotide repeat number and disease duration in frontotemporal degeneration. *Acta Neuropathologica* 1–10. doi: 10.1007/s00401-015-1445-9

- Taylor JP, Brown RH, Cleveland DW, 2016 Decoding ALS: from genes to mechanism. *Nature* 539, 197–206. doi:10.1038/nature20413 [PubMed: 27830784]
- Toledo JB, Van Deerlin VM, Lee EB, Suh E, Baek Y, Robinson JL, Xie SX, McBride J, Wood EM, Schuck T, Irwin DJ, Gross RG, Hurtig H, McCluskey L, Elman L, Karlawish J, Schellenberg G, Chen-Plotkin A, Wolk D, Grossman M, Arnold SE, Shaw LM, Lee VMY, Trojanowski JQ, 2014 A platform for discovery: The University of Pennsylvania Integrated Neurodegenerative Disease Biobank. *Alzheimers Dement* 10, 477–84.e1. doi:10.1016/j.jalz.2013.06.003 [PubMed: 23978324]
- Tombaugh TN, Kozak J, Rees L, 1999 Normative data stratified by age and education for two measures of verbal fluency: FAS and animal naming. *Arch Clin Neuropsychol* 14, 167–177. [PubMed: 14590600]
- Tustison NJ, Avants BB, Cook PA, Zheng Y, Egan A, Yushkevich PA, Gee JC, 2010 N4ITK: Improved N3 Bias Correction. *IEEE Trans Med Imaging* 29, 1310–1320. doi:10.1109/TMI.2010.2046908 [PubMed: 20378467]
- Tustison NJ, Cook PA, Klein A, Song G, Das SR, Duda JT, Kandel BM, van Strien N, Stone JR, Gee JC, Avants BB, 2014 Large-scale evaluation of ANTs and FreeSurfer cortical thickness measurements. *NeuroImage* 99, 166–179. doi:10.1016/j.neuroimage.2014.05.044 [PubMed: 24879923]
- Van Deerlin VM, Sleiman PMA, Martinez-Lage M, Chen-Plotkin A, Wang L-S, Graff-Radford NR, Dickson DW, Rademakers R, Boeve BF, Grossman M, Arnold SE, Mann DMA, Pickering-Brown SM, Seelaar H, Heutink P, van Swieten JC, Murrell JR, Ghetti B, Spina S, Grafman J, Hodges J, Spillantini MG, Gilman S, Lieberman AP, Kaye JA, Woltjer RL, Bigio EH, Mesulam M, al-Sarraj S, Troakes C, Rosenberg RN, White CL, Ferrer I, Lladó A, Neumann M, Kretschmar HA, Hulette CM, Welsh-Bohmer KA, Miller BL, Alzualde A, de Munain AL, McKee AC, Gearing M, Levey AI, Lah JJ, Hardy J, Rohrer JD, Lashley T, Mackenzie IRA, Feldman HH, Hamilton RL, Dekosky ST, van der Zee J, Kumar-Singh S, Van Broeckhoven C, Mayeux R, Vonsattel JPG, Troncoso JC, Kiri J, Kwok JBJ, Halliday GM, Bird TD, Ince PG, Shaw PJ, Cairns NJ, Morris JC, McLean CA, DeCarli C, Ellis WG, Freeman SH, Frosch MP, Growdon JH, Perl DP, Sano M, Bennett DA, Schneider JA, Beach TG, Reiman EM, Woodruff BK, Cummings J, Vinters HV, Miller CA, Chui HC, Alafuzoff I, Hartikainen P, Seilhean D, Galasko D, Masliah E, Cotman CW, Tuñón MT, Martínez MCC, Munoz DG, Carroll SL, Marson D, Riederer PF, Bogdanovic N, Schellenberg GD, Hakonarson H, Trojanowski JQ, Lee VM-Y, 2010 Common variants at 7p21 are associated with frontotemporal lobar degeneration with TDP-43 inclusions 42, 234–239. doi:10.1038/ng.536
- van Es MA, Veldink JH, Saris CGJ, Blauw HM, van Vught PWJ, Birve A, Lemmens R, Schelhaas HJ, Groen EJN, Huisman MHB, van der Kooij AJ, de Visser M, Dahlberg C, Estrada K, Rivadeneira F, Hofman A, Zwarts MJ, van Doormaal PTC, Rujescu D, Strengman E, Giegling I, Muglia P, Tomik B, Slowik A, Uitterlinden AG, Hendrich C, Waibel S, Meyer T, Ludolph AC, Glass JD, Purcell S, Cichon S, Nöthen MM, Wichmann H-E, Schreiber S, Vermeulen SHM, Kiemeny LA, Wokke JHJ, Cronin S, McLaughlin RL, Hardiman O, Fumoto K, Pasterkamp RJ, Meininger V, Melki J, Leigh PN, Shaw CE, Landers JE, Al-Chalabi A, Brown RH, Robberecht W, Andersen PM, Ophoff RA, van den Berg LH, 2009 Genome-wide association study identifies 19p13.3 (UNC13A) and 9p21.2 as susceptibility loci for sporadic amyotrophic lateral sclerosis. 41, 1083–1087. doi: 10.1038/ng.442
- van Rheenen W, Shatunov A, Dekker AM, McLaughlin RL, Diekstra FP, Pulit SL, van der Spek RAA, Vösa U, de Jong S, Robinson MR, Yang J, Fogh I, van Doormaal PT, Tazelaar GHP, Koppers M, Blokhuis AM, Sproviero W, Jones AR, Kenna KP, van Eijk KR, Harschnitz O, Schellevis RD, Brands WJ, Medic J, Menelaou A, Vajda A, Ticozzi N, Lin K, Rogelj B, Vrabec K, Ravník-Glava M, Koritnik B, Zidar J, Leonardis L, Grošelj LD, Milicamps S, Salachas F, Meininger V, de Carvalho M, Pinto S, Mora JS, Rojas-García R, Polak M, Chandran S, Colville S, Swingle R, Morrison KE, Shaw PJ, Hardy J, Orrell RW, Pittman A, Sidle K, Pietro Fratta, Malaspina A, Topp S, Petri S, Abdulla S, Drepper C, Sendtner M, Meyer T, Ophoff RA, Staats KA, Wiedau-Pazos M, Lomen-Hoerth C, Van Deerlin VM, Trojanowski JQ, Elman L, McCluskey L, Basak AN, Tunca C, Hamzeiy H, Parman Y, Meitinger T, Lichtner P, Radivojkov-Blagojevic M, Andres CR, Maurel C, Bensimon G, Landwehrmeyer B, Brice A, Payan CAM, Saker-Delye S, Dürr A, Wood NW,

- Tittmann L, Lieb W, Franke A, Rietschel M, Cichon S, Nöthen MM, Amouyel P, Tzourio C, Dartigues J-F, Uitterlinden AG, Rivadeneira F, Estrada K, Hofman A, Curtis C, Blauw HM, van der Kooij AJ, de Visser M, Goris A, Weber M, Shaw CE, Smith BN, Pansarasa O, Cereda C, Del Bo R, Comi GP, D'Alfonso S, Bertolin C, Sorarù G, Mazzini L, Pensato V, Gellera C, Tiloca C, Ratti A, Calvo A, Moglia C, Brunetti M, Arcuti S, Capozzo R, Zecca C, Lunetta C, Penco S, Riva N, Padovani A, Filosto M, Muller B, Stuit RJ, Blair I, Zhang K, McCann EP, Fifita JA, Nicholson GA, Rowe DB, Pamphlett R, Kiernan MC, Grosskreutz J, Witte OW, Ringer T, Prell T, Stubendorff B, Kurth I, Hübner CA, Leigh PN, Casale F, Chiò A, Beghi E, Pupillo E, Tortelli R, Logroscino G, Powell J, Ludolph AC, Weishaupt JH, Robberecht W, Van Damme P, Franke L, Pers TH, Brown RH, Glass JD, Landers JE, Hardiman O, Andersen PM, Corcia P, Vourc'h P, Silani V, Wray NR, Visscher PM, de Bakker PIW, van Es MA, Pasterkamp RJ, Lewis CM, Breen G, Al-Chalabi A, van den Berg LH, Veldink JH, 2016 Genome-wide association analyses identify new risk variants and the genetic architecture of amyotrophic lateral sclerosis 48, 1043–1048. doi: 10.1038/ng.3622
- Vance C, Rogelj B, Hortobágyi T, De Vos KJ, Nishimura AL, Sreedharan J, Hu X, Smith B, Ruddy D, Wright P, Ganesalingam J, Williams KL, Tripathi V, Al-Saraj S, Al-Chalabi A, Leigh PN, Blair IP, Nicholson G, de Belleruche J, Gallo J-M, Miller CC, Shaw CE, 2009 Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323, 1208–1211. doi:10.1126/science.1165942 [PubMed: 19251628]
- Vass R, Ashbridge E, Geser F, Hu WT, Grossman M, Clay-Falcone D, Elman L, McCluskey L, Lee VMY, Van Deerlin VM, Trojanowski JQ, Chen-Plotkin AS, 2010 Risk genotypes at TMEM106B are associated with cognitive impairment in amyotrophic lateral sclerosis. *Acta Neuropathol* 121, 373–380. doi:10.1007/s00401-010-0782-y [PubMed: 21104415]
- Verstraete E, Veldink JH, Hendrikse J, Schelhaas HJ, van den Heuvel MP, van den Berg LH, 2012 Structural MRI reveals cortical thinning in amyotrophic lateral sclerosis. *Journal of Neurology, Neurosurgery & Psychiatry* 83, 383–388. doi:10.1136/jnnp-2011-300909
- Verstraete E, Veldink JH, van den Berg LH, van den Heuvel MP, 2014 Structural brain network imaging shows expanding disconnection of the motor system in amyotrophic lateral sclerosis. *Hum Brain Mapp* 35, 1351–1361. doi:10.1002/hbm.22258 [PubMed: 23450820]
- Vidal-Taboada JM, Lopez-Lopez A, Salvado M, Lorenzo L, Garcia C, Mahy N, Rodríguez MJ, Gamez J, 2015 UNC13A confers risk for sporadic ALS and influences survival in a Spanish cohort. *Journal of Neurology* 262, 2285–2292. doi:10.1007/s00415-015-7843-z [PubMed: 26162714]
- Wang D, Guo Y, Wrighton SA, Cooke GE, Sadee W, 2011 Intronic polymorphism in *CYP3A4* affects hepatic expression and response to statin drugs. *The Pharmacogenomics Journal* 11:4 11, 274–286. doi:10.1038/tpj.2010.28
- Wechsler D, 1945 A Standardized Memory Scale for Clinical Use. *The Journal of Psychology* 19, 87–95. doi:10.1080/00223980.1945.9917223
- Weintraub S, Salmon D, Mercaldo N, Ferris S, Graff-Radford NR, Chui H, Cummings J, DeCarli C, Foster NL, Galasko D, Peskind E, Dietrich W, Beekly DL, Kukull WA, Morris JC, 2009 The Alzheimer's Disease Centers' Uniform Data Set (UDS): The Neuropsychological Test Battery. *Alzheimer Dis Assoc Disord* 23, 91–101. doi:10.1097/WAD.0b013e318191c7dd [PubMed: 19474567]
- Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE, 2014 Permutation inference for the general linear model. *NeuroImage* 92, 381–397. doi:10.1016/j.neuroimage.2014.01.060 [PubMed: 24530839]
- Wood EM, Falcone D, Suh E, Irwin DJ, Chen-Plotkin AS, Lee EB, Xie SX, Van Deerlin VM, Grossman M, 2013 Development and validation of pedigree classification criteria for frontotemporal lobar degeneration. *JAMA Neurol* 70, 1411–1417. doi:10.1001/jamaneurol.2013.3956 [PubMed: 24081456]
- Zhang M, Zhao L-J, Zhou Y, Badr R, Watson P, Ye A, Zhou B, Zhang J, Deng H-W, Recker RR, Lappe JM, 2017 SNP rs11185644 of RXRA gene is identified for dose-response variability to vitamin D3 supplementation: a randomized clinical trial. *Sci Rep* 7, 40593. doi:10.1038/srep40593 [PubMed: 28079136]

Highlights

- rs12608932 in *UNC13A* may modify frontotemporal disease in sporadic ALS
- minor allele associated with reduced *ante mortem* frontotemporal cortical thickness
- minor allele associated with greater frontotemporal TDP-43 pathology at autopsy

Figure 1. Reduced cortical thickness associated with rs12608932 genotype.

In an analysis restricted to regions of reduced cortical thickness identified in sporadic ALS (N=109) relative to healthy controls (N= 113) (light blue regions), patients with sporadic ALS who are carriers of the minor allele (C) exhibited greater reduction of cortical thickness in dorsal prefrontal, ventromedial prefrontal, anterior temporal, and middle temporal cortices (regions indicated in red-yellow heatmap). The heatmap indicates the associated T-statistic for each voxel, with yellow representing the highest value.

Figure 2. Greater burden of pTDP-43 pathology associated with rs12608932 genotype. Box and whisker plots of pTDP-43 pathologic burden in our sporadic ALS autopsy cohort (N=102) show that minor allele homozygotes (CC) and heterozygotes (AC) feature greater burden of pathology in the middle frontal cortex. Minor allele homozygotes but not minor allele heterozygotes also feature greater burden of pathology in the superior/middle temporal cortex and motor cortex.

Table 1.

Demographic, clinical, and genetic information for neuroimaging and ADNI cohorts (Abbreviations: Alzheimer's Disease Neuroimaging Initiative – ADNI; ALS Functional Rating Scale-Revised - ALSFRS-R; Mini-Mental State Exam – MMSE)

	ALS	Controls	ADNI Controls
N (F)	109 (44)	113 (52)	84
Age, Y	59.54 (10.96)	61.54 (8.73)	75.44 (6.40)
Education, Y	15.26 (2.99)	15.35 (2.43)	-
Race, N			-
White	100	90	
Black	6	22	
Multi-racial	1	1	
Other	1	-	
Unknown	1	-	
Ethnicity, N			-
Latino	2	1	
Non-Latino	105	111	
Other	1	-	
Unknown	1	1	
Disease duration, M	38.67 (34.25)	-	-
ALSFRS-R	33.81 (7.39)	-	-
Forward digit span, % impaired	6.53 (1.45), 4.54	-	-
Reverse digit span, % impaired	4.21 (1.70), 13.63	-	-
Letter fluency, % impaired	11.29 (5.41), 23.86	-	-
Visual Verbal Test, % impaired	7.8 (4.13), 9.09	-	-
MMSE, % correct, % impaired	91.82 (14.11), 25.00	-	-
rs12608932 genotypes, N			
AA	43	-	41
AC	46	-	34
CC	20	-	9

Table 2.

Demographic, clinical, and genetic information for autopsy cohort.

	ALS
N (F)	102 (40)
Age at Death, Y	65.01 (10.83)
Race, N	
White	71
Black	3
Multi-racial	-
Other	1
Unknown	27
Ethnicity, N	
Latino	2
Non-Latino	72
Other	-
Unknown	28
Education, Y	15.18 (3.48)
Disease duration at Death, Y	4.83(4.98)
rs12608932 genotypes, N	
AA	38
AC	47
CC	17

Table 3.

Regions of reduced cortical thickness for ALS relative to controls and associated with rs12508932 genotype in ALS.

Neuroanatomic region (BA)	L / R	MNI coordinates			T statistic	p value	Voxels
		x	y	z			
Reduced cortical thickness in ALS relative to controls ¹ :							25627
Insular Cortex (13)	R	40	22	0	6.72	0.0006	
Inferior Frontal Gyrus (45)	L	−32	28	2	6.18	0.0001	
Dorsolateral Prefrontal Cortex (9)	-	0	46	18	6.12	0.0001	
Inferior Frontal Gyrus (45)	R	34	22	8	5.96	0.0007	
Ventral Medial Orbital Frontal Cortex (11)	R	12	14	−20	5.91	0.0001	
Ventral Medial Orbital Frontal Cortex (11)	L	−12	14	−18	5.79	0.0001	
Middle Temporal Gyrus (21)	R	64	−32	10	3.54	0.004	
Reduced cortical thickness associated with rs12608932 minor allele in ALS:							
Anterior Temporal Lobe (38)	R	34	10	−40	3.08	0.002	56
Ventromedial Prefrontal Cortex (11)	L	−20	26	−24	3.21	0.001	27
Middle Temporal Gyrus (22)	L	−50	−10	−12	2.88	0.003	24
Dorsal Prefrontal Cortex (10)	L	−28	44	28	3.56	0.001	23
Ventromedial Prefrontal Cortex (11)	R	16	16	−24	3.66	0.001	21

¹Note. Cortical regions identified from peak voxel coordinates in an effort to describe sub-peaks within a larger, contiguous cluster.