Molecular and cellular characterization of nAChR subtypes in the arcuate nucleus of the mouse hypothalamus

Cali A. Calarco¹, Zhiying Li², Seth R. Taylor¹, Somin Lee¹, Wenliang Zhou¹, Jeffrey M. Friedman², Yann S. Mineur¹, Cecilia Gotti³, and Marina R. Picciotto¹

¹Department of Psychiatry and Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven, CT 06508, USA

²The Rockefeller University

³CNR, Institute of Neuroscience Milan

Abstract

Nicotine, acting through nicotinic acetylcholine receptors (nAChRs), increases the firing rate of both orexigenic agouti-related peptide (AgRP) and anorexigenic pro-opiomelanocortin (POMC) neurons in the arcuate nucleus of the hypothalamus (ARC), yet nicotine and other nAChR agonists decrease food intake in mice. Viral-mediated knockdown of $\beta 4$ nAChR subunit in all neuronal cell types in the ARC prevents the nicotinic agonist cytisine from decreasing food intake, but it is not known whether the $\beta 4$ subunit is selectively expressed in anorexigenic neurons or how other nAChR subtypes are distributed in this nucleus. Using translating ribosome affinity purification (TRAP) on ARC tissue from mice with ribosomes tagged in either AgRP or POMC cells, we examined nAChR subunit mRNA levels using real-time PCR. Both AgRP and POMC cells express a comparable panel of nAChR subunits with differences in $\alpha 7$ mRNA levels and a trend for difference in $\alpha 4$ levels, but no differences in $\beta 4$ expression. Immunoprecipitation of assembled nAChRs revealed that the $\beta 4$ subunit forms assembled channels with $\alpha 3$, $\beta 2$ and $\alpha 4$, but not other subunits found in the ARC. Finally, using cell type-selective, virally-delivered small hairpin RNAs targeting either the $\beta 4$ or $\alpha 7$ subunit, we examined the contribution of each subunit in either AgRP or POMC cells to the behavioral response to nicotine, refining the understanding of nicotinic

Correspondence should be addressed to: Marina R. Picciotto, Dept. of Psychiatry, Yale University School of Medicine, 34 Park Street – 3rd floor research, New Haven, CT 06508, Phone: 203-737-2041; Fax: 203-737-2043; marina.picciotto@yale.edu. DR. MARINA PICCIOTTO (Orcid ID : 0000-0002-4404-1280)

Conflict of Interest Statement

The authors declare that there are no conflicts of interest.

Author Contributions

C.C. wrote the manuscript, performed all PCR experiments beyond initial sample validation, collected tissue for immunoprecipitations, validated the $\alpha 7$ viruses, and performed the behavior with the cell type selective viruses. Z.L. oversaw breeding of AgRP and POMC bacTRAP mice, collected tissue, and performed TRAP immunoprecipitations. Z.L. also created cDNA from these samples and validated the quality of samples. TRAP experiments were run in the lab of and with the support of J.F. S.L. developed the acute feeding paradigm, collected the wild-type feeding data, and aided in data collection for the AgRP-Cre and POMC-Cre behavioral data. S.L. also validated some of the PCR primers. C.G. supervised and ran all immunoprecipitation experiments including hypothalamic, arcuate, and $\beta 2$ knockout tissue. S.T. designed and validated the $\beta 4$ knockdown viruses and the SCR virus. W.Z. with Y.M. designed the $\alpha 7$ knockdown viruses. Y.M. contributed to project design, purified all the viruses used, collected tissue for immunoprecipitations, and provided general insight into the experiments. M.R.P. contributed to study design, interpretation and analyses as well as manuscript writing and editing.

Data Accessibility Statement

Raw data will be shared with the research community upon request.

regulation of this feeding circuit. These experiments identify a more complex set of nAChRs expressed in ARC than in other hypothalamic regions. Thus, ARC appears to be a particular target of nicotinic modulation.

Graphical Abstract

Nicotinic acetylcholine receptor (nAChR) signaling on both anorexogenic and orexogenic neurons in the arcuate nucleus of the hypothalamus (ARC) regulates the effects of nicotine on food intake. nAChR subunit expression, is not segregated by cell type, the $\beta 4$ subunit assembles into nAChRs with a different composition than in other $\beta 4$ containing brain regions, and cell type selective knockdown of the $\beta 4$ subunit in either anorexogenic and orexogenic cells in the ARC blocks the reduction in feeding due to nicotine.

Keywords

ARC; nicotine; food intake; nAChR subunits; AgRP; POMC

Introduction

Despite the well documented and well known health risks of cigarette smoking, many smokers report that they continue to smoke in part because of nicotine's association with appetite suppression and leaner body mass (Albanes *et al.*, 1987; Klesges *et al.*, 1997; Chiolero *et al.*, 2008). Like smoking in humans, nicotine reduces food intake and weight gain in rodent models (Grunberg *et al.*, 1986; Winders & Grunberg, 1990; Frankish *et al.*, 1995; Mineur *et al.*, 2011). In both the brain and the periphery nicotine acts on nicotinic acetylcholine receptors (nAChR), cation channels made up of a combination α - and β -type subunits arranged as a pentamer around a central pore (Albuquerque *et al.*, 2009; Morales-Perez *et al.*, 2016). The identity of the individual subunits and the stoichiometry of α and β subunits within an assembled nAChR determine many channel properties, including sensitivity to both endogenous and exogenous ligands, kinetics of ligand binding, the amplitude and identity of ionic currents that flow through the channel, and channel open time (Stokes & Papke, 2012; Lucero *et al.*, 2015; Mazzaferro *et al.*, 2016; George *et al.*, 2017). In addition, each nicotinic subunit is expressed in distinct, though overlapping,

patterns in many cell types throughout the brain (Sargent, 1993). Accordingly, specific nAChR subunits play dissociable roles in different behavioral paradigms. For example, mice lacking the $\beta 2$ subunit demonstrate abnormal avoidance learning and will not self-administer nicotine, unlike their wild type siblings (Picciotto *et al.*, 1995, 1998), $\beta 4$ knockout mice show greatly decreased symptoms of nicotine withdrawal compared to wild type mice (Salas *et al.*, 2004), and knockdown of the $\alpha 5$ subunit induces higher levels of nicotine self-administration, likely by reducing aversive consequences of nicotine intake (Fowler *et al.*, 2011).

Nicotine affects food intake and fat storage through actions of nAChRs at many levels in the brain and periphery (Jo *et al.*, 2002; Zoli & Picciotto, 2012; Stojakovic *et al.*, 2017). The $\alpha 3\beta 4^*$ nAChR full agonist cytosine, which also acts as a low affinity agonist of $\alpha 7$ nAChRs and a partial agonist of $\alpha 4\beta 2^*$ nAChRs (Papke & Heinemann, 1994; Stokes & Papke, 2012; Peng *et al.*, 2013), acutely reduces food intake. (* notation indicates the presence of variable additional subunits). This effect is abolished by viral-mediated knockdown of the $\beta 4$ nAChR subunit in the arcuate nucleus of the hypothalamus (ARC) (Mineur *et al.*, 2011), a brain area known to affect appetite. The ARC is a central node in the hypothalamus regulating food intake and its two main cell populations, agouti-related peptide (AgRP) and proopiomelanocortin (POMC) expressing neurons, induce hunger and satiety, respectively (Aponte *et al.*, 2011; Krashes *et al.*, 2011; Cansell *et al.*, 2012; Zhan *et al.*, 2013; Sternson & Atasoy, 2014; Kim *et al.*, 2015; Fenselau *et al.*, 2016). The reduction in food intake and weight gain due to $\beta 4$ agonism have been supported by additional studies using cytosine (Greibenstein *et al.*, 2013) and another $\beta 4$ agonist, levamisole (Lewis *et al.*, 2017). Knockout of POMC or viral knockdown of melanocortin receptors in the paraventricular nucleus also diminishes the ability of cytosine to decrease food intake (Mineur *et al.*, 2011), suggesting that activation of $\beta 4$ -containing nAChRs in response to nicotinic agonists leads to activation of the melanocortin pathway, signaling satiety and thus yielding reduced food intake and long term reductions in weight gain. One possibility is that $\beta 4$ nAChR subunits could be selectively or preferentially expressed on POMC neurons, regulating the increase in activity of POMC neurons in response to cytosine and nicotine (Huang *et al.*, 2011; Mineur *et al.*, 2011; Rubinstein & Low, 2011). However $\beta 4$ nAChR subunit expression on other cell types cannot yet be ruled out as the nAChR topography of the different cell-types in ARC has not yet been examined. Further, as cytosine is also a full agonist at $\alpha 7$ receptors with a lower EC_{50} than for $\beta 4^*$ receptors (Papke & Porter Papke, 2002; Stokes & Papke, 2012), these receptors could be playing a synergistic role in controlling food intake, and the role of this receptor class has not been examined specifically in the ARC. Finally, signaling through either $\alpha 7$ or $\beta 4^*$ containing nAChRs may regulate the feeding response to the broader agonist nicotine.

The current study was designed to characterize nAChR subunit expression in the ARC comprehensively, with a specific focus on how nAChRs on AgRP and POMC neurons might contribute to reductions in food intake in response to nicotinic agonists, including nicotine and cytosine. We used a genetic method to perform translating ribosome affinity purification (bacTRAP) to isolate actively translated mRNAs from AgRP or POMC cells, to analyze enrichment of nAChR subunit translation in either cell type compared to the hypothalamus in general, as well as to compare relative subunit mRNA translation between AgRP and

POMC cell types directly. To identify which nAChR subunits form assembled pentameric receptors in the hypothalamus overall and in ARC specifically, we used subunit-selective antibodies to immunoprecipitate nAChR subtypes, and further sequential immunoprecipitation to determine which subunits assemble with the $\beta 4$ subunit to form nAChRs. We determined the expression of the cholinergic cell markers choline acetyltransferase (ChAT) and vesicular acetylcholine transporter (VACHT) in AgRP and POMC cells to determine whether neurons within the ARC show a cholinergic phenotype and may be engaged in local acetylcholine (ACh) release. Finally, with the use of novel cell-type-selective knockdown viruses targeting the $\beta 4$ or $\alpha 7$ nAChR subunits, we determined whether expression of these subunits in AgRP or POMC cells affected the acute feeding response to nicotine and cytosine.

Methods

Animals

For bacTRAP experiments, AgRP and POMC-eGFP-RPL10a transgenic mice were made in the same way as Pdyn-eGFP-RPL10a mice (Allison *et al.*, 2015), except that pronuclear injection was done with C57BL/6J embryos. Briefly, eGFP-RPL10a as a fusion gene was cloned immediately downstream of the ATG of the AgRP and POMC genomic sequence in the BAC backbone (RP24-297L5 and RP23-168O9 respectively). Resulting N0 pups were bred to C57BL/6J to produce N1 pups which were screened for eutopic transgene expression. Selected founders were maintained on C57BL/6J background and inter-crossed to produce pups for the study.

For immunoprecipitation experiments and immunohistochemistry adult male C57BL/6J (>8wks) mice were purchased from Jackson Laboratories (Bar Harbor, ME) and group housed. Hypothalamus samples for IPs, $\beta 2$ KO mice for IPs were a generous gift of Dr. Michael J Marks (Institute for Behavioral Genetics, University of Colorado, Boulder, USA). For viral mediated knockdown validation experiments, GAD65-IRES-Cre mice (Jackson Labs Bar Harbor, ME; Stock #010802) and ChAT-IRES-Cre mice (Jackson Labs strain B6;129S6-Chattm2(cre)Lowl/J, Stock:006410; Bar Harbor, ME) were bred in house to C57BL/6J mice. AgRP-IRES-Cre mice ((AgRptm1-cre)Lowl, Jax #012899) and POMC-Cre (Tg (Pomc1-cre)16Lowl/J) mice were bred in house to C57BL/6J mice, and used for behavioral experiments. AgRP-Cre wild-type littermates were used for pharmacological feeding tests.

All animals were maintained in a temperature controlled vivarium (21–23°C) on a 12 hour light-dark cycle (lights on 7:00 AM) and housed 5 per cage with food and water available *ad libitum*, unless otherwise specified. All experiments were performed according to protocols approved by the Yale University, Rockefeller University and/or University of Colorado, Boulder Animal Care and Use Committees.

Drugs

All drug solutions were prepared in sterile phosphate buffered saline and administered intraperitoneally (IP). Nicotine hydrogen tartrate salt was provided through the NIDA drug

supply program. All nicotine doses are reported as free base (Matta *et al.*, 2007). Cytisine was purchased from Sigma Aldrich (St. Louis, MO).

Tissue Collection

BacTRAP—10 adult bacTRAP mice (>8wks) of the same sex were pooled for each sample. Ten AgRP-TRAP and 10 POMC-TRAP samples (5 male and 5 female for each) were collected from EGFPL10a transgenic mice, and tissue was dissected and processed as previously described (Heiman *et al.*, 2008, 2014). Briefly, mice were rapidly decapitated and the hypothalami of 10 bacTRAP transgenic mice of the same line and sex were manually dissected in dissection buffer (2.5 mM HEPES-KOH [pH 7.4], 35 mM Glucose, 4 mM NaHCO₃ in HBSS). Pooled tissue was immediately homogenized in a Teflon-glass homogenizer (300 rpm, 6 strokes, 900 rpm, 6 strokes) with ice-cold homogenization buffer (10 mM HEPES-KOH [pH 7.4], 150 mM KCl, 5 mM MgCl₂, 0.5 mM dithiothreitol, protease inhibitors, Rnasin, and 100 µg/mL cycloheximide). 20 µL of homogenate was saved at this stage to be used for total RNA analysis. The remaining homogenate was centrifuged for 10 min (2,000 × g) at 4°C. NP-40 (EMD Biosciences, San Diego, CA) and 1,2-Diheptanoyl-sn-Glycero-3-Phosphocholine (DHPC; Avanti Polar Lipids, Alabaster, AL) were added to the supernatant at a final concentration of 1% and 30 mM, respectively. Samples were incubated on ice for 5 min then centrifuged at 20,000 × g for 15 min.

The resulting supernatant was transferred to mouse anti-GFP (custom)-coated MyOne magnetic Dynabeads (Invitrogen) and incubated on a slow rotator at 4°C for 16–18hrs. Dynabeads were washed three times with wash buffer (10 mM HEPES-KOH [pH 7.4], 5 mM MgCl₂, 350 mM KCl, 1% NP-40, 0.5 mM dithiothreitol, Rnasin, 100 mg/ml cycloheximide, RNase inhibitors, 3% BSA and 500 ng/ml poly[d(I-C)]). Beads were resuspended in 100 µl lysis buffer (Stratagene Absolutely RNA NanoRNA kit) or RLT buffer (Qiagen RNeasy kit) with β-ME added, vortexed, and incubated at RT for 10 min. Beads were separated on a magnet and the supernatant was collected. 200 ng poly[d(I-C)] was added to the collected supernatant. RNA was quantified and checked for quality on a Bioanalyzer 2100 using RNA Picochip (Agilent Technologies, Santa Clara, CA). Remaining RNA was frozen at –80°C until it was converted to cDNA.

Immunoprecipitation and Qualitative PCR—Adult C57BL/6J mice were sacrificed via rapid decapitation and brains were removed. For immunoprecipitation experiments using the whole hypothalamus, each experiment was conducted on the pooled tissue from 10 mice (3–4 replicate experiments). For specific ARC dissections, brains were removed after rapid decapitation, the hypothalamic disc was removed from the ventral surface of the brain, and a 1.2 mm punch surrounding the 3rd ventricle containing the ARC was taken and immediately frozen on dry ice (Supplemental Fig. 1 for diagram). For these dissections, ARC punches were pooled before immunoprecipitation, resulting in tissue from 20 mice being used for each of two immunoprecipitation replicates. For PCR each animal was one biological replicate, two animals total. After collection, tissue was stored at –80°C until it was used for immunoprecipitation or for mRNA extraction.

cDNA Preparation

For qualitative PCR, TRAP, and viral validation experiments RNA was extracted from tissue using an RNeasy Lipid Tissue Mini Kit following the manufacturer's protocol (Qiagen, Valencia, CA). The concentration of extracted mRNA was measured with a spectrophotometer (NanoDrop ND-1000; Thermo Scientific, Waltham, MA). RNA extracted using the RNeasy kit or isolated via the TRAP protocol was converted to double stranded cDNA using Qiagen Quantitech Reverse Transcription Kit according to the manufacturer's instructions. cDNA was generated using the maximum possible RNA for qualitative PCR (1.2 µg), bacTRAP input and bound samples (4–89 ng), 20 ng RNA for medial habenula (MHb) samples, and 125 ng RNA for GAD65-Cre vHipp samples. Converted cDNA was stored at –20°C or –80°C until PCR analysis.

Qualitative PCR

ARC cDNA was diluted 1:20 and 3 µl of this working solution was used in a 20 µL PCR reaction mixture using Taq polymerase (Roche) and primers for the α2–7 and β2–4 subunits listed in Table 1. PCR mixtures were incubated in a DNAEngine Peltier Thermal Cycler (Bio-Rad) under the following conditions: 95°C 1 min, (95°C 30 sec, 58°C 30 sec, 72°C 30 sec) × 40 cycles, 72°C 5 min. PCR products were run via electrophoresis on a 1.5% agarose gel containing ethidium bromide at 150 V for 45 min. Gels were visualized with a UV box and images were taken using Foto Analyst PC Image software (Fotodyne Inc, Hartland, WI). One representative band is shown for each subunit.

Real Time qPCR

Data collection—For bacTRAP samples, extraction quality was assessed by real-time qPCR to ensure enrichment of GFP and AgRP or POMC depending on the mouse line. Taqman reagents, primers: TaqMan™ Fast Advanced Master Mix, AgRP Taqman Assay (Mm00475829_g1, Cat# 4453320), POMC Taqman Assay (Mm00435874_m1, Cat# 4453320), GFP Taqman Assay (Mr03989638_mr, Cat# 4448892), HPRT Taqman Assay (Mm03024075_m1, Cat# 4453320). For all other PCR reactions relative expression of target genes was measured on a 96-well StepOne Plus qPCR machine with Power SYBR Green Mix (Applied Biosystems, Foster City, CA). All primers were designed to yield only one PCR product, had ~100% efficiency in a standard curve, and resulted in single melt peaks (see Table 1 for primer pair sequences and notation of which primers were used for which samples). All primers were ordered from Integrated DNA Technologies and HPRT primers were part of a commercially available assay (Mm.PT.39a.22214828). PCR reactions were run in duplicate and were averaged to yield one Ct value per animal. If duplicate samples did not run within one Ct value of each other that sample was excluded from analysis for that particular gene, resulting in n as noted on figures.

Analysis—For bacTRAP samples, fold change, was calculated using the 2^{-Ct} method (Goni *et al.*, 2009) with HPRT used as the reference gene. To determine fold enrichment, the total hypothalamic homogenate was used as a reference group to which the TRAP purified samples were compared. To determine differences in expression between AgRP and POMC cells, the POMC group was used as the reference group to which AgRP samples were

compared. The resulting fold-change measures were graphed and used for statistical analysis between groups. Individual t-tests were run for each gene analyzed.

Efficiency of $\beta 4$ knockdown was quantified using the 2^{-Ct} method with *tbp1* as the reference gene, and Cre- mice as the reference group. For analysis of $\alpha 7$ knockdown, the 2^{-Ct} method was used with *HPRT* as the reference gene, and AAV-dsRed-Sico-Scr shRNA as the reference group. The resulting fold-change measure was graphed and used for statistical analysis.

Immunoprecipitation

Antibody production and characterization—Affinity-purified, subunit-specific, polyclonal antibodies were used for the detection of nAChR subunits. The production of antibodies respected the guidelines established by the Italian Council on Animal Care, and was approved by Italian Government Decree No. 2/2010.

Subunit-specific polyclonal antibodies were produced in rabbits against peptides derived from or intracytoplasmic loop (CYT) regions of the $\alpha 2$, $-\alpha 3$, $-\alpha 4$, $-\alpha 5$, $\alpha 6$, $-\beta 2$, $-\beta 3$, $\beta 4$ rodent (mouse or rat) or human subunit sequences, and affinity purified and characterized as previously described (Gotti *et al.*, 2008; Grady *et al.*, 2009). For the $\alpha 3$ and $\beta 4$ subunits, polyclonal antibodies directed against the rodent COOH $\alpha 3$ (CQPLMARDDT) and the rodent $\beta 4$ (CGLPPLFQIHAPSKDS) peptides were also used.

Tissue homogenates and Triton X-100 extracts—The dissected frozen whole hypothalami or ARC nuclei were homogenized using a glass potter and washed in 50 mM Tris HCl pH 7, 120 mM NaCl, 5 mM KCl, 1 mM $MgCl_2$, 2.5 mM $CaCl_2$ and 2 mM PMSF by centrifugation twice. The resulting pellets were suspended in the same buffer containing a mixture of 20 $\mu g/ml$ of each of the following protease inhibitors: leupeptin, bestatin, pepstatin A and aprotinin. Triton X-100 at a final concentration of 2% was added to the washed membranes and extraction was performed for 2 hours at 4 °C as previously described (Gotti *et al.*, 2008).

Protein content of the membranes and 2% Triton X-100 extracts was measured using the bicinchonic acid protein assay (Pierce, Rockford, IL, USA) with bovine serum albumin as the standard.

[3H]-Epibatidine binding studies—[3H]-Epibatidine binds to heteromeric nAChR with high affinity (pM), and to homomeric $\alpha 7$ nAChRs with lower affinity (nM). In order to ensure that the $\alpha 7$ nAChRs did not contribute to [3H]-Epibatidine binding to solubilized receptors (present in the extract and immunoprecipitation experiments) the binding was performed in the presence of 1 μM α -Bungarotoxin, which specifically binds to $\alpha 7$ -nAChR (and thus prevents [3H]-Epibatidine binding to these sites). Triton X-100 extracts were incubated with cold α -bungarotoxin for two hours and binding experiments were carried out by incubating 50 μl of the extracts overnight with 1 nM [3H]-Epibatidine at 4°C. Following overnight incubation, nAChRs were captured using DEAE-Sepharose™ Fast flow (GE Healthcare, Uppsala, Sweden). Bound receptors were eluted with 1N NaOH and counted after addition of the scintillation mixture (filter count, GE Healthcare, Uppsala, Sweden) in a

Tri-Carb 2100TR β -counter (Perkin Elmer) counter. Non-specific binding (averaging 5–10% of total binding) was determined in parallel samples containing 100 nM unlabeled epibatidine.

Immunoprecipitation of [3 H]-Epibatidine-labelled nAChRs by subunit-specific antibodies

—Extracts (100 μ l), were incubated with 1 μ M α -bungarotoxin followed by incubation with 1 nM [3 H]-Epibatidine, then incubated overnight with a saturating concentration (10 μ g) of affinity purified anti-subunit IgG (anti- α 2, - α 3, - α 4, - α 5, - α 6, - β 2, - β 3, - β 4) or control IgG bound to Sepharose-ProteinA (GE). Immunoprecipitates were recovered by centrifugation (5 min at 2,400 $\times$ g).

[3 H]- Epibatidine binds to assembled heteromeric nAChRs, but it does not bind to subunits that are not associated with a subunit partner. Therefore, once the presence of a particular nAChR subunit is established with an antibody in the first immunoprecipitation assay, the subunit(s) it is associated with can be determined by performing a second immunoprecipitation on the resultant supernatant with a different antibody. The rationale for this sequential procedure is that if two (or more) different subunits are part of the same receptor, then initial immunoprecipitation of that receptor with an antibody directed at one subunit will decrease the amount of the receptor available in the remaining supernatant of the sample for immunoprecipitation with a subsequent antibody.

For the sequential immunoprecipitation assays, a fixed amount of the clear [3 H]-Epibatidine labeled supernatant remaining after immunoprecipitation with the first antibody (anti- α 2, - α 3, - α 4, - α 5, - α 6, - β 2, - β 3, - β 4) was incubated with a saturating concentration (10 μ g) of anti- β 4 antibody bound to protein A and the immunoprecipitation steps then repeated, as described above. In control studies, the number of solubilized nAChRs measured with [3 H]-Epibatidine was stable when incubated in the presence of control IgG over the time course of the sequential immunoprecipitation procedure.

Plasmid Design and Viral Packaging

Short-hairpin RNAs targeting the β 4 and α 7 nAChR subunits were designed using siRNA Wizard (InvivoGen, www.invivogen.com; San Diego, CA) (previously described in Mineur et al., 2015, 2013, 2011) and ligated into a pAAV-dsRed-Sico plasmid designed to restrict shRNA expression to cells that express Cre-recombinase (described in Wohleb *et al.*, 2016). Target sequences used were (5'-3'): β 4 RNAi1: GAGGAGAAGCTGATGGATGATCTC; β 4 RNAi 2: GTTGTACAACAATGCCGATGGGAC; α 7 RNAi 1: GGCTGTACAAGGAGCTGGTCAAGA; α 7 RNAi 2: AGCCCTGGCTTTGCTGGTATTCTT; α 7 RNAi 3: GCCAGGATGAGAGTGAGGTGATCT. A scrambled RNAi sequence not targeting any murine mRNAs (SCR) was also used: GCCAGATTTCTCAGGTGATAA. For β 4shRNAs, single-stranded oligonucleotides featuring a T, shRNA sense, 9 base pair hairpin (TTCAAGAGA), shRNA antisense, and a 3' overhang AGCT for XhoI cloning were ordered (Integrated DNA Technologies) and annealed. For α 7 shRNAs single stranded oligonucleotides featuring the shRNA sense, 10-basepair hairpin (CTTCCTGTCA), shRNA antisense, and 3' overhang ATTTTT (shRNA 1 and 3) or GTTTTT (shRNA 2) were ordered

(Integrated DNA Technologies) and annealed. The annealed oligonucleotides were ligated into pAAV-dsRed-Sico using HpaI and AbsI (for SCR shRNA), HpaI and XhoI ($\beta 4$ and $\alpha 7$ shRNAs). Proper ligation was visualized by PCR amplification over the ligation region using the forward primer (ACATGGTCCTGCTGGAGTTC) and reverse primer (CCAGCTTGGTTCCCAATAGA). The same primers were used to sequence the plasmid to ensure proper ligation.

After ligation of shRNAs into pAAV-dsRed-Sico, plasmids were packaged into adeno-associated viruses (AAV2) as previously described (Mineur *et al.*, 2011, 2013, 2016). This resulted in 100–200 μ l of each purified virus, which were stored at 4° C.

Stereotactic Viral Infusions

For validation of cre-specific $\beta 4$ viral knockdown, a cocktail of AAV-dsRed-Sico- $\beta 4$ 1 and 2 (1:1) was infused bilaterally into the MHb of adult (>8wks) ChAT-IRES-Cre mice or wild type littermates. For validation of Cre-specific $\alpha 7$ viral knockdown a cocktail of AAV-dsRed-Sico- $\alpha 7$ shRNA1, 2 and 3 (1:1:1) or AAV-dsRed-Sico-SCR shRNA were infused bilaterally into the ventral hippocampus (vHipp) of GAD65-Cre mice. For behavioral feeding experiments a cocktail of AAV-dsRed-Sico- $\beta 4$ shRNA1 and 2 (1:1), a cocktail of AAV-dsRed-Sico- $\alpha 7$ shRNA1, 2 and 3 (1:1:1) or AAV-dsRed-Sico-SCR were infused bilaterally into the ARC of adult (>8wks) POMC-Cre or AgRP-Cre mice.

Surgery was performed using isoflurane anesthesia, and all infusions were performed using stereotaxic coordinates empirically derived from pilot surgeries guided by a mouse brain atlas (Paxinos & Franklin, 2004) using a stereotaxic surgical apparatus (Kopf Instruments, Tijuana, CA). Coordinates (from bregma) for the MHb were: (2 infusions were done on each side) AP –1.4 and –1.8 mm, ML 0.25 mm on first side, 0.3 mm on the second side, DV –2.6 mm; for vHipp: AP –3.0 mm, ML $\pm$ 3.0 mm, DV –3.2 mm; for ARC: AP –0.9 mm, ML $\pm$ 0.3 mm, DV –6.0 mm. All infusions were pressure injections done with pulled glass pipettes (inner diameter of 10–40 μ m) and 0.5 – 0.7 μ l of virus was infused at each site. The pipette was left in place for 5–6 min before being withdrawn.

Animals were monitored until completely alert, returned to their home cages, and monitored post-operatively for 72 hrs. Animals were administered post-operative analgesia via intraperitoneal injection on the day of surgery and once per day for the next 48 hrs.

Viral Knockdown Validation

$\beta 4$ Knockdown—To verify knockdown, a cocktail of the two shRNA constructs targeting the $\beta 4$ subunit were injected bilaterally into the MHb of ChAT-Cre mice, as the $\beta 4$ subunit is highly expressed in the MHb and a large proportion of the ventral MHb cells express ChAT. ChAT-IRES-Cre mice and their wild-type littermates infused with either AAV-dsRed-Sico- $\beta 4$ shRNAs or AAV-dsRed-Sico-Scr shRNA were sacrificed 12 wks after surgery by rapid decapitation, and brains were removed and immediately frozen in isopentane chilled in a dry ice-ethanol bath. 200- μ m thick GFP-containing sections were taken using a NIGHTSEA BlueStar flashlight and filter goggles (Electron Microscopy Sciences, Hatfield, PA) to visualize GFP. Sections were mounted on glass slides, and 1-mm diameter punches of GFP-

enriched MHb from the frozen sections were taken under LED visualization. 4–6 punches were obtained from each animal. Four animals with off-target GFP expression were excluded from analysis. Punches were stored on dry ice and subsequently at -80°C until RNA extraction.

For $\beta 4$ knockdown, expression of dsRed (as determined by qPCR Ct value) was used to determine viral infection, and samples with greater than 2 cycle differences (after normalizing to *tbp1*) from the group mean were excluded. This resulted in $n = 3$ for wild-type mice, $n = 4$ for ChAT-IRES-Cre mice. In Cre+ mice, expression levels of $\beta 4$ mRNA detected by real time qPCR were significantly reduced compared to Cre- animals (Supplemental Fig. 2: $t(5) = 6.488$, $p = 0.0013$). In the same animals, $\beta 2$ expression was not significantly altered, indicating the specificity of the construct.

$\alpha 7$ Knockdown—To validate the $\alpha 7$ knockdown constructs *in vivo*, a cocktail of the AAVs containing the three shRNAs was injected bilaterally into the ventral hippocampus of GAD-Cre mice, as GABAergic cells in this region express high levels of the $\alpha 7$ subunit. GAD-Cre mice infused with either AAV-dsRed-Sico- $\alpha 7$ shRNAs or AAV-dsRed-Sico-Scr shRNA were allowed to recover for 5 wks and were sacrificed by rapid decapitation. Brains were removed and sectioned using a brain block (Zivic Instruments), and slices containing the hippocampus were transferred to ice cold phosphate buffered saline. Virally infected regions of the hippocampus were identified using a NIGHTSEA BlueStar flashlight and filter goggles (Electron Microscopy Sciences, Hatfield, PA) and 1.2 mm diameter punches of GFP-enriched hippocampus were taken under LED visualization. 6 punches were obtained from each animal, 3 from each hemisphere. Left and right hippocampi were kept separate and treated as individual replicates. Punches were stored on dry ice and subsequently at -80°C .

Samples that did not yield visible PCR products for GFP and dsRed, when run on a 1.5% agarose gel were consistent with samples in which GFP was not visualized during tissue dissection. These samples were considered to be off target and excluded from analysis (2 hippocampi). This resulted in $n = 6$ GAD-Cre hippocampi with SCR shRNA, and $n = 8$ GAD-Cre hippocampi with $\alpha 7$ shRNA. Compared to GAD-Cre mice injected with a SCR virus not targeting any specific mRNA, expression of $\alpha 7$ subunit mRNA was reduced by approximately 50% in the hippocampus of GAD-Cre mice that received the viral knockdown cocktail (Supplemental Fig. 3: $t(12) = 4.43$, $p = 0.0008$). As with the $\beta 4$ knockdown, expression levels of $\beta 2$ mRNA were not significantly changed compared to controls.

Assessment of Food Intake

Mice were single housed one week before experimental manipulations or at the time of surgery. Body weight and food weight were measured daily for the duration of behavioral experiments. Mice were allowed to recover for 6 weeks after surgery to ensure receptor turnover and effective nAChR subunit knockdown. Mice were acclimated to saline injection until animals consistently ate a minimum of 0.7 g of food in the 2 hr window as described in the feeding paradigm below. After acclimation, mice were randomized to receive saline,

nicotine (0.5 mg/kg) or cytosine (1.5 mg/kg) in one of six possible presentation orders, and the feeding paradigm was repeated in successive rounds until all animals received each drug. Following completion of all successive rounds, mice were again randomized and the procedure was repeated for higher doses of nicotine (1.0 mg/kg) and cytosine (3.0 mg/kg). If control mice receiving saline did not eat an average of 1.0 g of food in the 2 hr window as a group, all data from that round were excluded from analysis and repeated after the completion of remaining planned rounds of the sequence at that dose.

The feeding paradigm consisted of a 48-hour window beginning with a 0.1 mL IP saline injection, and removal of food at 5pm. Twenty-four hrs later, animals received an IP injection of drug or saline. Fifteen min after injection, food was returned to the cage. Food weight was measured 2 hrs and 24 hrs later. Mice were allowed to recover for 5 days to regain their initial free-feeding body weight before any changes in food intake were tested again.

Tissue Processing and Immunohistochemistry

Wild type animals used for ARC immunohistochemistry were deeply anesthetized via IP injection of an overdose of pentobarbital (**130 mg/kg**; Fatal Plus, Patterson Veterinary), transcardially perfused with ice-cold PBS followed by 4% paraformaldehyde. Brains were removed, post-fixed overnight in the same fixative, then cryoprotected in 30% sucrose. 40- μ m-thick sections were cut on a sliding microtome and collected in six series. Slices containing the ARC (one section series) were stained for POMC and ChAT expression. Sections were washed three times for 5 min in PBS, then blocked in 3% normal donkey serum (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) in PBS with 0.3% Triton X-100 (PBST). Sections were then incubated in primary antibody (rabbit anti-POMC precursor, Phoenix Pharmaceuticals H-029-30, 1:3,000; Goat anti-Choline Acetyltransferase, Millipore Sigma AB144P, 1:1,000) overnight at room temperature in PBST. After three 8 min washes, sections were incubated in secondary antibody (Donkey anti-Rabbit IgG AlexaFluor 555, ThermoFisher Scientific A-31572, 1:1,000; Donkey anti-Goat AlexaFluor 647, Life Technologies A-21447, 1:1,000) in PBST for 2 hr at room temperature, then washed three times, mounted and cover slipped with Fluoromount G (EMS, Hatfield, PA).

After completion of all behavioral experiments, tissue was collected from POMC-Cre and AgRP-Cre mice as described above. Three to four slices containing the ARC (one section series) from each mouse was stained for the virally-expressed fluorophores dsRed and GFP to verify proper targeting. Sections were stained using the same protocol as above with rabbit polyclonal anti-dsRed (Clontech 632496, 1:1000) and goat polyclonal anti-GFP (Abcam ab6673, 1:1000) primary antibodies followed by donkey anti-Rabbit IgG AlexaFluor 555 (ThermoFisher Scientific A-31572, 1:1000) and donkey anti-Goat AlexaFluor 488 (ThermoFisher Scientific A-11055, 1:1000) secondary antibodies.

Sections were visualized on a confocal microscope (Fluoview FV10i, Olympus). AgRP- and POMC-Cre animals with viral expression in the ARC and appropriate viral recombination indicated by the presence of red-only cells were considered to be appropriately targeted. All

animals that did not meet this criterion, independent of which virus was infused, were combined to generate an “off-target control” group for analyses.

Statistics

Data are presented as box plots representing the min, max, and quartile values of the group or as bar graphs showing the mean and SEM. Enrichment for bacTRAP qPCR samples was compared with paired t-tests, and differences between AgRP and POMC TRAP mRNA expression was compared with unpaired t-tests. Relative protein expression in immunoprecipitation experiments were compared with unpaired t-tests. A one-way ANOVA with Holm-Sidak post hoc correction of the planned comparisons was used to determine $\beta 4$ binding partners with sequential immunoprecipitation. For food intake of wild type animals a one-way, repeated measures ANOVA with planned comparisons was used to compare effects of nicotine and cytosine to saline control. A two-way, repeated measures ANOVA with planned comparisons of active drug (nicotine or cytosine) to saline, within viral treatment group, was used for viral knockdown feeding experiment. AgRP and POMC mice were separate for these analyses. Holm-Sidak *post-hoc* corrections were used to determine significance. P or adjusted p of 0.05 was regarded as significant. Statistical analyses were carried out using Graph Pad Prism 8.

Results

nAChR subunit mRNA representation in ARC

As an initial step to identify the nAChR subunits expressed in ARC we used reverse transcription, qualitative PCR to determine whether the mRNAs encoding each of the neuronal nAChR subunits were expressed in the ventral hypothalamus. All nAChR subunits except the $\alpha 2$ subunit were detected in this experiment (Fig. 1A). For cell type specific mRNA expression we used quantitative real time PCR analysis (qPCR) of mRNA isolated from whole hypothalamus and mRNA isolated using TRAP from either AgRP or POMC cells. We determined whether mRNA for individual nAChR subunits were enriched in the pool of actively transcribed mRNA from either cell type over the concentration of all mRNAs found in the hypothalamus in general, and if there were differences in relative concentration of translating mRNA for any subunit between these two cell populations.

As with the qualitative PCR from mRNA obtained from ventral hypothalamic punches (Fig. 1A), the majority of nAChR subunits were detectable in the hypothalamic input fraction, although transcripts encoding the $\beta 3$ subunit were not consistently detected in either the input fraction or the TRAP-isolated mRNA from either cell type (Fig. 1). Further, while the $\alpha 2$ subunit was not detectable in the PCR samples derived from a single mouse, it was identified in the hypothalamic tissue pooled from multiple animals. However, like the $\beta 3$ subunit, the $\alpha 2$ and $\alpha 5$ subunits were not detected reliably in TRAP-isolated mRNA from either AgRP or POMC cells (data not shown), indicating that while mRNAs encoding these three fairly low abundance subunits are expressed in the hypothalamus, they are not represented in the pool of actively transcribed mRNA in AgRP or POMC cells of the ARC.

The majority of detectable subunits were enriched in both AgRP and POMC cell types when the TRAP fractions were normalized to mRNA concentrations in the whole hypothalamic homogenate (input fraction), which would include other neuronal subtypes as well as multiple glial and endothelial cell types (Fig. 1B–D). The $\alpha 4$ and $\alpha 6$ subunits showed the most robust enrichment over the hypothalamic input fraction in both AgRP and POMC cells with group median values between 20 and 30 fold enrichment (Fig. 1B) ($\alpha 4$: AgRP, $t(6) = 6.453$, $p = 0.0001$; POMC, $t(9) = 5.841$, $p = 0.0002$; $\alpha 6$: AgRP, $t(6) = 2.951$, $p = 0.0256$; POMC, $t(8) = 8.564$, $p < 0.0001$). Both the $\beta 4$ and $\alpha 3$ subunits were significantly enriched in TRAP mRNA from both POMC and AgRP cells as well (Fig. 1C) ($\beta 4$: AgRP, $t(3) = 4.016$, $p = 0.0277$; POMC, $t(4) = 4.788$, $p = 0.0087$; $\alpha 3$: AgRP, $t(6) = 3.09$, $p = 0.0214$; POMC, $t(7) = 10.42$, $p < 0.0001$). Finally, the $\beta 2$ subunit was also enriched about 3 fold in both AgRP and POMC cells (Fig. 1D: AgRP, $t(7) = 6.618$, $p = 0.0003$; POMC, $t(9) = 4.812$, $p = 0.0010$). Unlike the other detectable subunits, the $\alpha 7$ subunit was not enriched in the AgRP or POMC cell types. While qPCR CT values indicated high levels of expression in both the TRAP and input fractions (Ct values around 18), TRAP isolated $\alpha 7$ mRNA was significantly reduced in AgRP neurons compared to the input and there was a trend for reduction of this mRNA in POMC cells (Fig. 1E: AgRP, $t(7) = 5.227$, $p = 0.0012$; POMC, $t(8) = 2.089$, $p = 0.0701$). This result suggests that $\alpha 7$ mRNA is highly expressed in other cell types in ARC in addition to the AgRP and POMC cells.

When concentrations of mRNAs encoding nAChR subunits were directly compared between AgRP and POMC cells directly, only transcripts encoding two of the six detected subunits differed between the cell types. As suggested by the enrichment comparison, mRNA for the $\alpha 7$ subunit is more highly expressed in POMC cells compared to AgRP cells (Fig. 1F; $t(16) = 2.488$, $p = 0.0243$). Further, there was a trend toward a difference in expression of mRNA encoding the $\alpha 4$ subunit, with higher expression in AgRP cells (Fig. 1G; $t(18) = 2.092$, $p = 0.0509$). No differences in transcript levels between POMC and AgRP cells were detected for $\alpha 3$, $\alpha 6$, $\beta 2$ or $\beta 4$ (data not shown).

Levels of nAChR subunit proteins in ARC

Although mRNA levels are informative about proteins transcribed in neurons with cell bodies in ARC, mRNA levels do not always correspond well with protein levels (Carlyle *et al.*, 2017), and not all protein subunits may be incorporated into assembled receptors. To understand which nAChR subunits are incorporated into assembled nAChRs expressed in the ARC itself and not trafficked out to terminals, we performed immunoprecipitations from hypothalamic homogenates after isolating assembled heteromeric nAChRs by [^3H]-Epibatidine binding. All the tested nAChR subunits ($\alpha 2$ – 6 , $\beta 2$ – 4) were detectable in a crude dissection of hypothalamic tissue, and all but the $\alpha 2$ subunit were detectable above background in a more specific punch of hypothalamus containing the ARC (Table 2). As there was more starting material in the hypothalamic samples, normalizing the raw [^3H]-Epibatidine counts obtained by subunit-specific antibody immunoprecipitation with the total [^3H]-Epibatidine binding present in the samples allowed for a relative comparison of how much of the total signal is accounted for by each subunit. When compared directly, signal from the $\alpha 3$ subunit makes up a greater percentage of the total ARC signal compared to the hypothalamus as a whole ($t(4) = 3.52$, $p = 0.0245$). Similarly, signal for the $\beta 4$ subunit

trended in the same direction ($t(5) = 2.216$, $p = 0.0775$). Conversely, the $\alpha 4$ and $\beta 2$ subunits appeared to account for a smaller proportion of the signal in the ARC compared to the hypothalamus overall ($\alpha 4$, $t(3) = 2.0279$, $p = 0.0720$; $\beta 2$, $t(4) = 2.077$, $p = 0.1064$). The other nAChR subunits represented approximately the same proportion of the signal in both the total hypothalamus and the more specific ARC sample.

Evidence that the $\beta 4$ subunit is expressed in the hypothalamus is clear from examining its expression in the hypothalamus of $\beta 2$ nAChR subunit knockout mice (Table 3). As expected, the [^3H]-Epibatidine signal is much lower in homogenates from the $\beta 2$ knockout mice compared to homogenates from wild type mice because of the loss of the more abundant $\beta 2$ nAChR subunit. However, when raw radioactivity counts are normalized to total [^3H]-Epibatidine binding in each sample, $\beta 2$ signal is reduced to background levels in homogenates from knockout mice ($t(4) = 20.82$, $p < 0.0001$) and relative $\alpha 4$ subunit incorporation into assembled nAChRs is dramatically reduced ($t(5) = 14.87$, $p < 0.0001$). Incorporation of the $\alpha 5$ subunit is also reduced in nAChRs isolated from $\beta 2$ knockout mice ($t(3) = 3.419$, $p = 0.0419$). Further, both the $\alpha 3$ and $\beta 4$ subunits make up a significantly larger percentage of the total nAChR signal in the hypothalamus from $\beta 2$ knockout mice ($\alpha 3$ $t(4) = 4.128$, $p = 0.0145$; $\beta 4$ $t(5) = 5.737$, $p = 0.0023$).

Since the $\beta 4$ nAChR subunit is more highly expressed in ARC compared to the rest of the hypothalamus, sequential immunoprecipitations were performed to understand which nAChR subunits incorporate with this subunit to form assembled nAChRs in the ARC.

To perform this experiment, an initial immunoprecipitation was performed with the antibody for the nAChR subunit of interest, then the supernatant from this immunoprecipitation was used for the second immunoprecipitation with the $\beta 4$ antibody. These results are summarized in Table 4. As an internal control, two sequential immunoprecipitations were performed with the $\beta 4$ antibody, which reduced the signal in the second immunoprecipitation to near zero, indicating that the majority of the $\beta 4$ subunits in the sample had been bound and removed in the first immunoprecipitation. For comparison, if a sequential immunoprecipitation was not significantly different from this internal control, then this was considered evidence for subunit interactions and binding in assembled channels. A one-way ANOVA indicated that signal from some subunits did differ from the $\beta 4$ - $\beta 4$ control (Table 4; $F_{(8,7)} = 7.753$, $p = 0.0074$).

Based on Holm-Sidak adjusted post hoc comparisons, immunoprecipitation for the $\alpha 3$ subunit followed by $\beta 4$ immunoprecipitation was not different from the $\beta 4$ - $\beta 4$ control, suggesting that these two subunits co-assemble, as expected (Grady *et al.*, 2009). $\beta 4$ signal was also reduced when the first immunoprecipitation was performed with antibodies against $\alpha 4$ or $\beta 2$ nAChR subunits, but signal was not reduced and remained significantly higher than $\beta 4$ - $\beta 4$ control levels after first immunoprecipitation with antisera against $\alpha 2$ (adjusted $p = 0.0284$), $\alpha 5$ (adjusted $p = 0.0141$), $\alpha 6$ (adjusted $p = 0.0147$), or $\beta 3$ (adjusted $p = 0.087$). Based on these data, $\beta 4$ subunits assemble into nAChRs with $\alpha 3$, $\alpha 4$, and $\beta 2$ subunits, but not $\alpha 2$, $\alpha 5$, $\alpha 6$ or $\beta 3$ subunits in the ARC. In a small follow up experiment ($n = 1$), sequential immunoprecipitation with the $\beta 2$ subunit antibody after either the $\alpha 3$ or $\beta 4$ subunit showed that the percent of total [^3H]-Epibatidine binding is greatly reduced,

consistent with the described interaction with $\beta 4$ and suggesting that $\beta 2$ also combines with the $\alpha 3$ in assembled nAChRs in ARC, as has been shown in other brain areas (Grady *et al.*, 2009). This analysis indicates that either $\beta 4$ or $\beta 2$ may be acting as an auxiliary subunit in $\alpha 4\beta 2^*$ or $\alpha 3\beta 4^*$ channels respectively.

Expression of cholinergic markers in the ARC

In addition to testing the levels of translated nAChR subunits in TRAP mRNA samples, we also measured expression of translated choline acetyltransferase (ChAT) and vesicular acetylcholine transporter (VACHT) mRNAs, two enzymes in the ACh synthesis and packaging pathway commonly used as markers for cholinergic neurons. Previous studies have ascribed a cholinergic phenotype specifically to POMC neurons in this region (Meister *et al.*, 2006), but we found that translated mRNAs encoding ChAT and VACHT were enriched in both AgRP and POMC cells when compared to expression in the hypothalamic input (Fig. 2A: AgRP, $t(6) = 6.972$, $p = 0.0004$; POMC, $t(9) = 5.346$, $p < 0.0005$). In addition, when compared to POMC neurons directly, AgRP neurons express these mRNAs at a 2–3 fold higher concentration (Fig. 2B: ChAT $t(15) = 3.561$, $p = 0.0028$; VACHT $t(14) = 7.549$, $p < 0.0001$). Consistent with mRNA measurements, immunohistochemistry for ChAT protein in the ARC revealed that ChAT protein is expressed not only in POMC cells in ARC (white arrowheads), but also in other cells in the ARC that do not express POMC (white arrows; Fig. 2C).

Relative role of $\beta 4$ and $\alpha 7$ nAChR subunits in the ARC in the feeding response to nicotine and cytosine

To examine the effect of nAChR agonists on food intake, we used a repeated measures feeding paradigm in which animals were food deprived for 24 hr then injected with saline, nicotine or cytosine in a randomized order before returning food to their home cage. Food intake was measured 2 and 24 hr after each treatment. Both nicotine and cytosine reduced food intake at 2 hr compared to saline administration in this acute feeding paradigm (Fig. 3A; repeated measures ANOVA $F(2,20) = 4.273$, $p = 0.0285$, *post hoc* nicotine v. saline adjusted $p = 0.0317$, cytosine v. saline adjusted $p = 0.0269$). There were no differences in 24 hr food intake (data not shown).

To gain a cell-type specific understanding of nAChR subunit control of food intake in response to these nicotinic drugs, we developed cell type selective shRNA viral constructs designed to target either the $\beta 4$ or $\alpha 7$ nAChR subunit (Fig. 3B). After verification, viral knockdown cocktails were injected bilaterally into the ARC of AgRP- or POMC-Cre animals to selectively knockdown either subunit in one cell type, rather than all neuronal cell types in the ARC. In the absence of Cre, both dsRed and GFP are expressed, while recombination in the presence of Cre leads to excision of GFP and expression of the shRNA construct. A representative image of bilateral ARC injection and the resulting recombination are shown in Fig. 3C. Virally injected mice were singly housed and weighed daily for 6 wk before the start of behavioral experiments to allow for viral expression and nAChR protein turn over. Body weight between groups did not differ at the time of surgery or after the 6 wk viral incubation period in either AgRP- or POMC-Cre mice (data not shown). After habituation to the feeding paradigm, animals were food deprived for 24 hours then injected

with saline, nicotine or cytosine in a randomized order for a within-subjects comparison of food intake. A repeated measures ANOVA indicated a main effect of drug in both POMC and AgRP animals (POMC: $F_{(2,124)} = 16.91$, $p < 0.0001$; AgRP: $F_{(2,118)} = 9.894$, $p = 0.001$), but no significant main effects of knockdown condition or interaction were observed. Planned comparisons of drug (cytosine or nicotine) to saline within knockdown condition and cell type revealed significant differences in food intake at the 2 hr time point (Fig. 3D), but not at 24 hrs (data not shown).

Knockdown of the $\beta 4$ subunit in POMC neurons blunted the ability of cytosine and nicotine to decrease food intake compared to saline treatment as animals with $\beta 4$ knockdown did not eat significantly less after cytosine or nicotine than they did with saline (adjusted $p > 0.05$). For both drugs, the reduction in food intake following $\beta 4$ subunit knockdown in POMC neurons was not statistically significant, indicating at least a partial blockade of the effects of cytosine and nicotine by $\beta 4$ knockdown in POMC cells of the ARC. Interestingly, $\beta 4$ subunit knockdown in AgRP neurons was also sufficient to block the decrease in feeding caused by cytosine and nicotine (adjusted $p > 0.05$). In contrast to $\beta 4$ knockdown, knockdown of the $\alpha 7$ subunit in POMC cells did not decrease the ability of nicotine or cytosine, as both drugs significantly reduced food intake compared to saline intake (cytosine, adjusted $p = 0.0066$, nicotine, adjusted $p = 0.0010$). $\alpha 7$ knockdown in AgRP cells also did not block the decrease in feeding due to nicotine (adjusted $p = 0.0079$). There was a trend for two-hour food intake in response to cytosine to be reduced compared to saline food intake, although this decrease did not reach significance (adjusted $p = 0.0567$).

In addition to $\beta 4$ and $\alpha 7$ knockdown, a subset of AgRP- and POMC-Cre animals were infused with AAV2 carrying a scrambled (SCR) shRNA (Supplemental Fig. 4). There was a trend for cytosine and nicotine to decrease food intake in these groups. There was a trend for cytosine and nicotine to decrease food intake in these groups. In POMC animals, cytosine trended to decrease intake (adjusted $p = 0.0636$), while the effect of nicotine was not as robust (adjusted $p = 0.1392$). In AgRP animals, those with the SCR KD had the lowest saline intake of any group, and there was not a robust further decrease in intake after cytosine or nicotine, potentially indicating a floor effect in this group. Because of the small size of the ARC, in all groups there were numerous animals in which the injected virus was off-target and viral expression was observed in other parts of the hypothalamus, with no expression or recombination in the ARC (Supplemental Fig. 4B). These animals were pooled and analyzed as an off-target control group. Nicotine and cytosine significantly decreased food intake compared to saline in both AgRP- and POMC-Cre off-target animals (POMC adjusted $p < 0.0001$ for both drugs, AgRP adjusted $p < 0.0001$ for both drugs), indicating that both nicotine and cytosine so reduce intake at baseline in this paradigm in these mouse lines.

Discussion

The data presented here suggest that the effects of nAChR signaling in the ARC are significantly more complex than was originally hypothesized, particularly in the control of feeding regulation by nicotinic compounds. Numerous nAChR subunits are expressed in hypothalamus ($\alpha 2$ – $\alpha 7$, $\beta 2$ – $\beta 4$) at the mRNA level, with a subset of these selectively enriched in ARC AgRP and POMC neurons ($\alpha 3$, $\alpha 4$, $\alpha 6$ and $\beta 2$, $\beta 4$). While high levels of

the $\alpha 7$ subunit mRNA were detected, this subunit was not significantly enriched in either AgRP or POMC cells above levels detected in the rest of the hypothalamus. Surprisingly, there was not dramatic cellular segregation in nAChR expression, with no differences between cell types detected in expression of $\alpha 3$, $\alpha 6$, $\beta 2$ or $\beta 4$. mRNA levels for $\alpha 4$ trended toward higher levels in AgRP compared to POMC neurons and $\alpha 7$ mRNA levels were significantly higher in POMC neurons. At the protein level, the same set of nAChR subunits were detected (excluding $\alpha 7$, which was not tested because specific antisera are not available) in both the hypothalamus overall and in an ARC-enriched fraction of hypothalamus. Compared to the hypothalamus overall, a higher proportion of nAChR signal included the $\alpha 3$ subunit in the ARC, with a trend toward ARC enrichment for the $\beta 4$ subunit as well. In contrast, the $\beta 2$ and $\alpha 4$ subunits showed a trend to make up less of the nAChR signal in the ARC compared to the overall hypothalamus, indicating region-selective expression of particular nAChR subunits. Further, consistent with a previous report (Picciotto *et al.*, 1995) deletion of $\beta 2$ subunit eliminates approximately 90% of the high affinity [^3H]-Epibatidine binding in the hypothalamus, and a higher percentage of the remaining receptors contain the $\alpha 3$ and $\beta 4$ subunits thus confirming that the $\alpha 3$ and $\beta 4$ signal observed in wild type animals is specific.

A subset of ARC neurons express the enzymes necessary to synthesize and package ACh, indicating the potential for local ACh synthesis and release in this brain area. Both AgRP and POMC neurons express mRNA for ChAT and VAcHT, with AgRP neurons expressing comparatively more than POMC neurons. At the protein level, both POMC-positive and POMC-negative neurons express ChAT. At the level of food intake, while the knockdown of the $\beta 4$ subunit appears to diminish the behavioral feeding response to nicotine and cytosine relative to knockdown of the $\alpha 7$ subunit, the behavioral responses appear to be similar when these subunits are knocked down in AgRP versus POMC cells. Knockdown of the $\beta 4$ subunit in AgRP-Cre mice blocks the decrease in food intake caused by nicotine and cytosine, and similarly, when the $\beta 4$ subunit is knocked down in POMC neurons the decrease in food intake is blunted, though not completely abolished. Following knockdown of the $\alpha 7$ subunit in either cell type, the decrease in feeding due to cytosine or nicotine is significantly more robust than in either line after knockdown of the $\beta 4$ subunit.

The distinct function of AgRP and POMC cells in the control of food intake has been confirmed clearly in recent optogenetic studies, which identify rapid and dramatic increase in food intake following stimulation of AgRP neurons and decreased food intake following stimulation of POMC cells (Aponte *et al.*, 2011; Krashes *et al.*, 2011; Atasoy *et al.*, 2012; Betley *et al.*, 2013; Fenselau *et al.*, 2016). A number of recent studies have evaluated broad gene expression profiles of AgRP or POMC neurons using RNA-Seq (Henry *et al.*, 2015; Lam *et al.*, 2017) or more targeted gene profiling techniques (Jeong *et al.*, 2016), although these studies did not isolate actively translated mRNAs, like the TRAP method used here, nor did they focus on nAChRs. Although not the explicit focus of the analysis, Henry *et al.* (2015) did identify a number of nAChR subunit mRNAs in both AgRP and POMC cells. As in the current study, they also identified mRNAs encoding the $\alpha 3$, $\alpha 4$, and $\alpha 7$ and $\beta 2$ and $\beta 4$ subunits, and even in this genome wide study, identified a difference in $\alpha 4$ and $\alpha 7$ mRNA levels between AgRP and POMC cells in the same directions observed here when animals were in a fed state. Consistent with the current TRAP study, they saw no differences in

levels of mRNAs encoding the other nAChR subunits between AgRP and POMC cells. However, in the genome-wide study they did not identify mRNAs encoding $\alpha 2$, $\alpha 5$, $\alpha 6$ or $\beta 3$ nAChR subunits, likely because these transcripts are present at lower levels and the experiment was not designed to detect this population of molecules specifically. Interestingly, this study suggested that transcripts encoding the $\alpha 4$ nAChR subunit are decreased in AgRP neurons when animals are food deprived compared to a fed state, indicating potential dynamic regulation of nAChR mRNA levels, although it is not known whether the proportion of actively translated mRNA differs. Notably, difference in nAChR stoichiometry can impact receptor function such that $(\alpha 4\beta 2)_2\alpha 4$ receptors have higher channel conductance, shorter open times and lower agonist sensitivity than $(\alpha 4\beta 2)_2\beta 2$ receptors (Papke *et al.*, 2010; Lucero *et al.*, 2015; Mazzaferro *et al.*, 2016), thus differences in $\alpha 4$ subunit levels could alter nAChR channel dynamics.

In a single-cell RNA-Seq study that focused on POMC neurons (Lam *et al.*, 2017), only $\alpha 4$ was listed among nAChR subunit mRNAs in the supplemental list of transcripts. Interestingly, in the 4 clusters of POMC cells they defined based on gene expression profiles, the $\alpha 4$ subunit ranged in expression from 3.7% of cells up to 20% of cells, indicating that $\alpha 4$ mRNAs are not homogenous in all cells that express POMC. This experiment also showed that a portion of POMC cells also expresses low levels of AgRP transcript, and vice versa. Thus, the differences in translated nAChR subunit mRNAs between AgRP and POMC cells identified here may be due to a difference in the relative proportion of AgRP or POMC cells actively transcribing these subunits, and a single-cell approach could provide necessary insights into sub-populations within these broad peptide-defined categories of cells. Another single-cell qPCR study confirms this heterogeneity within cells of the ARC (Jeong *et al.*, 2016). Of the 26 ARC cells tested, all expressed mRNA encoding ChAT, whereas expression of mRNAs encoding $\alpha 4$, $\alpha 7$, and $\beta 2$ nAChR subunits varied across cells, with some cells expressing all three subunits, some a combination of one or two subunits, and some cells did not express any nAChR subunit mRNAs at all. Further, of these ChAT positive cells, only about half expressed mRNA encoding POMC, consistent with the data presented here that POMC cells and non-POMC cells express ChAT protein. This confirms previous reports describing POMC neurons as cholinergic in the ARC (Meister *et al.*, 2006; Meister, 2007), but shows that the translation of cholinergic markers is not confined to this cell type. While the current study shows expression of ChAT and VACHRT mRNA in POMC and AgRP cells and ChAT protein in POMC and non-POMC cells, Meister, *et al.* (2006), showed that ChAT protein did not overlap with NPY protein in the ARC. It is possible that post-translational modification in AgRP neurons results in translational blockade or degradation of ChAT or VACHT proteins, and that POMC-negative/ChAT-positive cells in ARC represent another cell type entirely. Therefore, these POMC-negative/ChAT-positive cells shown in Fig. 2 may represent an entirely different neuronal cell type in the ARC, such as tyrosine hydroxylase-expressing neurons, which have been described in this region (Zhang & van den Pol, 2015, 2016).

While not every AgRP or POMC cell may express each nAChR subunit, the cells that do express these subunits form receptors in combinations that are unique compared to other brain areas. Although $\alpha 3\beta 4^*$ receptors are the primary nAChR in autonomic ganglion neurons, the expression of this type of nAChR in the brain is significantly more limited, with

highest expression in the MHb and its output nucleus the interpeduncular nucleus (IPN) (Dineley-Miller & Patrick, 1992). Despite its presence at lower levels, the current study and others (Mineur *et al.*, 2011; Henry *et al.*, 2015; Herman *et al.*, 2016) show expression of $\beta 4$ -containing nAChRs in the ARC. Interestingly, the composition of $\beta 4$ -containing nAChRs in the ARC differs from subtypes in the MHb and IPN. In the MHb, $\beta 4^*$ receptors make up about half of the nAChR population, while in the IPN, there are more $\beta 4^*$ receptors than $\beta 2^*$ receptors (Grady *et al.*, 2009). In the hypothalamus, $\beta 4$ subunits represent 6% of the nAChR population, and this percentage increases to 14% in the ARC, indicating an anatomically selective expression in this part of the hypothalamus. In all these brain regions, as in the autonomic ganglion, the predominant partner of $\beta 4$ in assembled nAChRs is the $\alpha 3$ subunit.

The current study shows that the $\beta 4$ subunit also forms nAChRs with the $\alpha 4$ and $\beta 2$ subunits in ARC. While the $\beta 2$ and $\beta 4$ subunits contribute to independent nAChR populations in MHb and IPN, $\alpha 3\beta 4\beta 2^*$ receptors have been described in the autonomic ganglia (Stokes & Papke, 2012). Unlike in the MHb or IPN, the $\beta 4$ subunit does not form channels with the $\alpha 2$ or $\alpha 6$ subunit in ARC. Further, $\beta 4$ subunits in the ARC do not form channels with the auxiliary subunits $\alpha 5$ and $\beta 3$, which are very common partners in the autonomic ganglia (Conroy & Berg, 1995) and MHb/IPN (Grady *et al.*, 2009) respectively. In the IPN, $\alpha 3\beta 4^*$ and $\alpha 3\beta 4\beta 3^*$ nAChRs regulate ACh release, and the $\beta 3$ subunit may play an important role in regulating nAChR trafficking to axon terminals in the IPN. Further, and in contrast to $\beta 4^*$ receptors, $\alpha 4\beta 2^*$ receptors can regulate the release of other neurotransmitters, potentially glutamate, in the IPN (Grady *et al.*, 2009). As we have no evidence for $\alpha 3\beta 4\beta 3^*$ nAChRs in the ARC, $\alpha 3\beta 4^*$ nAChRs could induce ACh release in the ARC, rather than out at distal terminal regions of these neurons. It is important to note that these immunoprecipitation experiments identifying co-assembled nAChR subunits cannot differentiate between nAChRs on neurons with cell bodies in the hypothalamus and nAChRs on axons from other brain areas terminating in the ARC. However, because there is consistency between translating mRNA and protein data in these experiments, at least a large subset of identified nAChRs are made and expressed by cells within the ARC.

The precise subunit combinations that make up each nAChR subtype dramatically affects channel properties and function, but also confers selectivity for different nicotinic drugs. Nicotine alters the electrophysiological excitability (Huang *et al.*, 2011; Mineur *et al.*, 2011), and neuropeptide expression levels (Frankish *et al.*, 1995; Fornari *et al.*, 2007; Hur *et al.*, 2010; Chen *et al.*, 2012; Calarco *et al.*, 2017) in both AgRP and POMC neurons in ARC. Nicotine application results in depolarization of both AgRP and POMC cells with components that are sensitive to both the $\beta 2^*$ -selective antagonist Dh β E- and the $\alpha 7$ -selective antagonist MLA (Huang *et al.*, 2011). Nicotine increases the firing frequency in both AgRP and POMC cells, and cytosine increases the firing frequency of POMC cells, whereas effects on AgRP neurons have not been evaluated. While nicotine is an agonist at all nAChR subtypes, cytosine is a full agonist at both $\alpha 3\beta 4^*$ and $\alpha 7$ nAChRs, and a low-efficacy partial agonist at $\alpha 4\beta 2^*$ nAChRs. In POMC neurons, both nicotine and cytosine cause a comparable increase in cell firing and both drugs result in a similar decrease in feeding, which is dependent on signaling through the MC4 receptor, the downstream target of both AgRP and α -MSH, a POMC derivative (Mineur *et al.*, 2011). Knockdown of the $\beta 4$

receptor in the ARC was sufficient to block the reductions in food intake caused by cytosine (Mineur *et al.*, 2011), and in the current study knockdown in either AgRP or POMC cells blunted the reduction in feeding due to cytosine and nicotine. Conversely, knockdown of the $\alpha 7$ subunit in either cell type did not alter the feeding response to cytosine or nicotine. While $\alpha 7$ is highly expressed in the hypothalamus and may regulate other circuits controlling food intake (Jo *et al.*, 2005; McFadden *et al.*, 2014), based on these data, it appears to be unlikely that $\alpha 7$ regulates food intake in response to nicotine through AgRP or POMC neurons. Notably, our TRAP studies showed that $\alpha 7$ mRNA was enriched in other cell types in ARC compared to AgRP or POMC neurons, suggesting that signaling through this nAChR subtype occurs on as-yet-unidentified cell types. Interestingly, the $\alpha 7$ subunit induces calcium waves in hippocampal astrocytes (Shen & Yakel, 2012), suggesting that nicotinic control of glia function could alter ARC function. Since glia are important regulators of neuronal energy balance (Yang *et al.*, 2015; Djogo *et al.*, 2016; Argente-Arizón *et al.*, 2017), and activity of ARC neurons is responsive to overall energy demands (Cansell *et al.*, 2012; Kohno & Yada, 2012; Sternson & Atasoy, 2014), this could be an additional function for nAChR signaling that could be evaluated.

The data presented here add a new layer of complexity to this ARC feeding circuit. Not only are the comparatively rare $\beta 4^*$ nAChRs sufficient to affect feeding behavior in response to nicotine, their role in AgRP and POMC neurons have similar downstream results despite the fact that AgRP and POMC cells have well described opposing effects on food intake. As study of this brain circuit has increased with increasingly specific tools, it is becoming more apparent that AgRP cells and POMC cells are not uniform categories. Both AgRP and POMC neurons project to many locations, and for both cell types a given neuron is likely to project to only one downstream target (King & Hentges, 2011; Betley *et al.*, 2013), indicating anatomical specificity. Further, single cell genetic profiling studies (Jeong *et al.*, 2016; Lam *et al.*, 2017) show that subsets of POMC and AgRP cells have gene expression profiles that could correlate with and define their anatomical specificity. This single cell specificity becomes even more relevant when considering that some POMC neurons express and release GABA, while others signal using glutamate, and other still have neither fast neurotransmitter but may signal exclusively through neuropeptide release (Hentges *et al.*, 2004, 2009; Dicken *et al.*, 2012; Marshall *et al.*, 2017). Although the majority of AgRP neurons express and release GABA, neurotransmitter variety has also been observed in AgRP neurons (Horvath *et al.*, 1997). Further, some AgRP and POMC neurons may also release ACh in addition to other neurotransmitters, as they express ChAT and VAcHT, although no experiments to date have demonstrated ACh release from these neurons directly. Because of the unique qualities of individual AgRP and POMC cells with respect to their projections, gene expression, and neurotransmitter profile, it is plausible that $\beta 4^*$ nAChRs may play a role in a specific subset of AgRP and POMC cells.

Conclusion

These experiments demonstrate that the $\beta 4$ subunit is expressed on both POMC and AgRP neurons in ARC, identify a broad range of nAChRs that are differentially present in ARC as compared to the hypothalamus as a whole, and indicate a more complex interaction in the ARC with respect to nAChR signaling than was predicted. These studies are important in

developing a more complete model of how nAChR activation in the ARC regulates food intake. Future experiments examining the heterogeneity of these populations are needed to understand this interaction fully. Determining the unique features of nAChR-mediated modulation of these cells with respect their gene expression, connectivity, and pattern of activity will provide insight into the mechanistic control of food intake and potentially provide therapeutic targets for clinical interventions to control weight.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH grant DA037566. The authors would like to thank Dr. Milena Moretti for running immunoprecipitation experiments with Dr. Cecelia Gotti. Thanks to Angela Lee and Richard Crouse for help with food intake measurements.

Abbreviations

3V	3 rd ventricle
AgRP	agouti-related peptide
ARC	arcuate nucleus
ChAT	choline acetyltransferase
DMH	dorsomedial hypothalamus
IPN	interpeduncular nucleus
ME	median eminence
MHb	medial habenula
nAChR	nicotinic acetylcholine receptor
POMC	pro-opiomelanocortin
VAcHT	vesicular acetylcholine transporter
vHipp	ventral hippocampus

References

- Albanes D, Jones DY, Micozzi MS, Mattson ME. Associations between smoking and body weight in the US population: Analysis of NHANES II. *Am. J. Public Health.* 1987; 77:439–444. [PubMed: 3493709]
- Albuquerque EX, Pereira EFR, Alkondon M, Rogers SW. Mammalian nicotinic acetylcholine receptors: From structure to function. *Physiol. Rev.* 2009; 89:73–120. [PubMed: 19126755]
- Allison MB, Patterson CM, Krashes MJ, Lowell BB, Myers MG, Olson DP. TRAP-seq defines markers for novel populations of hypothalamic and brainstem LepRb neurons. *Mol. Metab.* 2015; 4:299–309. [PubMed: 25830093]

- Aponte Y, Atasoy D, Sternson SM. AGRP neurons are sufficient to orchestrate feeding behavior rapidly and without training. *Nat. Neurosci.* 2011; 14:351–355. [PubMed: 21209617]
- Argente-Arizón P, Guerra-Cantera S, Garcia-Segura LM, Argente J, Chowen JA. Glial cells and energy balance. *J. Mol. Endocrinol.* 2017; 58:R59–R71. [PubMed: 27864453]
- Atasoy D, Betley JN, Su HH, Sternson SM. Deconstruction of a neural circuit for hunger. *Nature.* 2012; 488:172–177. [PubMed: 22801496]
- Betley JN, Cao ZFH, Ritola KD, Sternson SM. Parallel, redundant circuit organization for homeostatic control of feeding behavior. *Cell.* 2013; 155:1337–1350. [PubMed: 24315102]
- Calarco CA, Lee S, Picciotto MR. Access to nicotine in drinking water reduces weight gain without changing caloric intake on high fat diet in male C57BL/6J mice. *Neuropharmacology.* 2017; 123:210–220. [PubMed: 28623168]
- Cansell C, Denis RGP, Joly-Amado A, Castel J, Luquet S. Arcuate AgRP neurons and the regulation of energy balance. *Front. Endocrinol. (Lausanne).* 2012; 3:169. [PubMed: 23293630]
- Carlyle BC, Kitchen RR, Kanyo JE, Voss EZ, Pletikos M, Sousa AMM, Lam TKT, Gerstein MB, Sestan N, Nairn AC. A multiregional proteomic survey of the postnatal human brain. *Nat. Neurosci.* 2017; 20:1787–1795. [PubMed: 29184206]
- Chen H, Saad S, Sandow SL, Bertrand PP. Cigarette smoking and brain regulation of energy homeostasis. *Front. Pharmacol.* 2012; 3:1–8. [PubMed: 22291651]
- Chiolero A, Faeh D, Paccaud F, Cornuz J. Consequences of smoking for body weight, body fat distribution, and insulin resistance. *Am. J. Clin. Nutr.* 2008; 87:801–809. [PubMed: 18400700]
- Conroy WG, Berg DK. Neurons can maintain multiple classes of nicotinic acetylcholine receptors distinguished by different subunit compositions. *J. Biol. Chem.* 1995; 270:4424–4431. [PubMed: 7876208]
- Dicken MS, Tooker RE, Hentges ST. Regulation of GABA and glutamate release from proopiomelanocortin neuron terminals in intact hypothalamic networks. *J. Neurosci.* 2012; 32:4042–4048. [PubMed: 22442070]
- Dineley-Miller K, Patrick J. Gene transcripts for the nicotinic acetylcholine receptor subunit, beta4, are distributed in multiple areas of the rat central nervous system. *Brain Res. Mol. Brain Res.* 1992; 16:339–344. [PubMed: 1337943]
- Djogo T, Robins SC, Schneider S, Kryzskaya D, Liu X, Mingay A, Gillon CJ, Kim JH, Storch KF, Boehm U, Bourque CW, Stroh T, Dimou L, Kokoeva MV. Adult NG2-glia are required for median eminence-mediated leptin sensing and body weight control. *Cell Metab.* 2016; 23:797–810. [PubMed: 27166944]
- Fenselau H, Campbell JN, Verstegen AMJ, Madara JC, Xu J, Shah BP, Resch JM, Yang Z, Mandelblat-Cerf Y, Livneh Y, Lowell BB. A rapidly acting glutamatergic ARC→PVH satiety circuit postsynaptically regulated by α -MSH. *Nat. Neurosci.* 2016; 20:42–51. [PubMed: 27869800]
- Fornari A, Pedrazzi P, Lippi G, Picciotto MR, Zoli M, Zini I. Nicotine withdrawal increases body weight, neuropeptide Y and Agouti-related protein expression in the hypothalamus and decreases uncoupling protein-3 expression in the brown adipose tissue in high-fat fed mice. *Neurosci. Lett.* 2007; 411:72–76. [PubMed: 17052838]
- Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ. Habenular $\alpha 5$ nicotinic receptor subunit signalling controls nicotine intake. *Nature.* 2011; 471:597–601. [PubMed: 21278726]
- Frankish HM, Dryden S, Wang Q, Bing C, MacFarlane IA, Williams G. Nicotine administration reduces neuropeptide Y and neuropeptide Y mRNA concentrations in the rat hypothalamus: NPY may mediate nicotine's effects on energy balance. *Brain Res.* 1995; 694:139–146. [PubMed: 8974638]
- George AA, Bloy A, Miwa JM, Lindstrom JM, Lukas RJ, Whiteaker P. Isoform-specific mechanisms of $\alpha 3\beta 4^*$ -nicotinic acetylcholine receptor modulation by the prototoxin lynx1. *FASEB J.* 2017; 31:1398–1420. [PubMed: 28100642]
- Goni R, García P, Foissac S. The qPCR data statistical analysis. *Integromics White Pap.* 2009; 1:1–9.
- Gotti C, Moretti M, Meinerz NM, Clementi F, Gaimarri A, Collins AC, Marks MJ. Partial deletion of the nicotinic cholinergic receptor $\alpha 4$ or $\beta 2$ subunit genes changes the acetylcholine sensitivity of receptor-mediated 86Rb⁺ efflux in cortex and thalamus and alters relative expression of $\alpha 4$ or $\beta 2$ subunits. *Mol. Pharmacol.* 2008; 73:1796–1807. [PubMed: 18337473]

- Grady SR, Moretti M, Zoli M, Marks MJ, Zanardi A, Pucci L, Clementi F, Gotti C. Rodent habenulo-interpeduncular pathway expresses a large variety of uncommon nAChR subtypes, but only the $\alpha 3\beta 4^*$ and $\alpha 3\beta 3\beta 4^*$ subtypes mediate acetylcholine release. *J. Neurosci.* 2009; 29:2272–2282. [PubMed: 19228980]
- Grebenstein PE, Harp JL, Rowland NE. The effects of noncontingent and self-administered cytosine on body weight and meal patterns in male Sprague-Dawley rats. *Pharmacol. Biochem. Behav.* 2013; 110:192–200. [PubMed: 23876236]
- Grunberg NE, Bowen DJ, Winders SE. Effects of nicotine on body weight and food consumption in female rats. *Psychopharmacology (Berl)*. 1986; 90:101–105. [PubMed: 3094051]
- Heiman M, Kulicke R, Fenster RJ, Greengard P, Heintz N. Cell type-specific mRNA purification by translating ribosome affinity purification (TRAP). *Nat. Protoc.* 2014; 9:1282–1291. [PubMed: 24810037]
- Heiman M, Schaefer A, Gong S, Peterson JD, Day M, Ramsey KE, Suárez-Fariñas M, Schwarz C, Stephan DA, Surmeier DJ, Greengard P, Heintz N. A translational profiling approach for the molecular characterization of CNS cell types. *Cell.* 2008; 135:738–748. [PubMed: 19013281]
- Henry FE, Sugino K, Tozer A, Branco T, Sternson SM. Cell type-specific transcriptomics of hypothalamic energy-sensing neuron responses to weight-loss. *Elife.* 2015; 4:e09800.
- Hentges ST, Nishiyama M, Overstreet LS, Stenzel-Poore M, Williams JT, Low MJ. GABA release from proopiomelanocortin neurons. *J. Neurosci.* 2004; 24:1578–1583. [PubMed: 14973227]
- Hentges ST, Otero-Corchon V, Pennock RL, King CM, Low MJ. Proopiomelanocortin expression in both GABA and glutamate neurons. *J. Neurosci.* 2009; 29:13684–13690. [PubMed: 19864580]
- Herman AM, Ortiz-Guzman J, Kochukov M, Herman I, Quast KB, Patel JM, Tepe B, Carlson JC, Ung K, Selever J, Tong Q, Arenkiel BR. A cholinergic basal forebrain feeding circuit modulates appetite suppression. *Nature.* 2016; 538:253–256. [PubMed: 27698417]
- Horvath TL, Bechmann I, Naftolin F, Kalra SP, Leranth C. Heterogeneity in the neuropeptide Y-containing neurons of the rat arcuate nucleus: GABAergic and non-GABAergic subpopulations. *Brain Res.* 1997; 756:283–286. [PubMed: 9187344]
- Huang H, Xu Y, van den Pol AN. Nicotine excites hypothalamic arcuate anorexigenic proopiomelanocortin neurons and orexigenic neuropeptide Y neurons: similarities and differences. *J. Neurophysiol.* 2011; 106:1191–1202. [PubMed: 21653710]
- Hur Y-NN, Hong G-HH, Choi S-HH, Shin K-HH, Chun B-GG. High fat diet altered the mechanism of energy homeostasis induced by nicotine and withdrawal in C57BL/6 mice. *Mol. Cells.* 2010; 30:219–226. [PubMed: 20803089]
- Jeong JH, Woo YJ, Chua S, Jo Y-H. Single-cell gene expression analysis of cholinergic neurons in the arcuate nucleus of the hypothalamus. *PLoS One.* 2016; 11:e0162839. [PubMed: 27611685]
- Jo Y-H, Talmage DA, Role LW. Nicotinic receptor-mediated effects on appetite and food intake. *J. Neurobiol.* 2002; 53:618–632. [PubMed: 12436425]
- Jo Y-H, Wiedl D, Role LW. Cholinergic modulation of appetite-related synapses in mouse lateral hypothalamic slice. *J. Neurosci.* 2005; 25:11133–11144. [PubMed: 16319313]
- Kim ER, Wu Z, Sun H, Xu Y, Mangieri LR, Xu Y, Tong Q. Hypothalamic non-AgRP, non-POMC GABAergic neurons are required for postweaning feeding and NPY hyperphagia. *J. Neurosci.* 2015; 35:10440–10450. [PubMed: 26203139]
- King CM, Hentges ST. Relative number and distribution of murine hypothalamic proopiomelanocortin neurons innervating distinct target sites. *PLoS One.* 2011; 6:e25864. [PubMed: 21991375]
- Klesges RC, Winders SE, Meyers AW, Eck LH, Ward KD, Hultquist CM, Ray JW, Shadish WR. How much weight gain occurs following smoking cessation? A comparison of weight gain using both continuous and point prevalence abstinence. *J. Consult. Clin. Psychol.* 1997; 65:286–291. [PubMed: 9086692]
- Kohno D, Yada T. Arcuate NPY neurons sense and integrate peripheral metabolic signals to control feeding. *Neuropeptides.* 2012; 46:315–319. [PubMed: 23107365]
- Krashes MJ, Koda S, Ye C, Rogan SC, Adams AC, Cusher DS, Maratos-flier E, Roth BL, Lowell BB. Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. *J. Clin. Invest.* 2011; 121:2–6.

- Lam BYH, Cimino I, Poley-Wolf J, Nicole Kohnke S, Rimmington D, Iyemere V, Heeley N, Cossetti C, Schulte R, Saraiva LR, Logan DW, Blouet C, O'Rahilly S, Coll AP, Yeo GSH. Heterogeneity of hypothalamic pro-opiomelanocortin-expressing neurons revealed by single-cell RNA sequencing. *Mol. Metab.* 2017; 6:383–392. [PubMed: 28462073]
- Lewis JA, Yakel JA, Pandya AA. Levamisole: A positive allosteric modulator for the $\alpha 3\beta 4$ nicotinic acetylcholine receptors prevents weight gain in the CD-1 mice on a high fat diet. *Curr. Pharm. Des.* 2017; 23:1869–1872. [PubMed: 27908269]
- Lucero LM, Weltzin MM, Eaton JB, Cooper JF, Lindstrom JM, Lukas RJ, Whiteaker P. Differential $\alpha 4(+)/(-)\beta 2$ agonist binding site contributions to $\alpha 4\beta 2$ nicotinic acetylcholine receptor function within and between isoforms. *J. Biol. Chem.* 2015; 291:2444–2459. [PubMed: 26644472]
- Marshall CJ, Desrozier E, McLennan T, Campbell RE. Defining subpopulations of arcuate nucleus GABA neurons in male, female, and prenatally androgenized female mice. *Neuroendocrinology.* 2017; 105:157–169. [PubMed: 27710963]
- Matta SG, Balfour DJ, Benowitz NL, Boyd RT, Buccafusco JJ, Caggiula AR, Craig CR, Collins AC, Damaj MI, Donny EC, Gardiner PS, Grady SR, Heberlein U, Leonard SS, Levin ED, Lukas RJ, Markou A, Marks MJ, McCallum SE, Parameswaran N, Perkins KA, Picciotto MR, Quik M, Rose JE, Rothenfluh A, Schafer WR, Stolerman IP, Tyndale RF, Wehner JM, Zirger JM. Guidelines on nicotine dose selection for in vivo research. *Psychopharmacology (Berl).* 2007; 190:269–319. [PubMed: 16896961]
- Mazzaferro S, Bermudez I, Sine SM. $\alpha 4\beta 2$ nicotinic acetylcholine receptors: Relationships between subunit stoichiometry and function at the single channel level. *J. Biol. Chem.* 2016; 292:2729–2740. [PubMed: 28031459]
- McFadden KL, Cornier M-A, Tregellas JR. The role of $\alpha 7$ nicotinic receptors in food intake behaviors. *Front. Psychol.* 2014; 5:1–7. [PubMed: 24474945]
- Meister B. Neurotransmitters in key neurons of the hypothalamus that regulate feeding behavior and body weight. *Physiol. Behav.* 2007; 92:263–271. [PubMed: 17586536]
- Meister B, Gömüç B, Suarez E, Ishii Y, Dürr K, Gillberg L. Hypothalamic proopiomelanocortin (POMC) neurons have a cholinergic phenotype. *Eur. J. Neurosci.* 2006; 24:2731–2740. [PubMed: 17156199]
- Mineur YS, Abizaid A, Rao Y, Salas R, DiLeone RJ, Gündisch D, Diano S, De Biasi M, Horvath TL, Gao X-B, Picciotto MR. Nicotine decreases food intake through activation of POMC neurons. *Science.* 2011; 332:1330–1332. [PubMed: 21659607]
- Mineur YS, Fote GM, Blakeman S, Cahuzac ELM, Newbold SA, Picciotto MR. Multiple nicotinic acetylcholine receptor subtypes in the mouse amygdala regulate affective behaviors and response to social stress. *Neuropsychopharmacology.* 2016; 41:1579–1587. [PubMed: 26471256]
- Mineur YS, Obayemi A, Wigstrand MB, Fote GM, Calarco CA, Li AM, Picciotto MR. Cholinergic signaling in the hippocampus regulates social stress resilience and anxiety- and depression-like behavior. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110:3573–3578. [PubMed: 23401542]
- Morales-Perez CL, Noviello CM, Hibbs RE. X-ray structure of the human $\alpha 4\beta 2$ nicotinic receptor. *Nature.* 2016; 538:411–415. [PubMed: 27698419]
- Papke R, Wecker L, Stitzel J. Activation and inhibition of mouse muscle and neuronal nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J. Pharmacol. Exp. Ther.* 2010; 333:501–518. [PubMed: 20100906]
- Papke RL, Heinemann SF. Partial agonist properties of cytosine on neuronal nicotinic receptors containing the $\beta 2$ subunit. *Mol. Pharmacol.* 1994; 45:142–149. [PubMed: 8302273]
- Papke RL, Porter Papke JK. Comparative pharmacology of rat and human $\alpha 7$ nAChR conducted with net charge analysis. *Br. J. Pharmacol.* 2002; 137:49–61. [PubMed: 12183330]
- Paxinos G, Franklin K. *The Mouse Brain in Stereotaxic Coordinates.* Elsevier Academic Press; Boston: 2004.
- Peng C, Stokes C, Mineur YS, Picciotto MR, Tian C, Eibl C, Tomassoli I, Guendisch D, Papke RL. Differential modulation of brain nicotinic acetylcholine receptor function by cytosine, varenicline, and two novel bispidine compounds: Emergent properties of a hybrid molecule. *J. Pharmacol. Exp. Ther.* 2013; 347:424–437. [PubMed: 23959137]

- Picciotto MR, Zoli M, Lena C, Bessis A, Lallmand Y, LeNovere N, Vincent P, Pich EM, Brulet P, Changeux J-P. Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature*. 1995; 374:65–67. [PubMed: 7870173]
- Picciotto MR, Zoli M, Rimondini R, Léna C, Marubio LM, Pich EM, Fuxe K, Changeux JP. Acetylcholine receptors containing the $\beta 2$ subunit are involved in the reinforcing properties of nicotine. *Nature*. 1998; 391:173–177. [PubMed: 9428762]
- Rubinstein M, Low MJ. The smoking gun in nicotine-induced anorexia. *Cell Metab*. 2011; 14:145–147. [PubMed: 21803282]
- Salas R, Pieri F, De Biasi M. Decreased signs of nicotine withdrawal in mice null for the $\beta 4$ nicotinic acetylcholine receptor subunit. *J. Neurosci*. 2004; 24:10035–10039. [PubMed: 15537871]
- Sargent P. The Diversity of neuronal nicotinic acetylcholine receptors. *Annu. Rev. Neurosci*. 1993; 16:403–443. [PubMed: 7681637]
- Shen JX, Yakel JL. Functional $\alpha 7$ nicotinic ACh receptors on astrocytes in rat hippocampal CA1 slices. *J. Mol. Neurosci*. 2012; 48:14–21. [PubMed: 22351110]
- Sternson SM, Atasoy D. Agouti-related protein neuron circuits that regulate appetite. *Neuroendocrinology*. 2014; 100:95–102. [PubMed: 25402352]
- Stojakovic A, Espinosa EP, Farhad OT, Lutfy K. Effects of nicotine on homeostatic and hedonic components of food intake. *J. Endocrinol*. 2017; 235:R13–R31. [PubMed: 28814527]
- Stokes C, Papke RL. Use of an $\alpha 3\beta 4$ nicotinic acetylcholine receptor subunit concatamer to characterize ganglionic receptor subtypes with specific subunit composition reveals species-specific pharmacologic properties. *Neuropharmacology*. 2012; 63:538–546. [PubMed: 22580377]
- Winders SE, Grunberg NE. Effects of nicotine on body weight, food consumption and body composition in male rats. *Life Sci*. 1990; 46:1523–1530. [PubMed: 2355798]
- Wohleb ES, Wu M, Gerhard DM, Taylor SR, Picciotto MR, Alreja M, Duman RS. GABA interneurons mediate the rapid antidepressant-like effects of scopolamine. *J. Clin. Invest*. 2016; 126:2482–2494. [PubMed: 27270172]
- Yang L, Qi Y, Yang Y. Astrocytes Control Food Intake by Inhibiting AGRP Neuron Activity via Adenosine A1 Receptors. *Cell Rep*. 2015; 11:798–807. [PubMed: 25921535]
- Zhan C, Zhou J, Feng Q, Zhang J-E, Lin S, Bao J, Wu P, Luo M. Acute and long-term suppression of feeding behavior by POMC neurons in the brainstem and hypothalamus, respectively. *J. Neurosci*. 2013; 33:3624–3632. [PubMed: 23426689]
- Zhang X, van den Pol AN. Dopamine/Tyrosine Hydroxylase Neurons of the Hypothalamic Arcuate Nucleus Release GABA, Communicate with Dopaminergic and Other Arcuate Neurons, and Respond to Dynorphin, Met-Enkephalin, and Oxytocin. *J. Neurosci*. 2015; 35:14966–14982. [PubMed: 26558770]
- Zhang X, van den Pol AN. Hypothalamic arcuate nucleus tyrosine hydroxylase neurons play orexigenic role in energy homeostasis. *Nat. Neurosci*. 2016; 19:1341–1347. [PubMed: 27548245]
- Zoli M, Picciotto MR. Nicotinic regulation of energy homeostasis. *Nicotine Tob. Res*. 2012; 14:1270–1290. [PubMed: 22990212]

Figure 1. Expression of nAChR subunit mRNA in the hypothalamus and actively translated mRNAs isolated from AgRP or POMC neurons

A. Ethidium bromide gel visualization of nAChR subunit PCR products from an ARC enriched punch of ventral hypothalamus. B–E. TRAP-isolated nAChR subunit mRNA levels from AgRP (white) or POMC (grey) neurons normalized to the respective whole hypothalamic input fraction. Input and TRAP fractions were compared with paired t-tests. F–G. Direct comparison of TRAP-isolated α7 (F) and α4 (G) mRNA from AgRP and POMC cells. AgRP and POMC TRAP expression were compared with unpaired t-tests. Data are expressed as fold change from the reference group (Input B–E or POMC F–G), which is mathematically defined as 1 (noted with a dashed line). Box plots represent min, max, and quartiles, n=6–10, as noted. *p < 0.05, **p < 0.01, ***p < 0.001

Figure 2. Both AgRP and POMC cells in the ARC express cholinergic markers

A. TRAP-isolated ChAT and VACHT mRNA levels from AgRP (white) or POMC (grey) neurons normalized to the respective whole hypothalamic input fraction. Input and TRAP fractions were compared with paired t-tests. AgRP and POMC TRAP expression were compared with unpaired t-tests. B. Direct comparison of TRAP-isolated ChAT and VACHT mRNA from AgRP (white) and POMC (grey) cells. AgRP and POMC TRAP expression were compared with unpaired t-tests. Data are expressed as fold change from the reference group (Input for A or POMC for B), which is mathematically defined as one (noted with a dashed line). Box plots represent min, max, and quartiles, $n=7-10$, as noted. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

< 0.01 , *** $p < 0.001$ C. 60× image of neurons in the ARC; immunostaining for ChAT in red, immunostaining for POMC is in green, and a merge of these two channels to show overlap. Cells in which ChAT and POMC co-localize are marked with white arrowheads and cells that are ChAT positive, but POMC negative are marked with white arrows. 3V=3rd ventricle, scale bar = 50μm.

Figure 3. The effect of nicotine and cytosine on food intake in wild type mice and after cell type-selective knockdown of either the $\beta 4$ or $\alpha 7$ nAChR subunit

A. Two-hour food intake in wild type mice following saline (white), cytosine (1.5 mg/kg, grey), or nicotine (0.5 mg/kg, pink) IP injection. Drug order was randomized for a within-subjects analysis. One-way ANOVA compared cytosine and nicotine intake to saline intake. Box plots represent min, max, and quartiles; nicotine = pink, cytosine = grey, n = 11. B. Diagram of the AAV plasmid vector used to deliver shRNA knockdown constructs (top) for either $\beta 4$ or $\alpha 7$ nAChR subunit into the ARC (bottom). In the absence of Cre-recombinase cells will express both dsRed and eGFP, while in the presence of Cre-recombinase, the GFP coding DNA is excised, allowing the U6 promoter to drive shRNA expression in red only cells. C. Representative image of bilateral AAV expression in the ARC at 10× (top) and individual fluorophore expression (bottom) at 60×. Cells in which successful infection and recombination have taken place express dsRed, but not GFP (white arrows). Cells which are infected, but do not express Cre-recombinase express both dsRed and GFP and appear yellow in the merged image. 3V = 3rd Ventricle, ARC = Arcuate, ME = Median Eminence, 10× scale bar = 100μm, 60× scale bar = 50μm. D. Two-hour food intake in POMC-Cre (left) or AgRP-Cre (right) mice with shRNA knockdown of either $\beta 4$ or $\alpha 7$ nAChR subunit following saline (white), cytosine (3.0mg/kg, grey), or nicotine (1.0mg/kg, pink) IP injection. Drug order was randomized for a within subjects analysis. Two-way ANOVAs with planned

post hoc comparisons of cytosine and nicotine to saline intake within knockdown group were used. Box plots represent min, max, and quartiles; $n=7-12$, as noted. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Primer sequences used for quantitative and qualitative PCR

Target	Forward Primer 5' to 3'	Reverse Primer 5' to 3'
ChAT ^{\$}	CCATTGTGAAGCGGTTTGGG	GAGGCTGGCTGCAAACCTTA
Chn2 ^{*\$}	TCTACCCCGACGTCACCTAC	TAGAACACGAGCACGGTGAG
Chn3 [*]	ACTCCAAAAGCTGCAAGGAA	CAGAGCAGACAGGGACAACA
Chn3 ^{\$}	TGTCCTGTCTGTCTGTACCA	GCGAGCCTTGGCGAACAGGT
Chn4 [*]	CCAACTGGACTTCTGGGAAA	GGCGTAGGTGATGTCAGGAT
Chn4 ^{\$}	CCTGACTTGAAGACATCA	CACAGTACTGGATACTCC
Chn5 ^{*\$}	CTCCATGAAATTCGGCTCAT	CGCTCATGATTCCCATTTCT
Chn6 [*]	TGATCTTTGTACGCTGTCC	CTCCTGCCTCCTTTGTCTTG
Chn6 ^{\$}	ATTGGCAATCACGCAACTGG	AAGTCGCCGACAGCATTGTT
Chn7 ^{*\$#}	GCCTAAGTGGACCAGGATCA	CTCGGAAGCCAATGTAGAGC
Chnb2 ^{*\$#}	ATCCCTGCGTACTCATCAC	GTACTTCCCACCAGCGGTA
Chnb2 [^]	GTGGACGGGTGTACGCTTCATTG	GGTCGTGGCAGTGTAGTTCTGG
Chnb3 [*]	TGCACGCGATGACTAGTGTCTTCA	AGGAGTACAGGCCCAACAATGACA
Chnb3 ^{\$}	CCTACGGGTCTTCTTGCTC	TGGTCTGTCCATTCCACATCT
Chnb4 ^{*\$^}	CCTGCTCCTCGTCTCTGT	CAGGTTGTGTAGCGGGTTT
dsRed ^{^#}	TGAACGCCACTACTTCGAG	TGGAGCCGTACTGGAAGT
GFP ^{^#}	CAGAAGAACGGCATCAAGGT	TGGGTGCTCAGGTAGTGGTT
HPRT ^{\$#}	AACAAAGTCTGGCCTGTATCC	CCCCAAAATGGTTAAGGTTGC
Tbp1 [^]	AAAGGGAGAATCATGGACCAGAACAA	TGGACTAAAGATGGGAATTCCAGGAG
VChT ^{\$}	CGTTTCTCGAGCCACCATA	GTACGACGCAAGAGCTCACT

* Qualitative PCR

^{\$} bacTRAP[^] β 4 knockdown validation[#] α 7 knockdown validation

Table 2

Immunoprecipitation analysis of the nAChR subunit content in extracts obtained from whole hypothalamic or from more specific ARC-enriched hypothalamic punches.

	Hypothalamus (fmol/mg of protein)		Arcuate (fmol/mg of protein)		Hypothalamus (%Immunoprecipitation)		Arcuate (%Immunoprecipitation)		Significance HYP v ARC
	Mean $\pm$ SEM	n (pools of 10)	Mean $\pm$ SEM	n (pools of 20)	Mean $\pm$ SEM	n (pools of 10)	Mean $\pm$ SEM	n (pools of 20)	
[³ H]-Epi	54.03 $\pm$ 2.8	3	23.99 $\pm$ 0.47	2					
$\alpha 2$	2.60 $\pm$ 0.36	3	1.10 $\pm$ 0.30	2	4.90 $\pm$ 0.60	3	4.50 $\pm$ 1.30	2	
$\alpha 3$	5.20 $\pm$ 0.15	3	4.17 $\pm$ 0.58	3	9.70 $\pm$ 0.10	3	17.10 $\pm$ 2.10	3	*
$\alpha 4$	48.03 $\pm$ 1.4	3	18.05 $\pm$ 1.05	2	90.00 $\pm$ 2.20	3	75.20 $\pm$ 6.20	2	p=0.0720
$\alpha 5$	6.23 $\pm$ 0.83	3	2.80 $\pm$ 0.30	2	11.60 $\pm$ 1.20	3	11.80 $\pm$ 1.60	2	
$\alpha 6$	8.13 $\pm$ 1.19	3	2.15 $\pm$ 0.45	2	13.00 $\pm$ 2.60	4	9.00 $\pm$ 2.00	2	
$\beta 2$	49.93 $\pm$ 0.55	3	18.55 $\pm$ 0.75	2	90.40 $\pm$ 3.80	4	76.90 $\pm$ 5.10	2	p=0.1064
$\beta 3$	8.77 $\pm$ 0.90	3	4.70 $\pm$ 0.20	2	16.40 $\pm$ 1.70	3	19.70 $\pm$ 1.20	2	
$\beta 4$	3.03 $\pm$ 0.46	3	3.50 $\pm$ 1.05	3	6.10 $\pm$ 0.70	4	14.10 $\pm$ 4.20	3	p=0.0775

Unpaired t-tests comparing normalized hypothalamus and ARC signal

*
p < 0.05

Table 3

Immunoprecipitation analysis of the nAChR subunit content in samples obtained from whole hypothalamic extracts of wild type and $\beta 2$ knockout mice

	WT Hypothalamus (fmol/mg of protein)		KO Hypothalamus (fmol/mg of protein)		WT Hypothalamus (% Immunoprecipitation)		KO Hypothalamus (% Immunoprecipitation)		Significance WT v KO
	Mean $\pm$ SEM	n (pools of 10)	Mean $\pm$ SEM	n (pools of 10)	Mean $\pm$ SEM	n (pools of 10)	Mean $\pm$ SEM	n (pools of 10)	
^3H -Epi	54.37 $\pm$ 2.01	4	4.29 $\pm$ 1.02	3					
$\alpha 2$	2.60 $\pm$ 0.36	3	0.45 $\pm$ 0.25	2	4.77 $\pm$ 0.51	3	14.38 $\pm$ 8.13	2	
$\alpha 3$	5.20 $\pm$ 0.15	3	1.33 $\pm$ 0.29	3	9.52 $\pm$ 0.18	3	31.83 $\pm$ 5.40	3	*
$\alpha 4$	47.225 $\pm$ 1.28	4	0.63 $\pm$ 0.34	3	86.68 $\pm$ 3.19	4	13.07 $\pm$ 3.83	3	***
$\alpha 5$	6.23 $\pm$ 0.83	3	0.20 $\pm$ 0.00	2	11.40 $\pm$ 0.96	3	6.88 $\pm$ 0.63	2	*
$\alpha 6$	6.975 $\pm$ 1.43	4	0.30 $\pm$ 0.10	3	12.89 $\pm$ 2.66	4	7.25 $\pm$ 0.52	3	
$\beta 2$	49.93 $\pm$ 0.55	3	0.09 $\pm$ 0.05	3	92.23 $\pm$ 4.02	3	2.92 $\pm$ 1.50	3	***
$\beta 3$	8.77 $\pm$ 0.90	3	0.70 $\pm$ 0.20	2	16.23 $\pm$ 1.93	3	20.65 $\pm$ 5.65	2	
$\beta 4$	3.35 $\pm$ 0.46	4	1.5 $\pm$ 0.10	3	6.02 $\pm$ 0.73	4	38.27 $\pm$ 6.64	3	**

Unpaired t-tests comparing normalized WT and KO signal

* $p < 0.05$,

** $p < 0.01$,

*** $p < 0.001$

Table 4

Percent of total binding after sequential immunoprecipitations, first with a target subunit antibody (listed first), then $\beta 4$ antibody (listed second)

	Mean $\pm$ SEM	n (pools of 20)	Significance v $\beta 4$ - $\beta 4$
$\beta 4$ - $\beta 4$	1.2 $\pm$ 0.00	2	
$\alpha 2$ - $\beta 4$	7.25 $\pm$ 1.95	2	*
$\alpha 3$ - $\beta 4$	3.08 $\pm$ 0.00	1	
$\alpha 4$ - $\beta 4$	3.81 $\pm$ 1.09	2	
$\alpha 5$ - $\beta 4$	8.69 $\pm$ 1.12	2	*
$\alpha 6$ - $\beta 4$	8.39 $\pm$ 0.81	2	*
$\beta 2$ - $\beta 4$	2.60 $\pm$ 1.10	2	
$\beta 3$ - $\beta 4$	9.60 $\pm$ 1.00	2	*

One way ANOVA with Holm-Sidak post-hoc comparisons between $\beta 4$ - $\beta 4$ and other conditions.

*
p < 0.05