Figure 2: Mean plasma itraconazole concentration–time profile (day 15).

Disclosures. J. Lindsay, Mayne Pharma: Consultant, Consulting fee. S. Mudge, Mayne Pharma: Employee, Salary.

1355. Global Activity of Imipenem–Relebactam and Comparators Against Clinical Gram-Negative Pathogens – SMART 2017

Sibylle Lob, PhD¹; Krystyna Kazmierczak, PhD¹; Daryl Hoban, PhD¹; Meredith Hackel, PhD¹; Katherine Young, MS²; Mary Motyl, PhD² and Dan Sahm, PhD³; ¹IHMA, Inc., Schaumburg, Illinois, ²Merck & Co., Inc., Kenilworth, New Jersey, ³International Health Management Associates, Inc., Schaumburg, Illinois

Session: 144. Novel Agents

Friday, October 5, 2018: 12:30 PM

Background. Relebactam (REL), formerly MK-7655, is a β -lactamase inhibitor of class A and C β -lactamases that is in development in combination with imipenem (IMI). In this study, we evaluated the activity of IMI/REL against recent clinical isolates of Gram-negative bacilli (GNB) collected globally as part of the SMART surveillance program.

Methods. In 2017, 188 hospitals in 54 countries each collected up to 100 consecutive Gram-negative aerobic or facultatively anaerobic pathogens from lower respiratory tract infections, 75 from intra-abdominal infections, and 75 from urinary tract infections. MICs were determined for 41,319 GNB, including 30,864 *Enterobacteriaceae* and 6,933 *P. aeruginosa* isolates, using CLSI broth microdilution and interpreted with CLSI breakpoints; for comparison purposes, IMI susceptible breakpoints were applied to IMI/REL.

Results. Susceptibilities to IMI/REL and comparators of the 10 most commonly found *Enterobacteriaceae* species and *P. aeruginosa* are shown below.

Species	n	IMI/REL	IMI	FEP	CAZ	P/T	ATM	CIP	AMK	CST
<i>E. coli</i>	14176	99.5	99.0	75.8	79.0	91.0	76.1	64.4	99.0	99.6
<i>K. pneumoniae</i>	7208	92.5	86.9	59.0	58.2	68.6	58.5	60.4	93.8	95.1
<i>P. mirabilis</i>	1670	63.1	63.6	88.3	90.7	97.5	94.3	65.9	96.1	0.4
<i>E. cloacae</i>	1601	96.7	93.7	75.3	62.2	72.1	63.2	81.8	97.7	93.1
<i>S. marcescens</i>	1129	70.2	51.6	91.1	91.6	92.2	90.0	89.6	98.1	5.2
<i>K. oxytoca</i>	1012	99.2	98.5	94.3	93.9	86.8	86.9	94.1	99.5	99.2
<i>K. aerogenes</i>	828	97.5	88.3	92.6	72.1	75.0	75.4	93.8	98.9	98.6
<i>C. freundii</i>	568	98.6	96.7	90.7	70.8	79.6	71.8	87.0	99.1	99.8
<i>M. morganii</i>	538	32.0	5.6	94.4	84.4	97.4	94.1	76.8	99.3	0.6
<i>C. koseri</i>	402	98.5	96.0	98.3	97.8	97.3	96.5	98.5	99.8	100
<i>P. aeruginosa</i>	6933	88.8	68.6	74.6	73.4	68.3	63.9	74.8	90.7	99.5

IMI, imipenem; REL, relebactam; FEP, cefepime; CAZ, ceftazidime; P/T, piperacillin-tazobactam; ATM, aztreonam; AMK, amikacin; CIP, ciprofloxacin; CST, colistin; NA, breakpoint not available

IMI/REL showed activity >90% against seven of the top 10 *Enterobacteriaceae* species, typically ~5 to 35 percentage points higher than the β -lactam comparators, and it was active against 89% of *P. aeruginosa*, ~15 to 25 percentage points higher than the β -lactam comparators. Only amikacin and colistin showed similar or higher activity for most species, with colistin showing little activity against Proteaceae and *Serratia*.

Conclusion. IMI/REL could provide an important treatment option against infections with Gram-negative pathogens, especially since amikacin and colistin are associated with significant morbidity, including nephrotoxicity and ototoxicity, and amikacin is typically used in combination with another antibiotic.

Disclosures. S. Lob, IHMA, Inc.: Employee, Salary. Merck: Consultant, Consulting fee. K. Kazmierczak, Merck: Consultant, Consulting fee. IHMA, Inc.: Employee, Salary. D. Hoban, IHMA, Inc.: Employee, Salary. Merck: Consultant, Consulting fee. M. Hackel, IHMA, Inc.: Employee, Salary. Merck: Consultant, Consulting fee. K. Young, Merck: Employee and Shareholder, Dividends and Salary. M. Motyl, Merck: Employee and Shareholder, Dividends and Salary. D. Sahm, IHMA, Inc.: Employee, Salary. Merck: Consultant, Consulting fee.

1356. Improvement in Quality of Life for Adults With Acute Bacterial Skin and Skin Structure Infections Following Treatment With Omadacycline or Linezolid

Evan Tzanis, BA¹; Surya Chitra, PhD, MBA¹; Marla Curran, DrPH²; Paul McGovern, MD¹; Jordan Hinahara, BA³ and Thomas Goss, PharmD⁴; ¹Paratek Pharmaceuticals, Inc., King of Prussia, Pennsylvania, ²Paratek Pharmaceuticals, King of Prussia, Pennsylvania, ³Boston Healthcare Associates, Boston, Massachusetts, ⁴Boston Healthcare Associates, Inc., Boston, Massachusetts

Session: 144. Novel Agents

Friday, October 5, 2018: 12:30 PM

Background. The appearance of multidrug-resistant Gram-positive bacteria is a major challenge in clinical care. Omadacycline is the first aminomethylcycline antibiotic (semisynthetic compounds related to tetracyclines) in late-stage clinical development for acute bacterial skin and skin structure infections (ABSSSI), and demonstrates potent *in vitro* activity against many pathogens.

Methods. Seven hundred thirty-five patients were enrolled in the OASIS-2 randomized controlled trial comparing omadacycline and linezolid for the treatment of adult subjects with ABSSSI known or suspected to be due to a Gram-positive pathogen, with 368 and 367 enrolled in each group, respectively. Subjects completed the 36-Item Short Form Health Survey Version 2 (SF-36v2), a validated questionnaire on physical and mental health, at both screening and post-treatment evaluation. Results of the SF-36v2 were analyzed in accordance with established norm-based standards for the survey (Ware 2000) for the intention-to-treat population.

Results. Subjects who received omadacycline experienced a 3.25 point mean improvement in overall physical health ($P < 0.001$, Figure 1) and reported significant improvements across all but one component parameter of overall physical and mental health, including physical functioning, bodily pain, role physical, vitality, role emotional, mental health, and social functioning (Figure 2). In contrast, while overall physical health improved for subjects who received linezolid, the improvement in vitality, role emotional, mental health, and general health was not significant (Figure 2). Although omadacycline achieved greater increase from baseline than linezolid across all domains analyzed, the difference in scores was not statistically significant at the $P < 0.05$ level (Figure 1).

Conclusion. Omadacycline provides significant improvement in the physical component of quality of life over baseline for adult subjects with ABSSSI known or suspected to be due to a Gram-positive pathogen. Although the OASIS-2 trial was neither designed nor powered to measure differences in quality of life following treatment, trends identified in this analysis merit further investigation.

References

- Ware JE. SF-36 Health Survey Update. SPINE 2000; 25(24): 3130–3139.

Figure 1.

Figure 2.

Disclosures. E. Tzanis, Paratek Pharmaceuticals: Employee, Salary. M. Curran, Paratek Pharmaceuticals: Employee, Salary. P. McGovern, Paratek Pharmaceuticals: Employee, Salary. J. Hinahara, Paratek Pharmaceuticals: Consultant, Consulting fee. T. Goss, Paratek Pharmaceuticals: Consultant, Consulting fee.

1357. A Combination of Itraconazole and Amiodarone Is Highly Effective Against *Trypanosoma cruzi* Infection of Human Stem Cell-Derived Cardiomyocytes

Gabriele Sass, PhD¹; Roy Madigan, DVM²; Adriana Bozzi, PhD^{1,3,4,5,6}; Nazish Sayed, MD, PhD^{3,4,5}; Joseph Wu, MD^{3,4,5} and David Stevens, MD^{1,7}; ¹California Institute for Medical Research, San Jose, California, ²Animal Hospital of Smithson Valley,

Smithson Valley, Texas, ³Department of Medicine, Division of Cardiology, Stanford University, Stanford, California, ⁴Department of Radiology, Department of Medicine, Stanford University, Stanford, California, ⁵Institute of Stem Cell Biology and Regenerative Medicine, Stanford University, Stanford, California, ⁶Centro de Pesquisas René Rachou, FIOCRUZ, Belo Horizonte, Brazil, ⁷Division of Infectious Diseases and Geographic Medicine, Stanford University, Stanford, California

Session: 144. Novel Agents
Friday, October 5, 2018: 12:30 PM

Background. *Trypanosoma cruzi* is the etiologic agent of Chagas disease, which can result in severe cardiomyopathy. *Trypanosoma cruzi* is endemic to the Americas, and of particular importance in Latin America. In the United States and other nonendemic countries, rising case numbers have been observed. The only drugs available so far are benznidazole and nifurtimox, which have limited efficacy during chronic infection. We repurposed itraconazole, originally an antifungal, in combination with amiodarone, an antiarrhythmic, with the goal to interfere with Tc infection. Both drugs inhibit sterol synthesis, while amiodarone also inhibits calcium metabolism of *Trypanosoma cruzi*.

Methods. Human pluripotent stem cells (HiPSC) were differentiated to cardiomyocytes (HiPSC-CM). Vero cells or HiPSC-CM were infected with the *T. cruzi* trypomastigotes Y strain in the presence of itraconazole and/or amiodarone. After 48 hours, infection and multiplication were evaluated by Giemsa stain. Benznidazole was used as a reference compound. Cell viability was verified by XTT assay.

Results. Itraconazole and amiodarone showed dose-dependent interference with *T. cruzi* infection of Vero cells or HiPSC-CM. The combination of itraconazole and amiodarone was more potent than the single substances, or benznidazole at therapeutic concentrations, without affecting host cell metabolism. In addition to effects on infection, itraconazole, or amiodarone affected *T. cruzi* multiplication. Here, itraconazole/amiodarone combinations were more potent than either alone, both, in Vero cells, and HiPSC-CM.

Conclusion. Our *in vitro* data suggest that a combination of itraconazole and amiodarone might serve as an effective new treatment option for Chagas disease, particularly cardiac involvement, in human and animal patients.

Disclosures. All authors: No reported disclosures.

1358. *In vitro* Activity of Ceftazidime-Avibactam and Comparator Agents Against *Pseudomonas aeruginosa* Causing Intra-Abdominal, Lower Respiratory, and Urinary Tract Infections Collected in Latin America as Part of the INFORM Global Surveillance Program, 2012–2016

Mark Wise, PhD¹; Krystyna Kazmierczak, PhD¹; Gregory G. Stone, PhD² and Dan Sahm, PhD³; ¹IHMA, Inc., Schaumburg, Illinois, ²Pfizer, Inc., New York, New York, ³International Health Management Associates, Inc., Schaumburg, Illinois

Session: 144. Novel Agents
Friday, October 5, 2018: 12:30 PM

Background. The non-β-lactam β-lactamase inhibitor avibactam (AVI) is active against class A, C, and some class D β-lactamases, in combination with ceftazidime (CAZ) has been approved by the FDA and EMA for treatment of intra-abdominal infections (IAI), lower respiratory tract infections (LRTI), and urinary tract infections (UTI). This study reports on the *in vitro* activity of (CAZ-AVI) and comparators vs. *P. aeruginosa* collected from IAIs, LRTIs, and UTIs in Latin America as part of the INFORM surveillance study from 2012 to 2016.

Methods. For INFORM surveillance over 2012–2016 in Latin America, 1,595 nonduplicate *P. aeruginosa* isolates linked to IAIs, LRTIs, and UTIs were collected from 26 clinical sites in six countries. Susceptibility testing was done using broth microdilution according to CLSI guidelines and using CLSI 2018 breakpoints. CAZ was tested with AVI at a fixed concentration of 4 mg/mL. Meropenem (MEM) nonsusceptible organisms were screened for β-lactamase genes by PCR.

Results. Among the full collection of *P. aeruginosa*, CAZ-AVI showed consistently higher % susceptibilities than all comparators except for colistin (CST) for all infection sources. The addition of AVI to CAZ resulted in an increase in susceptibility ranging from 14.2% (IAI) to 19.5% (UTI). Against the non-metallo-β-lactamase (MBL) harboring subset, CAZ-AVI showed extremely potent activity (MIC₉₀, 8 mg/mL) for all infection sources. In this subset, the activity of CAZ-AVI approached that of colistin for IAIs (susceptibility of 93.3% vs. 96.4%, respectively).

Organism/Phenotype/Infection Source (n)	Drug (MIC90 [μg/mL]/% Susceptible)							
	CAZ-AVI	CAZ	MEM	CST	CAZ-AVI	CAZ	MEM	CST
<i>P. aeruginosa</i> , All sources (1595) ^a	16	86.0	64	68.7	>8	61.9	2	96.2
IAI (295)	32	85.1	64	70.9	>8	62.4	2	96.7
LRTI (942)	16	86.8	128	69.3	>8	60.7	2	96.3
UTI (358)	32	84.6	64	65.1	>8	64.8	2	95.8
<i>P. aeruginosa</i> MBL-negative, All sources (1488) ^b	8	91.9	64	73.5	>8	66.1	2	96.1
IAI (269)	8	93.3	64	77.7	>8	68.0	2	96.4
LRTI (889)	8	91.8	64	73.3	>8	64.0	2	96.0
UTI (330)	8	91.2	64	70.3	>8	70.3	2	96.2

% susceptible defined using CLSI 2018 breakpoints.

^aCST not tested in 2012 or 2013; totals for IAI, n = 215; LRTI, n = 721; UTI, n = 286.

^bCST not tested in 2012 or 2013; totals for IAI, n = 196; LRTI, n = 677; UTI, n = 260.

Conclusion. CAZ-AVI demonstrated very good *in vitro* activity against *P. aeruginosa* isolates, especially those without MBLs. More isolates were susceptible to CAZ-AVI than to MEM for all infection types.

Disclosures. M. Wise, Pfizer Inc.: Consultant, Consulting fee. IHMA, Inc.: Employee, Salary. K. Kazmierczak, Pfizer Inc.: Consultant, Consulting fee. IHMA, Inc.: Employee, Salary. G. G. Stone, Pfizer Inc.: Employee, Salary. AstraZeneca: Former Employee and Shareholder, Salary. D. Sahm, Pfizer Inc.: Consultant, Consulting fee. IHMA, Inc.: Employee, Salary.

1359. Activity of Meropenem/Nacubactam Combination Against Gram-Negative Clinical Isolates: ROSCO Global Surveillance 2017

Rusudan Okujava, PhD¹; Fernando Garcia-Alcalde, PhD¹; Andreas Haldemann, PhD²; Claudia Zampaloni, PhD³; Ian Morrissey, PhD²; Sophie Magnet, PhD²; Nimmi Kothari, MSc²; Ian Harding, PhD³ and Kenneth Bradley, PhD³; ¹Roche Pharma Research and Early Development, Immunology, Inflammation and Infectious Diseases, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland, ²IHMA Europe Sàrl, Monthey, Switzerland, ³Micron Research Ltd., Ely, Cambs, UK

Session: 144. Novel Agents
Friday, October 5, 2018: 12:30 PM

Background. Nacubactam (NAC, OP0595, RG6080) is a novel member of the diacyclicocloctane inhibitor family with a dual mode of action, acting as a β-lactamase inhibitor and an antibacterial agent by means of PB2 inactivation. NAC restores and extends the activity of β-lactam antibiotics, such as meropenem (MEM), when used in combination against a variety of carbapenem-resistant *Enterobacteriaceae* (CRE). The first year results of the ROSCO surveillance study for MEM/NAC against contemporary clinical isolates are presented here.

Methods. Isolates (n = 4,695) collected in 2017 from 50 sites in the United States and European hospitals included 30 different species of *Enterobacteriaceae* (EB, n = 3,306), *Pseudomonas* spp. (n = 960) and *Acinetobacter* spp. (n = 429). The predominant species of EB are shown in figure below. MICs were determined by broth microdilution following CLSI methodology for MEM/NAC at a fixed 1:1 ratio (w/w) and by titrating MEM with a constant concentration of NAC at 4 mg/L. Results were compared with MIC values of MEM and NAC alone and standard of care antibiotics, including ceftazidime/avibactam (CAZ/AVI).

Results. MIC_{50/90} for MEM, NAC, and MEM/NAC against all EB isolates and by species are shown in the figure below. NAC alone displayed a bimodal MIC distribution for EB, with a prominent separation at ≤4 mg/L. MEM/NAC 1:1 inhibited 99.5, 99.7, and 99.9% of the 3,306 EB isolates tested, at ≤2, ≤4, and ≤8 mg/L, respectively; while MEM inhibited 96.5, 96.8, and 97.3% of the isolates at the same concentrations. Of 117 (3.5% of total EB) MEM nonsusceptible (by EUCAST) and multidrug resistant (MDR, by Magiorakos AP, et al., 2012) EB, 87.2, 92.3, and 96.6% were inhibited by MEM/NAC 1:1 at ≤2, ≤4, and ≤8 mg/L, respectively. Additionally, MEM/NAC 1:1 displayed MIC ≤8 mg/L for 33 out of 37 CAZ/AVI-resistant MDR EB isolates. MEM/NAC had a similar activity to MEM alone against *Pseudomonas* spp. and *Acinetobacter* spp.

Species	MIC _{50/90} (mg/L)				
	MEM	MEM/NAC 1:1	MEM/NAC (4 mg/L)	CAZ/AVI (4 mg/L)	NAC
<i>Enterobacteriaceae</i> , n=3306	0.03/0.06	0.03/0.06	≤0.004/0.03	0.12/0.5	2/>32
<i>E. coli</i> , n=956	0.03/0.03	0.015/0.03	≤0.004/≤0.004	0.08/0.25	1/2
<i>K. pneumoniae</i> , n=892	0.03/4	0.03/0.5	≤0.004/0.015	0.12/0.5	4/>32
<i>S. marcescens</i> , n=414	0.06/0.12	0.06/0.12	0.03/0.06	0.25/0.5	>32/>32
<i>E. cloacae</i> , n=268	0.03/0.12	0.03/0.12	≤0.004/≤0.004	0.25/1	2/4
<i>C. freundii</i> , n=108	0.03/0.06	0.03/0.06	≤0.004/0.008	0.12/0.5	2/32
<i>Pseudomonas</i> spp., n=960	0.5/8	0.5/8	0.25/8	2/8	>32/>32
<i>Acinetobacter</i> spp., n=429	1/>32	1/>32	1/>32	16/>32	>32/>32

MIC for MEM/NAC combination is recorded as MEM concentration
MEM = meropenem, NAC = nacubactam, CAZ/AVI = ceftazidime/avibactam (avibactam fixed at 4 mg/L)

Conclusion. MEM/NAC combination shows excellent *in vitro* activity against current clinical EB isolates and the potential to extend MEM activity to MDR, MEM nonsusceptible and CAZ/AVI-resistant isolates, which supports the continued clinical development of MEM/NAC for infections caused by CREs. This project has been funded in part under HHS BARDA Contract HHSO100201600038C.

Disclosures. R. Okujava, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Employee, Salary. F. Garcia-Alcalde, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Employee, Salary. A. Haldemann, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Employee, Salary. C. Zampaloni, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Employee, Salary. I. Morrissey, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Research Contractor, Contracting fee to IHMA. S. Magnet, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Research Contractor, Contracting fee to IHMA. N. Kothari, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Research Contractor, Contracting fee to IHMA. I. Harding, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Research Contractor, Contracting fee to Micron. K. Bradley, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., Basel, Switzerland: Employee, Salary.

1360. Antimicrobial Activity of Cefepime in Combination with VNRX-5133 Against a Global Collection of *Enterobacteriaceae* Including Resistant Phenotypes
Meredith Hackel, Ph.D¹ and Dan Sahm, PhD²; ¹IHMA, Inc., Schaumburg, Illinois, ²International Health Management Associates, Inc., Schaumburg, Illinois

Session: 144. Novel Agents
Friday, October 5, 2018: 12:30 PM

Background. VNRX-5133 is a novel cyclic boronate-based broad-spectrum β-lactamase inhibitor with potent and selective direct inhibitory activity against both serine- and