Furthermore, FEP HSR + 1/8th VNRX-5133 HSR resulted in ≥ 1 -log reduction in the initial bacterial burden in 16 out of 24 isolates.

Conclusion. FEP/VNRX-5133 combination showed potent *in vivo* efficacy against serine β -lactamase-producing Gram-negative isolates. The extent of bacterial killing achieved with 1/8th VNRX-5133 HSR attested to the robustness of the inhibitor activity. These data support the consideration of FEP/VNRX-5133 combination for the treatment of serious infections due to these organisms in clinical trials.

Disclosures. D. P. Nicolau, VenatoRx Pharmaceuticals, Inc.: Grant Investigator, Research grant

1406. Augmented Renal Clearance Using Aminoglycoside Population-Based Pharmacokinetic Modeling with Bayesian Estimation in Children in the Pediatric Intensive Care Unit

Sean Avedissian, PharmD.¹; Nathaniel Rhodes, PharmD, MSc²; Yuna Kim, BS³; Josh Valdez, BS²; John Bradley, MD, FAAP⁴ and Jennifer Le, PharmD, MAS, FCCP, FCSHP, BCPS-ID⁵; ¹Pharmacy Practice, Midwestern University Chicago College of Pharmacy/Northwestern Memorial Hospital, Downers Grove, Illinois, ²Department of Pharmacy, Northwestern Medicine, Chicago, Illinois, ³University of California San Diego Skaggs School of Pharmacy, San Diego, California, ⁴Pediatric Infectious Disease, University of California San Diego, San Diego, California, ⁵Pharmacy/Infectious Diseases, University of California, San Diego Skaggs School of Pharmacy, La Jolla, California

Session: 145. PK/PD Studies

Friday, October 5, 2018: 12:30 PM

Background. Augmented renal clearance (ARC) in critically ill pediatric patients has been evaluated in limited studies. We evaluated ARC using clearance of aminoglycosides (CL_{AMINO}) derived from population-based pharmacokinetic modeling.

Methods. A retrospective, cohort study was conducted at two pediatric hospitals in patients who received aminoglycosides from 1999 to 2016. ARC was defined as a CL_{AMINO} of ≥ 130 mL/minute/1.73 m² within the first 24 hours of therapy. Pharmacokinetic (PK) models with nonparametric parameter estimation were constructed using Pmetrics in R, with the ultimate model selected by Akaike score and rule of parsimony. Covariate modifiers considered included: age, total body weight (TBW), serum creatinine (SCr) and sex. Noncompartmental analysis was performed on the Bayesian posteriors from the first dose to generate CL_{AMINO} within the first 24 hours and other PK exposure metrics (i.e., area under the curve for first 24 hours [AUC₂₄], maximum concentration [C_{MAX}]). Summary of patient demographics and statistical analysis were performed using GraphPad Prism version 7.

Results. ARC was identified in 34 of 117 (29%) subjects using 275 aminoglycoside serum concentrations. A two-compartment model fit the data well (See Figure 1: Population [a], Bayesian [b]). Allometric scaling of CL_{AMINO} utilized a fixed exponent of 0.75 and volume of distribution (VD) scaling utilized a fixed exponent of 1 in the final model. The final population model for CL_{AMINO} (L/hour) was $3.45 \times (TBW/40)^{0.75} + 0.05 \times 10^{(SCr/AGE)}$ and VD was $10.64 \times (TBW/40)^1$. Median age and baseline SCr were similar in those with and without ARC (13 [IQR 10–16] vs. 11.0 [5.0–15.0] years, $P = 0.11$, and 0.37 [0.27–0.49] vs. 0.38 [0.28–0.50] mg/dL, $P = 0.67$, respectively). Median TBW was found to be significantly higher in those with vs. without ARC (44.9 [26.9–61.7] vs. 34 [17.6–54.9] kg $P = 0.04$). Median 24 hours CL_{AMINO} was also found to be significantly higher in those with vs. without ARC (147.3 [138.7–163.9] vs. 94.5 [79.4–112.9], mL/minute/1.73 m², $P < 0.001$). Patients with vs. without ARC had significantly lower AUC₂₄ and C_{MAX} (40.7 [33.3–54.4] vs. 55.7 [46.7–66.4] mg hour/L, $P \leq 0.001$ and 5.06 [4.11–6.76] vs. 6.32 [5–7.44], μ g/mL, $P = 0.01$).

Conclusion. The incidence of ARC observed was similar to adult studies. Patients that exhibited ARC had lower AUC₂₄ and C_{MAX}; thus, higher doses may be warranted.

Disclosures. All authors: No reported disclosures.

1407. Disproportionality Analysis of Safety with Nafcillin and Oxacillin with the FDA Adverse Event Reporting System (FAERS)

Tristan T. Timbrook, PharmD, MBA, BCPS¹; Jesse Sutton, PharmD² and Emily Spivak, MD, MHS³; ¹University of Utah Health Care, Salt Lake City, Utah,

²George E. Wahlen Department of Veterans Affairs Medical Center, Salt Lake City, Utah and ³Internal Medicine, University of Utah Health, Salt Lake City, Utah

Session: 145. PK/PD Studies

Friday, October 5, 2018: 12:30 PM

Background. Antistaphylococcal penicillins including oxacillin and nafcillin are among the drugs of choice for severe and invasive MSSA infections. While alternative agents such as cefazolin are associated with improved safety compared with antistaphylococcal penicillins, comparative safety data between individual antistaphylococcal penicillins is limited and has shown possible improved safety with oxacillin among adults. The objective of this study was to determine the relative adverse events (AEs) reporting for these agents among the FDA Adverse Event Reporting System (FAERS) database.

Methods. We reviewed adverse events reports from the FAERS database from Q4/2003-Q1/2018 and performed a disproportionality analysis of safety events for nafcillin and oxacillin including ADEs related to acute kidney injury (AKI), rash, hypokalemia, and hepatotoxicity. Measures of association evaluated included reporting odds ratio (ROR) and proportion reporting ratio (PRR).

Results. Reports of AKI were substantially more common with nafcillin (PRR 23.2, 95% CI 18.4–29.3) than oxacillin (PRR 10.9, 95% CI 7.6–15.5). Rash was slightly higher with nafcillin than oxacillin (PRR 3.7, 95% CI 2.8–5.1 vs. 2.6, 95% CI 1.8–3.8). Hypokalemia was substantially more common with nafcillin than oxacillin (PRR 11.8, 95% CI 7.1–19.4 vs. 1.6, 95% CI 0.4–6.4). Hepatotoxicity was slightly higher reported among nafcillin than oxacillin (PRR 25.3, 95% CI 14.5–44.3 vs. 12.9, 95% CI 5.8–28.6). Similar observations were seen with RORs (Figure 1).

Figure 1. Reporting Odds Ratios of AEs with Nafcillin and Oxacillin

Conclusion. Oxacillin may be associated with overall improved safety compared with nafcillin based on reporting signals from FAERS. Our results support previous limited observational data. With the likely equal efficacy of these agents, clinicians may want to consider prescribing oxacillin over nafcillin if an antistaphylococcal penicillin is indicated for an invasive MSSA infection. However, given the limitations of reporting systems, further evaluation is warranted.

Disclosures. All authors: No reported disclosures.

1408. Population Pharmacokinetic (PK) Model to Describe Epithelial Lining Fluid (ELF) Penetration of ASN-1 and ASN-2 after ASN100 Administration to Healthy Subjects

Scott A. Van Wart, PhD, MS¹; Christopher Stevens, MD²; Zoltan Magyarics, MD, PhD³; Steven A. Luperchio, PhD, CMPP²; Christopher M. Rubino, PharmD¹ and Paul G. Ambrose, PharmD, FIDSA¹; ¹ICPD, Schenectady, New York, ²Arsanis, Inc., Waltham, Massachusetts, ³Arsanis Biosciences GmbH, Vienna, Austria

Session: 145. PK/PD Studies

Friday, October 5, 2018: 12:30 PM

Background. ASN100 is a combination of two co-administered fully human monoclonal antibodies (mAbs), ASN-1 and ASN-2, that together neutralize the six cytotoxins critical to *S. aureus* pneumonia pathogenesis. ASN100 is in development for prevention of *S. aureus* pneumonia in mechanically ventilated patients. A population PK model was developed to characterize the time-course of ASN-1 and ASN-2 in ELF following intravenous administration of ASN100 in healthy subjects.

Methods. A total of 42 healthy subjects received a single dose of ASN-1 or ASN-2 alone (200–4,000 mg) or ASN100 (3,600 or 8,000 mg; 1:1 ratio of ASN-1:ASN-2). All subjects contributed 13–17 serum samples for ASN-1/ASN-2 assay. Twelve subjects contributed 2 bronchoalveolar lavage (BALF) samples each for ELF concentration assay (Day 1 or 2 and Day 8 or 30 after dosing). A previously reported, linear, two-compartment population PK model for serum [ID Week 2017, Poster #1849] was expanded and fit to the ELF concentration–time data. Sequential analysis was used to fix serum PK as the driver for ELF PK; only those parameters controlling transfer into and out of the ELF were fit.

Results. An effect-site model adequately described the time-course of ELF concentrations. To allow for estimation of interindividual variability in the elimination from ELF, residual variability in ELF was fixed to that previously estimated for the serum PK data. Separate rate constants for transfer from serum to ELF were estimated for the 3,600 and 8,000 mg ASN100 dose groups to reflect the less than dose-proportional increase in ELF concentrations for both ASN-1 and ASN-2. Goodness-of-fit plots did not reveal any appreciable biases. A visual predictive check indicated that the model could adequately capture the observed data (Figure 1). Predicted ELF