Republic of (South), ³Pediatrics, Seoul National University College of Medicine, Seoul, Korea, Republic of (South), ⁴Pediatrics, Seoul National University Bundang Hospital, Seongnam, Korea, Republic of (South), ⁵Department of Pediatrics, Seoul National University Children's Hospital, Seoul, Korea, Republic of (South)

Session: 247. Pediatric Bacterial Infections

Saturday, October 6, 2018: 12:30 PM

Background. This study investigated the genetic structure of *Streptococcus pneumoniae* isolates from invasive pneumococcal disease (IPD) in Korean children after national immunization program (NIP) of extended-valency pneumococcal conjugate vaccines (PCVs) in Korea from 2014 to 2017.

Methods. Invasive isolates were collected from 23 hospitals throughout Korea. IPD cases were identified by isolating pneumococci from normally sterile sites. Each isolate was analyzed using standard microbiological techniques, Quellung reaction, multilocus sequence typing, and antimicrobial susceptibility testing. eBURST v3 software was used to estimate the relationships among the isolates and to assign the strains to a clonal complex (CC).

Results. Ninety-two pneumococcal isolates were analyzed. The source of isolates were blood (77), cerebrospinal fluid (7), pleural fluid (2), joint fluid (2), deep tissue abscess (2), and peritoneal fluid (2). A total of 38 STs and 17 singletons were assigned. Ten clonal complexes were identified: CC320, CC81, CC166, CC439, CC558, CC880, CC3280, CC9395, CC180, and CC310. New STs were assigned: ST13552, ST13553, ST13554, and ST13602. The serotypes were mostly non-vaccine type (NVTs) (82.6%). The most prevalent STs were ST11189 (17.4%, $n = 16$, all serotype 10A), ST6945 (10.9%, $n = 10$, all serotype 12F), ST166 (9.8%, $n = 9$, serotype 11A [22.2%, $n = 2$], 13 [22.2%, $n = 2$], 15B/C [22.2%, $n = 2$], and 23A [33.3%, $n = 3$]), and ST320 (6.5%, $n = 6$, all serotype 19A). Major CCs identified were CC166 (11.9%), CC320 (10.9%), and CC81 (10.9%). Serotypes constituting CC81 and CC166 were all NVTs except 6A ($n = 1$) and 23F ($n = 1$) in CC81. CC320 consisted of 19A ($n = 9$) and 19F ($n = 1$). The relative proportion of NVTs was 61.3% in major CCs. All major three CCs showed multi-drug resistance. Nonsusceptibility to penicillin (80%) and cefotaxime (100%) was the highest in CC320 (serotype 19A [$n = 9$, 90.0%] and 19F [$n = 1$, 10.0%]).

Conclusion. The introduction of extended-valency PCVs has resulted in the change of genetic structure of isolates from IPD of Korean children. In particular, two common CCs (CC81 and CC166), which previously contained vaccine types, were replaced with the NVTs, while CC320 remains unchanged.

Disclosures. All authors: No reported disclosures.

2339. Perianal Infections in Children With Acute Myeloid Leukemia: A Report From the Canadian Infection in Acute Myeloid Leukemia Research Group

Samuele Renzi, MD¹; Jack Bartram, MD, PhD¹; Salah Ali, MD¹; Carol Portwine, PhD²; David Mitchell, MD³; David Dix, MD⁴; Victor Lewis, MD⁵; Victoria Price, MMed⁶; Donna Johnston, MD⁷ and Lillian Sung, MD PhD^{1,8}; ¹Division of Haematology/Oncology, The Hospital for Sick Children, Toronto, ON, Canada, ²McMaster Children's Hospital at Hamilton Health Sciences, Hamilton, ON, Canada, ³Montreal Children's Hospital, Montreal, QC, Canada, ⁴British Columbia Children's Hospital, Vancouver, BC, Canada, ⁵Alberta Children's Hospital, University of Calgary, Calgary, AB, Canada, ⁶IWK Health Centre, Halifax, NS, Canada, ⁷Children's Hospital of Eastern Ontario, Ottawa, ON, Canada, ⁸Program in Child Health Evaluative Sciences, The Hospital for Sick Children, Toronto, ON, Canada

Session: 247. Pediatric Bacterial Infections

Saturday, October 6, 2018: 12:30 PM

Background. Little is known about the epidemiology of perianal infection in pediatric cancer patients. Objectives were to describe the characteristics, treatment and outcome of perianal infection and describe features of those with and without definite abscess in pediatric patients with acute myeloid leukemia (AML).

Methods. We performed a retrospective analysis of two multi-center cohort studies investigating risk factors for infection in children with AML. We included children with de novo AML ≤ 18 years of age with a perianal infection prior to the completion of AML treatment or stem cell transplantation

Results. Of 235 patients with AML, 17 (7%) experienced 19 perianal infections. Median age at perianal infection was 8.2 (range 0.6–16.1) years. Local bacterial cultures were positive in 6 (32%) episodes, but none matched bacteremia isolates ($n = 5$). Enterobacteriaceae were the most common pathogen. The 19 episodes were stratified by definite abscess ($n = 12$) and cellulitis/phlegmon ($n = 7$). All patients presented with local pain, erythema and induration or swelling. Fever was a frequent finding ($n = 17$, 89.4%). Among the patients with abscess, 9 (75%) were severely neutropenic at diagnosis and surgical intervention was required in 8 (42%). All patients received antibiotics; Metronidazole ($n = 14$) and Piperacillin/Tazobactam ($n = 10$) were the drugs most frequently used for treatment. Imaging was commonly performed ($n = 16$). Diagnostic yield was similar between computerized tomography of pelvis (5/10) and ultrasound (3/5). Severe complications occurred including fistula ($n = 1$), skin necrosis ($n = 2$) and mortality ($n = 1$).

Conclusion. Perianal infections occurred in 7% of pediatric patients with AML, with many consisting of definite abscess. Diagnostic yield were similar regardless of imaging modality and therefore, ultrasound may be considered for initial evaluation. Future research should develop consistent management approaches to perianal infection in order to improve outcomes.

Disclosures. All authors: No reported disclosures.

2340. Pneumococcal Colonization in Pediatric Patients Undergoing Bone Marrow Transplantation

Liset Olarte, MD, MSc¹; Jennifer Schuster, MD¹ and Kristina G. Hulten, PhD²;

¹Children's Mercy Hospital, Kansas City, Missouri, ²Baylor College of Medicine and Texas Children's Hospital, Houston, Texas

Session: 247. Pediatric Bacterial Infections

Saturday, October 6, 2018: 12:30 PM

Background. Hematopoietic cell transplant (HCT) recipients are at a significantly greater risk for invasive pneumococcal disease than the general population. However, pneumococcal colonization in pediatric HCT recipients has not been widely studied. We evaluated the dynamics of pneumococcal colonization in pediatric patients undergoing HSCT from conditioning regimen to 100 days post HCT.

Methods. Mid-turbinate samples obtained from pediatric patients undergoing HCT at Children's Mercy from September 2015 to January 2017 were tested for *Streptococcus pneumoniae* colonization via real-time PCR using *lytA* primer (autolysin-A-encoding gene). A cycle threshold value ≤ 35 was considered positive. First sample was obtained during conditioning regimen (week 1), second sample after HCT (week 2). Then, weekly samples were obtained for the first 100 days after HSCT.

Results. Twenty-two patients were included, representing 266 mid-turbinate samples. The median age at the time of HSCT was 9.5 years (IQR 3–16), and 14 patients were male (63.6%). The indication for HSCT was oncologic (15, 68.2%), hematologic (5, 22.7%) and immune deficiency (2, 9.1%). Fourteen patients (63.6%) underwent allogeneic HSCT. Six patients had documentation in our electronic medical record system of receiving ≥ 1 pneumococcal conjugate vaccine prior to conditioning regimen. Nine (40.9%) of 22 patients were colonized with *S. pneumoniae*, their median age was 14 years (IQR 2.5–16). Pneumococcal colonization during conditioning regimen was 9% (2/22). Pneumococcal colonization from week 14 to week 16 was 42% (5/12).

Conclusion. A third of pediatric HCT recipients were colonized with *S. pneumoniae*. Pneumococcal colonization was mainly identified either at the time of conditioning regimen or toward the end of the first 100 days after HCT, with the latter being the most common.

Disclosures. All authors: No reported disclosures.

2341. Trends in Adenovirus Infections in Singapore Children and Outcomes of Cidofovir Treatment in the Severely Ill

Valerie Xue Fen Seah, BSc(Pharm)(Hons), PharmD¹; Yirong Chew²; Koh Cheng Thoon, MBBS, MMed (PAEDS), MRCPCH³; Nancy Wen Sim Tee, MBBS, FRCPA⁴; Yelen⁵; Lin Cui⁵; Chia Yin Chong, MBBS, M. Med, FRCPCH³; Chee Fu Yung, MFPHM (UK), FFPHM (UK)³; Matthias Maiwald, MD, FRCPA⁶; Raymond Reinaldo Tanugroho, MBBS⁷ and Natalie Woon Hui Tan, MBBS, MRCPCH³; ¹Pharmacy, KK Women's and Children's Hospital, Singapore, Singapore, ²Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore, ³Infectious Disease Service, Department of Pediatrics, KK Women's and Children's Hospital, Singapore, Singapore, ⁴Microbiology, KK Women's and Children's Hospital, Singapore, Singapore, ⁵National Public Health Laboratory, Singapore, Singapore, ⁶Microbiology, KK Women's and Children's Hospital, Singapore, Singapore, ⁷Pediatrics, KK Women's and Children's Hospital, Singapore, Singapore

Session: 248. Pediatric Viral Infections

Saturday, October 6, 2018: 12:30 PM

Background. An increase in human adenovirus (HAdV) infections among hospitalized children in Singapore was observed since 2013. Cidofovir is often used to treat severe HAdV infections despite limited data. This study describes the epidemiology and outcomes of children with severe HAdV disease requiring high dependency (HD) or intensive care unit (ICU) admission in our hospital (KKH).

Methods. This is a retrospective cohort study of HAdV-infected children admitted to HD and ICU in KKH from January 2013 to September 2017. Characteristics and outcomes of those who received IV cidofovir was also reviewed.