including antibiotic use and offending organisms. We compared the cumulative incidence of reinfection between those with or without suppression using a competing risk model, with death and revision for mechanical failure as competing risks.

Results. The median age was 77 years (range, 48–94). Gender distribution was equal. The median Charlton Comorbidity Index (CCI) was 3 (1–11), while median BMI was 30.1 (17.8–59.5). Eleven of 51 patients received antibiotic suppression after spacer retention. A history of prior antibiotic suppressive therapy was the only variable associated with being placed back on antibiotic suppression after spacer retention [OR 18 (95% CI 3.2–100)]. During the median follow-up period of 31.3 months, there were five re-infections. The cumulative incidence of re-infection was not significantly different between suppressed and unsuppressed groups ($P = 0.89$). The re-infecting pathogens were different from the index offending organisms. Only the presence of preoperative draining sinus was significantly associated with re-infection [OR 10 (95% CI 1–99.6)].

Conclusion. In selected patients where a second-stage prosthesis re-implantation is not an option, and retention of “temporary” antibiotic loaded spacer is surgically preferred, the risk of re-infection was not prevented by prolonged antibiotic suppression. The presence of a draining sinus was significantly associated with re-infection, often with new pathogens.

Disclosures. All authors: No reported disclosures.

323. Childhood Outcomes Following Parechovirus Central Nervous System Infection in Young Infants at a US Children's Hospital

Rangaraj Selvarangan, PhD¹; Howard Kilbride, MD² and Mary Anne Jackson, MD, FIDSA, FPIDS³; ¹Microbiology Laboratory, Children's Mercy Hospital and Clinics, Kansas City, Missouri, ²Children's Mercy, Kansas City, Missouri, ³Pediatrics, Children's Mercy Hospital, Kansas City, Missouri

Session: 55. CNS Infections

Thursday, October 4, 2018: 12:30 PM

Background. Parechovirus (PeV), specifically *Parechovirus A* type 3 (PeV-A3), is a picornavirus associated with severe infection in young infants, with disease manifestations ranging from undifferentiated fever, to sepsis like illness, and meningoencephalitis. There are limited data regarding long-term outcomes of infected infants. The objective of this study was to describe early childhood outcomes following infantile PeV-CNS infection

Methods. Families of Infants hospitalized during 2014 with laboratory confirmed PeV-CNS infection were contacted for neurodevelopmental follow-up. Testing included medical history, standard neurologic examination, parental completion of Ages and Stages questionnaire (ASQ) and determination of Bayley III cognitive, motor, and language quotients. Neurodevelopmental impairment (NDI) was considered present if cognitive, motor, or language quotients were >1 standard deviation (mild) or >2 SD (severe) below the testing norms, the presence of cerebral palsy (CP), or sensory (vision/hearing) impairment. Relationship of children's outcomes to severity of PeV disease (uncomplicated febrile illness [mild], disseminated disease [moderate] or advanced disease requiring intensive care [severe]) was assessed by chi-square analysis.

Results. Nineteen children were available for testing at approximately 3 years of age (31–38 months), 12 (63%) with mild, five (26%) moderate, and two (11%) with severe disease. Mean Bayley quotients were within normal limits (see table), one infant had mild CP (5%) and two (11%) had mild NDI. There was no apparent relationship of NDI with infant PeV clinical presentation. ASQ results included 11% at referral level and 32% suspect, and were unrelated to severity of the viral illness. However, all parents of children with moderate or severe presentations of infantile PeV disease had medical or behavior concerns at 3 years of age compared with 25% of those with mild presentation ($P = 0.007$).

Bayley Quotients

	Mean	Range
IQ	98	85–125
Motor	108	91–130
Language	103	89–118

Conclusion. Neurodevelopmental impairments may be seen following infant PeV disease, but may not correlate with severity of clinical disease. Longitudinal monitoring of developmental status through early childhood following PeV infantile disease is warranted.

Disclosures. All authors: No reported disclosures.

324. Lack of Accuracy of the International Classification of Disease, Ninth (ICD-9) Codes in Identifying Patients With Encephalitis

Mohammed Samannodi, MD¹; Michael Hansen, MD² and Rodrigo Hasbun, MD, MPH³; ¹Division of Infectious Diseases, University of Texas Health Science Center at Houston, McGovern Medical School, Houston, Texas, ²Family and Community Medicine, Baylor College of Medicine, Houston, Texas, Houston, Texas

Session: 55. CNS Infections

Thursday, October 4, 2018: 12:30 PM

Background. ICD-9 codes have been widely used in studies utilizing large national databases that evaluate the clinical epidemiology of encephalitis in the United States. Many studies have showed that ICD-9 codes have poor accuracy in stroke, multiple sclerosis and pulmonary fibrosis but their utility in encephalitis is unknown.

Methods. Retrospective study of all adults with a discharge diagnosis of encephalitis by an ICD-9 code. The study was performed in 17 hospitals from the Memorial Hermann Hospital and Harris Health Hospital system in the Greater Houston area from March 2010 until July 2015. Medical records were reviewed and a case was considered accurately classified as encephalitis if they met the definition established by the international encephalitis consortium.

Results. A total of 1,241 cases were identified by a discharge diagnosis of ICD-9 code as having encephalitis. The most common cause identified was not having a central nervous system infection in 580 (46.7%) patients. A total of 244 (19.6%) patients were correctly identified as having encephalitis. Other causes identified were nosocomial meningitis (11.9%), community-acquired bacterial meningitis (8.1%), aseptic meningitis (5.8%), fungal meningitis (5.4%), tuberculosis (2.0%), and parasitic meningitis (0.2%).

Conclusion. ICD-9 codes have poor reliability in identifying patients with encephalitis questioning the accuracy of large nationwide studies that utilize them to identify patients.

Table 1: Correct Clinical Diagnosis in 1,241 Patients with a Discharge Diagnosis of Encephalitis by ICD-9 Codes.

Diagnosis	Number of Patients (%)
Non-CNS infection	580 (46.7)
Encephalitis	244 (19.6)
Nosocomial meningitis	148 (11.9)
Bacterial meningitis	101 (8.1)
Aseptic meningitis	72 (5.8)
Fungal meningitis	68 (5.4)
Tuberculous meningitis	25 (2.0)
Parasitic meningitis	3 (0.2)

Abbreviation: ICD-9, an International Classification of Disease, Ninth; CNS, Central Nervous System.

Disclosures. All authors: No reported disclosures.

325. Neurosyphilis Management in the Post-Procaine Penicillin Era

Chase Cannon, MD¹; Alison Beiler, PA-C, MS²; Meena Ramchandani, MD, MPH³; Roxanne Kerani, PhD⁴ and Shireesha Dhanireddy, MD¹; ¹Allergy and Infectious Diseases, University of Washington, Seattle, Washington, ²Harborview Medical Center, Seattle, Washington, ³King County Public Health STD Clinic / University of Washington, Seattle, Washington, ⁴King County Public Health/University of Washington, Seattle, Washington

Session: 55. CNS Infections

Thursday, October 4, 2018: 12:30 PM

Background. Neurosyphilis (NS) is an infection of the central nervous system caused by *Treponema pallidum*. Intramuscular (IM) penicillin (PCN) G procaine is a treatment option for those who cannot receive or decline intravenous (IV) therapy. Since August 24, 2016, it has been unavailable from the manufacturer, necessitating the use of IV PCN for NS. Our institutions organized a multidisciplinary, coordinated care system to expedite outpatient treatment of NS upon diagnosis. We report successful management of NS at an urban safety-net hospital in the post-procaine PCN era.

Methods. We identified patients with suspected NS from the King County Public Health STD and Harborview Infectious Disease clinics from October 2016 to February 2018. Demographics, clinical symptoms, diagnostics, treatment, and outcomes were collected by chart review. Successful NS treatment was defined as resolution of cerebrospinal fluid (CSF) pleocytosis or elevated protein, improvement in neurologic symptoms or appropriate decrease in serum rapid plasma reagin (RPR) or CSF Venereal Disease Research Laboratory (VDRL) titers.

Table 1: Demographic and Socioeconomic Characteristics

	Total = 43 n (%)
Gender	
Male	39 (91)
Race	
White	29 (67)
Black	3 (7)
Asian	3 (7)
Homeless	5 (12)
Insurance status ^a	
Private	15
Medicaid	20
Medicare	8
Charity care	11
Substance use disorder	15 (35)
HIV Positive	22 (51)
Viral load suppressed (< 200 copies/mL)	13 (59)

^aRepresents more than one payer per patient.