Published in final edited form as:

Cell Tissue Res. 2018 January ; 371(1): 171–179. doi:10.1007/s00441-017-2728-3.

Single cell analysis of diversity in human stem cell-derived neurons

Lise J. Harbom[#], Nadine Michel[#], and Michael J. McConnell

Department of Biochemistry and Molecular Genetics, Neuroscience Graduate Program,
University of Virginia School of Medicine, Charlottesville, VA, 22908

[#] These authors contributed equally to this work.

Abstract

Neural stem and progenitor cells produce one of the most remarkable organs in nature, the human brain. Among neural stem cell progeny, post-mitotic neurons are likewise remarkably diverse. Single cell transcriptomic approaches are now cataloguing a long-sought-after molecular taxonomy of neuronal diversity in the brain. Contemporary single cell ‘omic classifications of neuronal diversity build from electrophysiological approaches that for decades have measured and catalogued diverse biophysical properties of single neurons. With the widespread application of human pluripotent stem cell (hPSC)-based models of neurogenesis to investigate disease pathology and to develop new drugs, a high-resolution understanding of neuronal diversity in vivo is essential to benchmark the state of in vitro models of human neurological disease.

Introduction

Neural stem and progenitor cells (NSCs, NPCs) give rise to diverse neuronal and glial progeny during brain development (S. Lodato & Arlotta, 2015; Masland, 2004). While Golgi and Cajal first glimpsed the extraordinary morphological diversity of neurons in the cerebral cortex over a century ago (Cajal, 1890, 1891; Golgi, 1886), careful accounting of neural diversity at the molecular level, continues today. A complete census of neural subtypes and circuit composition will further our understanding of how complex behaviors and neurological diseases arise from neural circuits.

Tissue-wide measurement of any brain region obscures the underlying molecular diversity so rare cell types go unmeasured and diversity among common cell types becomes population-averaged (Figure 1). Thus, measurement of single cells is essential to define neuronal diversity (Eberwine & Kim, 2015). The ongoing development of in vitro hPSC-based models of neurological disease adds urgency to the need for a high-resolution understanding of diversity among primary neurons (K. J. Brennand et al., 2015). The limitations of current hPSC-based neurogenesis approaches are best evaluated by comparison to primary human neurons; innovative in vitro approaches can then focus on clear obstacles with well-defined measures of success.

Single cell analysis has been fundamental to neuroscience since the patch-clamp technique allowed one to record the electrical activity of single neurons (Neher & Sakmann, 1976). Patch-clamp has been used to functionally classify neurons based on their response to synaptic inputs, ion channel composition, and intrinsic firing properties (Contreras, 2004; Migliore & Shepherd, 2005). Following from these studies, the maturation of hPSC-derived neurons is typically validated based on the presence of sodium and potassium currents that underlie action potential generation. More careful characterization of hPSC-derived neuronal subtypes is assessed based on the extent to which their electrophysiological response matches the target cell type in slice recordings.

Today single cell transcriptome measurement is routine (Macaulay & Voet, 2014). Efforts to map neuronal diversity in mouse and human cortex have produced several thousand single cell transcriptomic profiles which are sorted into known cell-types based on the expression of one or a few marker genes (Lake et al., 2016; Linnarsson, 2013; Tasic et al., 2016). As expected, further analysis of gene expression revealed sub-types within the known types, and ongoing analysis of these data provides evidence of additional diversity. Current approaches to generate hPSC-derived neurons produce neurons that express known marker genes, but neuronal diversity in vitro is still limited. Herein, we review single cell approaches to define neuronal diversity in primary tissue and assess current hPSC-based models of neurogenesis.

Single Neuron Morphology and Physiology

Microscopy and differential staining methods revealed exquisitely distinct morphological diversity among neurons; however, measurements of the fundamental electrical properties of single neurons required the patch clamp technique. Patch clamp approaches yield information about the resting membrane potential, membrane resistance, and intrinsic excitability of single neurons, as well as synaptic function. In turn, these differences reflect electrical diversity among NSC progeny.

In the neocortex, neurons are broadly defined as excitatory or inhibitory based on what neurotransmitters are released from the cell upon activation. However, apart from a cell's synaptic output, neurons also differ in their response to activation, and neural circuits are composed of cells with varying electrophysiological properties and firing patterns (Connors & Gutnick, 1990; Mountcastle, Talbot, Sakata, & Hyvarinen, 1969). The firing properties of neocortical neurons can be divided broadly into five categories: regular spiking, intrinsically bursting, fast spiking, fast repetitive bursting, and cells with low threshold spikes (Connors & Gutnick, 1990; Contreras, 2004), though a cell's firing class is flexible and may change based on changes in ion conductance in response to surrounding activity (Steriade, 2004). Firing patterns are highly related to neuronal morphology (Connors & Gutnick, 1990; Mason & Larkman, 1990), particularly the number and branching of dendritic spines (Mainen & Sejnowski, 1996). Additionally, patching cells followed by single-cell sequencing has led to the discovery that electrophysiological subtypes are often correlated with aspects of the cells' transcriptomic profiles (Cadwell et al., 2016; Foldy et al., 2016; Fuzik et al., 2016).

The question arises whether cortical neuron classification is entirely driven by NPC fate decisions, or whether electrophysiological subtypes are determined as the cells form into circuits. There is some evidence that the electrical properties of neurons are in part pre-determined while they are intermediate progenitors (Tyler, Medalla, Guillamon-Vivancos, Luebke, & Haydar, 2015); however, it has been shown that transplanted neural stem cells can also integrate into an already established circuit (Avaliani et al., 2014; Espuny-Camacho et al., 2013), indicating that circuit activity surrounding the cell is at least partially responsible for cell fate. Additionally, morphological constraints during development can affect a neuron's firing properties, again showing that the local environment contributes to the development of electrical diversity (Zhu, Du, Jin, & Offenhausser, 2016).

Fluorescent biomarkers and live or time-lapse imaging have been extensively used in animal models to track the fate of stem cells during development, including neural stem cells both during development and the role of neurogenesis in the adult brain (Gobel, Kampa, & Helmchen, 2007; Mizrahi, 2007; Schroeder, 2011; Yokota et al., 2007). Although it is not usually feasible to perform imaging of this extent in humans because there is usually a requirement for pre-loading stem cells with a dye or label, proton magnetic resonance spectroscopy has been used to visualize levels of neurogenesis in the human hippocampus at different ages (Manganas et al., 2007). Some morphological features of cortical neuron diversity are accurately reflected in hPSC-based neurogenesis. For example, neurons expressing superficial versus deep cortical layer markers can be found in culture (Handel et al., 2016; Mariani et al., 2012). In 3D cultures, imaging has shown that newly differentiating neurons undergo a process similar to interkinetic nuclear migration (Pasca et al., 2015). The cortical spheroids in this study also show a clear lamination pattern, with deep layer markers concentrated towards the center of the spheroid and upper-layer markers towards the outside, mimicking in vivo cortical lamination. While it seems that cortical diversity is being represented in this way, functional characterization distinguishing cells from different regions has been limited.

Different morphological subtypes are observed in in vitro cultures; however, the extent to which this correlates with in vivo data is unclear. In a study on Dravet syndrome patient-derived neurons, Liu et al. showed that bipolar-shaped cells and pyramidal-shaped cells show very limited differences in basal membrane properties, despite a slightly decreased membrane resistance in pyramidal cells (Y. Liu et al., 2013). However, other studies in hPSC-derived neuron cultures have shown a correlation between properties such as dendritic branching with more mature neurons hPSC (Bardy et al., 2016).

Physiological Diversity In Vitro

Although there is evidence that hPSC-derived neuron cultures may mimic diversity found in primary cortex, technical variation is a large source of heterogeneity in hPSC-derived neuron cultures. There is a wide discrepancy in the literature in reported electrophysiological properties of hPSC-derived neurons. Some of this diversity can be explained by differing culture methods. Factors such as astrocyte co-culture (Kuijlaars et al., 2016; Rushton, Mattis, Svendsen, Allen, & Kemp, 2013; Tang et al., 2013) and altered cell culture medium (Bardy et al., 2015; Kemp et al., 2016) have been shown to alter the time scale of neuron

maturation. However, even among neurons of the same origin cultured the same way, there is a range of neuron function, including neurons that do not exhibit firing behavior at all alongside neurons with robust spiking activity. This makes it difficult to analyze spiking using traditional measures, such as a plot of current injection versus frequency, since the high rate of non-responders skews the data. To overcome this, many studies have categorized neurons into firing ‘types,’ and analyzed spiking metrics within those categories (Bardy et al., 2015; Belinsky et al., 2014; Kemp et al., 2016; Lam, Topfer, Wood, Busskamp, & Bamberg, 2017; Song, Mohamad, Chen, & Yu, 2013).

In a study by Bardy and colleagues in 2016, neurons were separated into five ‘functional states’ based on firing activity, which was subsequently correlated with transcriptome data from the single cells (Bardy et al., 2016). While there was a general increase in the more mature functional states with increased culture duration, immature cells were present at all time points. Similar to findings in in vivo studies, a cell’s firing activity was shown to be correlated to the cell’s transcriptome; in this case, the expression of certain genes involved in higher level neuron function, such as synaptic plasticity. This indicates that functional variability is in fact reflecting differences in maturity as assessed by transcriptome analysis. Thus, different electrophysiological subtypes found in these cultures are more likely to reflect neurons in various stages of development, and unlikely to reflect the kind of variability that is present within an adult cortex. Because of this, hPSC-derived neurons are currently not an ideal method to study functional subtypes in the adult cortex.

While a difference in maturity may be the root cause of variability in neuron function within a single culture, this does not necessarily negate the notion that stem cell-derived neurons accurately reflect some aspects of in vivo diversity. The field standard for functional maturity in these cultures is that of an immature neuron in vivo (Randall, 2016). One indicator of this is the membrane resistance, which is on the order of 1–4 G Ω in hPSC-derived neurons as opposed to less than 1 G Ω in an average adult neuron in animal studies (Livesey, Magnani, Hardingham, Chandran, & Wyllie, 2016). However, this aligns with membrane resistance values from fetal cortex tissue, which similarly show values up to 5 G Ω , and also show immature, abortive action potentials similar to those seen in stem cell-derived cultures (Moore et al., 2009). With this in mind, it is possible that the heterogeneity of neuron cultures is accurately mimicking in vivo developmental timelines, as all neurons do not reach functional maturity at the same rate.

While having such wide variability can make using hPSC-derived neurons as a valid model of mature neural circuits difficult, it also offers an opportunity to study factors that contribute to electrophysiological development, including why some neurons seem to mature faster than others and what relevance this may have to later function. It is also important to note that hPSC-derived neurons do accurately reflect properties of immature cortical neurons, and as techniques improve to culture these neurons long-term or accelerate maturation, it is possible that circuitry more reminiscent of a mature cortex will develop. This will be a valuable tool for investigating what factors contribute to circuitry formation and how electrophysiological diversity is established.

Single Neuron Transcriptome Analysis

Transcriptomic analysis is routinely employed to validate pluripotency and lineage potential in hPSC lines (Muller et al., 2011; Tsankov et al., 2015). Likewise, transcriptomic analysis has become a central tool for phenotyping patient-derived hPSC lines. Several groups have taken this approach to study hPSCs derived from patients with neuropsychiatric disorders such as schizophrenia (K. Brennand et al., 2015; Brennand et al., 2011). Studies looking at the transcriptome of hPSC-derived neurons are typically performed on large cell populations, which necessarily obscure neuron-to-neuron diversity. With the development of single-cell transcriptomic approaches, fine-grained maps of neuronal diversity are emerging rapidly.

Transcriptome Diversity In Vivo

Various single cell approaches have been used to better understand cellular diversity within known subpopulations of primary NPCs and neurons. Specific neural subtypes are purified from bulk tissue using fluorescence activated cell sorting (FACS). Single cell transcriptomes can then be measured on targeted collections of hundreds of specific genes using multiplexed qPCR and microfluidic devices (e.g. Fluidigm Biomark) or comprehensively on RNA libraries by next generation sequencing. Two examples of these approaches are highlighted from human cerebral cortex; however, several similar studies have also examined other populations of NPCs and neurons (Dulken, Leeman, Boutet, Hebestreit, & Brunet, 2017; S. J. Liu et al., 2016; Tasic et al., 2016).

Johnson et al. (Johnson et al., 2015) isolated two cortical progenitor cell populations (apical and outer (i.e., non-apical) radial glia from fetal human brain based on expression of known proteins (i.e., GLAST and LexA) and differential abundance of the apical marker prominin. Diversity in these human cortical progenitor cell types is of particular interest because the development of large, complex germinal zones is associated with evolutionary expansion of the human neocortex. Multiplexed qPCR was performed on 546 single progenitor cells to query the expression levels of dozens of genes. In contrast to previous bulk analysis of microdissected germinal zones, this single cell study identified new markers of outer radial glia and found diverse transcriptional states involving co-expression of both classic radial glia markers and pro-neural transcription factors in both apical and non-apical cell types.

A second single cell study by the Zhang and Chun labs used NeuN expression to separate neuronal nuclei from non-neuronal nuclei and investigated cortical neuron diversity in 6 discrete regions of the human cerebral cortex (Lake et al., 2016). Notably, they found that sequencing of nuclear RNA (Grindberg et al., 2013; Lacar et al., 2016) was superior to laser microdissection of intact neurons. A total of 3,227 neuronal data sets were classified based on 10-fold differentially expressed genes using an iterative clustering approach, then excitatory and inhibitory populations were identified based on expression of known marker genes. Neuronal diversity was classified into 8 excitatory and 8 inhibitory subtypes with additional diversity associated with regional identities. However, the authors note that even within one region there is still heterogeneity in gene expression among single cells that remains to be explored.

An unbiased alternative to these FACS-based approaches is to perform RNAseq analysis on all single cells in a defined brain region. Zeisel et al. (Zeisel et al., 2015) performed RNAseq on 3,005 single cells from mouse somatosensory cortex and hippocampal CA1 region. Cluster-based analysis identified pyramidal neurons and interneurons, as well as 6 non-neuronal subtypes including oligodendrocytes, astrocytes, microglia, vascular endothelial cells, mural cells, and ependymal cells. Distinct subclasses of cells were identified retrospectively based on expression of cell marker genes. Sixteen subclasses of interneurons were discovered and the authors noted that interneurons within the same region had the most similar transcriptomic profiles.

Transcriptome Diversity In Vitro

Single cell transcriptome analysis of primary neurons provides an essential benchmark for hPSC-derived neurons. Handel et al. (Handel et al., 2016) conducted a critical study assessing the similarities in transcriptomes of primary tissue and hPSC-derived cortical neurons. They discovered that hPSC-derived neurons had mixed layer identities based on the expression of specific upper and deep layer markers. However, this heterogeneity in cell layer marker expression was also present in primary fetal and adult neurons, suggesting that there is neuronal diversity throughout neurogenesis and that hPSC-derived cortical neurons may provide us with a reliable model to examine how that diversity is formed. Indirectly, this study also shows why transcriptional profiling will be a key method to identify cell subtypes moving forward. In both primary tissue and hPSC-derived cultures there were mixed layer identity cells; thus, transcriptional signatures may for the foreseeable future be the best tool to disambiguate those cortical layer identities.

The direct comparison between in vivo and in vitro single-cell transcriptomic analysis has been examined in mice as well. Dulken et al. specifically looked at transcriptional dynamics in adult neural stem cell lineage through single cell RNA-seq of the adult mouse subventricular zone in transgenic GFAP-GFP mice. They observed four cell populations: astrocytes, quiescent neural stem cells (qNSCs), activated neural stem cells (aNSCs), and NPCs. Through PCA analysis they found a continuum of quiescence, activation, and differentiation in aNSCs. Further investigation of this subpopulation revealed differential gene expression depending on the state of the NSC. Earlier aNSCs expressed cell cycle genes while later aNSCs expressed neuronal genes. When the authors compared the in vivo NSCs to in vitro, they found a greater enrichment for neuronal genes in vivo while the in vitro NSCs had enrichment for astrocyte genes (Dulken et al., 2017). This study highlights one of the key issues in the study of in vitro neurogenesis: determining real versus artificial heterogeneity. The in vitro condition showed more heterogeneity than the in vivo condition in that there was a greater spectrum of neural stem cells in various stages of activation. The study did not follow the NSCs far into the differentiation process, however, so it is still unclear if the heterogeneity in the NSCs translated into more diverse hPSC-derived neurons compared to primary tissue.

Single Neuron Genome Analysis

Single cell analysis has also revealed surprising diversity in neuronal genomes (McConnell et al., 2017). Brain somatic mosaicism is brought about by single nucleotide variants (SNVs), small insertion/deletion mutations (Hazen et al., 2016; M. A. Lodato et al., 2015), structural variants (e.g., copy number variants (CNVs)) (Cai et al., 2014; Knouse, Wu, & Amon, 2016; McConnell et al., 2013), and mobile genetic element insertions (Baillie et al., 2011; Erwin et al., 2016; Evrony et al., 2012; Upton et al., 2015). Collectively, the variety and abundance of somatic mutations indicates that every neuron in a circuit may be expected to have a different genome.

Cell proliferation during development is a prominent source of somatic mutations. Given about 20 billion neurons in human cortex, and more than 50% cell death during neurodevelopment, cortical NPCs must undergo at least 40 billion cell divisions (Lui, Hansen, & Kriegstein, 2011). Errors in DNA replication and repair are estimated to lead to 1 – 3 mutations per cell division (Lynch, 2010); thus, estimates of greater than 1,000 somatic SNVs per neuronal genome (M. A. Lodato et al., 2015) are quite reasonable. However, it is interesting to consider that the distribution of diverse genomes in the cortex is expected to be orthogonal to the arrangement of diverse cell types.

Clonally-related neurons of different cell types are born through a series of asymmetric cell divisions by an individual NPC (Namba & Huttner, 2017). Surviving progeny then reside in functionally connected radial columns (Yu, Bultje, Wang, & Shi, 2009) with early born neurons in deep cortical layers and subsequently born neurons migrating to increasingly superficial layers. Neuronal cell type diversity is often described in the context of layer 6 neurons expressing different genes than layer 5 neurons, and so forth. Likewise, neurons in different cortical layers have different morphological and physiological properties. Somatic mutations in NPCs, as a consequence of iterative cell fate decisions, will be shared among many of that NPCs progeny, while cortical columns that arise from other NPCs could have distinctly different genomes (Figure 2).

A central interest is whether somatic mutations are more prevalent in the brain than in other tissues. Although somatic mosaicism occurs in probably all organs, the long lifespan and direct influence on behavior of neural circuits suggest a direct consequence of somatic mosaicism in both neurotypical and diseased brains. In vitro models of human neurogenesis offer some insight into this question.

Single cell genomic analysis showed that NPCs derived from hPSCs exhibit few CNVs; whereas CNVs are abundant in 8-week differentiated neurons (McConnell et al., 2013). Using an L1 reporter system, L1 retrotransposon activity is likewise absent in hPSC-derived NPCs, but then observed during the first week of neuronal differentiation (Coufal et al., 2009). Additional study of the L1 promoter shows transcriptional regulation similar to that of the master neuronal transcription factor NeuroD. Hybrid Sox2/TCF-LEF motifs MeCP2-deficient phenotypes (Coufal et al., 2009) support a model wherein Sox2-binding and MeCP2 activity suppress L1 mobility in NPCs, and Sox2 de-repression coupled with Wnt signaling promote L1 transcription during neurogenesis (Kuwabara et al., 2009). If hPSC-

based neurogenesis is faithfully recapitulating the mechanisms that lead to primary brain somatic mosaicism, it will prove to be a valuable model system in defining the intersection between cell type diversity and genomic diversity in human neurons.

Future Directions

In addition to the single cell approaches highlighted herein, other advances in single cell epigenomic (Farlik et al., 2015; Gravina, Dong, Yu, & Vijg, 2016) and proteomic approaches (Budnik, 2017), as well as multi-omic approaches (Bock, Farlik, & Sheffield, 2016; Macaulay, Ponting, & Voet, 2017), a high-resolution molecular taxonomy of neuronal diversity is taking shape rapidly. Multiplex protein measurement approaches such as flow cytometry provide additional quantification of diverse neural types in specific brain regions. Pruszk et al used fluorescence-activated cell sorting to analyze heterogeneous hESC derived neural stem cells at various stages of differentiation and showed that specific surface antigens corresponded to the development or maintenance of neural precursor cells or neurons (Pruszk, Sonntag, Aung, Sanchez-Pernaute, & Isacson, 2007). Mass cytometry extends multiplex capability to at least 50 separate analytes on single cells by conjugating antibodies with heavy metals rather than fluorophores (Bandura et al., 2009; Bendall et al., 2014). Together with novel computational approaches, Zunder et. al used mass cytometry to track the dynamic process of iPSC reprogramming by looking at markers of pluripotency, differentiation, cell cycle status, and cell signaling (Zunder, Lujan, Goltsev, Wernig, & Nolan, 2015).

Development of improved approaches for hPSC-based neurogenesis is likewise proceeding with great pace. Three-dimensional culture approaches such as cerebral organoid technology (Lancaster et al., 2013) show improved tissue patterning over standard two-dimensional approaches; however, additional advances are required to fully recapitulate neuronal diversity in primary tissue. For example, neural vasculature is known to provide an important NPC niche, yet existing organoids are avascular. Without such a niche, limited diversity is not unexpected.

Very large single cell data sets from many individuals will ultimately be needed to establish a complete molecular taxonomy of neuronal diversity. While current single cell data sets number in the thousands, droplet-based approaches permit simultaneous capture, amplification, and barcoding of tens of thousands of single cells. Notably, the first application of DropSeq was to explore neural diversity in the retina using a data set of more than 10,000 cells (Macosko et al., 2015). Large data sets provide the statistical power not only to define a reference taxonomy, but also to begin to determine how neural diversity may vary among neurotypical individuals and in disease states. Moreover, a fine-grained map of primary brain tissue such as this will further inform our appreciation of the successes and limitations of hPSC-based models of neurogenesis.

Acknowledgements

We thank M. Beenhakker for helpful insights regarding the history of the patch-clamp technique and assessment of neuronal diversity by electrophysiological methods. We also acknowledge that the single cell field is rapidly evolving and apologize, in advance, to many colleagues whose advances in single cell measurement of neuronal

diversity will undoubtedly be reported after this review was prepared. This work was supported by NIMH U01 MH106882 to MJM, NIGMS T32 GM008328–24 to LJH and NM.

Literature cited

- Avaliani N, Sorensen AT, Ledri M, Bengzon J, Koch P, Brustle O, ... Kokaia M (2014). Optogenetics reveal delayed afferent synaptogenesis on grafted human-induced pluripotent stem cell-derived neural progenitors. *Stem Cells*, 32(12), 3088–3098. doi:10.1002/stem.1823 [PubMed: 25183299]
- Baillie JK, Barnett MW, Upton KR, Gerhardt DJ, Richmond TA, De Sapio F, ... Faulkner GJ (2011). Somatic retrotransposition alters the genetic landscape of the human brain. *Nature*, 479(7374), 534–537. doi:10.1038/nature10531 [PubMed: 22037309]
- Bandura DR, Baranov VI, Ornatsky OI, Antonov A, Kinach R, Lou X, ... Tanner SD (2009). Mass cytometry: technique for real time single cell multitarget immunoassay based on inductively coupled plasma time-of-flight mass spectrometry. *Anal Chem*, 81(16), 6813–6822. doi:10.1021/ac901049w [PubMed: 19601617]
- Bardy C, van den Hurk M, Eames T, Marchand C, Hernandez RV, Kellogg M, ... Gage FH (2015). Neuronal medium that supports basic synaptic functions and activity of human neurons in vitro. *Proc Natl Acad Sci U S A*, 112(20), E2725–2734. doi:10.1073/pnas.1504393112 [PubMed: 25870293]
- Bardy C, van den Hurk M, Kakaradov B, Erwin JA, Jaeger BN, Hernandez RV, ... Gage FH (2016). Predicting the functional states of human iPSC-derived neurons with single-cell RNA-seq and electrophysiology. *Mol Psychiatry*, 21(11), 1573–1588. doi:10.1038/mp.2016.158 [PubMed: 27698428]
- Belinsky GS, Rich MT, Sirois CL, Short SM, Pedrosa E, Lachman HM, & Antic SD (2014). Patch-clamp recordings and calcium imaging followed by single-cell PCR reveal the developmental profile of 13 genes in iPSC-derived human neurons. *Stem Cell Res*, 12(1), 101–118. doi:10.1016/j.scr.2013.09.014 [PubMed: 24157591]
- Bendall SC, Davis KL, Amir el AD, Tadmor MD, Simonds EF, Chen TJ, ... Pe'er D (2014). Single-cell trajectory detection uncovers progression and regulatory coordination in human B cell development. *Cell*, 157(3), 714–725. doi:10.1016/j.cell.2014.04.005 [PubMed: 24766814]
- Bock C, Farlik M, & Sheffield NC (2016). Multi-Omics of Single Cells: Strategies and Applications. *Trends Biotechnol*, 34(8), 605–608. doi:10.1016/j.tibtech.2016.04.004 [PubMed: 27212022]
- Brennand K, Savas JN, Kim Y, Tran N, Simone A, Hashimoto-Torii K, ... Gage FH (2015). Phenotypic differences in hiPSC NPCs derived from patients with schizophrenia. *Mol Psychiatry*, 20(3), 361–368. doi:10.1038/mp.2014.22 [PubMed: 24686136]
- Brennand KJ, Marchetto MC, Benvenisty N, Brustle O, Ebert A, Izpisua Belmonte JC, ... Jaenisch R (2015). Creating Patient-Specific Neural Cells for the In Vitro Study of Brain Disorders. *Stem Cell Reports*, 5(6), 933–945. doi:10.1016/j.stemcr.2015.10.011 [PubMed: 26610635]
- Brennand KJ, Simone A, Jou J, Gelboin-Burkhart C, Tran N, Sangar S, ... Gage FH (2011). Modelling schizophrenia using human induced pluripotent stem cells. *Nature*, 473(7346), 221–225. doi:10.1038/nature09915 [PubMed: 21490598]
- Budnik BL,E; Slavov N (2017). Mass-spectrometry of single mammalian cells quantifies proteome heterogeneity during cell differentiation. *bioRxiv* doi:https://doi.org/10.1101/102681
- Cadwell CR, Palasantza A, Jiang X, Berens P, Deng Q, Yilmaz M, ... Tolias AS (2016). Electrophysiological, transcriptomic and morphologic profiling of single neurons using Patch-seq. *Nat Biotechnol*, 34(2), 199–203. doi:10.1038/nbt.3445 [PubMed: 26689543]
- Cai X, Evrony GD, Lehmann HS, Elhosary PC, Mehta BK, Poduri A, & Walsh CA (2014). Single-cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human brain. *Cell Rep*, 8(5), 1280–1289. doi:10.1016/j.celrep.2014.07.043 [PubMed: 25159146]
- Cajal S (1890). Textura de las circunvoluciones cerebrales de los mamiferos inferiores. *Gaceta Medica Catalana*, 22–31.
- Cajal S (1891). Sur la structure de l'écorce cerebrale de quelques mammiferes. *La Cellule*, 125–176.
- Connors BW, & Gutnick MJ (1990). Intrinsic firing patterns of diverse neocortical neurons. *Trends Neurosci*, 13(3), 99–104. [PubMed: 1691879]

- Contreras D (2004). Electrophysiological classes of neocortical neurons. *Neural Netw*, 17(5–6), 633–646. doi:10.1016/j.neunet.2004.04.003 [PubMed: 15288889]
- Coufal NG, Garcia-Perez JL, Peng GE, Yeo GW, Mu Y, Lovci MT, ... Gage FH (2009). L1 retrotransposition in human neural progenitor cells. *Nature*, 460(7259), 1127–1131. doi:10.1038/nature08248 [PubMed: 19657334]
- Dulken BW, Leeman DS, Boutet SC, Hebestreit K, & Brunet A (2017). Single-Cell Transcriptomic Analysis Defines Heterogeneity and Transcriptional Dynamics in the Adult Neural Stem Cell Lineage. *Cell Rep*, 18(3), 777–790. doi:10.1016/j.celrep.2016.12.060 [PubMed: 28099854]
- Eberwine J, & Kim J (2015). Cellular Deconstruction: Finding Meaning in Individual Cell Variation. *Trends Cell Biol*, 25(10), 569–578. doi:10.1016/j.tcb.2015.07.004 [PubMed: 26410403]
- Erwin JA, Paquola AC, Singer T, Gallina I, Novotny M, Quayle C, ... Gage FH (2016). L1-associated genomic regions are deleted in somatic cells of the healthy human brain. *Nat Neurosci*, 19(12), 1583–1591. doi:10.1038/nn.4388 [PubMed: 27618310]
- Espuny-Camacho I, Michelsen KA, Gall D, Linaro D, Hasche A, Bonnefont J, ... Vanderhaeghen P (2013). Pyramidal neurons derived from human pluripotent stem cells integrate efficiently into mouse brain circuits in vivo. *Neuron*, 77(3), 440–456. doi:10.1016/j.neuron.2012.12.011 [PubMed: 23395372]
- Evrony GD, Cai X, Lee E, Hills LB, Elhosary PC, Lehmann HS, ... Walsh CA (2012). Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell*, 151(3), 483–496. doi:10.1016/j.cell.2012.09.035 [PubMed: 23101622]
- Farlik M, Sheffield NC, Nuzzo A, Datlinger P, Schonegger A, Klughammer J, & Bock C (2015). Single-cell DNA methylome sequencing and bioinformatic inference of epigenomic cell-state dynamics. *Cell Rep*, 10(8), 1386–1397. doi:10.1016/j.celrep.2015.02.001 [PubMed: 25732828]
- Foldy C, Darmanis S, Aoto J, Malenka RC, Quake SR, & Sudhof TC (2016). Single-cell RNAseq reveals cell adhesion molecule profiles in electrophysiologically defined neurons. *Proc Natl Acad Sci U S A*, 113(35), E5222–E5231. doi:10.1073/pnas.1610155113 [PubMed: 27531958]
- Fuzik J, Zeisel A, Mate Z, Calvigioni D, Yanagawa Y, Szabo G, ... Harkany T (2016). Integration of electrophysiological recordings with single-cell RNA-seq data identifies neuronal subtypes. *Nat Biotechnol*, 34(2), 175–183. doi:10.1038/nbt.3443 [PubMed: 26689544]
- Gobel W, Kampa BM, & Helmchen F (2007). Imaging cellular network dynamics in three dimensions using fast 3D laser scanning. *Nat Methods*, 4(1), 73–79. doi:10.1038/nmeth989 [PubMed: 17143280]
- Golgi C (1886). *Sulla Fina Anatomia degli Organi Centrali del Sistema Nervosa*. Milano: U. Hoepli
- Gravina S, Dong X, Yu B, & Vijg J (2016). Single-cell genome-wide bisulfite sequencing uncovers extensive heterogeneity in the mouse liver methylome. *Genome Biol*, 17(1), 150. doi:10.1186/s13059-016-1011-3 [PubMed: 27380908]
- Grindberg RV, Yee-Greenbaum JL, McConnell MJ, Novotny M, O'Shaughnessy AL, Lambert GM, ... Lasken RS (2013). RNA-sequencing from single nuclei. *Proc Natl Acad Sci U S A*, 110(49), 19802–19807. doi:10.1073/pnas.1319700110 [PubMed: 24248345]
- Handel AE, Chintawar S, Lalic T, Whiteley E, Vowles J, Giustacchini A, ... Cader MZ (2016). Assessing similarity to primary tissue and cortical layer identity in induced pluripotent stem cell-derived cortical neurons through single-cell transcriptomics. *Hum Mol Genet*, 25(5), 989–1000. doi:10.1093/hmg/ddv637 [PubMed: 26740550]
- Hazen JL, Faust GG, Rodriguez AR, Ferguson WC, Shumilina S, Clark RA, ... Baldwin KK (2016). The Complete Genome Sequences, Unique Mutational Spectra, and Developmental Potency of Adult Neurons Revealed by Cloning. *Neuron*, 89(6), 1223–1236. doi:10.1016/j.neuron.2016.02.004 [PubMed: 26948891]
- Johnson MB, Wang PP, Atabay KD, Murphy EA, Doan RN, Hecht JL, & Walsh CA (2015). Single-cell analysis reveals transcriptional heterogeneity of neural progenitors in human cortex. *Nat Neurosci*, 18(5), 637–646. doi:10.1038/nn.3980 [PubMed: 25734491]
- Kemp PJ, Rushton DJ, Yarova PL, Schnell C, Geater C, Hancock JM, ... Telezhkin V (2016). Improving and accelerating the differentiation and functional maturation of human stem cell-derived neurons: role of extracellular calcium and GABA. *J Physiol*, 594(22), 6583–6594. doi:10.1113/JP270655 [PubMed: 27616476]

- Knouse KA, Wu J, & Amon A (2016). Assessment of megabase-scale somatic copy number variation using single-cell sequencing. *Genome Res*, 26(3), 376–384. doi:10.1101/gr.198937.115 [PubMed: 26772196]
- Kuijlaars J, Oyelami T, Diels A, Rohrbacher J, Versweyveld S, Meneghello G, ... Verheyen A (2016). Sustained synchronized neuronal network activity in a human astrocyte co-culture system. *Sci Rep*, 6, 36529. doi:10.1038/srep36529 [PubMed: 27819315]
- Kuwabara T, Hsieh J, Muotri A, Yeo G, Warashina M, Lie DC, ... Gage FH (2009). Wnt-mediated activation of NeuroD1 and retro-elements during adult neurogenesis. *Nat Neurosci*, 12(9), 1097–1105. doi:10.1038/nn.2360 [PubMed: 19701198]
- Lacar B, Linker SB, Jaeger BN, Krishnaswami S, Barron J, Kelder M, ... Gage FH (2016). Nuclear RNA-seq of single neurons reveals molecular signatures of activation. *Nat Commun*, 7, 11022. doi:10.1038/ncomms11022 [PubMed: 27090946]
- Lake BB, Ai R, Kaeser GE, Salathia NS, Yung YC, Liu R, ... Zhang K (2016). Neuronal subtypes and diversity revealed by single-nucleus RNA sequencing of the human brain. *Science*, 352(6293), 1586–1590. doi:10.1126/science.aaf1204 [PubMed: 27339989]
- Lam RS, Topfer FM, Wood PG, Busskamp V, & Bamberg E (2017). Functional Maturation of Human Stem Cell-Derived Neurons in Long-Term Cultures. *PLoS One*, 12(1), e0169506. doi:10.1371/journal.pone.0169506 [PubMed: 28052116]
- Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurles ME, ... Knoblich JA (2013). Cerebral organoids model human brain development and microcephaly. *Nature*, 501(7467), 373–379. doi:10.1038/nature12517 [PubMed: 23995685]
- Linnarsson S (2013). Single-cell biology meeting marks rebirth of an old science. *Genome Biol*, 14(4), 305. doi:10.1186/gb-2013-14-4-305 [PubMed: 23607893]
- Liu SJ, Nowakowski TJ, Pollen AA, Lui JH, Horlbeck MA, Attenello FJ, ... Lim DA (2016). Single-cell analysis of long non-coding RNAs in the developing human neocortex. *Genome Biol*, 17, 67. doi:10.1186/s13059-016-0932-1 [PubMed: 27081004]
- Liu Y, Lopez-Santiago LF, Yuan Y, Jones JM, Zhang H, O'Malley HA, ... Parent JM (2013). Dravet syndrome patient-derived neurons suggest a novel epilepsy mechanism. *Ann Neurol*, 74(1), 128–139. doi:10.1002/ana.23897 [PubMed: 23821540]
- Livesey MR, Magnani D, Hardingham GE, Chandran S, & Wyllie DJ (2016). Functional properties of in vitro excitatory cortical neurons derived from human pluripotent stem cells. *J Physiol*, 594(22), 6573–6582. doi:10.1113/JP270660 [PubMed: 26608229]
- Lodato MA, Woodworth MB, Lee S, Evrony GD, Mehta BK, Karger A, ... Walsh CA (2015). Somatic mutation in single human neurons tracks developmental and transcriptional history. *Science*, 350(6256), 94–98. doi:10.1126/science.aab1785 [PubMed: 26430121]
- Lodato S, & Arlotta P (2015). Generating neuronal diversity in the mammalian cerebral cortex. *Annu Rev Cell Dev Biol*, 31, 699–720. doi:10.1146/annurev-cellbio-100814-125353 [PubMed: 26359774]
- Lui JH, Hansen DV, & Kriegstein AR (2011). Development and evolution of the human neocortex. *Cell*, 146(1), 18–36. doi:10.1016/j.cell.2011.06.030 [PubMed: 21729779]
- Lynch M (2010). Rate, molecular spectrum, and consequences of human mutation. *Proc Natl Acad Sci USA*, 107(3), 961–968. doi:10.1073/pnas.0912629107 [PubMed: 20080596]
- Macaulay IC, Ponting CP, & Voet T (2017). Single-Cell Multiomics: Multiple Measurements from Single Cells. *Trends Genet*, 33(2), 155–168. doi:10.1016/j.tig.2016.12.003 [PubMed: 28089370]
- Macaulay IC, & Voet T (2014). Single cell genomics: advances and future perspectives. *PLoS Genet*, 10(1), e1004126. doi:10.1371/journal.pgen.1004126 [PubMed: 24497842]
- Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, ... McCarroll SA (2015). Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell*, 161(5), 1202–1214. doi:10.1016/j.cell.2015.05.002 [PubMed: 26000488]
- Mainen ZF, & Sejnowski TJ (1996). Influence of dendritic structure on firing pattern in model neocortical neurons. *Nature*, 382(6589), 363–366. doi:10.1038/382363a0 [PubMed: 8684467]
- Manganas LN, Zhang X, Li Y, Hazel RD, Smith SD, Wagshul ME, ... Maletic-Savatic M (2007). Magnetic resonance spectroscopy identifies neural progenitor cells in the live human brain. *Science*, 318(5852), 980–985. doi:10.1126/science.1147851 [PubMed: 17991865]

- Mariani J, Simonini MV, Palejev D, Tomasini L, Coppola G, Szekely AM, ... Vaccarino FM (2012). Modeling human cortical development in vitro using induced pluripotent stem cells. *Proc Natl Acad Sci U S A*, 109(31), 12770–12775. doi:10.1073/pnas.1202944109 [PubMed: 22761314]
- Masland RH (2004). Neuronal cell types. *Curr Biol*, 14(13), R497–500. doi:10.1016/j.cub.2004.06.035 [PubMed: 15242626]
- Mason A, & Larkman A (1990). Correlations between morphology and electrophysiology of pyramidal neurons in slices of rat visual cortex. II. Electrophysiology. *J Neurosci*, 10(5), 1415–1428. [PubMed: 2332788]
- McConnell MJ, Lindberg MR, Brennand KJ, Piper JC, Voet T, Cowing-Zitron C, ... Gage FH (2013). Mosaic copy number variation in human neurons. *Science*, 342(6158), 632–637. doi:10.1126/science.1243472 [PubMed: 24179226]
- McConnell MJ, Moran JV, Abyzov A, Akbarian S, Bae T, Cortes-Ciriano I, ... Brain Somatic Mosaicism N (2017). Intersection of diverse neuronal genomes and neuropsychiatric disease: The Brain Somatic Mosaicism Network. *Science*, 356(6336). doi:10.1126/science.aal1641
- Migliore M, & Shepherd GM (2005). Opinion: an integrated approach to classifying neuronal phenotypes. *Nat Rev Neurosci*, 6(10), 810–818. doi:10.1038/nrn1769 [PubMed: 16276357]
- Mizrahi A (2007). Dendritic development and plasticity of adult-born neurons in the mouse olfactory bulb. *Nat Neurosci*, 10(4), 444–452. doi:10.1038/nn1875 [PubMed: 17369823]
- Moore AR, Filipovic R, Mo Z, Rasband MN, Zecevic N, & Antic SD (2009). Electrical excitability of early neurons in the human cerebral cortex during the second trimester of gestation. *Cereb Cortex*, 19(8), 1795–1805. doi:10.1093/cercor/bhn206 [PubMed: 19015375]
- Mountcastle VB, Talbot WH, Sakata H, & Hyvarinen J (1969). Cortical neuronal mechanisms in flutter-vibration studied in unanesthetized monkeys. Neuronal periodicity and frequency discrimination. *J Neurophysiol*, 32(3), 452–484. [PubMed: 4977839]
- Muller FJ, Schuldt BM, Williams R, Mason D, Altun G, Papapetrou EP, ... Loring JF (2011). A bioinformatic assay for pluripotency in human cells. *Nat Methods*, 8(4), 315–317. doi:10.1038/nmeth.1580 [PubMed: 21378979]
- Namba T, & Huttner WB (2017). Neural progenitor cells and their role in the development and evolutionary expansion of the neocortex. *Wiley Interdiscip Rev Dev Biol*, 6(1). doi:10.1002/wdev.256
- Neher E, & Sakmann B (1976). Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature*, 260(5554), 799–802. [PubMed: 1083489]
- Pasca AM, Sloan SA, Clarke LE, Tian Y, Makinson CD, Huber N, ... Pasca SP (2015). Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D culture. *Nat Methods*, 12(7), 671–678. doi:10.1038/nmeth.3415 [PubMed: 26005811]
- Pruszk J, Sonntag KC, Aung MH, Sanchez-Pernaute R, & Isacson O (2007). Markers and methods for cell sorting of human embryonic stem cell-derived neural cell populations. *Stem Cells*, 25(9), 2257–2268. doi:10.1634/stemcells.2006-0744 [PubMed: 17588935]
- Randall AD (2016). Are stem cell-derived neural cells physiologically credible? *J Physiol*, 594(22), 6569–6572. doi:10.1113/JP273348 [PubMed: 27870124]
- Rushton DJ, Mattis VB, Svendsen CN, Allen ND, & Kemp PJ (2013). Stimulation of GABA-induced Ca²⁺ influx enhances maturation of human induced pluripotent stem cell-derived neurons. *PLoS One*, 8(11), e81031. doi:10.1371/journal.pone.0081031 [PubMed: 24278369]
- Schroeder T (2011). Long-term single-cell imaging of mammalian stem cells. *Nat Methods*, 8(4 Suppl), S30–35. doi:10.1038/nmeth.1577 [PubMed: 21451514]
- Song M, Mohamad O, Chen D, & Yu SP (2013). Coordinated development of voltage-gated Na⁺ and K⁺ currents regulates functional maturation of forebrain neurons derived from human induced pluripotent stem cells. *Stem Cells Dev*, 22(10), 1551–1563. doi:10.1089/scd.2012.0556 [PubMed: 23259973]
- Steriade M (2004). Neocortical cell classes are flexible entities. *Nat Rev Neurosci*, 5(2), 121–134. doi:10.1038/nrn1325 [PubMed: 14735115]
- Tang X, Zhou L, Wagner AM, Marchetto MC, Muotri AR, Gage FH, & Chen G (2013). Astroglial cells regulate the developmental timeline of human neurons differentiated from induced

pluripotent stem cells. *Stem Cell Res*, 11(2), 743–757. doi:10.1016/j.scr.2013.05.002 [PubMed: 23759711]

Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, ... Zeng H (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat Neurosci*, 19(2), 335–346. doi:10.1038/nn.4216 [PubMed: 26727548]

Tsankov AM, Akopian V, Pop R, Chetty S, Gifford CA, Daheron L, ... Meissner A (2015). A qPCR ScoreCard quantifies the differentiation potential of human pluripotent stem cells. *Nat Biotechnol*, 33(11), 1182–1192. doi:10.1038/nbt.3387 [PubMed: 26501952]

Tyler WA, Medalla M, Guillamon-Vivancos T, Luebke JI, & Haydar TF (2015). Neural precursor lineages specify distinct neocortical pyramidal neuron types. *J Neurosci*, 35(15), 6142–6152. doi: 10.1523/JNEUROSCI.0335-15.2015 [PubMed: 25878286]

Upton KR, Gerhardt DJ, Jesuadian JS, Richardson SR, Sanchez-Luque FJ, Bodea GO, ... Faulkner GJ (2015). Ubiquitous L1 mosaicism in hippocampal neurons. *Cell*, 161(2), 228–239. doi:10.1016/j.cell.2015.03.026 [PubMed: 25860606]

Yokota Y, Gashghaei HT, Han C, Watson H, Campbell KJ, & Anton ES (2007). Radial glial dependent and independent dynamics of interneuronal migration in the developing cerebral cortex. *PLoS One*, 2(8), e794. doi:10.1371/journal.pone.0000794 [PubMed: 17726524]

Yu YC, Bultje RS, Wang X, & Shi SH (2009). Specific synapses develop preferentially among sister excitatory neurons in the neocortex. *Nature*, 458(7237), 501–504. doi:10.1038/nature07722 [PubMed: 19204731]

Zeisel A, Munoz-Manchado AB, Codeluppi S, Lonnerberg P, La Manno G, Jureus A, ... Linnarsson S (2015). Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science*, 347(6226), 1138–1142. doi:10.1126/science.aaa1934 [PubMed: 25700174]

Zhu G, Du L, Jin L, & Offenhauer A (2016). Effects of Morphology Constraint on Electrophysiological Properties of Cortical Neurons. *Sci Rep*, 6, 23086. doi:10.1038/srep23086 [PubMed: 27052791]

Zunder ER, Lujan E, Goltsev Y, Wernig M, & Nolan GP (2015). A continuous molecular roadmap to iPSC reprogramming through progression analysis of single-cell mass cytometry. *Cell Stem Cell*, 16(3), 323–337. doi:10.1016/j.stem.2015.01.015 [PubMed: 25748935]

Figure 1. Single cell approaches are essential to measure neuronal diversity.

(A) In a morphologically diverse population of neurons (left), transcriptomic diversity present only in a rare population of neurons is lost in bulk measurement. Schematized gene expression measurements from bulk tissue and from single cells (right) are used for illustration. Bulk analysis indicates that all neurons express high levels of gene A (orange), and low levels of gene B (green); whereas, single cell measurement identifies neuronal diversity. (B) Diversity among morphologically similar neurons (left) is also obscured by bulk measurement. As in (A, right), bulk measurement indicates that all neurons express equal levels of genes A and B; whereas, single cell measurement can reveal distinct subpopulations.

Figure 2. Genomic diversity may be greater within single neuronal subtypes, than among clonally-related neurons.

The molecular taxonomy of excitatory cortical neurons is based on laminar residence. Morphological, physiological, and transcriptomic diversity is catalogued by comparing layer VI neurons, to layer V neurons, for example. However, genomic diversity is expected to expand in an orthogonal direction. Radial cortical columns contain neurons of each subtype that were born from the same NPC, these clonally related progeny are expected to share somatic mutations. Tangentially distributed columns, on the other hand, arise from different NPCs and are expected to accumulate distinct mutational profiles from other columns.