EPA Public Access

Author manuscript

Ecol Modell. Author manuscript; available in PMC 2018 November 28.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Ecol Modell. 2017 November 10; 363: 17–30. doi:10.1016/j.ecolmodel.2017.08.020.

Parameter sensitivity and identifiability for a biogeochemical model of hypoxia in the northern Gulf of Mexico*

Marcus W. Beck^{*},

USEPA National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, 1 Sabine Island Drive, Gulf Breeze, FL 32561

John C. Lehrter,

University of South Alabama, Dauphin Island Sea Lab, Dauphin Island, AL 36528

Lisa L. Lowe, and

CSRA, Inc. supporting the USEPA, Research Triangle Park, NC 27709

Brandon M. Jarvis

USEPA National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, 1 Sabine Island Drive, Gulf Breeze, FL 32561

Abstract

Local sensitivity analyses and identifiable parameter subsets were used to describe numerical constraints of a hypoxia model for bottom waters of the northern Gulf of Mexico. The sensitivity of state variables differed considerably with parameter changes, although most variables were responsive to changes in parameters that influenced planktonic growth rates and less sensitive to physical or chemical parameters. Variation in sensitivity had a direct correspondence with identifiability, such that only small subsets of the complete parameter set had unique effects on the model output. Selecting parameters by decreasing sensitivity demonstrated that only eight of 51 total parameters had a sufficiently unique effect on model output for accurate calibration. As a result, parameter selection heuristics were used to identify parameters for model calibration that depended on combined effects on output, relative sensitivity of each parameter, and ecological categories for the biogeochemical equations. The calibrated zero-dimensional (0-D) unit of the hypoxia model had improved fit to the observed data if sensitive phytoplankton parameters were included in an identifiable subset. Extension of results to a three-dimensional grid of the Gulf of Mexico showed that sensitive parameters for the 0-D model translated to non-trivial changes in the areal estimates of hypoxia.

Keywords

Coastal General Ecosystem Model (CGEM); Gulf of Mexico (GOM); Hypoxia; Identifiability; Sensitivity

★Version: Thu Aug 17 18:08:14 2017 -0500, [7e3894b946e67efce7a075d46c2169be58729981](https://doi.org/10.1016/j.ecolmodel.2017.08.020)

*Corresponding author beck.marcus@epa.gov.

(Marcus W. Beck), jlehrter@disl.org (John C. Lehrter), lowe.lisa@epa.gov (Lisa L. Lowe), jarvis.brandon@epa.gov (Brandon M. Jarvis)

1. Introduction

Hypoxia formation in bottom waters of coastal oceans occurs primarily from excess nutrient inputs from land-based sources (Justić et al., 1987; Diaz and Rosenberg, 1995; Howarth et al., 1996). These events are detrimental to aquatic organisms and have significant negative effects on economic resources derived from coastal ecosystems (Lipton and Hicks, 2003; Diaz and Rosenberg, 2011). An understanding of the biological, physical, and chemical processes that influence hypoxic areas is a critical concern for mitigating and preventing these negative impacts. Numerical ecosystem models are important tools that synthesize knowledge of ecosystem processes that contribute to hypoxia formation and for predicting the effects of proposed management activities or future scenarios (Scavia et al., 2004; Hagy and Murrell, 2007; Camacho et al., 2014b; Pauer et al., 2016). Unlike statistical models with more generic structures, simulation and process-based models include explicit descriptions of relevant processes that are constrained by empirical or observational data relevant to the system of interest (e.g. Omlin et al., 2001b; Eldridge and Roelke, 2010). These models are often coupled with hydrodynamic grids to provide spatially-explicit representations of patterns in three dimensions (Warner et al., 2005; Dortch et al., 2007; Zhao et al., 2010; Ganju et al., 2016). Combined hydrodynamic and bio-geo-chemical models have been developed specifically to describe hypoxic conditions on the Louisiana continental shelf (LCS) in the northern Gulf of Mexico (GOM) (Fennel et al., 2013; Obenour et al., 2015; Pauer et al., 2016; Lehrter et al., 2017). This area drains a significant portion of the continental United States through the Mississippi-Atchafalaya River Basin (MARB) and is the second largest hypoxic area in the world (Rabalais et al., 2002). Understanding processes that contribute to the frequency and duration of hypoxic events remains a critical research goal for the region.

The development of a model represents a tradeoff between achieving predictive accuracy and a realistic representation of environmental processes. An ideal model is sufficiently generalizable across systems, provides results that are accurate given the inputs, and includes components that are valid descriptions of actual processes (Levins, 1966; Ganju et al., 2016). Given that these characteristics cannot be simultaneously achieved, models are developed to balance predictive accuracy with environmental realism or often favor one at the expense of the other (Morrison and Morgan, 1999; Ganju et al., 2016). These challenges are analogous to the well-known bias-variance tradeoff in statistical models that balances the competing objectives of over- and under-fitting to an observed dataset. Process-based models are more commonly imbalanced between reality and theory, such that most are over-parameterized in an attempt to completely describe reality (Denman, 2003; Nossent and Bauwens, 2012; Petrucci and Bonhomme, 2014). Quantitative limitations of over-parameterization are analogous to degrees of freedom in standard statistical models as free parameters cannot be numerically estimated when constrained to an observed dataset (Kirchner, 2006). More importantly, over-parameterization can limit use across systems outside of the data domain and impose uncertainty in model predictions as realistic values for every variable may not be known or inaccurately applied from existing studies (Durand et al., 2002; Refsgaard et al., 2007; Wade et al., 2008).

Model fit can be evaluated relative to the effects of initial conditions or the observed data used for calibration, changes in parameter values, or variation in the structural components (i.e., observational, parameter, or structural uncertainty) (Beck, 1987). Evaluating effects of parameter changes is by far the most common and simplest approach. Although sensitivity analyses should be integrated with model development, parameters are often evaluated post-hoc as a form of ‘damage control’ for further calibration. This approach is sometimes called inverse modelling where results from sensitivity analyses are used to guide calibration or fit of the developed model to observations (Soetaert and Petzoldt 2010, or confronting models with data, *sensu* Hilborn and Mangel 1997). Parameter sensitivity analysis combined with inverse modelling necessarily involves questions of parameter ‘identifiability’.

Redundancies in parameter effects lead to unidentifiable models where calibration is empirically impossible (i.e., standard algorithms will not converge) or parameter values may be non-unique leading to the right answer for the wrong reason (Kirchner, 2006).

Unidentifiable parameter sets have effects on model output that can be undone or compensated for by alteration of other parameters. Identifiability issues are not foreign to hypoxia or eutrophication models (Omlin et al., 2001a; Estrada and Diaz, 2010; Mateus and Franz, 2015), although there is a clear need for greater integration of these concepts in practice (Fasham et al., 2006).

This study describes a sensitivity and identifiability analysis of a zero-dimensional (0-D) unit of a larger spatial-temporal model of hypoxia dynamics on the LCS. In general, the analysis makes a case for unit-testing with lower-dimensional models, particularly when computational limitations preclude the use of conventional optimization algorithms with larger three-dimensional models. The objectives were to provide a statistical approach that demonstrates numerical limitations of parameter sets for model calibration and provide a framework for selecting parameters within the identifiability constraints. The specific goals were to 1) identify the parameters that have the greatest influence on state variables using local sensitivity analysis, 2) quantify the identifiability of subsets of the total parameter space based on sensitivity, and 3) provide a set of heuristics for choosing parameters based on sensitivity, identifiability, and parameter categories. The 0-D model was calibrated with selected parameter subsets to demonstrate use of the selection heuristics for optimization. Sensitive parameters for the 0-D model were also varied in the larger 3-dimensional model to demonstrate scalability of the results. In addition to dissolved oxygen (O_2), other state variables that were evaluated included ammonium, chlorophyll *a* (chl-*a*), irradiance, nitrate, particulate organic matter (POM), dissolved organic matter (DOM), and phosphorus. In general, we provide empirical results to support the assumption that models are generally over-parameterized and only a finite and smaller subset of the complete parameter set can be optimized.

2. Materials and Methods

2.1. Model description

Hypoxic events, defined as $<2 \text{ mg L}^{-1}$ of O_2 ($< 64 \text{ mmol m}^{-3}$), occur seasonally in bottom waters in the northern GOM. The hypoxic area averages $15,540 \text{ km}^2$ annually (1993–2015) with minimum concentrations observed from late spring to early fall. Seasonal variation is

strongly related to carbon and nutrient export from the MARB (Lohrenz et al., 2008; Bianchiet al., 2010), whereas hydrologic variation, currents, and wind patterns can affect vertical salinity gradients that contribute to hypoxia formation (Wiseman et al., 1997; Paerl et al., 1998; Obenour et al., 2015). The Coastal General Ecosystem Model (CGEM) was developed to describe hypoxia dynamics on the LCS and includes elements from the Navy Coastal Ocean Model (Martin, 2000) for hydrodynamics and a biogeochemical model with multiple plankton groups, water-column metabolism, and sediment diagenesis (Eldridge and Roelke, 2010). The hydrodynamic component of CGEM provides a spatially-explicit description of hypoxia using an orthogonal grid with an approximate horizontal resolution of 1.9 km² and twenty equally-spaced vertical sigma layers on the shelf. The biogeochemical component includes equations for 36 state variables including six phytoplankton groups (with nitrogen and phosphorus quotas for each), two zooplankton groups, nitrate, ammonium, phosphate, dissolved inorganic carbon, oxygen, silica, and multiple variables for dissolved and particulate organic matter from different sources.

The core unit of CGEM is FishTank, a 0-D model that implements the biogeochemical equations in Eldridge and Roelke (2010) and does not include any advection, mixing, or sediment diagenesis (Fig. 1). A full description of the model structure, equations, and parameters is described in appendices A-F in Lehrter et al. (2017). Although FishTank was developed for specific application in CGEM, it can easily be applied to other hydrodynamic grids. Results are based on time-dependent differential equations that describe energy flow between phytoplankton and zooplankton groups given nutrient uptake rates, organic matter inputs and losses, inherent optical properties, and temperature (Penta et al. 2008; Eldridge and Roelke 2010, see appendices in Lehrter et al. 2017). A total of 108 equations are estimated at each time step to return values for each of the 36 state variables described by the model. In addition to the initial conditions, 251 parameter values for each of the equations are also supplied at model execution. Values for each of the parameters were based on estimates from the literature, field or laboratory-based measurements, or expert knowledge in absence of the former. As such, a sensitivity analysis of parameter values is warranted given that, for example, literature or field-based estimates may not apply under all scenarios or expert knowledge is not completely certain (Refsgaard et al., 2007).

The sensitivity of state variables to perturbations of all relevant parameters for the 108 equations was estimated using a five minute timestep with daily output from January 1st to December 31st, 2006. Irrelevant parameters were removed for several reasons; parameters were not relevant for the 0-D model (i.e., hydrodynamic parameters), were considered physical constants, or had no effect given initial conditions. Additionally, FishTank includes six phytoplankton and two zooplankton groups to add complexity in community structure and foodweb dynamics. To remove obvious redundancies, the sensitivity analyses were conducted using only one phytoplankton and one zooplankton group. The final set that was evaluated included 51 parameters that were further grouped into one of six categories based on applicable biogeochemical components of the model: optics ($n=4$ parameters), organic matter (12), phytoplankton (22), temperature (2), and zooplankton (11) (Table 1). A full description of the model parameters is available as an appendix in Lehrter et al. (2017).

2.2. Local sensitivity analysis

A local sensitivity analysis was performed by evaluating the change in state variables following perturbation of each parameter from its original value (Soetaert and Petzoldt, 2010; Camacho et al., 2014a; RDCT (R Development Core Team), 2017). This approach is described as a First Order Variance Analysis that linearly propagates changes from model parameters to model predictions (Camacho et al., 2014a). Parameters were individually perturbed by a 50% increase of the original values (Table 1) and sensitivity S was estimated for each time step i given a change for parameter j as:

$$S_{ij} = \frac{\partial y_i}{\partial \Theta_j} \cdot \frac{w_{\Theta_j}}{w_{y_i}} \quad (1)$$

where the estimate is based on the change in the predicted value for response variable y divided by the change in the parameter Θ_j multiplied by the quotient of scaling factors w for each. The scaling factors, w_{Θ_j} for the parameter Θ_j and w_{y_i} for response variable y_i , were set as the default value of the unperturbed parameter and the predicted value of y_i after perturbation (Soetaert and Petzoldt, 2010). Scaling makes the estimates unitless to compare model sensitivity to parameters and state variables that differ in relative magnitude. Sensitivity values for all j parameters were summarized as a single value across the time series from $i = 1$ to n as $L1$:

$$L1 = \sum |S_{ij}| / n \quad (2)$$

All parameters for each of the six equation categories (optics, organic matter, phytoplankton, temperature, and zooplankton) that had non-zero $L1$ were retained for identifiability analysis.

2.3. Identifiability and selecting parameter subsets

The collinearity index γ provides a measure of potential redundancies in the response of a state variable to changes in parameter values. The index measures the linear dependence between sensitivity functions (i.e., S_j for j parameters) described above for parameter subsets and was estimated from the minimum eigenvector of the cross-product of a selected sensitivity matrix (Brun et al., 2001; Omlin et al., 2001a):

$$\gamma = \frac{1}{\sqrt{\min(\text{EV}[\hat{S}^T \hat{S}])}} \quad (3)$$

where γ ranges from one to infinity for perfectly identifiable (orthogonal) or unidentifiable (perfectly collinear) parameter sets. The sensitivity functions were supplied as a matrix $\hat{S}$ with rows i and columns j (eq. (1)) that described deviations of predicted O_2 following perturbations of each parameter. Thus, γ can be estimated from results for any subset of

parameter combinations. Sensitivity matrices were first normalized by dividing by the square root of the summed residuals (Omlin et al., 2001a; Soetaert and Petzoldt, 2010). Estimates of γ greater than 10–15 suggest parameter sets are poorly identifiable (Brun et al., 2001; Omlin et al., 2001a), meaning parameter values that maximize fit on a calibration dataset are inestimable by conventional optimization algorithms. An intuitive interpretation of γ is provided by Brun et al. (2001), such that a change in a state variable caused by a change in one parameter can be offset by the fraction $1 - 1/\gamma$ by the remaining parameters. That is, $\gamma = 10$ suggests the relative change in O_2 for an arbitrary parameter in the selected set can be compensated for by 90% with changes in the other parameters.

Parameter selection for model calibration must consider the competing objectives of increased goodness of fit with parameter inclusion and reduced identifiability as it relates to optimization. An additional challenge is a large number of combinations of parameter sets, which complicates selection given sensitivity differences and desired ecological categories of each parameter (e.g., parameters for a group of related structural equations could be of interest). Fig. 2 provides a simple graphic of the unique number of combinations that are possible for different subsets of ‘complete’ parameter sets of different sizes (i.e., n choose k combinations, $n! / (k! (n - k)!)$). The number of unique combinations increases with the total parameters in the set and is also maximized for moderate selections (e.g., selecting half the total). For example, over 10^{14} combinations are possible by selecting 25 parameters from a set of 50.

A set of heuristics was developed that addresses the tradeoff in model complexity and identifiability given the challenges described above (see also Wagener et al., 2001). These rulesets were developed to select parameters with preference for those with high sensitivity and identifiability based on $\gamma < 15$ as an acceptable threshold for subsets (e.g., 93% accountability between parameters). Heuristics for parameter selection also recognized that parameter categories (i.e., optics, organic matter, phytoplankton, temperature, zooplankton) may have unequal preferences by users given desired application of the model. For all selection heuristics, parameters were selected by decreasing sensitivity starting with the most sensitive until identifiability did not exceed $\gamma = 15$ where selections were 1) blocked within each of five parameter categories, 2) independent of parameter category, 3) or considering all categories equally.

2.4. Model calibration and scalability

Selected parameter subsets from the seven heuristics were used to calibrate the 0-D model. This analysis demonstrated that different parameter sets can produce model output with differing goodness of fit and calibration effort given 1) differences in variable sensitivity to parameters in each subset, 2) differences in the parameters selected from different categories (heuristic difference), and 3) differences in identifiability values. Generally, this analysis was a proof of concept that the selection heuristics can lead to an ‘optimal’ model by maximizing the balance between sensitivity and identifiability.

Estimates of oxygen production from closed bottle experiments were used to calibrate the 0-D model. Surface water was collected from a fixed location in Pensacola Bay, Florida (30.35° N, 87.20° W) on September 25th, 2013 (full details in Murrell et al. in revision).

Water samples were placed into closed 300 mL glass bottles and transported back to the lab for treatments. Concentrations of O_2 and nutrients (NH_4^+ , NO_2^- , NO_3^- , PO_4^{3-} , and SiO_3^{2-}) were extracted from the samples within one hour of collection. Oxygen concentrations were established using Winkler reagents (Parsons et al., 1985) and titration using a Metrohm Titrando with thiosulfate titrant calibrated against an iodate standard and electrochemical endpoint detection. Ammonium was analyzed using a fluorometric method (Holmes et al., 1999); other nutrients were analyzed using either a Thermo Fisher Aquakem 200 discrete analyzer or an Astoria-Pacific continuous flow analyzer using standard colorimetric methods (APHA 2005).

Oxygen production after an approximate 12 hour period (morning to evening) was estimated within the bottles using light and dark treatments. All treatments were exposed to ambient outdoor conditions in a water bath at the laboratory. The dark treatments used opaque black bottles to ensure no exposure of ambient light to the samples. The exposure experiments began at 7am and were concluded at 7pm, after which the concentrations of O_2 in each bottle were measured. Each treatment had four replicates and HOBO[®] pendants were used to measure continuous temperature and light. Starting O_2 concentrations for the experiments were $200.78 (\pm 4.47) \text{ mmol } O_2 \text{ m}^{-3}$ and ending concentrations were $229.57 (\pm 2.66) \text{ mmol } O_2 \text{ m}^{-3}$ for the light treatments and $195.51 (\pm 1.9) \text{ mmol } O_2 \text{ m}^{-3}$ for the dark treatments. Initial nutrient concentrations were 0.71 mmol m^{-3} for NH_4^+ (single sample), 0.13 mmol m^{-3} for PO_4^{3-} , and 38.4 mmol m^{-3} for SiO_3^{2-} . Concentrations for NO_2^- and NO_3^- were below detection limit.

Each parameter subset for O_2 from the selection heuristics was calibrated to the ending oxygen concentration for each of the four replicates of each treatment. The initial nutrient and oxygen concentrations were used as starting conditions for Fish Tank. The corresponding continuous temperature and light measurements were also used as forcing conditions. Parameter values were searched using the optim function in R that implemented the limited memory modification of the BFGS quasi-Newton method (Byrd et al., 1995; Nocedal and Wright, 2006; RDCT (R Development Core Team), 2017). This optimization scheme searches for values within the user-defined lower and upper bounds of each parameter. We set the algorithm to search within a range of values for a subset that was chosen to best reflect expectations given professional judgment and reasonable bounds from the literature. While we recognize that ‘actual’ parameter values could possibly extend beyond these ranges, constraining the values was expected to produce output that was more interpretable within the context of the sensitivity functions defined above. Moreover, precise limits on each parameter were less of a concern for the current analysis as our primary goal was to demonstrate differences in fit as a function of sensitivity and identifiability. The optimal values for each parameter were identified based on convergence of the model results as a minimization of the root mean squared error (*RMSE*) between the observed and modelled data.

As a final analysis, effects of parameter changes on the larger 3-dimensional model were evaluated to demonstrate that results from the 0-D FishTank model are scalable. For simplicity, only the most sensitive parameter in each category was selected. The CGEM

model was run for one year using the default parameter conditions and by increasing each of the selected parameters by 50%. This produced six spatially-referenced time series (one default, five for each parameter change) from which changes in the daily hypoxic area (number of cells < 64 mmol O₂) were estimated.

3. Results

3.1. Local sensitivity analysis

Local sensitivity analyses showed that O₂ was sensitive to perturbations in 38 of the 51 (75% of total) parameters that were evaluated in FishTank (Fig. 3, Table 2). Within each parameter category, O₂ was sensitive to three parameters for optics (75% of all optic parameters), eight for organic matter (67%), 16 for phytoplankton (73%), one for temperature (50%), and 10 for zooplankton (91%). Although O₂ had the greatest sensitivity to parameters in the zooplankton category (as percentage of total), the relative effects varied. Among all parameters, average sensitivity was $L1 = 9.2e^{-3}$ with values ranging from $L1 = 8.34e^{-8}$ for *QminP* (phytoplankton) to 0.05 for *umax* (phytoplankton). Within categories (excluding temperature with one sensitive parameter), sensitivity ranged from $4.39e^{-5}$ (*astarOMA*) to $7.51e^{-4}$ (*astar490*) for optics, $4.17e^{-4}$ (*KNH4*) to $6.15e^{-3}$ (*KGI*) for organic matter, $8.34e^{-8}$ (*QminP*) to 0.05 (*umax*) for phytoplankton, and $3.69e^{-5}$ (*ZQp*) to 0.05 (*ZKa*) for zooplankton (Table 2). Average sensitivity values in each category were $L1 = 2.81e^{-4}$ for optics, $2.17e^{-3}$ for organic matter, 0.02 for temperature, 0.01 for phytoplankton, and 0.01 for zooplankton.

Local sensitivity analyses for the additional state variables (ammonium, chl-a, irradiance, nitrate, POM, DOM, and phosphorus) had similar results as O₂ with some exceptions (Fig. 3 and Tables S1 to S7). All variables were sensitive to the same parameters as O₂ (38 of 51 evaluated), although average sensitivity differed between variables. Average $L1$ ranged from 0.02 for irradiance (Table S3) to 0.71 for DOM (Table S6). All average sensitivity values for the state variables were higher than the average for O₂ ($L1 = 9.2e^{-3}$). For each variable, $L1$ ranged from $2.24e^{-6}$ (*QminP*) to 8.49 (*mA*) for ammonium (Table S1), $1.38e^{-6}$ (*QminP*) to 13.94 (*mA*) for chl-a (Table S2), $1.92e^{-7}$ (*QminP*) to 0.13 (*ZKa*) for irradiance (Table S3), $6.67e^{-7}$ (*QminP*) to 8.49 (*umax*) for nitrate (Table S4), $6.41e^{-5}$ (*KNH4*) to 7.22 (*mA*) for POM (Table S5), $7.41e^{-5}$ (*KNH4*) to 14.25 (*mA*) for DOM (Table S6), and $8.21e^{-7}$ (*QminP*) to 1.47 (*ZKa*) for phosphate (Table S7). For the parameter categories, ammonium was most sensitive to phytoplankton parameters (average $L1 = 0.8$ across all parameters in the category), chl-a to phytoplankton ($L1 = 1.14$), irradiance to zooplankton ($L1 = 0.03$), nitrate to zooplankton ($L1 = 1.06$), POM to temperature ($L1 = 0.86$), DOM to temperature ($L1 = 1.48$), and phosphate to zooplankton ($L1 = 0.31$). Finally, average sensitivity between parameter categories independent of the state variables ranged from $8.38e^{-3}$ for optics (average $L1$ across all variables) to 0.62 for phytoplankton.

3.2. Identifiability of parameter subsets and selection rules

The identifiability analyses suggested that many parameter subsets exceeded the thresholds of $\gamma = 10, 15$. Parameter identifiability for O₂ decreased (increasing γ) at different rates with increasing size of parameter subsets depending on the parameter category or the number of

top parameters that were selected (Fig. 4). By category, identifiability was lowest for all combinations of parameter subsets in the phytoplankton (60% of subsets less than $\gamma = 15$, 43% less than $\gamma = 10$) and zooplankton categories (53.1% less than $\gamma = 15$, 40% less than $\gamma = 10$), whereas all combinations were identifiable for optics (100% less than $\gamma = 15, 10$) and a majority identifiable for organic matter (91.9% less than $\gamma = 15$, 76.5% less than $\gamma = 10$). Identifiability for parameters in the temperature category was not evaluated because O₂ was sensitive to only one parameter (i.e., $\gamma = 1$). Parameter combinations for choosing from the top, top two, top three, and top four parameters in each category together had decreasing identifiability with the increasing size of the selection pool (e.g., top one versus top four parameters, Fig. 4). The percentage of parameter subsets that were below the acceptable thresholds for identifiability was 100% less than $\gamma = 15, 10$ for the top parameter in each category, 90.6% and 80.7% for the top two, 80.7% and 70.9% for the top three, and 55.8% and 45.7% for the top four. Results for the remaining state variables had similar patterns in identifiability with increasing size of parameter subsets and selection categories, although differences in identifiability between state variables was observed (Fig. 5). Most notably, nitrate was consistently the least identifiable variable (highest overall γ), whereas O₂ was most identifiable.

An evaluation of the effects of individual parameters on γ suggested that some parameters had disproportionate effects on identifiability. For example, Fig. 4 suggests that most parameter sets for organic matter were identifiable ($\gamma < 15$ for most), regardless of how many parameters were selected (i.e., two through eight). However, some subsets were not identifiable such that identification of one or more redundant parameters that are inflating γ values could provide useful information. Fig. 6 shows an alternative view of identifiability of O₂ with exclusion and inclusion of individual parameters in different sets for the organic matter category. As before, collinearity increases with more parameters in a subset, although the increase varies depending on which parameter was included or excluded from the set. For example, inclusion of *KNO3* in a parameter set almost always inflated γ . All parameter subsets that did not include *KNO3* were well below $\gamma = 15$, suggesting that exclusion of this parameter improves identifiability. Interestingly, the inclusion of some parameters caused a reduction in γ , which contradicts the general rule that more parameters caused reduced identifiability. For example, parameter sets that included *KGcdom* generally had lower γ values relative to those that excluded the parameter.

Results for each of the three selection heuristics (blocked by parameter category, independent of category, all categories equally) applied to each state variable differed in the number of selected parameters and distribution of parameters within each category (Tables 3 to 5). In general, a correspondence was observed between the number of parameters that were selected given the threshold of $\gamma = 15$ and relative identifiability between the state variables. As noted above, nitrate was the least identifiable variable (Fig. 5), whereas other variables (e.g., O₂, irradiance) were more identifiable. The constraints on identifiability between variables were demonstrated with the selection heuristics. For example, heuristics for nitrate typically selected only one or two parameters that met the criteria as compared to more identifiable variables that included several parameters. Overall, the first selection heuristic demonstrated that the number of parameters chosen by parameter category differed independently of the state variables (Table 3). The number of selected parameters averaged

across state variables in decreasing order was 4.25 parameters from the phytoplankton category, 3.5 from organic matter, 2.75 from optics, and 2.38 from zooplankton. The second and third selection heuristics (Tables 4 and 5) were similar, although more parameters were generally selected for the third heuristic given equal importance between categories.

3.3. Model calibration and scalability

Calibration of parameter values for O_2 for each of the seven selection heuristics (Tables 3 to 5) produced model output that varied in goodness of fit and computational effort required to identify optimal values (Table 6). As expected, the number of required iterations of the optimization algorithm to minimize *RMSE* of observed O_2 and model output varied in proportion to the number of parameters in a subset. For example, only 12 iterations were required to optimize the single parameter selected for the temperature category, whereas 615 iterations were required for the seven parameters selected for phytoplankton (Fig. 7). No association was observed between collinearity (γ) and number of iterations or calibration. Of the seven parameter subsets, the optimal phytoplankton parameters had the largest reduction in *RMSE* values from the initial output (starting *RMSE* 21.14 mmol O_2 m⁻³, reduction to 5.81). The identifiable parameters selected equally within categories (third heuristic) and independent of category (second heuristic) produced the second and third greatest reduction in *RMSE* (19.09, and 19.45, respectively), although much less than that observed for phytoplankton parameters. Overall, calibration to most of the parameter subsets had a small effect on *RMSE*. However, the magnitude of parameter changes for calibration varied up to several orders from the initial values (Table 7). The percent change in parameters from the initial values varied from no change (optics parameters *astarOMZ* and *astarOMA*) to 900% (organic matter parameter *KO2*). Many calibrated parameter values included those at the lower or upper bounds used for the optimization routine (as for phytoplankton, Fig. 7). There was no relationship between magnitude of parameter changes within a subset and overall reduction of *RMSE*.

The estimated areal extent of bottom-water hypoxia using initial parameter values for the 3-dimensional model peaked on August, 28th at 1.19e⁴ km² (Fig. 8a, August mean area 8988 ± 551 km², Fig. 8b). Increases in the top sensitive parameters for O_2 in each category (Table 2) increased the estimated areal extent of hypoxia (Figs. 8c, 8e), with the exception of the parameter for optimum temperature of growth (*Tref(nospA+nospZ)_{p1}*), which caused a decrease (mean difference from July default area -1070 ± 1161 km², Fig. 8d). The largest increase in area was caused by an increase in the maximum growth rate of phytoplankton (*umax*), with the greatest increase occurring in July (monthly mean increase of 4109 ± 1085 km², Fig. 8d). Overall, changes in hypoxia extent were largest for parameters with large sensitivity values (i.e., L1 = 0.05 for *ZKa*, L1 = 0.05 for *umax*).

4. Discussion

State variables were most sensitive to phytoplankton and zooplankton parameters, particularly the maximum growth rates (*umax* for phytoplankton, *Zumax* for zooplankton), mortality coefficient for phytoplankton (*mA*), and the zooplankton half saturation coefficient for grazing (*ZKa*). An increase in the growth rate of primary producers can increase oxygen

concentration through photosynthetic processes, although increased production of organic matter is balanced with respiration and bacterial decomposition that reduce O₂ in the water column. Similarly, increases in zooplankton abundance with increased growth rates can reduce phytoplankton biomass through grazing, which is expected to further deplete pools of organic matter from algal sources. Most variables were also sensitive to variation in the halfsaturation grazing coefficient which moderates nutrient concentrations that support half the maximum grazing rate. Although the tradeoff between abundance, grazing, and decomposition is complex, the sensitivity of model state variables to parameters that directly control the abundance of primary producers is in agreement with empirical observations of factors that influence hypoxia dynamics on the LCS (Fahnenstiel et al., 1995; Roelke, 2000; Eldridge and Roelke, 2010). The sensitivity of the model output to variation in other parameters that relate to physical and chemical properties of the system was of secondary importance. That is, state variables were sensitive to changes in light and temperature parameters, although to a lesser extent than biological parameters. As such, the differing sensitivities of state variables to parameters in each of the categories was not unexpected given ecological relationships that are well understood and described by the model.

A general conclusion from the identifiability analyses is that only limited subsets of parameters were identifiable within the constraints of local sensitivity analyses. These results support previous studies that have suggested similarly small subsets of parameters can be identified using traditional calibration schemes (e.g., Wheeler et al., 1986; Ye et al., 1997; Omlin et al., 2001a). In addition to CGEM, these conclusions have relevance for other biogeochemical models that include numerous parameters and structural equations to characterize processes in the model domain. A general conclusion is that less complex models are potentially beneficial given that only a small subset of parameters is identifiable and that ecosystem processes may in fact be sufficiently characterized with few parameters (Ye et al., 1997). Conversely, others have argued that model complexity is not in itself a disadvantage when parsimony is not the only determinant of model structure (Reichert and Omlin, 1997). Over-parameterization can be useful if processes have importance that were not evaluated during model identification. Single objective functions that maximize model fit with identifiable parameters may also provide an incomplete characterization of model worth, which has prompted the development of probability-based models of hypoxia that explicitly include uncertainty in model components (e.g. Obenour et al., 2015). Our results demonstrated that approximately 75% of the evaluated parameters had an effect on the eight state variables, whereas CGEM includes a total of 36 variables and multiple plankton groups, not all of which have immediate concern for understanding hypoxia. The redundancies identified with the sensitivity analyses are challenging only if the primary interest is, for example, O₂ dynamics. Moreover, the proposed selection heuristics provide flexibility for choosing different parameters with the assumption that those chosen depend on the research or management question.

Results from the identifiability analyses provided additional insight into the interactions of parameters in large biogeochemical models. First, identifiability of parameter subsets was not related to the sensitivity of individual variables. As noted above, an identifiable parameter is one that has a unique effect on model predictions that cannot be compensated for or undone by changing other parameters. The magnitude of the effect of a parameter has

no bearing on identifiability, which further complicates the selection of parameters for calibration. Although identifiability is the primary limiting factor in choosing a set, the relative sensitivities are more important for the decision to include or exclude individual parameters. For example, Table 6 and Fig. 8 demonstrated that parameter sets with sensitive parameters had improved model fit and that these parameters had effects that translated to the larger model. Our analysis addressed this challenge by presenting multiple selection criteria for identifiable parameter sets that prioritized the most sensitive parameters during the selection process. Similarly, identifiability was not always related to the number of parameters in a set. Although the general trend was decreasing identifiability with more parameters, the unique effects of including an individual parameter with an existing set often reduced the γ estimate. For example, Fig. 6 showed that including *KGcdom*, *KO2*, or *nitmax* in parameter sets more often reduced γ relative to sets that excluded the parameters. The selection criteria proposed above can facilitate parameter selection and also provide diagnostic tools to identify parameters with disproportionate effects on γ .

4.1. Recommendations and conclusions

Our results demonstrated that small parameter subsets relative to all sensitive parameters were within the identifiability thresholds described in the literature. The identifiable parameter subsets varied considerably between state variables and the method for parameter selection. Use of identifiable parameter subsets with sensitive parameters improved model fit, particularly for phytoplankton parameters. However, calibration to most parameter sets showed little improvement in model fit despite relatively large changes in parameter values. This result underscores the challenge of model optimization when considering many parameters. Although we demonstrated model sensitivity to individual parameters, the interactive effects are also important. For example, optimization to zooplankton parameters showed almost no change in model *RMSE* despite a relatively large sensitivity of the model to individual parameters. A more comprehensive approach that considers the interaction of zooplankton parameters with others, such as phytoplankton, is warranted given the structural dependencies between parameter categories. Indeed, the reduction in *RMSE* for selection heuristics that equally considered the parameter categories (third heuristic) showed the second largest reduction in model error. This subset included both phytoplankton and zooplankton parameters, suggesting more careful consideration of these sensitive parameters could further improve model calibration.

The effects of structural or observational uncertainty could be evaluated as an extension of the analyses presented above. For example, the sensitivity of O_2 to variation in the halfsaturation constants for phytoplankton (the concentration supporting half the maximum uptake rate of nutrients) could vary given the initial nutrient concentrations (Eppley et al., 1969). Further, changes in the ratio between nitrogen and phosphorus could affect the sensitivity of state variables to parameter changes depending on the limiting nutrient. A more challenging analysis is an evaluation of the effects of structural components on model output, which requires exclusion or inclusion of explicit biogeochemical equations. The FishTank model includes several ‘switches’ that allow users to change the governing equations that estimate state variables, such as switches that ‘turn on’ different structural equations for light attenuation in the water column. This design is uncommon in

biogeochemical models and could be leveraged for an evaluation of structural uncertainty. As such, our analysis of parameter sensitivity and identifiability could be combined with an evaluation of observational and structural uncertainty for a more complete characterization of the model, having further implications for understanding drivers of hypoxia in coastal waters.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the research staff of the USEPA Gulf Ecology Division, Environmental Modeling and Visualization Laboratory, and Naval Research Laboratory for their efforts developing the hydrodynamic and biogeochemical components of CGEM. We thank James Pauer, Yonghsan Wan, Susan Elizabeth George, and an anonymous reviewer for providing helpful comments on an earlier draft. We thank Michael Murrell for assistance with the bottle dataset.

Role of Funding Sources

This study was funded, reviewed, and approved for publication by the US EPA, National Health and Environmental Effects Research Laboratory. However, the views expressed in this paper are those of the authors and do not necessarily reflect the views or policies of the US EPA. Mention of trade names or commercial products does not constitute endorsement by the authors or the US EPA.

Acronyms:

chl-a	chlorophyll a
CGEM	Coastal General Ecosystem Model
O₂	dissolved oxygen
DOM	dissolved organic matter
GOM	Gulf of Mexico
LCS	Louisiana continental shelf
MARB	Mississippi-Atchafalaya River Basin
POM	particulate organic matter
RMSE	root mean squared error
0-D	zero-dimensional

References

- APHA (American Public Health Association), 2005 Standard methods for the examination of water and wastewater. Washington, DC.
- Beck MB, 1987 Water quality modeling: A review of the analysis of uncertainty. Water Resour. Res 23 (8), 1393–1442.

- Bianchi TS, DiMarco SF, Hetland RD, Chapman P, Day JW, Allison MA, 2010 The science of hypoxia in the Northern Gulf of Mexico: a review. *Sci. Total. Environ* 408 (7), 1471–1484. [PubMed: 20092873]
- Brun R, Reichert P, Künsch HR, 2001 Practical identifiability analysis of large environmental simulation models. *Water Resour. Res* 37 (4), 1015–1030.
- Byrd RH, Lu P, Nocedal J, Zhu C, 1995 A limited memory algorithm for bound constrained optimization. *SIAM J. Sci. Comput* 16 (5), 1190–1208.
- Camacho RA, Martin JL, Diaz-Ramirez J, McAnally W, Rodriguez H, Suscy P, Zhang S, 2014a Uncertainty analysis of estuarine hydrodynamic models: an evaluation of input data uncertainty in the Weeks Bay estuary, Alabama. *Appl. Ocean. Res* 47, 138–153.
- Camacho RA, Martin JL, Watson B, Paul MJ, Zheng L, Stribling JB, 2014b Modeling the factors controlling phytoplankton in the St. Louis Bay estuary, Mississippi and evaluating estuarine responses to nutrient load modifications. *J. Environ. Eng* 141 (3), 04014067.
- Denman KL, 2003 Modelling planktonic ecosystems: parameterizing complexity. *Prog. Oceanogr* 57 (3–4), 429–452.
- Diaz RJ, Rosenberg R, 1995 Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanogr. Mar. Biology* 33, 245–303.
- Diaz RJ, Rosenberg R, 2011 Introduction to environmental and economic consequences of hypoxia. *Int. J. Water Resour. Dev* 27 (1), 71–82.
- Dortch MS, Zakikhani M, Noel MR, Kim SC, 2007 Application of a water quality model to Mississippi Sound to evaluate impacts of freshwater diversions Tech. rep., Engineer Research and Development Center, Vicksburg, Mississippi.
- Durand P, Gascuel-Oudou C, Cordier MO, 2002 Parameterisation of hydrological models: a review and lessons learned from studies of an agricultural catchment (Naisin, France. *Agron* 22 (2), 217–228.
- Eldridge PM, Roelke DL, 2010 Origins and scales of hypoxia on the Louisiana shelf: importance of seasonal plankton dynamics and river nutrients and discharge. *Ecol. Model* 221 (7), 1028–1042.
- Eppley RW, Rogers JN, McCarthy J, 1969 Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton. *Limnol. Oceanogr* 14 (6), 912–920.
- Estrada V, Diaz M, 2010 Global sensitivity analysis in the development of first principle- based eutrophication models. *Environ. Model. Softw* 25, 1539–1551.
- Fahnenstiel GL, McCormick MJ, Lang GA, Redalje DG, Lohrenz SE, Markowitz M, Wagoner B, Carrick HJ, 1995 Taxon-specific growth and loss rates for dominant phytoplankton populations from the northern Gulf of Mexico. *Mar. Ecol. Prog. Ser* 117 (1–3), 229–239.
- Fasham MJR, Flynn KJ, Pondaven P, Anderson TR, Boyd PW, 2006 Development of a robust marine ecosystem model to predict the role of iron in biogeochemical cycles: A comparison of results for iron-replete and iron-limited areas, and the SOIREE iron-enrichment experiment. *Deep. Res I* 53, 333–366.
- Fennel K, Hu J, Laurent A, Marta-Almeida M, Hetland R, 2013 Sensitivity of hypoxia predictions for the northern Gulf of Mexico to sediment oxygen consumption and model nesting. *J. Geophys. Res. Ocean* 118 (2), 990–1002.
- Ganju NK, Brush MJ, Rashleigh B, Aretxabaleta AL, del Barrio P, Gear JS, Harris LA, Lake SJ, McCardell G, O'Donnell J, Ralston DK, Signell RP, Testa JM, Vaudrey JMP, 2016 Progress and challenges in coupled hydrodynamic- ecological estuarine modeling. *Estuaries Coasts* 39 (2), 311–332. [PubMed: 27721675]
- Hagy JD, Murrell MC, 2007 Susceptibility of a northern Gulf of Mexico estuary to hypoxia: An analysis using box models. *Estuar. Coast. Shelf Sci* 74, 239–253.
- Hilborn R, Mangel M, 1997 *The Ecological Detective: Confronting Models with Data*. Princeton University Press, Princeton, New Jersey.
- Holmes RM, Amino A, Kerouel R, Hooker BA, Peterson BJ, 1999 A simple precise method for measuring ammonium in marine and freshwater ecosystems. *J. Fish. Aquat. Sci* 56 (10), 1801–1808.
- Howarth RW, Billen G, Swaney D, Townsend A, Jaworski N, Lajtha K, Downing JA, Elmgren R, Caraco N, Jordan T, Berendse F, Freney J, Kudeyarov V, Murdoch P, Zhao-Liang Z, 1996 Regional

- nitrogen budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean: natural and human influences. *Biogeochem.* 35 (1), 75–139.
- Justić D, Legović T, Rottini-Sandrini L, 1987 Trends in oxygen content 1911–1984 and occurrence of benthic mortality in the northern Adriatic Sea. *Estuar. Coast. Shelf Sci* 4), 435–445.
- Kirchner JW, 2006 Getting the right answers for the right reasons: Linking measurements, analyses, and models to advance the science of hydrology. *Water Resour. Res* 42 (3), W03S04.
- Lehrter JC, Ko DS, Lowe LL, Penta B, 2017 Predicted effects of climate change on northern Gulf of Mexico hypoxia In: Justic et al. (Ed.), *Modeling Coastal Hypoxia: Numerical Simulations of Patterns, Controls, and Effect of Dissolved Oxygen Dynamics*. Springer, New York, pp. 173–204.
- Levins R, 1966 The strategy of model building in population biology. *Am. Sci* 54 (4), 421–431.
- Lipton D, Hicks R, 2003 The cost of stress: low dissolved oxygen and economic benefits of recreational striped bass (*Morone saxatilis*) fishing in the Patuxent River. *Estuaries* 26 (2A), 310–315.
- Lohrenz SE, Redalje DG, Cai WJ, Acker J, Dagg M, 2008 A retrospective analysis of nutrients and phytoplankton productivity in the Mississippi River plume. *Cont. Shelf Res* 28 (12), 1466–1475.
- Martin PJ, 2000 Description of the navy coastal ocean model version 1.0. Tech. Rep. NRL/FR/7322–00-9962, Naval Research Lab, Stennis Space Center, Mississippi.
- Mateus MD, Franz G, 2015 Sensitivity analysis in a complex marine ecological model. *Water* 7, 2060–2081.
- Morrison M, Morgan MS, 1999 Models as mediating agents In: Morgan MS, Morrison M (Eds.), *Models as Mediators*. Cambridge University Press, Cambridge, p. 401.
- Murrell MC, Caffrey JC, Marcovich DT, Beck MW, Jarvis BM, III, J. D. H., in revision. Seasonal oxygen dynamics in a warm temperate estuary: Effects of hydrologic variability on measurements of primary production, respiration, and net metabolism. *Estuaries Coasts* - (-), -.
- Nocedal J, Wright SJ, 2006 Numerical Optimization, 2nd Edition. Springer-Verlag, New York, New York.
- Nossent J, Bauwens W, 2012 Multi-variable sensitivity and identifiability analysis for a complex environmental model in view of integrated water quantity and water quality modeling. *Water Sci. Technol* 65 (3), 539–549. [PubMed: 22258687]
- Obenour DR, Michalak AM, Scavia D, 2015 Assessing biophysical controls on Gulf of Mexico hypoxia through probabilistic modeling. *Ecol. Appl* 25 (2), 492–505. [PubMed: 26263670]
- Omlin M, Brun R, Reichert P, 2001a Biogeochemical model of Lake Zürich: sensitivity, identifiability and uncertainty analysis. *Ecol. Model* 141 (1–3), 105–123.
- Omlin M, Reichert P, Forster R, 2001b Biogeochemical model of Lake Zürich: model equations and results. *Ecol. Model* 141 (1–3), 77–103.
- Paerl HW, Pinckney JL, Fear JM, Peierls BL, 1998 Ecosystem responses to internal and watershed organic matter loading: consequences for hypoxia in the eutrophying Neuse River Estuary, North Carolina, USA. *Mar. Ecol. Prog. Ser* 166, 17–25.
- Parsons TR, Maita Y, Lalli CM, 1985 A manual of chemical methods for seawater analysis. Pergamon Press, New York.
- Pauer JJ, Feist TJ, Anstead AM, DePetro PA, Melendez W, Lehrter JC, Murrell MC, Zhang X, Ko DS, 2016 A modeling study examining the impact of nutrient boundaries on primary production on the Louisiana continental shelf. *Ecol. Model* 328, 136–147.
- Penta B, Lee Z, Kudela RM, Palacios SL, Gray DJ, Jolliff JK, Shulman IG, 2008 An underwater light attenuation scheme for marine ecosystem models. *Opt. Express* 16 (21), 16581–16591. [PubMed: 18852767]
- Petrucci G, Bonhomme C, 2014 The dilemma of spatial representation for urban hydrology semi-distributed modelling: trade-offs among complexity, calibration and geographical data. *J. Hydrol* 517, 997–1007.
- Rabalais NN, Turner RE, Scavia D, 2002 Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi river. *Biosci.* 52 (2), 129–142.
- RDCT (R Development Core Team), 2017 R: A language and environment for statistical computing, v3.3.2. R Foundation for Statistical Computing, Vienna, Austria <http://www.R-project.org>.

- Refsgaard JC, van der Sluijs JP, Højberg AL, Vanrolleghem PA, 2007 Uncertainty in the environmental modelling process - a framework and guidance. *Environ. Model. Softw* 22 (11), 1543–1556.
- Reichert P, Omlin M, 1997 On the usefulness of over parameterized ecological models. *Ecol. Model* 95 (2), 289–299.
- Roelke DL, 2000 Copepod food-quality threshold as a mechanism influencing phytoplankton succession and accumulation of biomass, and secondary productivity: a modeling study with management implications. *Ecol. Model* 134, 245–274.
- Scavia D, Justic D, Bierman VJ, 2004 Reducing hypoxia in the Gulf of Mexico: Advice from three models. *Estuaries* 27 (3), 419–425.
- Soetaert K, Petzoldt T, 2010 Inverse modelling, sensitivity, and Monte Carlo analysis in R using package FME. *J. Stat. Softw* 33 (3), 1–28. [PubMed: 20808728]
- Wade AJ, Jackson BM, Butterfield D, 2008 Over-parameterised, uncertain ‘mathematical marionettes’ - how can we best use catchment water quality models? An example of an 80-year catchment-scale nutrient balance. *Sci. Total. Environ* 400 (1–3), 52–74. [PubMed: 18538825]
- Wagener T, Boyle DP, Lees MJ, Wheeler HS, Gupta HV, Sorooshian S, 2001 A framework for development and application of hydrological models. *Hydrol. Earth Syst. Sci* 5 (1), 13–26.
- Warner JC, Geyer WR, Lerczak JA, 2005 Numerical modeling of an estuary: a comprehensive skill assessment. *J. Geophys. Res. Ocean* 110 (C5), 13.
- Wheeler HS, Bishop KH, Beck MB, 1986 The identification of conceptual hydrological models for surface water acidification. *Hydrol. Process* 1 (1), 89–109.
- Wiseman WJ, Rabalais NN, Turner RE, Dinnel SP, MacNaughton A, 1997 Seasonal and interannual variability within the Louisiana coastal current: stratification and hypoxia. *J. Mar. Syst* 12 (1–4), 237–248.
- Ye W, Bates BC, Viney NR, Sivapalan M, Jakeman AJ, 1997 Performance of conceptual rainfall-runoff models in low yielding ephemeral catchments. *Water Resour. Res* 33 (1), 153–166.
- Zhao L, Chen C, Vallino J, Hopkinson C, Beardsley RC, Lin H, Lerczak J, 2010 Wetland-estuarine-shelf interactions on the Plum Island Sound and Merrimack River in the Massachusetts coast. *J. Geophys. Res* 115 (C10), 13.

Fig. 1:

Conceptual representation of biogeochemical components included in FishTank (complete equations in Eldridge and Roelke 2010, appendices A-F in Lehrter et al. 2017). The Coastal General Ecosystem Model couples FishTank with a hydrodynamic model that includes advection, mixing, and dispersion.

Fig. 2:

Examples of unique parameter combinations from different parameter sets and number of selected parameters. The number of combinations are shown for increasing numbers of selected parameters from the total in the set, where 50 parameter sets are shown each with one through 50 total parameters. Note that the number of unique combinations is shown as the natural-log.

Fig. 3:

Sensitivity values (L1, eq. (2)) of all state variables to changes in a 50% increase in parameter values. Parameters are grouped by category: optics, organic matter, phytoplankton, zooplankton, temperature, and zooplankton. See Table 2 for L1 values for O2 and Tables S1 to S7 for the other state variables.

Fig. 4:

Collinearity (γ as a measure of identifiability, eq. (3)) of parameter subsets for O₂. Plots in (a) show collinearity by parameter categories and (b) shows collinearity by selecting the top 1 through 4 parameters in all categories. Lines represent collinearity ranges for the possible combinations given the number of parameters in the set. The temperature category is not shown because O₂ was sensitive to only one parameter (i.e., $\gamma = 1$).

Fig. 5:

Average collinearity (γ as a measure of identifiability, eq. (3)) of parameter subsets for all state variables. Plots in (a) show collinearity by parameter categories and (b) shows collinearity by selecting the top 1 through 4 parameters in all categories. Collinearity was averaged for all combinations in a parameter set to evaluate relative differences between state variables. The temperature category is not shown because all state variables were sensitive to only one parameter (i.e., $\gamma = 1$).

Fig. 6: Collinearity (γ as a measure of identifiability, eq. (3)) for O2 of organic matter parameters for subset combinations in Fig. 4. Collinearity is evaluated for subsets that excluded and included the parameters at the top of each plot. Collinearity of including all eight parameters is in Fig. 4. Grey lines indicate potential thresholds at $\gamma = 10, 15$ for maximum acceptable collinearity.

Fig. 7:

Example of calibration steps for the phytoplankton parameter subset. The FishTank model was calibrated using data from an approximate 12-hour flask experiment of oxygen production. Model calibration adjusted parameters (left plots) to minimize the *RMSE* between experimental and modelled dissolved oxygen (top left). The phytoplankton parameters were those identifiable for dissolved oxygen (Table 3), arranged from most (top plot) to least sensitive (bottom plot) (Table 2). The relative parameter changes in the right

plots show the initial parameter values (black) and the selected parameter values (blue) after calibration.

Fig. 8:

Estimated hypoxic area and monthly averages (km² × 1000) using the default parameter settings (a, b) and the differences using a 50% increase in each parameter (c, d). The selected parameters were the top sensitive in each parameter category for O₂ in Fig. 3 and Table 2. Error bars are 95% confidence intervals for the monthly averages in (b) and the difference in monthly averages (d) from the perturbed and default results. Areal extent of estimated hypoxia on August 28th using the default parameters (grey) and estimated changes with parameter perturbations is shown in (e).

Table 1:

FishTank parameters evaluated for sensitivity and identifiability. Sensitivity analyses were based on a 50% increase ($\text{Value} \cdot 1.5$) from the initial value of each parameter. Dashed units are dimensionless. Parameters are grouped by categories as optics, temperature, phytoplankton, zooplankton, and organic matter. A full description of the model structure, equations, and parameters is described in Lehrter et al. (2017).

Description	Parameter	Units	Value	Value - 1.5
Optics				
Chla specific absorption at 490 nm	<i>astar490</i>	$\text{m}^{-1} (\text{mg Chla m}^{-3})^{-1}$	0.04	0.06
OMA specific absorption at 490 nm	<i>astarOMA</i>	$\text{m}^{-1} (\text{mg OMA m}^{-3})^{-1}$	0.1	0.15
OMZ specific absorption at 490 nm	<i>astarOMZ</i>	$\text{m}^{-1} (\text{mg OMZ m}^{-3})^{-1}$	0.1	0.15
sinking rate	<i>sink CDOM</i>	m d^{-1}	0	0
Temperature				
Optimum temperature for growth	<i>Tref(nospA+nospZ)_{pl}</i>	C	22	33
Optimum temperature for growth	<i>Tref(nospA+nospZ)_{pl}</i>	C	22	33
Phytoplankton				
initial slope of photosynthesis v irradiance	<i>alpha</i>	$10^{-16} \text{ cm}^2 \text{ s quanta}^{-1} \text{ d}^{-1}$	$8.42e^{-17}$	$1.26e^{-16}$
coefficient for non-limiting nutrient	<i>aN</i>	-	1	1.5
phytoplankton threshold for grazing	<i>Athresh</i>	$10^7 \text{ cells m}^{-3}$	$1.72e^8$	$2.58e^8$
edibility vector for Z1	<i>ediblevector(Z1)</i>	-	0.25	0.38
edibility vector for Z2	<i>ediblevector(Z2)</i>	-	0.25	0.38
half-saturation constant for N	<i>Kn</i>	mmol m^{-3}	4.51	6.76
half-saturation constant for P	<i>Kp</i>	mmol m^{-3}	2.86	4.29
Qn constant for Flynn growth model	<i>KQn</i>	-	5	7.5
Qp constant for Flynn growth model	<i>KQp</i>	-	0.2	0.3
half-saturation constant for Si uptake	<i>Ksi</i>	mmol m^{-3}	4.51	6.76
mortality coefficient	<i>mA</i>	d^{-1}	0.1	0.15
phytoplankton carbon/cell	<i>Qc</i>	$10^{-7} \text{ mmol C cell}^{-1}$	$1.35e^{-6}$	$2.03e^{-6}$
minimum N cell-quota	<i>QminN</i>	$10^{-9} \text{ mmol N cell}^{-1}$	$6.08e^{-9}$	$9.12e^{-9}$
minimum P cell-quota	<i>QminP</i>	$10^{-9} \text{ mmol P cell}^{-1}$	$6.19e^{-10}$	$9.29e^{-10}$
phytoplankton basal respiration coefficient	<i>respb</i>	d^{-1}	0.02	0.03
phytoplankton growth respiration coefficient	<i>respg</i>	-	0.1	0.15
sinking rate of phytoplankton cells	<i>sink A</i>	m d^{-1}	1.49	2.23
maximum growth rate	<i>umax</i>	d^{-1}	0.41	0.62
N-uptake rate measured at umax	<i>vmaxN</i>	$10^{-8} \text{ mmol cell}^{-1} \text{ d}^{-1}$	$4.1e^{-8}$	$6.15e^{-8}$
P-uptake rate measured at umax	<i>vmaxP</i>	$10^{-8} \text{ mmol cell}^{-1} \text{ d}^{-1}$	$2.68e^{-8}$	$4.02e^{-8}$
Si-uptake rate measured at umax	<i>vmaxSi</i>	$10^{-8} \text{ mmol cell}^{-1} \text{ d}^{-1}$	$4.1e^{-8}$	$6.15e^{-8}$
phytoplankton volume/cell	<i>volcell</i>	μm^3	$3.37e^4$	$5.05e^4$
Zooplankton				
assimilation efficiency as a fraction of ingestion	<i>Zeffic</i>	-	0.4	0.6
half saturation coefficient for grazing	<i>ZKa</i>	$\mu\text{m}^3 \text{ m}^{-3}$	$1.12e^{12}$	$1.68e^{12}$
quadratic mortality constant	<i>Zm</i>	$\text{m}^6 \text{ ind}^{-2} \text{ d}^{-1}$	$7.2e^{-4}$	0

Description	Parameter	Units	Value	Value - 1.5
zooplankton carbon/individual	<i>ZQc</i>	mmol C ind ⁻¹	$3.13e^{-4}$	$4.7e^{-4}$
zooplankton nitrogen/individual	<i>ZQn</i>	mmol N ind ⁻¹	$6.95e^{-5}$	$1.04e^{-4}$
zooplankton phosphorus/individual	<i>ZQp</i>	mmol P ind ⁻¹	$3.77e^{-6}$	$5.66e^{-6}$
zooplankton biomass-dependent respiration factor	<i>Zrespb</i>	d ⁻¹	0.1	0.15
zooplankton growth-dependent respiration factor	<i>Zrespg</i>	-	0.2	0.3
proportion of phytoplankton lost to sloppy feeding	<i>Zslop</i>	-	0.25	0.38
maximum growth rate of zooplankton	<i>Zumax</i>	$\mu\text{m}^3 \text{ ind}^{-1} \text{ d}^{-1}$	$9.45e^7$	$1.42e^8$
zooplankton volume/individual	<i>Zvolcell</i>	$\mu\text{m}^3 \text{ ind}^{-1}$	$2.98e^7$	$4.47e^7$
Organic Matter				
turnover rate for OM1A and OM1Z	<i>KG1</i>	y ⁻¹	50	75
turnover rate for OM2A and OM2Z	<i>KG2</i>	y ⁻¹	50	75
decay rate of CDOM	<i>KGcdom</i>	d ⁻¹	0.01	0.02
NH4 rate constant for nitrification	<i>KNH4</i>	mmol m ⁻³	1	1.5
half-saturation concentration for denitrification	<i>KNO3</i>	mmol m ⁻³	10	15
half-saturation concentration for O2 utilization	<i>KO2</i>	mmol m ⁻³	10	15
O2 concentration that inhibits denitrification	<i>KstarO2</i>	mmol m ⁻³	10	15
maximum rate of nitrification per day	<i>nitmax</i>	mmol m ⁻³ d ⁻¹	0.52	0.78
phytoplankton sinking rate	<i>sink OM1_A</i>	m d ⁻¹	10	15
zooplankton sinking rate	<i>sink OM1_Z</i>	m d ⁻¹	10	15
phytoplankton sinking rate	<i>sink OM2_A</i>	m d ⁻¹	0	0
zooplankton sinking rate	<i>sink OM2_Z</i>	m d ⁻¹	0	0

Table 2:

Sensitivity of O₂ to perturbations of individual parameters. Sensitivities are based on a 50% increase from the initial parameter value, where *L1* summarizes differences in model output (see eq. (2)). Parameters that did not affect O₂ are not shown. Parameters are grouped by categories as optics, temperature, phytoplankton, zooplankton, and organic matter.

Description	Parameter	L1
Optics		
Chla specific absorption at 490 nm	<i>astar490</i>	$7.51e^{-4}$
OMZ specific absorption at 490 nm	<i>astarOMZ</i>	$4.92e^{-5}$
OMA specific absorption at 490 nm	<i>astarOMA</i>	$4.39e^{-5}$
Organic Matter		
turnover rate for OM1A and OM1Z	<i>KG1</i>	$6.15e^{-3}$
turnover rate for OM2A and OM2Z	<i>KG2</i>	$3.14e^{-3}$
O ₂ concentration that inhibits denitrification	<i>KstarO2</i>	$3.04e^{-3}$
decay rate of CDOM	<i>KGcdom</i>	$2.98e^{-3}$
half-saturation concentration for O ₂ utilization	<i>K02</i>	$5.85e^{-4}$
half-saturation concentration for denitrification	<i>KN03</i>	$5.8e^{-4}$
maximum rate of nitrification per day	<i>nitmax</i>	$4.99e^{-4}$
NH ₄ rate constant for nitrification	<i>KNH4</i>	$4.17e^{-4}$
Phytoplankton		
maximum growth rate	<i>umax</i>	0.05
mortality coefficient	<i>mA</i>	0.02
initial slope of photosynthesis v irradiance	<i>alpha</i>	0.02
edibility vector for Z1	<i>ediblevector(Z1)</i>	0.02
phytoplankton carbon/cell	<i>Qc</i>	0.01
phytoplankton growth respiration coefficient	<i>respg</i>	$8.36e^{-3}$
N-uptake rate measured at umax	<i>vmaxN</i>	$8.12e^{-3}$
phytoplankton basal respiration coefficient	<i>respb</i>	$6.94e^{-3}$
phytoplankton threshold for grazing	<i>Athresh</i>	$4.57e^{-3}$
minimum N cell-quota	<i>QminN</i>	$4.32e^{-3}$
P-uptake rate measured at umax	<i>vmaxP</i>	$4.27e^{-3}$
coefficient for non-limiting nutrient	<i>aN</i>	$4.23e^{-3}$
phytoplankton volume/cell	<i>volcell</i>	$4.13e^{-3}$
half-saturation constant for P	<i>Kp</i>	$2.9e^{-3}$
half-saturation constant for N	<i>Kn</i>	$2.77e^{-4}$
minimum P cell-quota	<i>QminP</i>	$8.34e^{-8}$
Temperature		
Optimum temperature for growth	<i>Tref(nospA+nospZ)_{pt}</i>	0.02
Zooplankton		
half saturation coefficient for grazing	<i>ZKa</i>	0.05
zooplankton nitrogen/individual	<i>ZQn</i>	0.02

Description	Parameter	L1
quadratic mortality constant	Zm	0.02
maximum growth rate of zooplankton	Z_{max}	0.02
assimilation efficiency as a fraction of ingestion	Z_{effic}	0.01
proportion of phytoplankton lost to sloppy feeding	Z_{slop}	$7.78e^{-3}$
zooplankton growth-dependent respiration factor	Z_{resp_g}	$5.32e^{-3}$
zooplankton biomass-dependent respiration factor	Z_{resp_b}	$2.96e^{-3}$
zooplankton carbon/individual	ZQ_c	$9.38e^{-5}$
zooplankton phosphorus/individual	ZQ_p	$3.69e^{-5}$

* Temperature parameters apply separately to phytoplankton ($p1$, one group) or zooplankton ($z1$, one group), denoted by subscripts

Table 3:

Parameter identifiability (as γ , eq. (3)) by category for relevant state variables. Selections followed the first heuristic where parameters were selected within categories from most to least sensitive until $\gamma > 15$. Rank describes the relative parameter sensitivity in each category for each state variable. Duplicate parameters and ranks in the first two columns apply only to γ values in the same row (i.e., parameter ranks vary for each variable).

Parameter	Rank	Ammonium	Chl-a	O2	Irradiance	Nitrate	POM	DOM	Phosphate
Optics									
<i>astar490</i>	1	1	1	1	1	1	1	1	1
<i>astarOMA</i>	2	7.33	5.42	-	5.36	-	7.78	7.87	-
<i>astarOMZ</i>	2	-	-	1.39	-	-	-	-	4.73
<i>astarOMA</i>	3	-	-	3.87	-	-	-	-	10.04
<i>astarOMZ</i>	3	7.58	5.51	-	6.02	-	7.91	7.87	-
Organic Matter									
<i>KG1</i>	1	-	-	1	-	-	1	-	1
<i>KG2</i>	1	-	-	-	-	-	-	1	-
<i>KGcdom</i>	1	-	1	-	1	-	-	-	-
<i>KstarO2</i>	1	-	-	-	-	1	-	-	-
<i>nitmax</i>	1	1	-	-	-	-	-	-	-
<i>KG1</i>	2	-	1.12	-	1.93	-	-	-	-
<i>KG2</i>	2	-	-	6	-	-	-	-	13.43
<i>KGcdom</i>	2	-	-	-	-	-	1.47	1.39	-
<i>KNH4</i>	2	4.03	-	-	-	-	-	-	-
<i>KG1</i>	3	4.09	-	-	-	-	-	-	-
<i>KG2</i>	3	-	-	-	8.19	-	-	-	-
<i>KGcdom</i>	3	-	-	-	-	-	-	-	13.75
<i>KO2</i>	3	-	-	-	-	-	14.07	11.96	-
<i>KstarO2</i>	3	-	-	6.04	-	-	-	-	-
<i>KGcdom</i>	4	4.19	-	6.12	-	-	-	-	-
<i>KO2</i>	4	-	-	-	-	-	-	-	14.68
<i>KstarO2</i>	4	-	-	-	10.65	-	14.08	-	-
<i>KO2</i>	5	9.47	-	8.61	-	-	-	-	-
Phytoplankton									
<i>mA</i>	1	1	1	-	-	-	1	1	-
<i>umax</i>	1	-	-	1	1	1	-	-	1
<i>ediblevector(Z1)</i>	2	1.13	1.17	-	-	-	1.15	-	-
<i>mA</i>	2	-	-	1.19	1.29	-	-	-	-
<i>Qc</i>	2	-	-	-	-	11.57	-	-	-
<i>umax</i>	2	-	-	-	-	-	-	1.21	-
<i>vmaxP</i>	2	-	-	-	-	-	-	-	7.45
<i>alpha</i>	3	-	-	1.44	1.98	-	-	-	-
<i>ediblevector(Z1)</i>	3	-	-	-	-	-	-	2.9	-

Parameter	Rank	Ammonium	Chl-a	O2	Irradiance	Nitrate	POM	DOM	Phosphate
<i>umax</i>	3	2.73	2.11	-	-	-	3.26	-	-
<i>alpha</i>	4	3.55	4.57	-	-	-	-	-	-
<i>ediblevector(ZI)</i>	4	-	-	2.09	4.09	-	-	-	-
<i>Qc</i>	4	-	-	-	-	-	4.98	-	-
<i>vmaxN</i>	4	-	-	-	-	-	-	4.9	-
<i>alpha</i>	5	-	-	-	-	-	10.11	-	-
<i>Qc</i>	5	-	-	2.9	-	-	-	-	-
<i>vmaxN</i>	5	8.14	-	-	-	-	-	-	-
<i>Athresh</i>	6	11.27	-	-	-	-	-	-	-
<i>respg</i>	6	-	-	3.41	-	-	-	-	-
<i>vmaxN</i>	7	-	-	3.97	-	-	-	-	-
Zooplankton									
<i>ZKa</i>	1	-	-	1	1	1	-	-	1
<i>Zumax</i>	1	1	1	-	-	-	1	1	-
<i>ZKa</i>	2	-	4.31	-	-	-	7.3	5.43	-
<i>ZQn</i>	2	-	-	3.18	6.32	9.76	-	-	8.54
<i>Zm</i>	3	-	-	4.57	-	-	-	-	-
<i>Zumax</i>	3	-	-	-	6.93	-	-	-	-
<i>Zm</i>	4	-	-	-	11.86	-	-	-	-
<i>Zumax</i>	4	-	-	5.2	-	-	-	-	-

Table 4:

Parameter identifiability (as γ , eq. (3)) for relevant state variables. Selections followed the second heuristic where parameters were selected independent of category from most to least sensitive (L1, eq. (2)), until $\gamma > 15$. Rank describes the relative parameter sensitivity in each category for each state variable (O: optics, OM: organic matter, P: phytoplankton, T: temperature, Z: zooplankton).

Selections by state variable	Parameter	L1	Rank	γ
Ammonium				
1	<i>mA</i>	8.49	1 _P	1
2	<i>nitmax</i>	1.54	1 _{OM}	1.16
3	<i>Zumax</i>	1.42	1 _Z	2.9
Chlorophyll				
1	<i>mA</i>	13.94	1 _P	1
2	<i>Zumax</i>	1.02	1 _Z	1.18
Dissolved Oxygen				
1	<i>umax</i>	0.05	1 _P	1
2	<i>ZKa</i>	0.05	1 _Z	2.17
3	<i>mA</i>	0.02	2 _P	2.31
4	<i>Tref(nospA+nospZ)_{pl}</i>	0.02	1 _T	2.37
5	<i>ZQn</i>	0.02	2 _Z	4.69
6	<i>alpha</i>	0.02	3 _P	4.91
7	<i>Zm</i>	0.02	3 _Z	6.73
8	<i>Zumax</i>	0.02	4 _Z	6.81
DOM				
1	<i>mA</i>	14.25	1 _P	1
2	<i>Tref(nospA+nospZ)_{pl}</i>	1.48	1 _T	1.05
3	<i>umax</i>	1.11	2 _P	2.46
4	<i>Zumax</i>	1.01	1 _Z	2.91
Irradiance				
1	<i>ZKa</i>	0.13	1 _Z	1
2	<i>umax</i>	0.09	1 _P	4.41
3	<i>ZQn</i>	0.06	2 _Z	7.54
4	<i>mA</i>	0.05	2 _P	8.17
5	<i>KGcdom</i>	0.05	1 _{OM}	9.44
6	<i>alpha</i>	0.04	3 _P	9.66
7	<i>Zumax</i>	0.04	3 _Z	10.79
Nitrate				
1	<i>umax</i>	8.49	1 _P	1
Phosphate				
1	<i>ZKa</i>	1.47	1 _Z	1

Selections by state variable	Parameter	L1	Rank	γ
2	u_{max}	0.78	1 _p	11.45
3	v_{maxP}	0.59	2 _p	11.48
4	ZQn	0.5	2 _z	13.74
POM				
1	mA	7.22	1 _p	1
2	Z_{umax}	0.96	1 _z	1.15
3	KGI	0.92	1 _{OM}	3.87

Table 5:

Parameter identifiability (as γ , eq. (3)) for relevant state variables. Selections followed the third heuristic where parameters were selected equally within each category from most to least sensitive (L1, eq. (2)), until $\gamma > 15$. Rank describes the relative parameter sensitivity in each category for each state variable (O: optics, OM: organic matter, P: phytoplankton, T: temperature, Z: zooplankton).

Selections by state variable	Parameter	L1	Rank	γ
Ammonium				
1	<i>mA</i>	8.49	1 _P	1
2	<i>nitmax</i>	1.54	1 _{OM}	1.16
3	<i>Zumax</i>	1.42	1 _Z	2.9
4	<i>Tref(nospA+nospZ)_{pl}</i>	0.79	1 _T	3.46
5	<i>astar490</i>	0.03	1 _O	4.25
Chlorophyll				
1	<i>mA</i>	13.94	1 _P	1
2	<i>Zumax</i>	1.02	1 _Z	1.18
3	<i>Tref(nospA+nospZ)_{pl}</i>	0.6	1 _T	2.62
4	<i>KGcdom</i>	0.07	1 _{OM}	3.24
5	<i>astar490</i>	0.02	1 _O	5.98
Dissolved Oxygen				
1	<i>umax</i>	0.05	1 _P	1
2	<i>ZKa</i>	0.05	1 _Z	2.17
3	<i>Tref(nospA+nospZ)_{pl}</i>	0.02	1 _T	2.29
4	<i>KG1</i>	$6.15e^{-3}$	1 _{OM}	3.85
5	<i>astar490</i>	$7.51e^{-4}$	1 _O	3.89
6	<i>mA</i>	0.02	2 _P	4.42
7	<i>ZQn</i>	0.02	2 _Z	5.22
DOM				
1	<i>mA</i>	14.25	1 _P	1
2	<i>Tref(nospA+nospZ)_{pl}</i>	1.48	1 _T	1.05
3	<i>Zumax</i>	1.01	1 _Z	2.61
4	<i>KG2</i>	0.94	1 _{OM}	3.39
5	<i>astar490</i>	0.04	1 _O	4.46
6	<i>umax</i>	1.11	2 _P	6.02
7	<i>ZKa</i>	0.88	2 _Z	9.21
Irradiance				
1	<i>ZKa</i>	0.13	1 _Z	1
2	<i>umax</i>	0.09	1 _P	4.41
3	<i>KGcdom</i>	0.05	1 _{OM}	4.5
4	<i>Tref(nospA+nospZ)_{pl}</i>	0.03	1 _T	4.5

Selections by state variable	Parameter	L1	Rank	γ
5	<i>astar490</i>	0.02	1 _O	6.9
6	<i>ZQn</i>	0.06	2 _Z	10.63
7	<i>mA</i>	0.05	2 _P	11.21
8	<i>KGI</i>	$3.96e^{-3}$	2 _{OM}	14.65
9	<i>astarOMA</i>	$1.47e^{-3}$	2 _O	14.72
Nitrate				
1	<i>umax</i>	8.49	1 _P	1
Phosphate				
1	<i>ZKa</i>	1.47	1 _Z	1
2	<i>umax</i>	0.78	1 _P	11.45
3	<i>Tref(nospA+nospZ)_{pl}</i>	0.16	1 _T	13.71
4	<i>KGI</i>	0.14	1 _{OM}	14.64
POM				
1	<i>mA</i>	7.22	1 _P	1
2	<i>Zumax</i>	0.96	1 _Z	1.15
3	<i>KGI</i>	0.92	1 _{OM}	3.87
4	<i>Tref(nospA+nospZ)_{pl}</i>	0.86	1 _T	3.93
5	<i>astar490</i>	0.03	1 _O	5.81

Table 6:

Results of model calibration to observed O2 for each of seven parameter subsets selected by sensitivity within each parameter category (first), independent of category (second), and equally within each category (third). The collinearity values (γ) and parameters of each subset are from Tables 3 to 5. Mean L1 is the average of sensitivity values (eq. (2)) for the subset. *RMSE* shows the final error comparing observed and model output after the specified number of iterations for optimization.

Selection heuristic	γ	<i>n</i> parameters	mean L1	Iterations	<i>RMSE</i> *
1					
Optics	3.87	3	$2.81e^{-4}$	14	21.14
Organic Matter	8.61	5	$3.18e^{-3}$	132	20.6
Phytoplankton	3.97	7	0.02	615	5.81
Temperature	1	1	0.02	12	21.14
Zooplankton	5.2	4	0.03	54	21.14
2					
Independent of category	6.81	8	0.03	221	19.45
3					
Equally within category	5.22	7	0.03	435	19.09

* Starting *RMSE* 21.14

Table 7:

Results of parameter changes from model calibration to observed O2 for each of seven parameter subsets selected by sensitivity within each parameter category (first), independent of category (second), and equally within each category (third). The parameters of each subset are from Tables 3 to 5. The L1 values are relative sensitivity of O2 to a parameter change. Initial and final parameter values after calibration are shown, including the percent change. The calibration routine was bounded by minimum and maximum values for each parameter.

Selection heuristic	Parameter	L1	Min/Max	Initial	Final	% change
1						
Optics	<i>astar490</i>	$7.51e^{-4}$	0.02/0.12	0.04	0.04	-0.3
	<i>astarOMZ</i>	$4.92e^{-5}$	0.01/0.15	0.01	0.01	nc
	<i>astarOMA</i>	$4.39e^{-5}$	0.01/0.15	0.01	0.01	nc
Organic Matter	<i>KG1</i>	$6.15e^{-3}$	5/100	30	5	-83.3
	<i>KG2</i>	$3.14e^{-3}$	5/100	30	52.31	+74.4
	<i>KstarO2</i>	$3.04e^{-3}$	1/100	10	10.01	+0.1
	<i>KGcdom</i>	$2.98e^{-3}$	0/1	0.01	0.01	+0.1
	<i>KO2</i>	$5.85e^{-4}$	1/100	10	100	+900
	<i>umax</i>	0.05	0.1/2.5	1.12	2.49	+121.9
Phytoplankton	<i>mA</i>	0.02	0.01/0.5	0.11	0.07	-38.9
	<i>alpha</i>	0.02	$1e^{-17}/6e^{-16}$	$3.96e^{-16}$	$6e^{-16}$	+51.5
	<i>ediblevector(Z1)</i>	0.02	0.05/1	0.5	0.37	-26.5
	<i>Qc</i>	0.01	$1e^{-9}/5e^{-7}$	$4.54e^{-8}$	$3.21e^{-7}$	+606.3
	<i>respg</i>	$8.36e^{-3}$	0.01/0.5	0.1	0.01	-89.6
	<i>vmaxN</i>	$8.12e^{-3}$	$1e^{-11}/1e^{-8}$	$1.33e^{-9}$	$1.35e^{-9}$	+1.8
Temperature	<i>Tref(nospA+nospZ)_{pl}</i>	0.02	0/40	17	16.5	-2.9
Zooplankton	<i>ZKa</i>	0.05	$1e^{11}/2.5e^{12}$	$1.12e^{12}$	$1.61e^{12}$	+43.6
	<i>ZQn</i>	0.02	$1e^{-6}/2.04e^{-4}$	$6.95e^{-5}$	$7.53e^{-5}$	+8.3
	<i>Zm</i>	0.02	$1e^{-4}/10$	$7.2e^{-4}$	$8.9e^{-4}$	+23.6
	<i>Zumax</i>	0.02	$5e^8/2e^9$	$9.45e^8$	$5.33e^8$	-43.6
2						
Independent of category	<i>Umax</i>	0.05	0.1/2.5	1.12	2.5	+123.2
	<i>ZKa</i>	0.05	$1e^{11}/2.5e^{12}$	$1.12e^{12}$	$2.5e^{12}$	+123.2
	<i>mA</i>	0.02	0.01/0.5	0.11	0.01	-90.9
	<i>Tref(nospA+nospZ)_{pl}</i>	0.02	0/40	17	5.82	-65.7
	<i>ZQn</i>	0.02	$1e^{-6}/2.04e^{-4}$	$6.95e^{-5}$	$7.04e^{-5}$	+1.2
	<i>alpha</i>	0.02	$1e^{-17}/6e^{-16}$	$3.96e^{-16}$	$6e^{-16}$	+51.5
	<i>Zm</i>	0.02	$1e^{-4}/10$	$7.2e^{-4}$	$1.07e^{-3}$	+48.9
	<i>Zumax</i>	0.02	$5e^8/2e^9$	$9.45e^8$	$5e^8$	-47.1
3						
Equally within category	<i>Umax</i>	0.05	0.1/2.5	1.12	2.5	+123.2
	<i>ZKa</i>	0.05	$1e^{11}/2.5e^{12}$	$1.12e^{12}$	$2.5e^{12}$	+123.2

Selection heuristic	Parameter	L1	Min/Max	Initial	Final	% change
	<i>mA</i>	0.02	0.01/0.5	0.11	0.01	-90.9
	<i>Tref(nospA+nospZ)_{pl}</i>	0.02	0/40	17	12.2	-28.2
	<i>ZQn</i>	0.02	$1e^{-6}/2.04e^{-4}$	$6.95e^{-5}$	$8.87e^{-5}$	+27.6
	<i>KG1</i>	$6.15e^{-3}$	5/100	30	5	-83.3
	<i>astar490</i>	$7.51e^{-4}$	0.02/0.12	0.04	0.02	-46.7