Occupational Health

AHR gene-dioxin interactions and birthweight in the Seveso Second Generation Health Study

Jennifer Ames,¹ Marcella Warner,¹ Paolo Mocarelli,² Paolo Brambilla,² Stefano Signorini,² Claudia Siracusa,² Karen Huen,¹ Nina Holland^{1†} and Brenda Eskenazi^{1*†}

¹Center for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, CA, USA and ²Department of Laboratory Medicine, University of Milan-Bicocca, Hospital of Desio, Desio-Milano, Italy

*Corresponding author. Center for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, 1995 University Ave Suite 265, Berkeley, CA 94720, USA. E-mail: eskenazi@berkeley.edu

†Drs Holland and Eskenazi are joint senior authors.

Editorial decision 2 July 2018; Accepted 17 July 2018

Abstract

Background: 2, 3, 7, 8-tetrachlorodibenzo-*p*-dioxin (TCDD) is proposed to interfere with fetal growth via altered activity of the aryl hydrocarbon receptor (protein: AHR; gene: *AHR*) pathway which regulates diverse biological and developmental processes including xenobiotic metabolism. Genetic variation in *AHR* is an important driver of susceptibility to low birthweight in children exposed to prenatal smoking, but less is known about these genetic interactions with TCDD, AHR's most potent xenobiotic ligand.

Methods: The Seveso Women's Health Study (SWHS), initiated in 1996, is a cohort of 981 Italian women exposed to TCDD from an industrial explosion in July 1976. We measured TCDD concentrations in maternal serum collected close to the time of the accident. In 2008 and 2014, we followed up the SWHS cohort and collected data on birth outcomes of SWHS women with post-accident pregnancies. We genotyped 19 single nucleotide polymorphisms (SNPs) in *AHR* among the 574 SWHS mothers.

Results: Among 901 singleton births, neither SNPs nor TCDD exposure alone were significantly associated with birthweight. However, we found six individual SNPs in *AHR* which adversely modified the association between maternal TCDD and birthweight, implicating gene-environment interaction. We saw an even stronger susceptibility to TCDD due to interaction when we examined the joint contribution of these SNPs in a risk allele score. These SNPs were all located in noncoding regions of *AHR*, particularly in proximity to the promoter.

Conclusions: This is the first study to demonstrate that genetic variation across the maternal *AHR* gene may shape fetal susceptibilities to TCDD exposure.

Key words: Dioxins, aryl hydrocarbon receptor, Seveso, birthweight, gene-environment, polymorphisms

Key Messages

- Some genetic subpopulations of *AHR*, a gene coding for a key transcription factor involved in xenobiotic metabolism, may be more susceptible to *in utero* TCDD exposure than others.
- We observed six polymorphisms in regulatory regions of *AHR*, particularly in proximity to the gene's promoter, that adversely modified the association between maternal TCDD exposure and child birthweight.
- Analyses using a risk allele score suggested that the combination of these risk variants may jointly influence susceptibility to dioxin.
- *AHR* polymorphisms and maternal TCDD levels were not significantly associated with birthweight when examined independently, supporting the presence of gene-environment interaction.

Introduction

In 1976, an industrial accident near Seveso Italy resulted in one of the highest residential exposures to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) in history.^{1–5} TCDD, a common by-product of industrial and combusive processes, is a highly lipophilic, persistent organic pollutant, and a known carcinogen⁶ and endocrine disruptor.⁷ The half-life of TCDD is relatively long, in the order of 7–10 years in humans.⁸ TCDD has been shown to cross the placenta^{9,10} and *in utero* exposure to TCDD and dioxin-like chemicals has been linked in animal studies to altered immune function, glucose regulation, steroidogenesis and neurobehavioural and bone development.^{11–19}

Animal studies have also suggested that TCDD may impact fetal growth,^{15,20–23} possibly via altered activity of the aryl hydrocarbon receptor (protein: AHR; gene: *AHR*), a nuclear receptor and transcription factor active in many tissues including the placenta.^{24–29} Upon binding TCDD in the cell cytoplasm, AHR moves to the nucleus, where it induces expression of several xenobiotic metabolizing enzymes such as cytochrome P450s, and exhibits cross-talk with pathways of hormone synthesis.³⁰ In addition to detoxification, the AHR pathway regulates a myriad of biological processes related to development, cell growth, apoptosis and immune function.³¹

Of the epidemiological studies that have examined biological markers of maternal dioxin exposures and birthweight, three found no association,^{32–34} another three found adverse associations^{35–37} and three had adverse associations that did not reach statistical significance.^{38–40} In the Seveso Women's Health Study (SWHS), a follow-up study of women exposed to high levels of TCDD from an industrial explosion in 1976, we previously reported a non-significant inverse association between serum TCDD concentrations and birthweight of first post-explosion births [adjusted- $\beta = -47.7$ g, 95% confidence interval (CI): $-107.3, 11.9$ for a 10-fold increase in serum TCDD concentration].^{38,41} The lack of consistency across the

literature examining the effects of TCDD on birthweight may be due to the wide variation in sample size, difference in exposure levels and, perhaps, genetic variation represented in these study populations.

Inter-strain and interspecies differences in AHR ligand-binding affinities suggest that genetic variation in *AHR* may influence susceptibility to TCDD.⁴² Human evidence supporting this hypothesis is drawn from two Japanese studies that found that a polymorphism in maternal *AHR* (rs2066853) conferred significant reductions in birthweight in pregnant women who smoked cigarettes (components of cigarette smoke, such as benzo(a)pyrene, also bind AHR).⁴³ A more recent study of Japanese infants ($n = 421$), which examined the relationship between this single nucleotide polymorphism (SNP) and prenatal dioxins toxic equivalency (TEQ), reported no relationship with birthweight;⁴⁴ consideration of additional *AHR* SNPs in larger cohorts with higher exposures to specifically TCDD, the most potent compound of the TEQ, is warranted.

In the present analysis, we investigate whether maternal *AHR* gene variation modifies the association between maternal exposure levels of TCDD and birthweight in the children born after the Seveso explosion to mothers who participated in SWHS, a follow-up study of women living in Seveso, Italy, at the time of the accident.

Methods

Study population and procedures

In 1996, 20 years after the explosion, the Seveso Women's Health Study was initiated. Eligible women were aged 40 years or younger on 10 July 1976, resided in the most contaminated areas and had blood samples collected soon after the explosion. A total of 981 women (80% of those eligible) participated.⁵ In 2008 and 2014, we followed up these participants. Details of the study procedure for the 2008 and 2014 studies are described elsewhere.^{45,46}

At each follow-up visit, women were interviewed in a private room at the Hospital of Desio by a nurse interviewer who was blinded to participant TCDD levels. Information was obtained on medical and reproductive history, with detailed information on each pregnancy and on demographic and lifestyle factors. Between 1976 and 2016, a total of 574 SWHS mothers reported 943 live births occurring after the 1976 accident. We obtained genetic information on 567 mothers (98.8%), corresponding with 929

births. Seven women (who had 14 live births) either did not consent to biobanking their blood specimens or did not have adequate amounts of blood specimen for DNA isolation. We excluded an additional 27 multiple births and one singleton with missing birthweight, leaving 901 singletons from 562 mothers for the primary analyses (Table 1). The study was approved by the institutional review boards of the participating institutions and we obtained written informed consent from all mothers before participation.

Table 1. Descriptive statistics of mothers with genetic data in the SWHS, 1996–2014

Characteristic	<i>n</i>	(%)	1976 serum TCDD (ppt) median (IQR)	
Total women	562	(100.0)	61.3	(29.0, 163.0)
Total live births	901	(100.0)		
Age at explosion (years)				
0–10	164	(29.2)	157.5	(51.8, 320.5)
11–20	239	(42.5)	53.4	(25.4, 105.0)
21–30	139	(24.7)	41.3	(22.0, 80.9)
31–40	20	(3.6)	39.0	(27.7, 66.8)
Menarche status at explosion				
Premenarche	211	(37.5)	131.0	(50.5, 286.0)
Postmenarche	351	(62.5)	44.4	(22.5, 86.7)
Pre-explosion parity				
0	451	(80.3)	71.1	(31.7, 192.0)
1	71	(12.6)	36.6	(21.1, 70.4)
≥2	40	(7.1)	35.6	(24.5, 69.1)
Maternal education at last follow-up				
<Required	105	(18.7)	42.5	(23.5, 73.4)
Required/high school	432	(76.9)	64.7	(30.0, 188.0)
University	25	(4.5)	75.7	(30.2, 180.0)
Age at pregnancy (years)				
<25	152	(16.9)	46.4	(21.5, 103.5)
25–29	309	(34.3)	55.5	(28.6, 131.0)
30–34	270	(30.0)	67.2	(29.9, 187.0)
≥35	170	(18.9)	64.3	(35.0, 176.0)
Smoking during pregnancy				
No	813	(90.2)	61.2	(29.3, 164.0)
Yes	88	(9.8)	50.0	(20.7, 102.0)
Weight gain during pregnancy (kg) ^a				
<10	190	(21.1)	67.9	(38.2, 141.0)
10–14	421	(46.7)	54.3	(26.7, 136.0)
15–19	171	(19.0)	62.8	(22.0, 214.0)
≥20	96	(10.7)	70.9	(30.0, 204.0)
Low birthweight (<2500 g)				
No	844	(93.7)	60.4	(28.4, 157.0)
Yes	57	(6.3)	64.7	(31.7, 131.0)
Preterm (<37 weeks)				
No	839	(93.1)	60.4	(28.4, 156.0)
Yes	62	(6.9)	76.2	(33.2, 206.0)
Infant sex				
Male	473	(52.5)	55.0	(27.2, 130.0)
Female	428	(47.5)	67.0	(29.9, 179.5)

^aMissing data on pregnancy weight gain for 23 live births.

Outcome assessment

Birthweights and gestational duration were based on maternal report. In a small sample ($n = 139$), we confirmed reported birthweights using hospital records. These data indicated that women slightly over-reported birthweight, by 22 g on average, but this was non-differential by TCDD exposure.⁴¹

TCDD analysis

TCDD was measured in archived maternal serum samples collected near the time of the explosion by high-resolution gas chromatography/high-resolution mass spectrometry methods at the Centers for Disease Control and Prevention (CDC).^{47,48} Details of the serum sample selection and TCDD concentrations are presented elsewhere.^{5,49} Before statistical analysis, maternal serum TCDD levels were adjusted for blood lipid concentrations by dividing TCDD on a whole-weight basis by total serum lipid content, estimated from measurements of triglycerides and total cholesterol.⁵⁰ Serum TCDD levels were reported in picograms per gram lipid or parts per trillion (ppt). The median lipid-adjusted limit of detection (LOD) for the full population was 18.8 ppt. Quantifiable results less than the method detection limits were reported when observed. Otherwise, samples below the LOD (9.4% in the full cohort) were assigned a value equal to one-half of the LOD.⁵¹ By considering the 1976 TCDD levels, we examine the hypothesis that the mother's primary dose permanently altered her reproductive system or oocytes, possibly

resulting in persistent epigenetic changes that could impact fetal growth.⁵²

SNP selection and genotyping

We used the HapMap browser⁵³ in the Caucasian population of European descent (CEPH) and the 1000 Genomes Toscani in the Italia population (TSI), to choose SNPs from *AHR* which were expected to have minor allele frequencies greater than 5% in our Italian Caucasian sample. In cases where SNPs were in linkage disequilibrium, we sought an appropriate tagging SNP representative of this group of co-varying SNPs, to conserve study power and resources. These candidate SNPs were further pared down to those with known or suspected functional relevance as reported in the literature or as listed in the open access Regulome SNP database (Stanford University).⁵⁴ We particularly prioritized SNPs linked to xenobiotic exposure and fetal development, though we considered *AHR* SNPs associated with any health outcome. Our final genotyping assay comprised 18 SNPs across 50 kb of the *AHR* gene as well as one SNP in *AHR*'s upstream intergenic region. Location of the SNPs and their physical distribution are presented in Table 2 and Supplementary Figure 1, available as Supplementary data at *IJE* online.

Maternal DNA for genotyping was isolated from archived blood using a QIAamp Blood DNA Maxi kit (QIAGEN, Valencia, CA, USA) with some modification as previously described.⁵⁵ High-throughput genotyping of selected SNPs was performed using the multiplex platform

Table 2. Description of *AHR* SNPs genotyped in SWHS mothers, 1996-2014

SNP	Location (chr: bp)	Alleles	MAF (%)	Functional consequence
rs6968865	7: 17 247 645	T/A	41.2	Upstream, intergenic region
rs3757824	7: 17 296 411	A/G	18.5	intron, regulatory region variant
rs10249788	7: 17 298 523	C/T	12.4	intron, regulatory region variant
rs713150	7: 17 300 533	C/G	25.7	intron, regulatory region variant
rs17722841	7: 17 303 970	G/C	14.2	intron
rs2282885	7: 17 305 990	T/C	35.3	intron
rs3802083	7: 17 309 283	G/A	38.9	intron
rs17779352	7: 17 310 002	T/C	8.1	exon, synonymous variant
rs1476080	7: 17 318 249	C/A	37.2	intron, regulatory region variant
rs2237297	7: 17 319 970	C/T	9.3	intron
rs17137566	7: 17 320 897	T/C	15.7	intron
rs4236290	7: 17 323 944	T/C	12.0	intron
rs6960165	7: 17 328 461	A/G	24.1	intron
rs2158041	7: 17 328 796	G/A	23.8	intron
rs3802082	7: 17 330 557	T/A	15.3	intron
rs7811989	7: 17 331 739	G/A	27.3	intron
rs2066853	7: 17 339 486	G/A	9.8	exon, missense variant
rs2040623	7: 17 341 038	T/G	19.4	intron
rs2106728	7: 17 342 116	A/G	35.2	intron

iPLEX (Sequenom, San Diego, CA) at the Genomics Center at the University of Minnesota. The main steps involved multiplex polymerase chain reaction (PCR), single-base primer extension and finally mass spectrometry to determine the genotype. Quality assurance procedures for genotyping included assessment of randomly distributed blank samples and duplicates of participant samples. Call rates were above 98% for all 19 SNPs. Samples with lower success rates were resolved with additional genotyping. All genotype distributions were in accordance with Hardy-Weinberg equilibrium assumptions.

Statistical analyses

TCDD measures were analysed as a $\log_{10}$ -transformed, continuous variable. We considered those covariates that were used in previous reports of birthweight in this cohort or identified a priori as confounders between *in utero* dioxin and birthweight in a directed acyclic graph (DAG) (Supplementary Figure 2, available as Supplementary data at *IJE* online). These covariates included maternal age at pregnancy (continuous, years), year of pregnancy, smoking during pregnancy (yes/no), parity (0, 1 or >2 pregnancies), maternal height (cm), pre-accident history of delivering a low birthweight infant, child sex and gestational age (maternal report in weeks). Model parameters were estimated with use of generalized estimating equations (GEE) with exchangeable correlation of the variance structure to account for siblings.⁵⁶ We first re-evaluated the association between TCDD and birthweight in this analytical sample for comparison with our previously reported result of an adverse but non-significant relationship.^{38,41}

Before model fitting, linkage disequilibrium between SNPs was assessed with r^2 and observed SNP distributions and correlations were compared with those in the TSI of the 1000 Genomes. Genotype analyses considered two penetrance models with reasonable power given our sample sizes: (i) additive allelic penetrance (i.e. groups inheriting 0, 1 or 2 minor alleles at each SNP); and (ii) dominant penetrance (i.e. inheriting 0 vs at least 1 minor allele). In models assuming dominant penetrance of the minor 'variant' allele, the genotype at each SNP was analysed as an indicator variable. In separate analyses assuming additive allelic penetrance, the genotype was analysed categorically and if a dose-response pattern was observed with the increasing number of variant alleles (0, 1, 2), an ordinal variable was used. In both scenarios, the reference group was the genotype with 0 variant alleles.

To evaluate the main effects of each SNP, we fitted multivariate models of birthweight regressed on maternal genotype controlling for the above covariates and TCDD levels. We then considered crude and adjusted models of

interaction between maternal genotype and TCDD on birthweight by constructing a cross-product term of genotype and TCDD (i.e. $\text{SNP}_{\text{Maternal}} \times \text{TCDD}$). TCDD was considered as a continuous variable ($\log_{10}$ -transformed) and as a categorized variable to explore GxE interaction allowing for nonlinear associations between TCDD and birthweight. The lowest TCDD category with no variant alleles served as the reference group. We examined cumulative associations across SNPs through calculation of a genetic risk allele score by summing, for each individual, the number of risk alleles (1 point if ≥ 1 risk allele present) across all SNPs that were significant ($p_{\text{int}} < 0.1$) in the individual SNP models and, in a sensitivity analysis, by weighting the SNP components by the size of their coefficients in the individual models. Genotypes associated with lowered birthweight in interaction with TCDD were designated as the 'risk' genotype and summed across all SNPs included in the score. The score was then examined in models with and without interaction with maternal TCDD levels. The score, a common method that conserves study power and potentially gives insight into underlying biological mechanisms, was examined as a continuous variable.⁵⁷

Multiple births were excluded from the initial models, but were added back in a sensitivity analysis. Additional sensitivity analyses considered models excluding preterm (<37 weeks' gestation)⁵⁸ births and models restricted to the mother's first birth after the explosion, using robust regression. All analyses were performed in Stata 13 (StataCorp LP, College Station, TX) with the exception of Hardy-Weinberg and linkage disequilibrium statistics and the Benjamini-Hochberg false-discovery rate (FDR) correction for multiple comparisons which were performed using the *SNP assoc* and *p.adjust* packages in R, version 3.4.3,⁵⁹ respectively.^{60,61}

Results

Descriptive statistics of study sample

Maternal 1976 serum TCDD concentrations across demographic and pregnancy characteristics are presented in Table 1. Mothers in this sample averaged 14.7 (± 7.7) years of age at the time of the accident. The average maternal age at pregnancy across all 901 births was 29.6 years (± 5.3) and 20% of women had a history of pregnancy preceding the accident. The majority of mothers (81.4%) had educational attainment beyond the compulsory education in Italy. Women who were younger in 1976 tended to have higher levels of serum TCDD, higher levels of educational attainment at follow-up and were slightly older at time of pregnancy than SWHS women who were adults at the time of the accident, a pattern described in previous studies of

the SWHS.^{4,49} The median TCDD level in maternal serum was 61.3 ppt [interquartile range (IQR): 29.0–163.0] in 1976. The mothers in our analytical sample ($n = 562$) did not differ in sociodemographic characteristics or medical history from the sample of all mothers with eligible children in the Seveso Second Generation Health Study ($n = 574$).

Among the 901 singleton births, the average birthweight was 3264 (± 526) g. The number of low birthweight (<2500 g)⁶² and preterm infants (<37 weeks)⁵⁸ was 57 (6.3%) and 62 (6.9%), respectively. Without considering genotype, a 10-fold increase in 1976 serum TCDD was associated with a non-significant reduction in birthweight ($\beta = -33.65$, 95% CI: -85.11, 17.80; $P = 0.20$), adjusting for covariates. These findings are similar to what has been observed in previous studies in this cohort, although marginally different due to the slightly different sample who had DNA available for analysis.^{38,41,46}

Minor allele frequencies for the 19 SNPs, shown in Table 2, ranged from 8.1% (rs17779352) to 41.2% (rs6968865). The allelic frequencies in the SWHS population were similar to those observed in the 1000 Genomes TSI.⁶³ Linkage disequilibrium was relatively low among the *AHR* SNPs, with 5% of pairwise LD comparisons exceeding an $r^2 > 0.8$ (see Supplementary Figure 1, available as Supplementary data at IJE online, for LD Plot).

Associations between SNPs and birthweight

We observed no association between any of the 19 SNPs in *AHR* on birthweight (Supplementary Table 1, available as Supplementary data at IJE online). The only SNP with a suggestive association was rs2066853. Under an additive model of penetrance, each additional risk allele (A) was associated with a -62 g reduction in birthweight relative to the wild-type GG mothers ($P_{\text{trend}} = 0.085$). However, this trend was based on seven mothers with the AA genotype, and therefore a dominant model (adjusted $\beta = -67.42$, 95% CI: -145.13, 10.30; $P = 0.089$) was used in subsequent sensitivity analyses to conserve power (Figure 1). The only other SNP (rs2237297) that was notably associated with birthweight (adjusted $\beta = -58.04$, 95% CI: -137.49, 21.42; $P = 0.152$) is in high LD with rs2066853 ($r^2 = 0.93$) in this population.

Gene-environment interactions

In crude models, the variant alleles of four of the 19 SNPs (rs6968865, rs3757824, rs10249788, rs2040623) exhibited interaction with 1976 maternal TCDD levels on birthweight (Supplementary Table 2, available as Supplementary data at IJE online). When these models were adjusted for covariates,

Figure 1. Main effects of maternal *AHR* SNP rs2066853 on birthweight under dominant and additive genetic models, SWHS, 1996–2014. Models adjusted for gestational age (weeks), history of low birthweight, maternal age at pregnancy, parity, maternal height, maternal smoking, birth year, child sex.

effect sizes were attenuated, but these four SNPs and two others (rs2282885, rs2106728) suggested gene-environment interaction (Table 3). Further, these six interactive SNPs were clustered around the promoter and first and last introns of *AHR*. We observed five SNPs (rs6968865, rs3757824, rs10249788, rs3802082, rs2040623) for which presence of the variant allele in mothers was associated with strong adverse associations between TCDD and birthweight relative to the reference group of homozygous wild-type mothers. For example, a 10-fold increase in TCDD was associated with -149.09 g (95% CI: -252.34, -45.83; $P = 0.015$) and -1.76 g (95% CI: -59.89, 56.37; $P = 0.953$) reductions in birthweight among children of mothers with the variant (CC/CT) and wild-type (TT) rs10249788 genotypes, respectively. This interactive association was particularly notable at the low and high ends of TCDD exposure where, continuing with the example of rs10249788, the variant allele was protective at low TCDD levels but crossed to adverse at higher TCDD exposures (Figure 2). We observed this pattern of GxE interaction, whereby the risk allele conferred fetal hypersusceptibility to TCDD, for three other SNPs (rs6968865, rs3757824, rs2040623). For two SNPs, rs2282885 and rs2106728, the variant allele exhibited a

Table 3. GEE regression models of 1976 log10TCDD exposure on child birthweight stratified by genotype at each locus using dominant model of inheritance, SWHS, 1996–2014

SNP	Genetic model	0 minor alleles			1 or 2 minor alleles			P_{int}	FDR
		β_{adj} (95% CI)		P	β_{adj} (95% CI)		P		
rs6968865	AA + TA vs TT	23.23	(-66.93, 113.40)	0.614	-62.34	(-124.58, -0.10)	0.050*	0.034	0.17
rs3757824	GG + AG vs AA	18.44	(-45.40, 82.28)	0.571	-126.75	(-210.00, -43.49)	0.003**	0.021	0.14
rs10249788	TT + CT vs CC	-1.76	(-59.89, 56.37)	0.953	-149.09	(-252.34, -45.83)	0.005**	0.015	0.14
rs713150	GG + CG vs CC	-46.10	(-119.31, 27.11)	0.217	-2.95	(-78.30, 72.40)	0.939	0.741	0.80
rs17722841	AA + GA vs GG	-50.31	(-109.15, 8.53)	0.094	-7.97	(-111.77, 95.83)	0.880	0.704	0.80
rs2282885	CC + TC vs TT	-86.66	(-164.08, -9.24)	0.028*	10.86	(-57.15, 78.87)	0.754	0.067	0.22
rs3802083	AA + GA vs GG	-36.32	(-131.41, 58.78)	0.454	-35.05	(-95.96, 25.86)	0.259	0.654	0.80
rs17779352	CC + TC vs TT	-37.35	(-90.59, 15.90)	0.169	15.29	(-150.22, 180.81)	0.856	0.618	0.80
rs1476080	AA + CA vs CC	-14.00	(-103.84, 75.83)	0.760	-46.64	(-109.23, 15.96)	0.144	0.360	0.62
rs2237297	TT + CT vs CC	-23.65	(-80.81, 33.51)	0.417	-79.12	(-191.05, 32.81)	0.166	0.369	0.62
rs17137566	CC + TC vs TT	-14.76	(-76.99, 47.46)	0.642	-84.39	(-172.78, 4.00)	0.061	0.409	0.62
rs4236290	CC + TC vs TT	-11.52	(-70.17, 47.13)	0.700	-103.61	(-210.38, 3.16)	0.057	0.161	0.45
rs6960165	GG + AG vs AA	-48.01	(-118.69, 22.67)	0.183	-9.59	(-83.44, 64.25)	0.799	0.772	0.80
rs2158041	AA + GA vs GG	-47.28	(-117.89, 23.32)	0.189	-16.02	(-90.12, 58.08)	0.672	0.861	0.85
rs3802082	AA + TA vs TT	4.04	(-59.70, 67.79)	0.901	-88.91	(-174.30, -3.53)	0.041*	0.221	0.53
rs7811989	AA + GA vs GG	-20.32	(-94.72, 54.09)	0.593	-46.31	(-117.47, 24.85)	0.202	0.370	0.62
rs2066853	AA + GA vs GG	-26.83	(-83.60, 29.94)	0.354	-59.57	(-175.78, 56.65)	0.315	0.416	0.62
rs2040623	GG + TG vs TT	36.74	(-29.19, 102.68)	0.275	-116.13	(-197.27, -34.99)	0.005**	0.021	0.14
rs2106728	GG + AG vs AA	-80.23	(-158.02, -2.45)	0.043*	6.88	(-60.90, 74.66)	0.842	0.061	0.22

Models adjusted for gestational age (weeks), history of low birthweight, maternal age at pregnancy, parity, maternal height, maternal smoking, birth year, child sex. $P_{\text{interaction}}$ from model with an interaction term between log10TCDD and genotype.

* P -value significant at $\alpha < 0.05$.

** P -value significant at $\alpha < 0.01$.

Figure 2. Interactive effects of maternal rs10249788*Maternal log10TCDD on birthweight under a dominant genetic model, SWHS, 1996–2014. Models adjusted for gestational age (weeks), history of low birthweight, maternal age at pregnancy, parity, maternal height, maternal smoking, birth year, child sex.

protective influence on the association between TCDD and birthweight.

We developed a genetic risk allele score to explore the joint association of multiple *AHR* SNPs in interaction with

TCDD on birthweight. We considered the six SNPs that were interactive with TCDD. Two of these SNPs (rs2040623 and rs2106728), located at the end of the *AHR* gene, were excluded from the score as they were in high LD with two other SNPs already in the score, rs2282885 and rs3757824 ($r^2 > 0.84$), respectively, and did not contribute appreciably to the association or variance of the models. Thus, the final risk allele score was based on the four SNPs located in the intergenic and promoter regions only (rs6968865, rs3757824, rs10249788, rs2282885). Considered independently of dioxin interaction, the risk allele score had no association with birthweight (adjusted $\beta = -3.84$, 95% CI: -27.20, 19.53; $P = 0.75$). However, the score adversely modified the association of TCDD and birthweight ($P_{\text{int}} = 0.001$) (Figure 3). For example, a 10-fold increase in TCDD was not associated with birthweight among 199 children whose mothers had the lowest risk allele score (adjusted $\beta = -2.41$, 95% CI: -119.65, 114.83; $P = 0.97$). In contrast, TCDD was associated with significant decreases in birthweight in children of mothers with scores greater than 1, with the largest reductions observed when mothers carried all four risk genotypes (adjusted $\beta = -192.93$, 95% CI: -314.85, -71.01; $P = 0.002$).

Figure 3. Association of maternal TCDD levels on birthweight, stratified by maternal risk allele score. Score based on four SNPs in *AHR*, SWHS, 1996–2014. Models adjusted for gestational age (weeks), history of low birthweight, maternal age at pregnancy, parity, maternal height, maternal smoking, birth year, child sex. *Model without consideration of risk allele score.

In sensitivity analyses, excluding preterm births did not appreciably change the findings of interaction between the *AHR* gene and TCDD exposure (Supplementary Table 3, available as Supplementary data at *IJE* online) nor did including multiple births (Supplementary Table 4, available as Supplementary data at *IJE* online). When we restricted to the first post-explosion birth, the GxE coefficients were also comparable but with diminished statistical precision due to the smaller sample sizes (Supplementary Table 5, available as Supplementary data at *IJE* online). When we accounted for multiple testing with FDR adjustment, the significance of the interactions for individual SNPs was attenuated (Table 3); nevertheless, the GxE interaction by the risk allele score, which mitigates issues of multiple-testing, remained statistically significant.

Discussion

In our previous research,^{38,41} we found that birthweight was suggestively related to maternal TCDD blood concentrations and hypothesized that there may be a susceptible subgroup that is at higher risk. In this study, we examined whether variation in maternal *AHR* genotypes, coding for a transcription factor with diverse functions including dioxin metabolism, could explain heterogeneity in the effects of *in utero* dioxin exposure on birthweight. We found

interactions between maternal serum TCDD levels and six SNPs in *AHR*'s regulatory regions, particularly in proximity to the gene's promoter. We observed an even stronger adverse interaction when we examined the joint contribution of these SNPs via a risk allele score. These novel results demonstrate for the first time an interactive association between *AHR* genetics and TCDD exposure on birthweight, supporting the existence of susceptible genetic subgroups, and add to previous evidence that variation in maternal *AHR* in interaction with smoking may influence fetal growth.^{43,64,65} The only extant report of maternal *AHR* genetics (specifically, a single SNP, rs2066853) on fetal susceptibility to dioxins suggested no interaction;⁴⁴ our study, the first report in a population of European ancestry, builds upon this work by examining additional SNPs across the *AHR* gene and interaction across a wider distribution of TCDD in a population with background exposures to other dioxin-like compounds. Average reductions in birthweight (60–140 g) associated with TCDD in our study were in some cases as large as those reported among maternal smokers within high-risk genotypes in previous studies.^{43,64,65} Susceptibility to TCDD culminated in an average 192-g reduction in birthweight among children of mothers carrying all four of the highest risk genotypes.

This finding is noteworthy, given that the independent associations of neither TCDD nor maternal *AHR* genotypes with birthweight were significant (though as we previously noted, the relationship between maternal TCDD and birthweight has trended in an adverse direction in earlier analyses). Specifically, we did not observe associations between 19 *AHR* SNPs and birthweight, though the variant allele at rs2066853 was associated with a large but statistically non-significant weight reduction. This lack of a relationship is consistent with what is known from genome-wide association studies (GWAS) of birthweight; *AHR* activity is integral to fetal development, but no variants have been linked to altered fetal growth in GWAS.^{66,67} Nevertheless, previous GWAS studies of birthweight have only considered child's genetics and not maternal genetics, as examined here. This affirms the methodological caution advanced by Humblet *et al.*⁶⁸ that only testing for interaction among marginally significant SNPs may screen out and miss potentially important interactive loci in epidemiological studies.

The SNPs in the *AHR* risk allele score in our study may be important for development and toxicant metabolism. Previous experimental and epidemiological research indicate that the three SNPs in proximity to *AHR*'s promoter region may influence expression of the *AHR* gene,^{69,70} suggesting a possible biological mechanism for the interaction we observed. For example rs6968865,

upstream of *AHR*, is a documented expression quantitative trait locus (eQTL) associated with expression of *AHR* in pancreas tissue (Genotype-Tissue Expression GTEx Project, Broad Institute). The minor allele (T) of rs10249788, a SNP in *AHR*'s promoter, has been associated with higher expression of *AHR* than the C allele in human blood,⁷⁰ and downstream genes such as interleukin factors may be more upregulated in the presence of the TT genotype, independently of dioxin exposure.⁷¹ A possible biological explanation for this expression difference is offered by a study in human endometrial cells, which observed that nuclear factor 1-C (NF1C), a suppressor of *AHR* expression, preferentially bound to promoters with the C-allele compared with the T-allele at rs10249788.⁷² To the best of our knowledge, no studies on *AHR* regulation by rs3757824, another significant promoter SNP in our study that is located about 2 kb away from rs10249788, have been published, but the heterozygous genotype of this locus has been linked to higher risk of cryptorchidism in Italian boys,⁷³ suggesting a role in fetal development.

The most widely studied *AHR* variant in humans is rs2066853, an Arg554Lys missense mutation in exon 10, the codon site for *AHR*'s transactivation domain (TAD). Insight into the possible biological mechanism of rs2066853 is offered by a study that observed significantly higher mRNA levels of *AHR* in human lymphocytes with the wild-type genotype compared with the homozygote variant, Lys554Lys (AA).⁷⁴ However, functional studies of the polymorphism linking variation at rs2066853 to altered *CYP1A1* or *AHR* expression are mostly inconclusive.^{75,76} Though this SNP had the largest main effect on birthweight in our study, the association was of borderline significance and we did not observe interaction of this SNP with dioxin exposure. This finding is consistent with a recent study in a Japanese cohort, which also did not report evidence of interaction between this SNP and dioxins/TEQ on birthweight.⁴⁴ However, in this same Japanese cohort, rs2066853 was associated with circulating levels of dioxins in pregnant mothers, suggesting that rs2066853 may still be related to efficiency of dioxin metabolism.⁷⁷

To date, there is limited evidence on the functional relevance of rs2106728, found in the gene's last exon, on *AHR* activity but one study reported a large but insignificant association of this SNP with endometriosis in Japanese women.⁷⁸ This SNP and others could be worthy of further investigation in the SWHS, where dioxin exposure was observed to be non-significantly associated with a doubling of endometriosis risk.⁷⁹

There are several possible pathways by which maternal xenobiotic-metabolizing genes such as *AHR* can alter

the intrauterine environment and affect birthweight; for example, their activity contributes to xenobiotic levels in maternal-fetal circulation, placental function, endocrine regulation, and/or indirectly through the fetus's detoxification capacity inherited from the mother's alleles.⁸⁰ The relative contributions of these pathways are difficult to disentangle, but our study corroborates previous reports that maternal variation in other detoxifying proteins, particularly in the promoter region, may influence fetal susceptibility to toxicants.^{81,82} For example, we previously found that maternal and child promoter variation in *PON1*, another multifunctional, detoxifying gene, has been linked to subsequent expression and detoxifying activity of organophosphate pesticides⁸³ and possibly modifies the relationship between prenatal organophosphate exposure and neurodevelopment.^{84,85} However, whether *AHR* exhibits analogous genetic-epigenetic regulation at the site of the promoter and the contribution of the child's genetics is not yet known.

This study has several notable strengths, including the long follow-up of the SWHS through reproductive years, and one of the largest sample sizes among analyses examining GxE with *AHR*. The wide exposure distribution of TCDD in our study population also allowed for higher resolution of significant gene-dioxin interactions, particularly at the low and high ends of exposure where the interaction was most prominent. In addition, a candidate gene approach focusing on the *AHR* gene, where the biological plausibility is strong, conserved study power by limiting the number of multiple tests. Last, the genetic homogeneity in the SWHS limits confounding by population stratification.

Our study has several limitations. First, this is the earliest report examining this particular GxE association in a Caucasian population, and its generalizability to other ancestral populations is uncertain and warrants replication in additional cohorts. Second, though its potency for dioxin metabolism has been widely studied, *AHR* is just one player in a complex network of genes and proteins that mediate dioxin toxicity. Further, we only considered 19 main *AHR* SNPs among approximately 125 SNPs with >1% MAF across human populations, and we prioritized those evidencing functional significance for fetal development or detoxification. Third, in the interest of conserving study power, we used a dominant model of inheritance for all the risk alleles and an additive count of risk alleles across SNPs in composing the risk allele score on birthweight, assumptions which may not accurately reflect underlying biology. Future studies will investigate gene-dioxin interactions with respect to additional outcomes in the Seveso Women's Health Study and in the Seveso Second

Generation and the roles of other genes in the AHR pathway as well as the child's genotype.

Conclusions

This is one of the first studies to evaluate gene-dioxin interactions in a large, homogeneous cohort with a broad range of exposure levels. We used the prospective study design of the SWHS, high-quality TCDD measurements collected close to the time of the exposure, and an assessment of multiple polymorphisms across the AHR gene. We found six SNPs in maternal AHR that significantly potentiated the adverse relationship between maternal TCDD concentrations and child birthweight. AHR genetics may explain variation in human sensitivity to dioxin exposure, particularly among the offspring of TCDD-exposed mothers. Replication in additional cohorts or confirmation in mechanistic studies is warranted.

Supplementary Data

Supplementary data are available at *IJE* online.

Funding

This work was supported by funding from the National Institutes of Health (grant number F06 TW02075-01); National Institute of Environmental Health Sciences (grant numbers R01ES007171, R01ES007171-17S1, 2P30-ESO01896-17 and R24 ES028529); U.S. Environmental Protection Agency (grant number R82471); and Regione Lombardia and Fondazione Lombardia Ambiente Milan, Italy (grant number 2896). J.A. was supported by the Ruth L. Kirschstein Predoctoral Individual National Research Service Award (NIEHS) F31 Fellowship (grant number F31ES026488).

Acknowledgements

We acknowledge the significant contributions in sample analyses made by D Patterson, L Needham and W Turner to the Seveso Women's Health Study. We gratefully acknowledge Nicole Gelpi and Giulia Del Giorno for coordinating data collection at the Hospital of Desio and Maria Escudero-Fung, and Toshnia Hossain and Michael Zhou for assistance in DNA isolations at the Children's Environmental Health Laboratory (Richmond Field Station, University of California, Berkeley). We also acknowledge the significant contributions to the iPlex assay design by Shalane Porter and her colleagues at the Genomics Center of the University of Minnesota.

Conflict of interest: None declared.

References

- Di Domenico A, Cerlesi S, Ratti S. A two-exponential model to describe the vanishing trend of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) in the soil of Seveso, Northern Italy. *Chemosphere* 1990;20:1559–66.
- Mocarelli P, Pocchiari F, Nelson N. Preliminary report: 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin exposure to humans—Seveso, Italy. *MMWR Morb Mortal Wkly Rep* 1988;37:733–36.
- Mocarelli P. Seveso: a teaching story. *Chemosphere* 2001;43:391–402.
- Warner M, Mocarelli P, Brambilla P *et al.* Serum TCDD and TEQ concentrations among Seveso women, 20 years after the explosion. *J Expo Sci Environ Epidemiol* 2014;24:588–94.
- Eskenazi B, Mocarelli P, Warner M *et al.* Seveso Women's Health Study: a study of the effects of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin on reproductive health. *Chemosphere* 2000;40:1247–53.
- IARC. Polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans. *IARC Monogr Eval Carcinog Risks Hum* 1997;69:33–342.
- Rysavy NM, Maaetoft-Udsen K, Turner H. Dioxins: diagnostic and prognostic challenges arising from complex mechanisms. *J Appl Toxicol* 2013;33:01–08.
- Milbrath M, Wenger Y, Chang C *et al.* Apparent half-lives of dioxins, furans, and polychlorinated biphenyls as a function of age, body fat, smoking status, and breast feeding. *Environ Health Perspect* 2009;117:417–25.
- Chao HR, Wang SL, Lin LY, Lee WJ, Pöpke O. Placental transfer of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in Taiwanese mothers in relation to menstrual cycle characteristics. *Food Chem Toxicol* 2007;45:259–65.
- Nau H, Bass R, Neubert D. Transfer of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) via placenta and milk, and postnatal toxicity in the mouse. *Arch Toxicol* 1986;59:36–40.
- Bruner-Tran KL, Osteen KG. Developmental exposure to TCDD reduces fertility and negatively affects pregnancy outcomes across multiple generations. *Reprod Toxicol* 2011;31:344–50.
- Jin MH, Ko HK, Hong CH, Han SW. In utero exposure to 2, 3, 7, 8-Tetrachlorodibenzo-p-Dioxin affects the development of reproductive system in mouse. *Yonsei Med J* 2008;49:843–50.
- Ding T, McConaha M, Boyd KL, Osteen KG, Bruner-Tran KL. Developmental dioxin exposure of either parent is associated with an increased risk of preterm birth in adult mice. *Reprod Toxicol* 2011;31:351–58.
- Gray LE Jr, Kelce WR, Monosson E, Ostby JS, Birnbaum LS. Exposure to TCDD during development permanently alters reproductive function in male Long Evans rats and hamsters: reduced ejaculated and epididymal sperm numbers and sex accessory gland weights in offspring with normal androgenic status. *Toxicol Appl Pharmacol* 1995;131:108–18.
- Gray LE Jr, Ostby JS. In utero 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) alters reproductive morphology and function in female rat offspring. *Toxicol Appl Pharmacol* 1995;133:285–94.
- Ikeda M, Tamura M, Yamashita J, Suzuki C, Tomita T. Repeated in utero and lactational 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin exposure affects male gonads in offspring, leading to sex ratio changes in F2 progeny. *Toxicol Appl Pharmacol* 2005;206:351–55.
- Arima A, Kato H, Ooshima Y *et al.* In utero and lactational exposure to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) induces a reduction in epididymal and ejaculated sperm number in rhesus monkeys. *Reprod Toxicol* 2009;28:495–502.
- Takeyama M, Sone H, Tohyama C. Perinatal exposure of female rats to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin induces central

- precocious puberty in the offspring. *J Endocrinol* 2008;**197**: 351–58.
19. Manikkam M, Guerrero-Bosagna C, Tracey R, Haque MM, Skinner MK. Transgenerational actions of environmental compounds on reproductive disease and identification of epigenetic biomarkers of ancestral exposures. *PLoS One* 2012;**7**: e31901.
20. Myllymaki SA, Haavisto TE, Brokken LJ, Viluksela M, Toppari J, Paranko J. In utero and lactational exposure to TCDD; steroidogenic outcomes differ in male and female rat pups. *Toxicol Sci* 2005;**88**:534–44.
21. Roman BL, Peterson RE. In utero and lactational exposure of the male rat to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin impairs prostate development. 1. Effects on gene expression. *Toxicol Appl Pharmacol* 1998;**150**:240–53.
22. Sommer RJ, Ippolito DL, Peterson RE. In utero and lactational exposure of the male Holtzman rat to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin: decreased epididymal and ejaculated sperm numbers without alterations in sperm transit rate. *Toxicol Appl Pharmacol* 1996;**140**:146–53.
23. Murray FJ, Smith FA, Nitschke KD, Humiston CG, Kociba RJ, Schwetz BA. Three-generation reproduction study of rats given 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) in the diet. *Toxicol Appl Pharmacol* 1979;**50**:241–52.
24. Poland A, Glover E, Kende AS. Stereospecific, high affinity binding of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. Evidence that the binding species is receptor for induction of aryl hydrocarbon hydroxylase. *J Biol Chem* 1976;**251**: 4936–46.
25. Abe Y, Sinozaki H, Takagi T *et al.* Identification of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD)-inducible genes in human amniotic epithelial cells. *Reprod Biol Endocrinol* 2006;**4**:27.
26. Okey AB, Giannone JV, Smart W *et al.* Binding of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin to Ah receptor in placentas from normal versus abnormal pregnancy outcomes. *Chemosphere* 1997; **34**:1535–47.
27. Peltier MR, Arita Y, Klimova NG *et al.* 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD) enhances placental inflammation. *J Reprod Immunol* 2013;**98**:10–20.
28. Stejskalova L, Vecerova L, Perez LM *et al.* Aryl hydrocarbon receptor and aryl hydrocarbon nuclear translocator expression in human and rat placentas and transcription activity in human trophoblast cultures. *Toxicol Sci* 2011;**123**:26–36.
29. Wu Y, Chen X, Zhou Q *et al.* ITE and TCDD differentially regulate the vascular remodeling of rat placenta via the activation of AhR. *PLoS One* 2014;**9**:e86549.
30. Safe S, Wang F, Porter W, Duan R, McDougal A. Ah receptor agonists as endocrine disruptors: antiestrogenic activity and mechanisms. *Toxicol Lett* 1998;**102–103**:343–47.
31. Mimura J, Fujii-Kuriyama Y. Functional role of AhR in the expression of toxic effects by TCDD. *Biochim Biophys Acta* 2003; **1619**:263–68.
32. Nishijo M, Tawara K, Nakagawa H *et al.* 2, 3, 7, 8-Tetrachlorodibenzo-p-dioxin in maternal breast milk and newborn head circumference. *J Expo Sci Environ Epidemiol* 2008; **18**:246–51.
33. Halldorsson TI, Thorsdottir I, Meltzer HM, Strøm M, Olsen SF. Dioxin-like activity in plasma among Danish pregnant women: dietary predictors, birth weight and infant development. *Environ Res* 2009;**109**:22–28.
34. Tawara K, Nishijo M, Honda R *et al.* Effects of maternal dioxin exposure on newborn size at birth among Japanese mother-infant pairs. *Environ Health Prev Med* 2009;**14**:88–95.
35. Konishi K, Sasaki S, Kato S *et al.* Prenatal exposure to PCDDs/PCDFs and dioxin-like PCBs in relation to birth weight. *Environ Res* 2009;**109**:906–13.
36. Papadopoulou E, Kogevinas M, Botsivali M *et al.* Maternal diet, prenatal exposure to dioxin-like compounds and birth outcomes in a European prospective mother-child study (NewGeneris). *Sci Total Environ* 2014;**484**:121–28.
37. Van Tung D, Kido T, Honma S *et al.* Low birth weight of Vietnamese infants is related to their mother's dioxin and glucocorticoid levels. *Environ Sci Pollut Res Int* 2016;**23**:10922–29.
38. Wesselink A, Warner M, Samuels S *et al.* Maternal dioxin exposure and pregnancy outcomes over 30 years of follow-up in Seveso. *Environ Int* 2014;**63**:143–48.
39. Vartiainen T, Jaakkola JJ, Saarikoski S, Tuomisto J. Birth weight and sex of children and the correlation to the body burden of PCDDs/PCDFs and PCBs of the mother. *Environ Health Perspect* 1998;**106**:61–66.
40. Tajimi M, Uehara R, Watanabe M, Oki I, Ojima T, Nakamura Y. Relationship of PCDD/F and Co-PCB concentrations in breast milk with infant birthweights in Tokyo, Japan. *Chemosphere* 2005;**61**:383–88.
41. Eskenazi B, Mocarelli P, Warner M *et al.* Maternal serum dioxin levels and birth outcomes in women of Seveso, Italy. *Environ Health Perspect* 2003;**111**:947–53.
42. Denison MS, Pandini A, Nagy SR, Baldwin EP, Bonati L. Ligand binding and activation of the Ah receptor. *Chem Biol Interact* 2002;**141**:3–24.
43. Sasaki S, Kondo T, Sata F *et al.* Maternal smoking during pregnancy and genetic polymorphisms in the Ah receptor, CYP1A1 and GSTM1 affect infant birth size in Japanese subjects. *Mol Hum Reprod* 2006;**12**:77–83.
44. Kobayashi S, Sata F, Miyashita C *et al.* Dioxin-metabolizing genes in relation to effects of prenatal dioxin levels and reduced birth size: the Hokkaido study. *Reprod Toxicol* 2017;**67**: 111–16.
45. Warner M, Mocarelli P, Samuels S, Needham L, Brambilla P, Eskenazi B. Dioxin exposure and cancer risk in the Seveso Women's Health Study. *Environ Health Perspect* 2011;**119**: 1700–05.
46. Eskenazi B, Warner M, Brambilla P, Signorini S, Ames J, Mocarelli P. The Seveso accident: A look at 40 years of health research and beyond. 2018. Submitted for publication.
47. Patterson DG, Hampton L, Lapeza CR *et al.* High-resolution gas chromatographic/high-resolution mass spectrometric analysis of human serum on a whole-weight and lipid basis for 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin. *Anal Chem* 1987;**59**: 2000–05.
48. Needham LL, Gerthoux PM, Patterson DG Jr *et al.* Serum dioxin levels in Seveso, Italy, population in 1976. *Teratog Carcinog Mutagen* 1997;**17**:225–40.
49. Eskenazi B, Mocarelli P, Warner M *et al.* Relationship of serum TCDD concentrations and age at exposure of female residents of Seveso, Italy. *Environ Health Perspect* 2004;**112**:22–27.

50. Akins JR, Waldrep K, Bernert JT Jr. The estimation of total serum lipids by a completely enzymatic 'summation' method. *Clin Chim Acta* 1989;184:219–26.
51. Hornung RW, Reed LD. Estimation of average concentration in the presence of non-detectable values. *Appl Occup Environ Hyg* 1990;5:46–51.
52. Walker SK, Hartwich KM, Robinson JS. Long-term effects on offspring of exposure of oocytes and embryos to chemical and physical agents. *Hum Reprod Update* 2000;6:564–77.
53. Thorisson GA, Smith AV, Krishnan L, Stein LD. The International HapMap Project Web site. *Genome Res* 2005;15:1592–93.
54. Boyle AP, Hong EL, Hariharan M *et al.* Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* 2012;22:1790–97.
55. Holland N, Furlong C, Bastaki M *et al.* Paraoxonase polymorphisms, haplotypes, and enzyme activity in Latino mothers and newborns. *Environ Health Perspect* 2006;114:985–91.
56. Liang K-Y, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika* 1986;73:13–22.
57. Chawla R, Badon SE, Rangarajan J *et al.* Genetic risk score for prediction of newborn adiposity and large-for-gestational-age birth. *J Clin Endocrinol Metab* 2014;99:E2377–86.
58. Lawn JE, Gravett MG, Nunes TM, Rubens CE, Stanton C; GAPPs Review Group. Global report on preterm birth and stillbirth (1 of 7): definitions, description of the burden and opportunities to improve data. *BMC Pregnancy Childbirth* 2010;10(Suppl 1):S1.
59. R Core Team R: *A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing, 2017.
60. Gonzalez JR, Armengol L, Sole X *et al.* SNPassoc: an R package to perform whole genome association studies. *Bioinformatics* 2007;23:654–55.
61. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B* 1995;57:289–300.
62. Wardlaw TM; World Health Organization; UNICEF. *Low Birthweight: Country, Regional and Global Estimates*. Geneva, New York: WHO, UNICEF, 2004.
63. Auton A, Brooks LD, Durbin RM *et al.* A global reference for human genetic variation. *Nature* 2015;526:68–74.
64. Kobayashi S, Sata F, Sasaki S *et al.* Combined effects of AHR, CYP1A1, and XRCC1 genotypes and prenatal maternal smoking on infant birth size: Biomarker assessment in the Hokkaido Study. *Reprod Toxicol* 2016;65:295–306.
65. Kobayashi S, Sata F, Sasaki S *et al.* Modification of adverse health effects of maternal active and passive smoking by genetic susceptibility: dose-dependent association of plasma cotinine with infant birth size among Japanese women - The Hokkaido Study. *Reprod Toxicol* 2017;74:94–103.
66. Horikoshi M, Beaumont RN, Day FR *et al.* Genome-wide associations for birth weight and correlations with adult disease. *Nature* 2016;538:248–52.
67. Freathy RM, Mook-Kanamori DO, Sovio U *et al.* Variants in ADCY5 and near CCNL1 are associated with fetal growth and birth weight. *Nat Genet* 2010;42:430–35.
68. Humblet O, Korrick SA, Williams PL *et al.* Genetic modification of the association between peripubertal dioxin exposure and pubertal onset in a cohort of Russian boys. *Environ Health Perspect* 2012;121:111–17.
69. Sulem P, Gudbjartsson DF, Geller F *et al.* Sequence variants at CYP1A1-CYP1A2 and AHR associate with coffee consumption. *Hum Mol Genet* 2011;20:2071–77.
70. Wang X, Li K, Liu L *et al.* AHR promoter variant modulates its transcription and downstream effectors by allele-specific AHR-SP1 interaction functioning as a genetic marker for vitiligo. *Sci Rep* 2015;5:13542.
71. Liu G, Asanoma K, Takao T *et al.* Aryl hydrocarbon receptor SNP -130 C/T associates with dioxins susceptibility through regulating its receptor activity and downstream effectors including interleukin 24. *Toxicol Lett* 2015;232:384–92.
72. Li D, Takao T, Tsunematsu R *et al.* Inhibition of AHR transcription by NF1C is affected by a single-nucleotide polymorphism, and is involved in suppression of human uterine endometrial cancer. *Oncogene* 2013;32:4950–59.
73. Qin X-Y, Kojima Y, Mizuno K *et al.* Association of variants in genes involved in environmental chemical metabolism and risk of cryptorchidism and hypospadias. *J Hum Genet* 2012;57:434–41.
74. Helmig S, Seelinger JU, Döhrel J, Schneider J. RNA expressions of AHR, ARNT and CYP1B1 are influenced by AHR Arg554Lys polymorphism. *Mol Genet Metab* 2011;104:180–84.
75. Wong JM, Harper PA, Meyer UA *et al.* Ethnic variability in the allelic distribution of human aryl hydrocarbon receptor codon 554 and assessment of variant receptor function in vitro. *Pharmacogenetics* 2001;11:85–94.
76. Daly AK, Fairbrother KS, Smart J. Recent advances in understanding the molecular basis of polymorphisms in genes encoding cytochrome P450 enzymes. *Toxicol Lett* 1998;102–103:143–47.
77. Kobayashi S, Sata F, Sasaki S *et al.* Genetic association of aromatic hydrocarbon receptor (AHR) and cytochrome P450, family 1, subfamily A, polypeptide 1 (CYP1A1) polymorphisms with dioxin blood concentrations among pregnant Japanese women. *Toxicol Lett* 2013;219:269–78.
78. Matsuzaka Y, Kikuti YY, Goya K *et al.* Lack of an association human dioxin detoxification gene polymorphisms with endometriosis in Japanese women: results of a pilot study. *Environ Health Prev Med* 2012;17:512–17.
79. Eskenazi B, Mocarelli P, Warner M *et al.* Serum dioxin concentrations and endometriosis: a cohort study in Seveso, Italy. *Environ Health Perspect* 2002;110:629–34.
80. Wells PG, Lee CJ, McCallum GP, Perstin J, Harper PA. Receptor- and reactive intermediate-mediated mechanisms of teratogenesis. *Handb Exp Pharmacol* 2010;196:131–62.
81. Li YF, Gauderman WJ, Conti DV, Lin PC, Avol E, Gilliland FD. Glutathione S-transferase P1, maternal smoking, and asthma in children: a haplotype-based analysis. *Environ Health Perspect* 2008;116:409–15.
82. Tekin D, Kayaaltı Z, Söylemezoğlu T. The effects of metallothionein 2A polymorphism on lead metabolism: are pregnant women with a heterozygote genotype for metallothionein 2A polymorphism and their newborns at risk of having higher blood lead levels? *Int Arch Occup Environ Health* 2012;85:631–37.

-
83. Huen K, Yousefi P, Street K, Eskenazi B, Holland N. PON1 as a model for integration of genetic, epigenetic, and expression data on candidate susceptibility genes. *Environ Epigenet* 2015;1 2015 Jan;1. pii: dvv003. [Epub 2015 Sep 11].
 84. Eskenazi B, Kogut K, Huen K *et al.* Organophosphate pesticide exposure, PON1, and neurodevelopment in school-age children from the CHAMACOS study. *Environ Res* 2014;134: 149–57.
 85. Eskenazi B, Huen K, Marks A *et al.* PON1 and neurodevelopment in children from the CHAMACOS study exposed to organophosphate pesticides in utero. *Environ Health Perspect* 2010; 118:1775–81.