HSP90 Contributes to Entrainment of the *Arabidopsis* Circadian Clock via the Morning Loop

Amanda M. Davis,^{*,†,1} James Ronald,^{†,1} Zisong Ma,^{*,†} Anthony J. Wilkinson,[†] Koumis Philippou,^{*} Takayuki Shindo,^{*} Christine Queitsch,[†] and Seth J. Davis^{*,†,2}

^{*}Max Planck Institute for Plant Breeding Research, Cologne, Germany, [†]University of York, UK, and [‡]University of Washington, Seattle, Washington

ORCID IDs: 0000-0002-8847-0378 (J.R.); 0000-0001-5928-9046 (S.J.D.)

ABSTRACT The plant circadian clock allows the synchronization of internal physiological responses to match the predicted environment. HSP90.2 is a molecular chaperone that has been previously described as required for the proper functioning of the *Arabidopsis* oscillator under both ambient and warm temperatures. Here, we have characterized the circadian phenotype of the *hsp90.2-3* mutant. As previously reported using pharmacological or RNA interference inhibitors of HSP90 function, we found that *hsp90.2-3* lengthens the circadian period and that the observed period lengthening was more exaggerated in warm–cold-entrained seedlings. However, we observed no role for the previously identified interactors of HSP90.2, GIGANTEA and ZEITLUPPE, in HSP90-mediated period lengthening. We constructed phase-response curves (PRCs) in response to warmth pulses to identify the entry point of HSP90.2 to the oscillator. These PRCs revealed that *hsp90.2-3* has a circadian defect within the morning. Analysis of the *cca1*, *lhy*, *prp9*, and *prp7* mutants revealed a role for CCA1, LHY, and PRR7, but not PRR9, in HSP90.2 action to the circadian oscillator. Overall, we define a potential pathway for how HSP90.2 can entrain the *Arabidopsis* circadian oscillator.

KEYWORDS *Arabidopsis*; circadian clock; HSP90

MANY organisms have evolved an internal timing mechanism called the circadian clock to anticipate predictable environmental changes. In *Arabidopsis*, the circadian clock regulates approximately one-third of genes and 36% of *Arabidopsis* promoters show circadian regulation by transcript accumulation (Covington and Harmer 2007; Covington *et al.* 2008; Staiger *et al.* 2013). During the process of entrainment, the internal circadian clock is reset by daily exogenous cues (*zeitgebers*) to maintain synchronization with the diurnal cycle. For most organisms, the dominant *zeitgebers* are light and temperature changes perceived at dawn (Oakenfull and Davis 2017). Light input to the clock occurs via multiple types of photoreceptors; for example, in plants, phytochrome and cryptochromes control red- and blue-light signaling to the

clock (Oakenfull and Davis 2017). However, for temperature, the *zeitgeber* input pathway leading to clock entrainment remains poorly understood (Boikoglou *et al.* 2011; Bujdoso and Davis 2013; Anwer *et al.* 2014).

One current model of the plant circadian clock consists of interlocked transcriptional–translational feedback loops (Bujdoso and Davis 2013; Ronald and Davis 2017). At the center of these loops are the morning-expressed *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) and *LATE ELONGATED HYPOCOTYL* (*LHY*), and the evening-expressed *TIMING OF CAB EXPRESSION 1* (*TOC1*, also called *PSEUDO RESPONSE REGULATOR 1*, *PRR1*) (Mizoguchi *et al.* 2002; Ding *et al.* 2007). Upon being expressed, CCA1/LHY bind to the evening-element within the *TOC1* promoter and directly repress *TOC1* expression (Nagel *et al.* 2015). At dusk, *TOC1* reciprocally represses the expression of *CCA1/LHY*, generating a negative feedback loop (Gendron *et al.* 2012).

Morning- and evening-phased regulators subsequently regulate CCA1/LHY and *TOC1* activity. Starting just after dawn, *PRR9/7/5* are sequentially expressed throughout the day and directly repress the expression of *CCA1/LHY* through the recruitment of the TOPLESS corepressor (Nakamichi

Copyright © 2018 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.118.301586>

Manuscript received September 7, 2018; accepted for publication October 11, 2018; published Early Online October 18, 2018.

Supplemental material available at Figshare: <https://doi.org/10.25386/genetics.7223579>.

¹These authors contributed equally to this work.

²Corresponding author: University of York, Wentworth Way, York, UK. E-mail: seth.davis@york.ac.uk

et al. 2010; Wang *et al.* 2013). In the evening, GIGANTEA (GI) interacts and stabilizes the F-box protein ZEITLUPPE (ZTL) in a blue-light-dependent manner to promote the degradation of TOC1 and PRR5 (Kim *et al.* 2007), which supports the role of protein–protein interactions in stabilizing circadian period (Schöning and Staiger 2005). Finally, the evening complex, composed of LUX ARRHYTHMO (LUX), EARLY FLOWERING3 (ELF3), and ELF4, represses the expression of *LUX*, *GI*, *PRR7*, and *PRR9* (Nusinow *et al.* 2011; Herrero *et al.* 2012).

HEAT SHOCK PROTEIN90 (HSP90) is a highly conserved and abundant protein in prokaryotes and eukaryotes. The HSP90 family of proteins is involved in the assembly, maturation, stabilization, and activation of proteins (Chen *et al.* 2005). *Arabidopsis* has seven HSP90 isoforms (HSP90.1–7). Of these, four display cytosolic localization (HSP90.1–4), and the remaining (HSP90.5–7) are predicted to be localized to the chloroplast, mitochondria, and endoplasmic reticulum, respectively (Krishna and Gloor 2001). HSP90.2 has been previously linked to the circadian clock through protein–protein interactions with GI and ZTL (Kim *et al.* 2011; Noren *et al.* 2016; Cha *et al.* 2017; Gil *et al.* 2017), and alleles at this locus have pathology phenotypes (Hubert *et al.* 2003). HSP90 and GI are reported to act as cochaperones to promote ZTL maturation and accumulation (Cha *et al.* 2017). Inhibition of global HSP90 activity by geldanamycin (GDA) application or through specific targeting of cytosolic HSP90 isoforms causes a lengthening of the circadian period (Kim *et al.* 2011). ZTL and HSP90 have recently been shown to impart thermotolerance to the circadian clock by acting as a protein quality control system at warmer temperatures (Gil *et al.* 2017). GI and ZTL also act as a hub in the plastid control of nuclear circadian rhythms in a PRR5 and HY5 signaling pathway (Noren *et al.* 2016). Therefore, HSP90 isoforms likely have multiple roles within the circadian oscillator.

In this study, we have characterized the circadian phenotype of the *hsp90.2-3* mutant. This specific allele at *HSP90.2* was chosen given its strong “poison pill” phenotype, as the null had no pathology phenotype (Hubert *et al.* 2003). We found that *hsp90.2-3* has a longer circadian period in both light–dark (LD)- and warm–cold (WC)-entrained plants. This period-lengthening effect did not require either of the previously identified circadian interacting partners of HSP90.2: ZTL or GI. Phase-response curves (PRCs) in response to warmth pulses were constructed and revealed that *hsp90.2-3* displayed a defect at the morning phase. Further analysis revealed that the period-lengthening effects of GDA were lost in the *cca1*, *lhy*, and *prp7* backgrounds. However, no genetic role of *PRR9* was observed, revealing functional independence between *PRR9* and *PRR7*. Thus, this work has revealed new insights regarding how HSP90 could contribute to clock function.

Materials and Methods

Plant lines

Ws-2 and Col-0 were used as wildtype (WT), either harboring *CCR2::LUC*, *CCA1::LUC*, or *CAB2::LUC* (Doyle *et al.* 2002;

Farré *et al.* 2005). The luciferase-containing *gi-11*, *ztl* alleles, *prp7-3*, *prp9-1*, *prp7-3/prp9-1*, *cca1-11*, *lhy-21*, *cca1-11/lhy-21*, and *hsp90.2-3* have all been described previously (Fowler *et al.* 1999; Hubert *et al.* 2003; Farré *et al.* 2005; Kevei *et al.* 2006; Ding *et al.* 2007). Before circadian phenotyping of *hsp90.2-3*, it was backcrossed six times to Ws-2 *CCR2::LUC*, and in the BC6F2 generation a homozygous *hsp90.2-3 CCR2::LUC* line was isolated and bulked for analyses.

Bioluminescent assays

Seeds were surface-sterilized and plated onto MS medium with 3% sucrose, and then stratified for ~3 days. After stratification, seedlings were entrained under either 12/12 LD cycles (with a constant temperature of 22°) or 12/12 cycles of 22°/16° (with constant light) for 7 days (Boikoglou *et al.* 2011; Anwer *et al.* 2014). On day 6, seedlings were transferred to black 96-well Microplates with MS medium containing 3% sucrose with DMSO and, where relevant, 2 μM of GDA. Plants were superficially treated with 15 μl 5 mM D-Luciferin. Seedlings were then reentrained for 1 day under the respective entrainment conditions before being transferred to the TOPCOUNT (Perkin-Elmer [Perkin-Elmer-Cetus], Norwalk, CT). All TOPCOUNT experiments were carried out under constant blue–red light and a constant temperature of 21°. Data were analyzed as previously described (Hanano *et al.* 2006, 2008; Kolmos *et al.* 2009). All experiments were replicated and provided consistent results.

PRC assays

For PRC assays, plants were grown as for luciferase assays, as described above, under 7 days LD conditions (12 hr in light and 12 hr in darkness), and then transferred to a TOPCOUNT under constant red and blue light for one full day before a 3-hr long 27° warmth pulse. This was respectively applied every 3 hr to a given 96-well plate beginning at (*zeitgeber* time) ZTO. Resultant data were then analyzed using *Peak Picker* in the Biological rhythms analysis software system (BRASS) (Southern and Millar 2005). Here, the first peak after the warmth treatment was chosen for both pulsed and nonpulsed plates, and the time difference of the timing of the peak between pulsed and nonpulsed populations was calculated (Covington *et al.* 2001).

Statistical analysis

All statistical analysis was completed using R (version 3.4.2) within the R studio software package (version 1.1). Unless stated otherwise, the sample size for determining period estimates was 48 plants.

Data availability

Seeds are available upon request. The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables. Supplemental material available at Figshare: <https://doi.org/10.25386/genetics.7223579>.

Figure 1 *hsp90.2-3* has a circadian period phenotype. Free-running profile of *CCR2::LUC* in Ws-2 and *hsp90.2-3* seedlings under constant red-blue light (LL) after plants were prior trained to 12:12 cycles of (A and C) light-dark (LD) or (B and D) warm-cold (WC). Plants were released into free-running conditions at *zeitgeber* time ZT36. (C and D) Mean period estimates of (A) and (B), respectively. Error bars represent SEM. *** $P < 0.001$. Significance determined using a Student's *t*-test. In each experiment, 48 WT and *hsp90.2-3* seedlings were analyzed for rhythms under each entrainment protocol. All experiments repeated at least once. CPS, counts per second.

Results

hsp90.2-3 causes a lengthening of circadian period

To establish if *hsp90.2-3* has a circadian phenotype, it (Hubert *et al.* 2003) was introgressed with Ws-2 WT plants harboring *CCR2::LUC* [also termed *GRP7* (Köster *et al.* 2014)] to generate the *hsp90.2-3 CCR2::LUC* line. The free-running period (FRP) of *CCR2::LUC* was then analyzed under constant light after plants were either entrained to LD or WC cycles. After either entrainment protocol, *hsp90.2-3* was found to lengthen *CCR2::LUC* FRP (Figure 1, A–D), but the magnitude of period lengthening was greater under WC entrainment compared to LD entrainment [$\Delta LD = 0.78 \pm 0.05$ hr, $\Delta WC = 0.92 \pm 0.05$ hr, ($P < 0.05$)]. *hsp90.2-3* plants also displayed a change in the amplitude of *CCR2::LUC* rhythms; the amplitude of *CCR2::LUC* rhythms increased in LD-entrained plants, while the amplitude decreased in WC-entrained plants (Figure 1, A and B).

As previously reported (O'Neill *et al.* 2011), GDA treatment of Ws-2 *CCR2::LUC* resulted in a similar lengthening of FRP as observed in the *hsp90.2-3* mutant (Figure 2A). To determine if multiple HSP90 isoforms signal redundantly to the circadian clock, *hsp90.2-3* mutants were treated with 2 μ M GDA. For both WC- and LD-entrained plants, GDA treatment resulted in further lengthening of *CCR2::LUC* FRP compared to non-treated *hsp90.2-3* seedlings (Figure 2B). However, unlike the *hsp90.2-3* mutant, *hsp90.2-3* in combination with 2 μ M GDA resulted in a more severe period lengthening under LD entrainment compared to WC entrainment ($\Delta LD = 1.42 \pm 0.05$ hr, $\Delta WC = 1.09 \pm 0.06$ hr, [$P < 0.001$]). This suggests that HSP90 isoforms act redundantly within the oscillator, but may contribute independently to different entrainment pathways.

To confirm the effects of GDA on periodicity, the FRP profile of Ws-2 *CAB2::LUC* was also tested. As seen with *CCR2::LUC*, *CAB2::LUC* FRP was longer when treated with 2 μ M GDA compared to WT regardless of the prior entrainment condition

(Supplemental Material, Figure S1A). The period-lengthening effect was also found to not be accession-dependent; Col-0 *hsp90.2-3 GI::LUC* lines also had a longer FRP than Col-0 WT *GI::LUC* (Figure S1B). Additionally, 2 μ M GDA treatment of Col-0 WT seedlings also lengthened the FRP of both morning (*CCA1::LUC*) and evening (*TOC1::LUC*) reporter genes (Figure S1, C and D). These results suggest that the *hsp90.2-3* phenotype and the effects of GDA on circadian periodicity are not dependent on the reporter gene or ecotype used.

HSP90 circadian period lengthening genetically does not require *GI* or *ZTL*

HSP90.2 protein has been previously reported to interact with the circadian component *GI* to stabilize *ZTL* (Kim *et al.* 2011). To determine if the observed effects of GDA on circadian periodicity required the activity of either *GI* or *ZTL*, the FRP of *CCR2::LUC* was analyzed in the previously described *gi-11* or *ztl-21* mutants (Gould *et al.* 2006; Kim *et al.* 2007). *gi-11 CCR2::LUC* or *ztl-21 CAB2::LUC* were entrained to either LD or WC cycles before being treated with 2 μ M GDA upon release into free-running conditions. Regardless of the prior entrainment condition, GDA treatment caused a lengthening of FRP in both the *gi-11* and *ztl-21* backgrounds (Figure 3). To confirm the *ztl* result, the FRPs of additional *ztl* mutants harboring *CAB2::LUC* were examined. As was observed with *ztl-21 CAB2::LUC*, in all but one instance the other *ztl* mutants displayed a lengthening of circadian period under both LD and WC entrainment when treated with GDA (Figure S2). The only *ztl* allele that did not show any period lengthening was *ztl-30*, but this response was only seen in LD-entrained plants (Figure S2A). Under WC conditions, GDA treatment in the *ztl-30* background caused the same period lengthening as observed for WT and the other *ztl* alleles treated with GDA (Figure S2B).

To identify if GDA had an additive effect in the *ztl* or *gi* mutant backgrounds, Δ period changes in response to

Figure 2 Geldanamycin (GDA) treatment has an additive effect on *hsp90.2-3* circadian period phenotype. Period estimates of the free-running period (FRP) of *CCR2::LUC* in the (A) *Ws-2* or (B) *hsp90.2-3* seedlings treated with either DMSO or with 2 μ M GDA under constant red-blue light (LL). For (A), plants were entrained under 12:12 light-dark cycles. Plants in (B) were entrained under the stated entrainment conditions. For both (A) and (B), GDA was applied upon transfer to free-running conditions. Error bars represent SEM. ** $P < 0.01$ and *** $P < 0.001$. Significance determined via a Student's *t*-test. In each experiment, the FRP of 48 seedlings was examined. Each experiment was repeated at least once.

GDA treatment were calculated (Tables S1 and S2). No significance difference in period lengthening was observed for *gi-11* under LD or WC entrainment, or the majority of the *ztl* alleles entrained under LD cycles compared to *Ws-2*. However, under WC cycles the reverse was seen. Most *ztl* alleles had a significantly greater period lengthening effect when treated with GDA compared to GDA-treated WT plants (Table S2). This therefore suggests that the general period lengthening effect caused by GDA treatment did not require the presence of GI or ZTL, but HSP90 and ZTL activity could converge in a temperature-entrainment pathway.

hsp90.2-3 has a morning-phase defect

To identify the point of entry of HSP90.2 to the circadian oscillator, a PRC was generated for the WT and the *hsp90.2-3* genotypes. PRCs test the sensitivity of the oscillator to resetting stimuli (*zeitgebers*) at different points of the day (Covington *et al.* 2001). Three-hour long warmth pulses of

Figure 3 Geldanamycin (GDA) lengthening of circadian period is not dependent on GI or ZTL. Period estimates of the free-running profile of (A) *gi-11* *CCR2::LUC* or (B) *ztl-21* *CAB2::LUC*. Plants were entrained under 12:12 cycles of light-dark (LD) or warm-cold (WC) before transfer to free-running conditions. Next, 2 μ M of GDA was applied upon transfer to free-running conditions. Error bars represent SEM. ** $P < 0.01$ and *** $P < 0.001$. Significance determined using a Student's *t*-test. In each experiment, 48 seedlings of WT and the respective mutant were examined under each entrainment condition, apart from *gi-11* LD (GDA) where $n = 30$. All experiments were repeated at least once.

27° were applied at respective 3-hr intervals to *Ws-2* and *hsp90.2-3* plants throughout the day, and then the phase change in *CCR2::LUC* expression was recorded. In *Ws-2*, these warmth pulses elicited a phase advance of ~4–7 hr during the period from dawn to late morning (Figure 4). *hsp90.2-3* showed reduced phase advances during the same period; there was no observable phase advance at dawn and only a maximum phase advance of ~2 hr by late morning (Figure 4). During the later afternoon, *hsp90.2-3* showed a subtler phase delay than was seen in *Ws-2* (~0.5 and 2 hr, respectively). No change in *CCR2::LUC* phase response was seen between *hsp90.2-3* and *Ws-2* across the subjective night. This indicates that the likely entry point of HSP90.2 to the circadian oscillator occurs within the morning.

CCA1/LHY and *PRR7* are a hub for HSP90 circadian activity

The morning loop of the *Arabidopsis* circadian clock is primarily composed of *CCA1/LHY* and *PRR9/7* arranged in a reciprocal repressive loop (Bujdosó and Davis 2013; Ronald and Davis 2017). To determine if the observed period

Figure 4 *hsp90.2-3* has a morning phase defect. *Ws-2* and *hsp90.2-3 CCR2::LUC* plants were entrained for 7 days under 12/12 light–dark cycles before being exposed to 3-hr long pulses of 27°. Phase-response curves were then constructed by plotting the observed phase shift in *CCR2::LUC* expression against the circadian time that heat pulses were administered. Positive values represent phase advances and negative values represent phase delays. *** $P < 0.001$; n.s., no significant difference. Significance determined via a Student's *t*-test. Error bars represent pooled SE.

lengthening effects of GDA required either *CCA1/LHY*, *PRR9*, or *PRR7*, the effects of GDA on the FRP profile of the respective mutant was analyzed after both LD- and WC-entrainment conditions, respectively. *cca1-11*, *lhy-21*, and *cca1-1/lhy-21* mutants harboring *CCR2::LUC* displayed no period lengthening when treated with 2 μ M GDA, regardless of the prior entrainment condition (Figure 5). Similarly to *cca1* and *lhy*, GDA treatment of *prp7-3 CCA1::LUC* and *prp7-3/prp9-1 CCA1::LUC* lines resulted in no observed period lengthening after either LD or WC entrainment (Figure 6). However, unlike *prp7*, *prp9-1 CCA1::LUC* did display a similar response to WT; GDA treatment caused a lengthening of *prp9-1* FRP and this occurred independently of the prior entrainment conditions (Figure 6, A and B). As was observed in the *hsp90.2-3* mutant and in WT treated with GDA, *prp9-1* displayed a greater period lengthening when WC entrained. Together, this suggests that HSP90 requires the presence of *CCA1*, *LHY*, and *PRR7*, but not *GI* nor *ZTL*, to lengthen circadian period.

Discussion

HSP90.2 is a molecular chaperone previously shown to be required for proper circadian rhythms (Kim *et al.* 2011; O'Neill *et al.* 2011; Gil *et al.* 2017). Here, we have confirmed this result by characterizing the *hsp90.2-3* mutant. This mutant was found to have a longer circadian period regardless of prior entrainment conditions, although there was a greater phenotypic defect after WC entrainment (Figure 1). Such phenotypes were not dependent on the *Arabidopsis* ecotype or reporter gene used (Figure S1). Our work also supports the notion that HSP90 isoforms function in a partially redundant manner to the circadian clock, as treating the *hsp90.2-3*

Figure 5 Geldanamycin (GDA) fails to lengthen circadian period in the *cca1* or *lhy* mutant. (A and B) Period estimates of *CCR2::LUC* profile under free-running conditions in *Ws-2* [wild-type (WT)], *cca1-11*, and *lhy-21* mutants treated with or without 2 μ M GDA. Plants were prior entrained to light–dark (A) or warm–cold (B) cycles before being released into free-running conditions. GDA treatment was applied upon transfer to free-running conditions. Error bars represent SEM. In each experiment, 48 WT and mutant seedlings were examined under each entrainment condition. n.s., no significance; *** $P < 0.001$. Significance determined by a Student's *t*-test. All experiments were repeated at least once.

mutant with GDA resulted in further lengthening of the circadian period (Figure 2). However, unlike the nontreated *hsp90.2-3* mutant, GDA treatment of *hsp90.2-3* resulted in a more pronounced period lengthening in LD- compared to WC-entrained plants. This suggests that HSP90 isoforms likely function redundantly in the general regulation of clock periodicity, and also that individual isoforms could contribute to separate light- and temperature-entrainment pathways.

Duality-of-function as both a core circadian component and separately as a contributor to the entrainment of the oscillator has now been described for *ELF3*, *PRR9*, *PRR7*, *CCA1/LHY*, and *GI* (Farré *et al.* 2005; Gould *et al.* 2006; Ding *et al.* 2007; Thines and Harmon 2010; Dalchau *et al.* 2011; Bujdosó and Davis 2013; Haydon *et al.* 2013). The contribution of HSP90 to either light or thermal entrainment of the clock could be dependent on the cellular localization of HSP90 isoforms. These isoforms in *Arabidopsis* are localized to different cellular compartments and would therefore have different client proteins, and indeed a host of protein interactions can be predicted in the regulation of periodicity (Schöning and Staiger 2005; Bujdosó and Davis 2013). For example, HSP90.5 is

Figure 6 The effect of geldanamycin (GDA) treatment on period length is disrupted in the *prr7* and *prr7/prr9* background. Period estimates of the free-running profile of *CCA1::LUC* in the Col-0, *prr7-3*, *prr9-1*, and *prr7-3/prr9-1* background. Plants were entrained under (A) light-dark or (B) warm-cold cycles before being released into free-running conditions. Plants were treated with or without 2 μ M GDA upon transfer to free-running conditions. Error bars represent SEM. ** $P < 0.01$ and *** $P < 0.001$; n.s., no significance. Significance determined via a Student's *t*-test. In each experiment, 48 WT and mutant seedlings were examined under each entrainment condition. All experiments were repeated at least once.

localized to the stroma, has a light-responsive transcript accumulation, and *hsp90.5* (*cr88*) has defects in red-light perception (Lin and Cheng 1997; Cao *et al.* 2003). The perception of red-light is critical for the entrainment of the clock (Oakenfull and Davis 2017). Therefore, HSP90 isoforms may contribute within separate light- and temperature-entrainment pathways in a cell localization-dependent manner.

HSP90 has been previously linked to the circadian oscillator through direct interactions with GI and ZTL (Kim *et al.* 2011). Applying 2 μ M of GDA to either the *gi* or *ztl* mutants resulted in a lengthening of circadian period, as seen with WT (Figure 3 and Figure S2), suggesting that HSP90 does not require either ZTL or GI to regulate FRP. HSP90 and ZTL have been recently shown to maintain circadian thermostability, and GI is also required to maintain circadian oscillations and the precision of these oscillations under warm temperatures (Gould *et al.* 2006; Gil *et al.* 2017). We did find that applying 2 μ M of GDA to either the *ztl* or *gi* mutant increased the

Figure 7 An expanding role of HSP90 within the *Arabidopsis* circadian oscillator. HSP90 has been previously shown to interact with ZTL to regulate both periodicity and, under heat stress, the stability of the oscillator. GI and HSP90 are thought to cooperatively stabilize ZTL activity. Here, we have found that Hsp90 also signals independently of GI and ZTL through the morning loop components CCA1/LHY and PRR7. We did not detect a direct effect of HSP90 on regulating *CCA1/LHY* expression, and HSP90 was also found previously to not regulate *PRR7* expression. Therefore, this indicates that HSP90 is signaling via an as yet unidentified protein to regulate *CCA1/LHY* and *PRR7* activity. Purple lines indicate an interaction (direct or indirect), red lines indicate a repressive interaction, and blue lines highlight the effect of the interaction on the oscillator.

variance of periodicity estimates, an effect not seen in *hsp90.2-3* or with GDA treatment in either the *Ws-2* or Col-0 background (data not shown). Therefore, the previously described HSP90/ZTL/GI module may act as a buffering agent to maintain clock precision through the regulation of proteostasis, while HSP90.2 could function independently of the GI and ZTL module to regulate clock periodicity (Figure 7).

PRCs for WT and *hsp90.2-3* plants exposed to warmth pulses were constructed to determine when HSP90.2 regulates circadian periodicity. These revealed a phase defect within the morning (Figure 4). We subsequently found that there was no lengthening of circadian period in *cca1*, *lhy*, and *cca1/lhy* plants treated with GDA regardless of the prior entrainment conditions (Figure 5). We also found that *prr7* and *prr7/9* failed to show any lengthening of FRP when treated with GDA after either LD or WC conditions, respectively (Figure 6). Genetically, one interpretation is that the timing of the entry point of HSP90.2 depends equally on *LHY*, *CCA1*, and *PRR7*. In contrast, *prr9* did show a longer circadian period when treated with GDA (Figure 6). This reveals a functional independence between *PRR9* and *PRR7* within the circadian oscillator. Together, these data suggest that *CCA1/LHY*, *HSP90*, and *PRR7* constitute a unique morning loop that regulates general circadian oscillations regardless of the prior entrainment condition (Figure 7), and that it is consistent with the morning PRC defects seen in *hsp90.2-3* plants (Figure 4).

It is not fully clear how HSP90.2 signals through *CCA1/LHY* and *PRR7*. HSP90 isoforms have a large client pool of proteins (Kadota and Shirasu 2012) and many genes become misregulated when *HSP90* function is inhibited, including *PRR9* and *CCA1* (Sangster *et al.* 2007). However, we observed

no role for PRR9 in GDA's period lengthening effect (Figure 6), and we also observed no change in *CCA1* or *LHY* expression levels or patterns of expression in LD-entrained plants (data not shown). Previous work also found no effect of GDA on PRR7's protein stability or gene-expression profile (Kim *et al.* 2011). This therefore suggests that the effects of HSP90 on *CCA1/LHY* and PRR7 may not be fully direct, and could occur upstream of these transcriptional regulators. The activity of *CCA1/LHY* and PRR7 is modulated by a range of morning-associated transcriptional activators and afternoon/evening-expressed transcriptional repressors (Ronald and Davis 2017). Further screens of the FRP of clock mutants when treated with GDA will provide further answers for how HSP90 regulates clock periodicity. In this, it is notable that physiologically and genetically, temperature and light set the clock in differing ways (Boikoglou *et al.* 2011; Anwer *et al.* 2014).

Acknowledgments

We thank Jeff Dangl for supplying the *hsp90.2-3* allele, Jane Parker for her encouragement of this work, Sarah Lock for her assistance with statistical analysis, and Paula Avello Fernández and Jon Pitchford for comments on the manuscript. This work is dedicated to the memory of Exakoustodios Philippou. Funding was through the Max-Planck-Gesellschaft, the International Max Planck Research Schools program, and Biotechnology and Biological Sciences Research Council award BB/N018540/1.

Literature Cited

- Anwer, M. U., E. Boikoglou, E. Herrero, M. Hallstein, A. M. Davis *et al.*, 2014 Natural variation reveals that intracellular distribution of ELF3 protein is associated with function in the circadian clock. *Elife* 3: e02206. <https://doi.org/10.7554/eLife.02206>
- Boikoglou, E., Z. Ma, M. von Korff, A. M. Davis, F. Nagy *et al.*, 2011 Environmental memory from a circadian oscillator: the *Arabidopsis thaliana* clock differentially integrates perception of photic vs. thermal entrainment. *Genetics* 189: 655–664. <https://doi.org/10.1534/genetics.111.131417>
- Bujdoso, N., and S. J. Davis, 2013 Mathematical modeling of an oscillating gene circuit to unravel the circadian clock network of *Arabidopsis thaliana*. *Front. Plant Sci.* 4: 3. <https://doi.org/10.3389/fpls.2013.00003>
- Cao, D., J. E. Froehlich, H. Zhang, and C. L. Cheng, 2003 The chlorate-resistant and photomorphogenesis-defective mutant *cr88* encodes a chloroplast-targeted HSP90. *Plant J.* 33: 107–118. <https://doi.org/10.1046/j.1365-313X.2003.016011.x>
- Cha, J. Y., J. Kim, T. S. Kim, Q. Zeng, L. Wang *et al.*, 2017 GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE in the *Arabidopsis* circadian clock. *Nat. Commun.* 8: 3. <https://doi.org/10.1038/s41467-016-0014-9>
- Chen, B., W. H. Piel, L. Gui, E. Bruford, and A. Monteiro, 2005 The HSP90 family of genes in the human genome: insights into their divergence and evolution. *Genomics* 86: 627–637. <https://doi.org/10.1016/j.ygeno.2005.08.012>
- Covington, M. F., and S. L. Harmer, 2007 The circadian clock regulates auxin signaling and responses in *Arabidopsis*. *PLoS Biol.* 5: e222. <https://doi.org/10.1371/journal.pbio.0050222>
- Covington, M. F., S. Panda, X. L. Liu, C. A. Strayer, D. R. Wagner *et al.*, 2001 ELF3 modulates resetting of the circadian clock in *Arabidopsis*. *Plant Cell* 13: 1305–1316. <https://doi.org/10.1105/tpc.13.6.1305>
- Covington, M. F., J. N. Maloof, M. Straume, S. A. Kay, and S. L. Harmer, 2008 Global transcriptome analysis reveals circadian regulation of key pathways in plant growth and development. *Genome Biol.* 9: R130. <https://doi.org/10.1186/gb-2008-9-8-r130>
- Dalchau, N., S. J. Baek, H. M. Briggs, F. C. Robertson, A. N. Dodd *et al.*, 2011 The circadian oscillator gene GIGANTEA mediates a long-term response of the *Arabidopsis thaliana* circadian clock to sucrose. *Proc. Natl. Acad. Sci. USA* 108: 5104–5109. <https://doi.org/10.1073/pnas.1015452108>
- Ding, Z., M. R. Doyle, R. M. Amasino, and S. J. Davis, 2007 A complex genetic interaction between *Arabidopsis thaliana* TOC1 and *CCA1/LHY* in driving the circadian clock and in output regulation. *Genetics* 176: 1501–1510. <https://doi.org/10.1534/genetics.107.072769>
- Doyle, M. R., S. J. Davis, R. M. Bastow, H. G. McWatters, L. Kozma-Bognár *et al.*, 2002 The ELF4 gene controls circadian rhythms and flowering time in *Arabidopsis thaliana*. *Nature* 419: 74–77. <https://doi.org/10.1038/nature00954>
- Farré, E. M., S. L. Harmer, F. G. Harmon, M. J. Yanovsky, and S. A. Kay, 2005 Overlapping and distinct roles of PRR7 and PRR9 in the *Arabidopsis* circadian clock. *Curr. Biol.* 15: 47–54. <https://doi.org/10.1016/j.cub.2004.12.067>
- Fowler, S., K. Lee, H. Onouchi, A. Samach, K. Richardson *et al.*, 1999 GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J.* 18: 4679–4688. <https://doi.org/10.1093/emboj/18.17.4679>
- Gendron, J. M., J. L. Pruneda-Paz, C. J. Doherty, A. M. Gross, S. E. Kang *et al.*, 2012 *Arabidopsis* circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proc. Natl. Acad. Sci. USA* 109: 3167–3172. <https://doi.org/10.1073/pnas.1200355109>
- Gil, K. E., W. Y. Kim, H. J. Lee, M. Faisal, Q. Saquib *et al.*, 2017 ZEITLUPE contributes to a thermoresponsive protein quality control system in *Arabidopsis*. *Plant Cell* 29: 2882–2894. <https://doi.org/10.1105/tpc.17.00612>
- Gould, P. D., J. C. W. Locke, C. Larue, M. M. Southern, S. J. Davis *et al.*, 2006 The molecular basis of temperature compensation in the *Arabidopsis* circadian clock. *Plant Cell* 18: 1177–1187. <https://doi.org/10.1105/tpc.105.039990>
- Hanano, S., M. A. Domagalska, F. Nagy, and S. J. Davis, 2006 Multiple phytohormones influence distinct parameters of the plant circadian clock. *Genes Cells* 11: 1381–1392. <https://doi.org/10.1111/j.1365-2443.2006.01026.x>
- Hanano, S., R. Stracke, M. Jakoby, T. Merkle, M. A. Domagalska *et al.*, 2008 A systematic survey in *Arabidopsis thaliana* of transcription factors that modulate circadian parameters. *BMC Genomics* 9: 182. <https://doi.org/10.1186/1471-2164-9-182>
- Haydon, M. J., O. Mielczarek, F. C. Robertson, K. E. Hubbard, and A. A. R. Webb, 2013 Photosynthetic entrainment of the *Arabidopsis thaliana* circadian clock. *Nature* 502: 689–692. <https://doi.org/10.1038/nature12603>
- Herrero, E., E. Kolmos, N. Bujdoso, Y. Yuan, M. Wang *et al.*, 2012 EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the *Arabidopsis* circadian clock. *Plant Cell* 24: 428–443. <https://doi.org/10.1105/tpc.111.093807>
- Hubert, D. A., P. Tornero, Y. Belkhadir, P. Krishna, A. Takahashi *et al.*, 2003 Cytosolic HSP90 associates with and modulates the *Arabidopsis* RPM1 disease resistance protein. *EMBO J.* 22: 5679–5689. <https://doi.org/10.1093/emboj/cdg547>
- Kadota, Y., and K. Shirasu, 2012 The HSP90 complex of plants. *Biochim. Biophys. Acta* 1823: 689–697. <https://doi.org/10.1016/j.bbamcr.2011.09.016>

- Kevei, É., P. Gyula, A. Hall, L. Kozma-Bognár, W.-Y. Kim *et al.*, 2006 Forward genetic analysis of the circadian clock separates the multiple functions of ZEITLUPE. *Plant Physiol.* 140: 933–945. <https://doi.org/10.1104/pp.105.074864>
- Kim, T. S., W. Y. Kim, S. Fujiwara, J. Kim, J. Y. Cha *et al.*, 2011 HSP90 functions in the circadian clock through stabilization of the client F-box protein ZEITLUPE. *Proc. Natl. Acad. Sci. USA* 108: 16843–16848. <https://doi.org/10.1073/pnas.1110406108>
- Kim, W. Y., S. Fujiwara, S. S. Suh, J. Kim, Y. Kim *et al.*, 2007 ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature* 449: 356–360. <https://doi.org/10.1038/nature06132>
- Kolmos, E., M. Nowak, M. Werner, K. Fischer, G. Schwarz *et al.*, 2009 Integrating ELF4 into the circadian system through combined structural and functional studies. *HFSP J.* 3: 350–366. <https://doi.org/10.2976/1.3218766>
- Köster, T., K. Meyer, C. Weinholdt, L. M. Smith, M. Lummer *et al.*, 2014 Regulation of pri-miRNA processing by the hnRNP-like protein at GRP7 in Arabidopsis. *Nucleic Acids Res.* 42: 9925–9936. <https://doi.org/10.1093/nar/gku716>
- Krishna, P., and G. Gloor, 2001 The Hsp90 family of proteins in Arabidopsis thaliana. *Cell Stress Chaperones* 6: 238–246. [https://doi.org/10.1379/1466-1268\(2001\)006<0238:THFOPI>2.0.CO;2](https://doi.org/10.1379/1466-1268(2001)006<0238:THFOPI>2.0.CO;2)
- Lin, Y., and C. L. Cheng, 1997 A chlorate-resistant mutant defective in the regulation of nitrate reductase gene expression in Arabidopsis defines a new HY locus. *Plant Cell* 9: 21–35. <https://doi.org/10.1105/tpc.9.1.21>
- Mizoguchi, T., K. Wheatley, Y. Hanzawa, L. Wright, M. Mizoguchi *et al.*, 2002 LHY and CCA1 are partially redundant genes required to maintain circadian rhythms in Arabidopsis. *Dev. Cell* 2: 629–641. [https://doi.org/10.1016/S1534-5807\(02\)00170-3](https://doi.org/10.1016/S1534-5807(02)00170-3)
- Nagel, D. H., C. J. Doherty, J. L. Pruneda-Paz, R. J. Schmitz, J. R. Ecker *et al.*, 2015 Genome-wide identification of CCA1 targets uncovers an expanded clock network in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 112: E4802–E4810. <https://doi.org/10.1073/pnas.1513609112>
- Nakamichi, N., T. Kiba, R. Henriques, T. Mizuno, N.-H. Chua *et al.*, 2010 PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in the Arabidopsis circadian clock. *Plant Cell* 22: 594–605. <https://doi.org/10.1105/tpc.109.072892>
- Norén, L., P. Kindgren, P. Stachula, M. Rühl, M. Eriksson *et al.*, 2016 HSP90, ZTL, PRR5 and HY5 integrate circadian and plastid signaling pathways to regulate CBF and COR expression. *Plant Physiol.* 171: 1392–1406. <https://doi.org/10.1104/pp.16.00374>
- Nusinow, D. A., A. Helfer, E. E. Hamilton, J. J. King, T. Imaizumi *et al.*, 2011 The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475: 398–402. <https://doi.org/10.1038/nature10182>
- Oakenfull, R. J., and S. J. Davis, 2017 Shining a light on the Arabidopsis circadian clock. *Plant Cell Environ.* 40: 2571–2585. <https://doi.org/10.1111/pce.13033>
- O'Neill, J. S., G. van Ooijen, L. E. Dixon, C. Troein, F. Corellou *et al.*, 2011 Circadian rhythms persist without transcription in a eukaryote. *Nature* 469: 554–558. <https://doi.org/10.1038/nature09654>
- Ronald, J., and S. Davis, 2017 Making the clock tick: the transcriptional landscape of the plant circadian clock. *F1000Res.* 6: 951. <https://doi.org/10.12688/f1000research.11319.1>
- Sangster, T. A., A. Bahrami, A. Wilczek, E. Watanabe, K. Schellenberg *et al.*, 2007 Phenotypic diversity and altered environmental plasticity in Arabidopsis thaliana with reduced Hsp90 levels. *PLoS One* 2: e648. <https://doi.org/10.1371/journal.pone.0000648>
- Schöning, J. C., and D. Staiger, 2005 At the pulse of time: protein interactions determine the pace of circadian clocks. *FEBS Lett.* 579: 3246–3252. <https://doi.org/10.1016/j.febslet.2005.03.028>
- Southern, M. M., and A. J. Millar, 2005 Circadian genetics in the model higher plant, Arabidopsis thaliana. *Methods Enzymol.* 393: 23–35. [https://doi.org/10.1016/S0076-6879\(05\)93002-4](https://doi.org/10.1016/S0076-6879(05)93002-4)
- Staiger, D., J. Shin, M. Johansson, and S. J. Davis, 2013 The circadian clock goes genomic. *Genome Biol.* 14: 208. <https://doi.org/10.1186/gb-2013-14-6-208>
- Thines, B., and F. G. Harmon, 2010 Ambient temperature response establishes ELF3 as a required component of the core Arabidopsis circadian clock. *Proc. Natl. Acad. Sci. USA* 107: 3257–3262. <https://doi.org/10.1073/pnas.0911006107>
- Wang, L., J. Kim, and D. E. Somers, 2013 Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc. Natl. Acad. Sci. USA* 110: 761–766. <https://doi.org/10.1073/pnas.1215010110>

Communicating editor: Y. Xue