Abnormal growth

Rita Slim¹ , Noel Aoun²

¹Department of Gastroenterology, Saint Joseph University of Beirut, Hotel-Dieu de France Hospital, Beirut, Lebanon

²Department of Radiology, Saint Joseph University of Beirut, Hotel-Dieu de France Hospital, Beirut, Lebanon

Cite this article as: Slim R, Aoun N. Abnormal growth. *Turk J Gastroenterol* 2018; 29: 509-10.

QUESTION

A 20-year-old man with a one-year history of severe ulcerative pancolitis had been receiving azathioprine for 8 months with low-dose prednisone to maintain clinical remission. The last 2 months were marked with episodes of abdominal pain and vomiting, and he subsequently presented to the emergency department for exacerbation

of abdominal pain with incoercible vomiting. Complete blood tests with CRP levels were within normal ranges. An abdominal and pelvic CT scan was then performed (Figure 1, 2).

What does-it show?

Figure 1. CT image of circumferential colonic thickening in axial plane

Figure 2. CT image of circumferential colonic thickening with irregular mucosal contours involving the splenic flexure and the descending colon in coronal plane

ORCID IDs of the authors: R.S. 0000-0001-5087-7407; N.A. 0000-0003-2907-1956.

Corresponding Author: **Slim Rita**; rslimkaram@hotmail.com

Received: **February 16, 2018** Accepted: **February 27, 2018** Available online date: **June 26, 2018**

© Copyright 2018 by The Turkish Society of Gastroenterology · Available online at www.turkjgastroenterol.org

DOI: **10.5152/tjg.2018.18130**

ANSWER

The CT scan revealed bowel obstruction and a circumferential colonic thickening with irregular mucosal contours involving the splenic flexure and the descending colon. On endoscopy, the rectum appeared normal, but the endoscope could not be inserted beyond the sigmoid colon because of the presence of a polypoid lesion. A subtotal colectomy was then performed with preservation of the

Figure 3. Thickened wall of the colon on macroscopic view

Figure 4. Cluster of pseudopolyps

rectum. Macroscopic examination of the excised portion showed a 2-cm thickening of the wall of the excised colon (50 cm) (Figure 3), with a continuous cluster of pseudopolyps all over the mucosa (Figure 4), reducing the colic lumen. Microscopic examination confirmed the diagnosis of obstructing postinflammatory pseudopolyps, composed of a layer of hyperplastic glandular epithelium, also including the mucosa muscularis. No dysplastic lesions were noted. The rest of the colon showed stigmata of quiescent colitis.

Pseudopolyps are a frequent finding in inflammatory bowel disease (IBD) accounting for approximately 10% to 20% of all cases (1). It is believed that they are due to hyperplastic healing after intense inflammatory flare. They can appear as early as 1 month after the diagnosis of the IBD and as late as many years after. The presence of numerous, relatively large polyps coalescing in a localized segment of the intestine are described as a giant inflammatory polyposis. This inflammatory polyposis can be misdiagnosed as tumor-like proliferation and be revealed by the development of intestinal obstruction or sub-obstruction, as in our case (2).

Informed Consent: Informed Consent: Informed consent was obtained from the patient who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - R.S.; Design - R.S.; Supervision - R.S.; Analysis and/or Interpretation - N.A.; Literature Search - R.S., N.A.; Writing Manuscript - R.S.; Critical Review - R.S., N.A.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Politis DS, Katsanos KH, Tsianos EV, Christodoulou DK. Pseudopolyps in inflammatory bowel diseases: Have we learned enough? *World J Gastroenterol* 2017; 23: 1541-51. [\[CrossRef\]](#)
2. Maggs JR, Browning LC, Warren BF, Travis SP. Obstructing giant post-inflammatory polyposis in ulcerative colitis: Case report and review of the literature. *J Crohns Colitis* 2008; 2: 170-80 [\[CrossRef\]](#)