Published in final edited form as:

AME Med J. 2018 October ; 3: . doi:10.21037/amj.2018.10.08.

Targeting ELK1: a wELKome addition to the prostate cancer armamentarium

Jelani C. Zarif^{1,2} and Emmanuel S. Antonarakis^{1,2}

¹Department of Oncology, The Sidney Kimmel Comprehensive Cancer Center

²The Bloomberg-Kimmel Institute for Cancer Immunotherapy, Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA

The androgen receptor (AR) is a critical steroid receptor required for prostatic luminal cell survival. In its unbound state, AR is initially localized to the cytoplasm and has been demonstrated to interact with a host of scaffold proteins and kinases such as c-Src (1) in a non-genomic fashion (2). Once bound by its natural ligand, dihydrotestosterone (DHT), AR forms a homo-dimer with itself, translocates into the nucleus, recruits transcriptional coactivators, and binds to DNA androgen-response elements (AREs) positively regulating prostate-specific antigen (PSA) and a variety of other AR target genes that control cellular division and survival (3). In prostate cancer, AR continues to remain a central target in both localized and castration-resistant disease states. Suppression of testosterone synthesis or inhibition of AR signaling using pharmacological modalities represents an effective approach for the treatment of hormone-sensitive and castration-resistant disease because tumor cells are still dependent on AR for their growth and survival. However, most men receiving these therapies will eventually become resistant to such hormonal manipulations, which is heralded by an increase in PSA level or a clinical worsening of metastases while on these therapies. This can be ascribed to a range of resistance mechanisms including activating mutations in the AR ligand-binding domain, hormonal independence via generation of mRNA splice variants of the AR (notably AR-V7), as well as induction of alternative steroid receptors such as the glucocorticoid and progesterone receptors which hijack promoters/enhancers of AR-responsive genes (3,4).

In the paper that accompanies this editorial, Rosati *et al.* 2018 (5) developed and tested a flavonoid agent that successfully disrupted AR signaling through an epigenetic mechanism involving ERK1. Using LNCaP, VCaP and 22RV1 castration-resistant prostate cancer cell lines, and by performing chromatin immunoprecipitation (ChIP) and competitive ligand-

Correspondence to: Emmanuel S. Antonarakis, MD. The Sidney Kimmel Comprehensive Cancer Center, Skip Viragh Building, 201 N. Broadway, Room 9129, Baltimore, MD 21287, USA., eantona1@jhmi.edu.

Conflicts of Interest: ES Antonarakis is a paid consultant/ advisor to Janssen, Astellas, Sanofi, Dendreon, Medivation, ESSA, AstraZeneca, Clovis, and Merck; he has received research funding to his institution from Janssen, Johnson & Johnson, Sanofi, Dendreon, Genentech, Novartis, Tokai, Bristol Myers- Squibb, AstraZeneca, Clovis, and Merck; and he is the co-inventor of a biomarker technology that has been licensed to Qiagen. JC Zarif declares no potential conflicts of interest.

Provenance: This is an invited Editorial commissioned by Section Editor Xiao Li (Department of Urologic Surgery, The Affiliated Cancer Hospital of Jiangsu Province of Nanjing Medical University, Nanjing, China).

Comment on: Rosati R, Polin L, Ducker C, et al. Strategy for Tumor-Selective Disruption of Androgen Receptor Function in the Spectrum of Prostate Cancer. Clin Cancer Res 2018. [Epub ahead of print].

binding assays, the authors were able to demonstrate that a lead small-molecule compound, KCI807, had the ability to bind directly to AR and block the recruitment of an ETS-family transcriptional coactivator, ELK1. AR/ELK1 complexes bound to specific sites of DNA enhancer/promoter regions, and KCI807 was able to reduce mRNA expression levels of downstream target genes regulated by the AR/ELK1 transcriptional complex. Importantly, KCI807 did not appear to impede the recruitment of AR to canonical ARE sites, for example those associated with expression of *PSA* or *TMPRSS2*. Knockdown experiments using siRNAs against ELK1 and AR further potentiated the action of KCI807; this resulted in increased levels of Fam112b relative to the ELK1 and AR knockdown experiments. Notably, in 22RV1 cells that harbor the AR-V7 splice variant, KCI807 was also effective at inhibiting promoter activation and transcription of AR-V7 target genes in these cells. Intriguingly, however, KCI807 did not suppress the level of protein expression of the wild-type AR or the AR-V7 splice variant. Finally, the authors also tested their agent at a concentration of 250 mg/kg and compared it against enzalutamide in an *in vivo* 22RV1 xenograft mouse model, and demonstrated the ability of KCI807 to delay tumorigenesis more so than enzalutamide *in vivo*. Altogether, the authors performed well-controlled experiments to evaluate KCI807 binding to AR, its effects on AR/ELK1-regulated genes, inhibition of cell proliferation, and suppression of tumorigenesis *in vivo*.

What are the potential clinical implications of this work? First, the development of therapeutic strategies capable of targeting AR-V7-expressing human castration-resistant prostate cancer (CRPC) represents one of the holy grails of advanced prostate cancer treatment. With the exception of taxane chemotherapies (6) and perhaps immune checkpoint blockade therapy (7), attempts to target AR-V7-positive prostate cancers have been discouraging (8,9) and this continues to represent an unmet medical need. To this end, if KCI807 (or its derivative) is capable of extinguishing AR-regulated gene expression at both the AR/ELK1 and the AR-V7/ELK1 transcriptional levels, then this strategy should theoretically be effective in many human CRPC tumors that usually co-express both wild-type and aberrant AR isoforms. Such an approach could either be used as a monotherapy in patients who have previously failed abiraterone and/or enzalutamide, or could be developed as a combination strategy together with abiraterone and/or enzalutamide in CRPC patients who have not yet received either agent. The combinatorial approach may serve to prolong the duration of responses to abiraterone/ enzalutamide, or to subvert secondary (i.e., acquired) resistance that invariably develops in most patients.

A second potential clinical utility of KCI807 (or its derivatives) would be for the treatment of neuroendocrine prostate cancer which is a rapidly lethal phenotype of CRPC that is usually refractory to most forms of androgen/ AR ablation, and for which novel therapeutic strategies are sorely needed (10). Interestingly, *ELK1* gene amplification and/or mRNA overexpression occurs in about 25–30% of human neuroendocrine CRPC cases (11) compared to <5% of conventional acinar CRPC cases, while the related ETS-family genes *ELK3* and *ELK4* are also amplified in many neuroendocrine prostate cancers. These findings might theoretically imply that disruption of the AR/ELK1 transcriptional axis in neuroendocrine prostate cancers might lead to effective therapies for a least a subset of these cases. This hypothesis could be further tested pre-clinically by administering KCI807 to mice harboring patient-derived xenografts (12) consisting of neuroendocrine CRPC,

especially those demonstrating amplification or high expression of *ELK1*. Finally, although only hypothesis-generating, it would be attractive to also explore potential synergy between ELK1 modulators and platinum agents in neuroendocrine prostate cancers, since the latter is currently being used commonly for the clinical management of neuroendocrine CRPC.

Acknowledgements

Funding: This work was partially supported by National Institutes of Health Cancer Center Support Grant P30 CA006973 (ES Antonarakis), Prostate Cancer Foundation Young Investigator award (JC Zarif), Maryland Cigarette Restitution Fund grant FHB33CRF (JC Zarif), The Patrick C. Walsh Prostate Cancer Research Fund (JC Zarif), and Department of Defense grant W81XWH-16-PCR-CCRS (ES Antonarakis).

References

1. Zarif JC, Lamb LE, Schulz VV, et al. Androgen receptor non-nuclear regulation of prostate cancer cell invasion mediated by Src and matriptase. *Oncotarget* 2015;6:6862–76. [PubMed: 25730905]
2. Zarif JC, Miranti CK. The importance of non-nuclear AR signaling in prostate cancer progression and therapeutic resistance. *Cell Signal* 2016;28:348–56. [PubMed: 26829214]
3. Nelson PS. Molecular states underlying androgen receptor activation: a framework for therapeutics targeting androgen signaling in prostate cancer. *J Clin Oncol* 2012;30:644–6. [PubMed: 22184375]
4. Antonarakis ES, Lu C, Wang H, et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N Engl J Med* 2014;371:1028–38. [PubMed: 25184630]
5. Rosati R, Polin L, Ducker C, et al. Strategy for Tumor-Selective Disruption of Androgen Receptor Function in the Spectrum of Prostate Cancer. *Clin Cancer Res* 2018 [Epub ahead of print].
6. Antonarakis ES, Lu C, Lubner B, et al. Androgen Receptor Splice Variant 7 and Efficacy of Taxane Chemotherapy in Patients With Metastatic Castration-Resistant Prostate Cancer. *JAMA Oncol* 2015;1:582–91. [PubMed: 26181238]
7. Boudadi K, Suzman DL, Anagnostou V, et al. Ipilimumab plus nivolumab and DNA-repair defects in AR-V7-expressing metastatic prostate cancer. *Oncotarget* 2018;9:28561–71. [PubMed: 29983880]
8. Antonarakis ES, Armstrong AJ, Dehm SM, et al. Androgen receptor variant-driven prostate cancer: clinical implications and therapeutic targeting. *Prostate Cancer Prostatic Dis* 2016;19:231–41. [PubMed: 27184811]
9. Taplin ME, Antonarakis ES, Ferrante KJ, et al. Clinical factors associated with AR-V7 detection in ARMOR3-SV, a randomized trial of galeterone (Gal) vs enzalutamide (Enz) in men with AR-V7+ metastatic castration-resistant prostate cancer. *J Clin Oncol* 2017;35:abstract 5005.
10. Rickman DS, Beltran H, Demichelis F, et al. Biology and evolution of poorly differentiated neuroendocrine tumors. *Nat Med* 2017;23:1–10.
11. Beltran H, Prandi D, Mosquera JM, et al. Divergent clonal evolution of castration-resistant neuroendocrine prostate cancer. *Nat Med* 2016;22:298–305. [PubMed: 26855148]
12. Navone NM, van Weerden WM, Vessella RL, et al. Movember GAP1 PDX project: An international collection of serially transplantable prostate cancer patient-derived xenograft (PDX) models. *Prostate* 2018 [Epub ahead of print].