Published in final edited form as:

Genesis. 2018 October ; 56(10): e23256. doi:10.1002/dvg.23256.

Generation of *MLC-2v-tdTomato* knock-in reporter mouse line

Zhentao Zhang^{1,2,3} and Young-Jae Nam^{1,2,3,*}

¹Department of Medicine, Division of Cardiovascular Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

²Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN, USA

³Vanderbilt Center for Stem Cell Biology, Vanderbilt University, Nashville, TN, USA

Abstract

MLC-2v is a myosin light chain regulatory protein which is specifically expressed in ventricular cardiomyocytes and slow twitch skeletal muscle cells. MLC-2v plays critical roles in ventricular maturation during heart development. Mice lacking *MLC-2v* are embryonic lethal due to heart failure associated with abnormal myofibrillar organization of ventricular cardiomyocytes. To study the development of ventricular cardiac muscle and slow twitch skeletal muscle, we generated a new *MLC-2v* reporter mouse line by knocking-in a *tdTomato* reporter cassette into 3' UTR of the *MLC-2v* gene without disrupting the endogenous gene. Our results demonstrated specific *MLC-2v-tdTomato* knock-in reporter expression in ventricular cardiomyocytes and slow twitch muscle during myogenesis, precisely recapitulating the spatiotemporal expression pattern of endogenous MLC-2v. No tdTomato expression was observed in the atria, fast twitch muscle or other organs throughout development into adulthood. Isolated neonatal and adult ventricular cardiomyocytes uniformly express tdTomato. Taken together, *MLC-2v-tdTomato* knock-in reporter mouse model described in this article will serve as a valuable tool to study cardiac chamber and skeletal muscle specification during development and regeneration by overcoming the pitfalls of transgenic strategies.

Keywords

MLC-2v; Heart; Ventricle; Muscle

INTRODUCTION

Striated muscle cells are defined by the presence of repeating contractile units of the sarcomere, which is composed of multiple sarcomeric proteins. Striated muscle is classified into cardiac (i.e. ventricular and atrial) and skeletal (i.e. slow twitch and fast twitch) muscles, based on its distinct anatomical location, physiological function, and molecular and

*To whom correspondence may be addressed: Young-Jae Nam, M.D., Ph.D., Department of Medicine, Division of Cardiovascular Medicine, Vanderbilt University Medical Center, Nashville, Tennessee 37232, USA, Phone: 615-936-5436, young-jae.nam@vanderbilt.edu.

CONFLICT OF INTEREST

The authors have no competing financial interests.

electrophysiological properties. For example, atrial cardiomyocytes which are exclusively located in the atria of the heart display a triangular shape action potential with a relatively short duration. By contrast, ventricular cardiomyocytes restricted to the ventricle of the heart exhibit an action potential with prominent plateau and longer duration, thereby generating greater contractile force (Spater, Hansson, Zangi, & Chien, 2014). Compared to fast twitch skeletal muscle cells, slow twitch skeletal muscle cells are more resistant to fatigue with their larger mitochondrial content and enriched oxidative enzymes, and characterized by a slower contraction-relaxation response (Schiaffino & Reggiani, 2011). Regardless of a classified type, striated muscle cells uniformly contain very similar molecular components and structures of the sarcomere. However, a distinct combinatorial expression of multiple isoforms of sarcomeric proteins allows each type of striated muscle to harness its unique physiological properties and to be distinguished from other types (Schiaffino & Reggiani, 1996).

Myosin light chain-2 (MLC-2 or MYL2) is a major component of the sarcomere in striated muscle cells. MLC-2 consists of three isoforms, MLC-2a, MLC-2f, and MLC-2v (Sheikh, Lyon, & Chen, 2015). Despite their high degree of homology, each MLC-2 isoform displays a distinct expression pattern during striated muscle development. MLC-2f is specifically expressed in fast twitch skeletal muscles (Nudel, Calvo, Shani, & Levy, 1984). MLC-2a expression is restricted to atrial cardiac muscle in the adult heart (Chen, Kubalak, Minamisawa, et al., 1998; Kubalak, Miller-Hance, O'Brien, Dyson, & Chien, 1994). MLC-2a mRNA starts to express in the entire heart tube at E7.5 during heart development. Its expression is selectively decreased in the ventricles at E11, while its atrial expression persists (Kubalak et al., 1994). At E13.5, MLC-2a mRNA expression becomes restricted to the atria. On the other hand, *MLC-2v* expression pattern is unique in that *MLC-2v* is expressed in both ventricular cardiac muscle and slow twitch skeletal muscle (Henderson, Xu, & Chien, 1988; Kumar, Cribbs, Delaney, Chien, & Siddiqui, 1986). *MLC-2v* is considered one of the earliest markers of ventricular chamber specification during heart development. MLC-2v mRNA is expressed in the ventricular portion of the heart tube at as early as E8.0 without detectable expression in the atrial or sinus venosus regions (O'Brien, Lee, & Chien, 1993). At E11.0, MLC-2v mRNA expression becomes restricted to the ventricular region and remains confined to the ventricular chamber throughout the rest of embryonic development into adulthood.

Multiple transgenic mice harboring a rat *MLC-2v* promoter fragment fused to a luciferase or *LacZ* reporter gene were generated (Lee et al., 1992; Ross, Navankasattusas, Harvey, & Chien, 1996). *MLC-2v* promoter driven *LacZ* activity became detectable as early as E7–7.5. From this linear heart tube stage during embryogenesis, the *LacZ* reporter expression remained within the heart and confined to the ventricular chamber. This ventricular restricted expression of the reporter continues throughout the embryonic development into adulthood. However, this reporter does not precisely represent endogenous expression of MLC-2v. While endogenous MLC-2v is uniformly expressed throughout both ventricles (Sheikh et al., 2015), the transgene expression was right ventricle dominant. Even in the right ventricle, only subset of cardiomyocytes expressed the transgene. These results suggest that the *MLC-2v cis*-regulatory element harbored in this mouse line contains only a part of the genetic elements responsible for chamber specification during heart development (Ross et

al., 1996). In addition, the LacZ reporter expression was not observed in slow twitch skeletal muscles in this mouse line. Taken together, the reporter driven by *MLC-2v cis*-regulatory element does not precisely recapitulate endogenous MLC-2v expression.

To study the biological and physiological roles of *MLC-2v* during development, *MLC-2v* null (*MLC-2v*^{-/-}) mice were generated by Cre-loxP recombination (Chen, Kubalak, Minamisawa, et al., 1998). Cre-recombinase coding sequences were inserted into the endogenous *MLC-2v* genomic locus by disrupting the exon 2 of the *MLC-2v* gene. Cre-recombinase activity was first shown in the heart at E8.75 (Chen, Kubalak, & Chien, 1998). *MLC-2v*^{-/-} mice were embryonic lethal at E12.5 (Chen, Kubalak, Minamisawa, et al., 1998). Although MLC-2a comparable to the level of MLC-2v protein in wild type mice was induced in the ventricles of *MLC-2v*^{-/-} embryos, it could not substitute for MLC-2v. *MLC-2v*^{-/-} embryos at E10.5-E12.5 demonstrated features of heart failure (i.e. enlarged cardiac chambers, thin ventricular wall, pleural effusion, and hepatic congestion) associated with abnormal sarcomere assembly in ventricular cardiomyocytes. These results indicate that MLC-2v plays irreplaceable roles in ventricular maturation during heart development.

In this study, we have generated a new *MLC-2v* reporter mouse line by knocking-in a *tdTomato* reporter cassette into 3' UTR of the *MLC-2v* gene. Although multiple transgenic mouse lines based on *cis*-regulatory sequences of the *MLC-2v* gene have been generated, the transgene expression in those mouse lines differs from endogenous MLC-2v expression (i.e. patchy and right ventricular dominant expression in the ventricles, and absence of slow twitch muscle expression). In contrast, *tdTomato* knock-in reporter expression in our new *MLC-2v* reporter mouse line faithfully represents endogenous MLC-2v expression. The reporter is uniformly expressed in both ventricles and selectively expressed in slow twitch muscles. Therefore, *MLC-2v-tdTomato* knock-in reporter mouse line that we generated will be a valuable new tool for studying cardiac and skeletal muscle specification and regeneration.

RESULTS AND DISCUSSION

Generation of *MLC-2v-tdTomato* knock-in reporter mice

We generated *MLC-2v-tdTomato* knock-in reporter mouse line by knocking-in *IRES-tdTomato-Neo^R* (Neomycin resistance) gene cassette into the 3'UTR immediately following the seventh exon of the *MLC-2v* gene. We designed a targeting construct containing an *IRES-tdTomato-FRT* (flippase recognition target)-*Neo^R-FRT* knock-in cassette flanked by homology arms as illustrated in Figure 1A. The linearized targeting construct was electroporated into embryonic stem cells (ESCs) of C57BL/6 mice. Neomycin resistant ESC clones were expanded and screened by Polymerase chain reaction (PCR). Homologous recombination at 3' was verified by amplifying 3.5 kb PCR product (Figure 1B). Then, homologous recombination of the PCR verified ESC clones was confirmed by Southern blotting (Figure 1C). The correctly targeted ESC clones were injected into blastocysts of C57BL/6 mice to generate mouse chimeras. By cross breeding the chimeras with FLP (flippase) transgenic mice, *Neo^R* cassette was removed to create the *MLC-2v-IRES-tdTomato* knock-in allele by FLP-FRT recombination (Figure 1A). We did not observe any

abnormality in viability and fertility in the generated heterozygote or homozygote *MLC-2v-tdTomato* knock-in reporter mice.

Characterization of *MLC-2v-tdTomato* knock-in reporter expression during development

We assessed the *MLC-2v-tdTomato* knock-in reporter expression during development. The reporter expression was first observed in the linear heart tube at E8.0 (Figure 2A). tdTomato expression was restricted to the ventricular portion of the forming heart (Figure 2A–C). It was not shown in either out flow tract or inflow tract (Figure 2G). At E10.5, tdTomato expression remained restricted to the ventricle of the embryonic heart (Figure 2D and 2H). Similar ventricular specific expression pattern of *MLC-2v-tdTomato* knock-in reporter was observed at E13.5 and E16.5 (Figure 2E, 2F, 2I, and 2J). The atria did not express tdTomato. Other areas outside the heart in E13.5 and E16.5 embryo did not show tdTomato expression (Figure 2E and 2F). Using immunostaining of frozen heart sections of the mouse embryos, we confirmed that *MLC-2v-tdTomato* knock-in reporter was exclusively expressed in the ventricles within the heart, while pan-cardiac marker α -actinin was expressed in both the atria and ventricles (Figure 3). At E10.5, only the heart in the entire chest area showed α -actinin and tdTomato expression, while other areas did not show any α -actinin or tdTomato expression (Figure 3A). tdTomato was expressed throughout the entire ventricle at E10.5, E13.5, and E16.5 (Figure 3B–3D). Both right and left ventricles uniformly expressed *tdTomato* knock-in reporter at E13.5 and E16.5, although the previous *MLC-2v-LacZ* transgenic reporter mouse line showed right ventricle dominant expression pattern (Ross et al., 1996) (Figure 3B and 3C). Every cardiomyocyte in the ventricle expressing α -actinin expressed tdTomato in the knock-in reporter mice (Figure 3E–3G). The symmetric and homogeneous ventricular expression pattern of *MLC-2v-tdTomato* knock-in reporter is consistent with endogenous MLC-2v expression (Sheikh et al., 2015).

At post-natal stages, *MLC-2v-tdTomato* knock-in reporter expression remained to be specific to the ventricles of the heart (Figure 4A). We examined if inserting *tdTomato* into the endogenous locus of *MLC-2v* alters the endogenous expression of MLC-2v by comparing the endogenous MLC-2v expression in a wild type littermate with the heterozygous or homozygous *MLC-2v-tdTomato* knock-in reporter mouse heart. There was no significant difference in the endogenous MLC-2v protein expression among those three distinct genotypes of mice, while each mouse showed different expression levels of tdTomato (Figure 4B). These results indicate that the endogenous *MLC-2v* gene expression is not affected by knocking-in *tdTomato* into the endogenous locus of *MLC-2v*. The adult homozygous mouse heart clearly demonstrated strong tdTomato expression in every ventricular cardiomyocyte expressing the endogenous MLC-2v (Figure 4C). Furthermore, we isolated neonatal ventricular cardiomyocytes at P1 and showed that tdTomato expression overlaps with endogenous MLC-2v expression using immunostaining (Figure 4D, left panel). We also isolated adult cardiomyocytes from *MLC-2v-tdTomato* knock-in reporter mice using Langendorff perfusion and demonstrated tdTomato expression in adult cardiomyocytes in a single cell level (Figure 4D, right panel).

We examined *tdTomato* knock-in reporter expression in skeletal muscles of *MLC-2v-tdTomato* knock-in reporter mice (Figure 5). Although the endogenous MLC-2v was

expressed in slow twitch muscles, the previous *MLC-2v* promoter fragment driven reporters did not express in slow twitch muscles (Lee et al., 1992; Ross et al., 1996). We found *MLC-2v-tdTomato* knock-in reporter expression in slow twitch muscle (soleus), but not in fast twitch muscle (extensor digitorum longus), recapitulating the expression pattern of endogenous *MLC-2v* in skeletal muscles (Lee et al., 1992). The *MLC-2v-tdTomato* knock-in reporter expression in slow twitch muscles first appeared at P0, and persisted throughout post-natal muscle development (Figure 5A). The reporter expression was significantly increased in the adult muscle, compared to the muscle at P0 or P1. We also demonstrated that *MLC-2v-tdTomato* knock-in reporter expression completely overlaps with slow twitch muscle-specific slow muscle myosin expression in slow twitch muscles, while it was not observed in fast twitch muscles expressing a fast twitch muscle marker, fast skeletal myosin (Figure 5B). Taken together, our study demonstrated temporal expression pattern of *MLC-2v* in slow twitch muscle during development.

In summary, we generated a new *MLC-2v-tdTomato* knock-in reporter mouse line without disrupting the endogenous *MLC-2v* gene, and characterized spatiotemporal expression of *MLC-2v-tdTomato* knock-in reporter during myogenesis. Its expression uniformly persists in both right and left ventricles throughout heart development. In skeletal muscle, *MLC-2v-tdTomato* knock-in reporter is selectively expressed in slow twitch muscle during post-natal development. Our results indicate that *MLC-2v-tdTomato* knock-in reporter faithfully recapitulates endogenous *MLC-2v* expression during pre- and post-natal development. Neither the heterozygote nor homozygote mice carrying the knock-in allele showed any developmental defect. Therefore, this new mouse line will be an important genetic tool for studying cardiac and skeletal muscle specification during development and regeneration after injury.

METHODS

Animals

All animal procedures were performed with the approval of Vanderbilt University Medical Center. *MLC-2v-tdTomato* knock-in reporter mice were generated with the aid of Cyagen Biosciences Inc. (Santa Clara, CA). In brief, the targeting vector was generated by inserting *IRES-tdTomato-FRT-Neo^R-FRT* flanked by 4.9 kb 5' homology arm and 2.9 kb 3' homology arm. The linearized targeting construct was electroporated into C57BL/6 ESCs. Neomycin (G418) resistant clones were expanded and screened by PCR for homologous recombination. Homologous recombination of the selected ESC clones by PCR was confirmed by Southern blotting. The following PCR and southern blotting primers were used: 1) PCR_F: GCTGACCGCTTCCTCGTGCTTTA, 2) PCR_R1: CGCAGAGAGCCGAGTTTGAATCT, 3) Southern_F: TCATCTCACCTTGCTCCTGC, and 4) Southern_R: AAGGCGATAGAAGGCGATGC. The correctly targeted ESC clones were expanded and then injected into C57BL/6 blastocysts to generate mouse chimeras. The chimera mice were cross-bred with FLP transgenic mice, obtained from Jackson Laboratory (Stock number 009086), to remove a *Neo^R* cassette. *MLC-2v-tdTomato* knock-in reporter mice will be available to the academic research community upon acceptance of the manuscript.

Whole mount epifluorescent microscopy and Immunohistochemistry

Mouse embryos and adult hearts were harvested, and washed with ice-cold PBS. Epifluorescent images of whole embryos, embryonic hearts, adult hearts, or neonatal skeletal muscles were acquired with Nikon Eclipse 800. The samples were then fixed in pre-chilled 4% paraformaldehyde (PFA) in PBS for 30 minutes, incubated in 30% sucrose in PBS overnight at 4°C and then embedded in Tissue Freezing Medium. The cryosections at 8 µm thickness were obtained. After 5–10 minutes air dry, the cryosection slides were washed with PBS three times, permeabilized in 0.1% Triton X-100 in PBS for 20 minutes. Following PBS washing three times, the cryosections were blocked with M.O.M mouse IgG blocking reagent (Vector Labs) for 1 hour and 5% goat serum in M.O.M protein diluent (Vector Labs) for 30 minutes at room temperature. The cryosections were incubated with a primary antibody against α -actinin (mouse monoclonal, Sigma, 1:200 dilution), cardiac troponin T (mouse monoclonal, Thermoscientific, 1:200 dilution), alpha skeletal muscle actin (rabbit polyclonal, GeneTex, 1:200 dilution), fast skeletal myosin (mouse monoclonal, GeneTex, 1:200 dilution), or slow muscle myosin (mouse monoclonal, EMD Millipore, 1:200 dilution) overnight at 4°C. After PBS washing three times, the cryosections were incubated with a secondary antibody (goat anti-mouse AlexaFluor 488 or 647, Thermofisher) for one hour at room temperature and mounted with Vectashield containing DAPI (VectorLabs). TdTomato fluorescence was visualized without immunostaining. Images were captured using Zeiss LSM 710 confocal microscope.

Isolation of neonatal mouse cardiomyocytes and immunocytochemistry

Neonatal mouse ventricular cardiomyocytes at P1 were isolated using the Neomyts kit (Cellutron) as per manufacturer's protocol. Isolated neonatal ventricular cardiomyocytes were plated on Poly-D-Lysine/Laminin coated coverslip (BD Biosciences). Following 48hr culture in NS media (Cellutron), isolated neonatal cardiomyocytes were fixed with 4% PFA for 15 min and permeabilized with permeabilization buffer (0.05% Triton-X in PBS) for 5 min three times at room temperature. Isolated neonatal cardiomyocytes were incubated with blocking buffer (Universal blocking buffer, BiogeneX) for 45 min and then with primary antibodies against α -actinin (Mouse monoclonal, Sigma, 1:400 dilution) and MLC-2v (rabbit polyclonal, Protein tech, 1:200 dilution) for overnight at 4°C. Following washing three times for 5 min with permeabilization buffer, cells were incubated with secondary antibodies (goat anti-mouse AlexaFluor 488 and goat anti-rabbit AlexaFluor 647, Thermofisher) at room temperature for 1hr. After another set of washing (5 min x3 with permeabilization buffer), cells were mounted with Vectashield with DAPI (VectorLabs) and images were captured using Zeiss LSM 710 confocal microscope.

Isolation of adult mouse cardiomyocytes

Adult mouse hearts were dissected and perfused retrogradely via aortic cannulation in a Langendorff apparatus as described previously (O'Connell, Rodrigo, & Simpson, 2007). Hearts were perfused with following buffers with the indicated order: 1) perfusion buffer (NaCl 120.4 mM, KCl 14.7 mM, Na₂HPO₄ 0.6 mM, K₂HPO₄ 0.6 mM, MgSO₄ 1.2 mM, Na-HEPES 10 mM, NaHCO₃ 4.6 mM, Taurine 30 mM, BDM 10 mM, Glucose 5.5 mM, pH 7.0), 2) digestion buffer without CaCl₂ (Collagenase II 2.4 mg/mL in perfusion buffer), and

3) digestion buffer with CaCl_2 (Collagenase II 2.4 mg/mL, and CaCl_2 40 μM in perfusion buffer). Hearts were then removed from the Langendorff apparatus. Ventricular cardiomyocytes were mechanically dissociated and triturated using a fine scalpel and scissors and resuspended in stopping buffer (CaCl_2 11.7 μM in calf serum 2 ml plus perfusion buffer 18 ml). Cells were centrifuged at low speed, and supernatant was removed. Adult cardiomyocytes were fixed with 4% PFA for 15 min. After PBS washing, a drop of cardiomyocyte containing solution was placed on glass slide and mounted with a coverslip and Vectashield with DAPI (VectorLabs). Images were captured using Zeiss LSM 710 confocal microscope.

ACKNOWLEDGEMENT

This work was supported by Gilead Sciences Research Scholars Program and NIH R03 HL140264 (Y.-J. N.).

Grant Numbers: Gilead Sciences Research Scholars Program, and NIH R03 HL140264 for Young-Jae Nam

References

- Chen J, Kubalak SW, & Chien KR (1998). Ventricular muscle-restricted targeting of the RXRalpha gene reveals a non-cell-autonomous requirement in cardiac chamber morphogenesis. *Development*, 125(10), 1943–1949. [PubMed: 9550726]
- Chen J, Kubalak SW, Minamisawa S, Price RL, Becker KD, Hickey R, ... Chien KR (1998). Selective requirement of myosin light chain 2v in embryonic heart function. *J Biol Chem*, 273(2), 1252–1256. [PubMed: 9422794]
- Henderson SA, Xu YC, & Chien KR (1988). Nucleotide sequence of full length cDNAs encoding rat cardiac myosin light chain-2. *Nucleic Acids Res*, 16(10), 4722. [PubMed: 3380697]
- Kubalak SW, Miller-Hance WC, O'Brien TX, Dyson E, & Chien KR (1994). Chamber specification of atrial myosin light chain-2 expression precedes septation during murine cardiogenesis. *J Biol Chem*, 269(24), 16961–16970. [PubMed: 8207020]
- Kumar CC, Cribbs L, Delaney P, Chien KR, & Siddiqui MA (1986). Heart myosin light chain 2 gene. Nucleotide sequence of full length cDNA and expression in normal and hypertensive rat. *J Biol Chem*, 261(6), 2866–2872. [PubMed: 3005270]
- Lee KJ, Ross RS, Rockman HA, Harris AN, O'Brien TX, van Bilsen M, ... et al. (1992). Myosin light chain-2 luciferase transgenic mice reveal distinct regulatory programs for cardiac and skeletal muscle-specific expression of a single contractile protein gene. *J Biol Chem*, 267(22), 15875–15885. [PubMed: 1379240]
- Nudel U, Calvo JM, Shani M, & Levy Z (1984). The nucleotide sequence of a rat myosin light chain 2 gene. *Nucleic Acids Res*, 12(18), 7175–7186. [PubMed: 6091059]
- O'Brien TX, Lee KJ, & Chien KR (1993). Positional specification of ventricular myosin light chain 2 expression in the primitive murine heart tube. *Proc Natl Acad Sci U S A*, 90(11), 5157–5161. [PubMed: 8506363]
- O'Connell TD, Rodrigo MC, & Simpson PC (2007). Isolation and culture of adult mouse cardiac myocytes. *Methods Mol Biol*, 357, 271–296. doi: 10.1385/1-59745-214-9:271 [PubMed: 17172694]
- Ross RS, Navankasattusas S, Harvey RP, & Chien KR (1996). An HF-1a/HF-1b/MEF-2 combinatorial element confers cardiac ventricular specificity and established an anterior-posterior gradient of expression. *Development*, 122(6), 1799–1809. [PubMed: 8674419]
- Schiaffino S, & Reggiani C (1996). Molecular diversity of myofibrillar proteins: gene regulation and functional significance. *Physiol Rev*, 76(2), 371–423. doi: 10.1152/physrev.1996.76.2.371 [PubMed: 8618961]
- Schiaffino S, & Reggiani C (2011). Fiber types in mammalian skeletal muscles. *Physiol Rev*, 91(4), 1447–1531. doi: 10.1152/physrev.00031.2010 [PubMed: 22013216]

- Sheikh F, Lyon RC, & Chen J (2015). Functions of myosin light chain-2 (MYL2) in cardiac muscle and disease. *Gene*, 569(1), 14–20. doi: 10.1016/j.gene.2015.06.027 [PubMed: 26074085]
- Spater D, Hansson EM, Zangi L, & Chien KR (2014). How to make a cardiomyocyte. *Development*, 141(23), 4418–4431. doi: 10.1242/dev.091538 [PubMed: 25406392]

Figure 1.

Generation of *MLC-2v-tdTomato* knock-in reporter mouse line. (A) Schematic illustration of the gene targeting strategy. *IRES-tdTomato-FRT-Neo^R-FRT* cassette was inserted into 3' UTR of the *MLC-2v* gene locus. The *Neo^R* gene cassette is flanked by two FRT sites. By homologous recombination, the mice carrying *IRES-tdTomato-FRT-Neo^R-FRT* cassette (targeted allele) were generated. By cross breeding the chimera mice with FLP transgenic mice, *Neo^R* cassette was removed and *MLC-2v-tdTomato* knock-in mice carrying constitutive knock-in allele were generated. (B) PCR screening for the targeted ESCs using the indicated PCR primers in (A) (i.e. PCR_F and PCR_R). Eleven expanded clones demonstrated expected size of PCR band (3.5 kb). A wild type (WT) clone did not show the band. (C) Southern blot analyses to confirm the correctly targeted ESC clones. Using six (2, 3, 4, 6, 7, and 8) out of eleven PCR positive ESC clones, restrictive digestions with the indicated enzymes and hybridizations with the indicated southern probe in (A) were performed. Targeted ESCs for *MLC-2v-tdTomato* knock-in displayed 11 kb (with PacI and EcoRV) and 6.2 kb (with HindIII) bands, while WT ESCs did not show any band. The

southern blot results showed that five (2, 3, 4, 7, and 8) of the expanded 6 clones (2, 3, 4, 6, 7, and 8) were correctly targeted.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2.
MLC-2v-tdTomato knock-in reporter expression during heart development. Whole mount embryos (A-F) and embryonic hearts (G-J) demonstrate specific tdTomato expression in the ventricle within the heart at the indicated developmental stages. A: atrium; V: ventricle; IFT: inflow tract; OFT: outflow tract.

Figure 3.

Expression of *MLC-2v-tdTomato* knock-in reporter in ventricular cardiomyocytes during heart development. Frozen sections of the whole embryo or hearts at E10.5, E13.5 and E16.5 were processed. The frozen sections were immunostained for a cardiomyocyte marker, α -actinin. tdTomato expression was visualized without immunostaining. The cross section of the entire chest area at E10.5 was shown in (A). tdTomato expression was shown in the hearts at the indicated developmental stages (B-D). Magnified views of the area indicated by the white inlet in B-D were shown (E-G). A: atrium; V: ventricle; S: septum; RA: right atrium; LA: left atrium; RV: right ventricle; LV: left ventricle.

Figure 4.

Expression of *MLC-2v-tdTomato* knock-in reporter in the post-natal hearts. The hearts of P1 and 6 weeks (wk) old *MLC-2v-tdTomato* knock-in reporter mice were dissected. *tdTomato* fluorescence was specifically shown in the ventricles (A). A: atrium; V: ventricle. The hearts of heterozygous, homozygous, and wild type littermate mouse for *MLC-2v-tdTomato* knock-in allele at P0 were processed for frozen sections. Immunostaining for *MLC-2v* and cardiac troponin T (*cTnT*) was performed to determine the expression level of *tdTomato* and endogenous *MLC-2v* in an each genotype of mouse (B). +/+ : homozygous; +/- : heterozygous; -/- : wild type. *tdTomato* was visualized without immunostaining. The heart of 6 week old *MLC-2v-tdTomato* knock-in homozygote mice was processed for frozen sections and immunostained for *MLC-2v* (C). The lower (top panel) and higher (bottom panel) magnification pictures are shown. Neonatal and adult ventricular cardiomyocytes were isolated. Isolated neonatal ventricular cardiomyocytes were immunostained for α -actinin and *MLC-2v* (D, left panel). *tdTomato* fluorescence was visualized in adult mouse ventricular cardiomyocytes without immunostaining (D, right panel).

Figure 5.

Expression of *MLC-2v-tdTomato* knock-in reporter in skeletal muscles. Slow twitch (soleus) and fast twitch (extensor digitorum longus) muscles were dissected from E20, P0, P1 and 6 week old *MLC-2v-tdTomato* knock-in mice. tdTomato fluorescence was specifically shown in soleus, not in extensor digitorum longus, at post-natal stages (A). The dissected soleus and extensor digitorum longus muscles at P1 were processed for frozen sections and immunostained for alpha skeletal muscle actin (a pan-skeletal muscle marker) and slow muscle myosin (a slow twitch muscle marker), or fast muscle myosin (a fast twitch muscle marker) (B). S: soleus; E: extensor digitorum longus