Published in final edited form as:

Org Lett. 2018 November 16; 20(22): 7345–7350. doi:10.1021/acs.orglett.8b03345.

Photocatalytic Oxyamination of Alkenes: Copper(II) Salts as Terminal Oxidants in Photoredox Catalysis

Nicholas L. Reed, Madeline I. Herman, Vladimir P. Miltchev, and Tehshik P. Yoon*

Department of Chemistry, University of Wisconsin-Madison, 1101 University Avenue, Madison WI 53706.

Abstract

A photocatalytic method for the oxyamination of alkenes using simple nucleophilic nitrogen atom sources in place of pre-functionalized electrophilic nitrogen atom donors is reported. Copper(II) is an inexpensive, practical, and uniquely effective terminal oxidant for this process. In contrast to oxygen, peroxides, and similar oxidants commonly utilized in non-photochemical oxidative methods, the use of copper(II) as a terminal oxidant in photoredox reactions avoids the formation of reactive heteroatom-centered radical intermediates that can be incompatible with electron-rich functional groups. As a demonstration of the generality of this concept, it has been shown that diamination and deoxygenation reactions can also be accomplished using similar photooxidative conditions.

Graphical Abstract

The oxyamination of alkenes remains an important problem in synthetic chemistry due to the prominence of amino alcohol-derived subunits in many important classes of bioactive natural products, pharmaceutical compounds, and chiral reagents for stereoselective synthesis.¹ However, the most extensively developed oxyamination methods, including the Sharpless aminohydroxylation and its derivatives, require pre-oxidized, electrophilic nitrogen donors such as chloramines, iminoiodinanes, or hydroxylamines that can be difficult to synthesize, are often explosive, and can be challenging to handle because of their chemical instability. One important objective in oxidative functionalization methodology has

*Corresponding Author Tehshik P. Yoon: tyoon@chem.wisc.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.
Experimental procedures and characterization data (PDF)

therefore been the development of new oxyamination protocols that can be conducted directly with simple nucleophilic nitrogen and oxygen atom donors.²

The design of effective net-oxidative transformations is a fundamental problem in catalysis, but it poses a particular challenge for photoredox chemistry.³ Although many powerful oxidative photoredox processes have been reported, the majority have exploited the reactivity of preoxidized electrophilic group-transfer reagents where some portion of the terminal oxidant is structurally incorporated into the product (e.g., halogenation,⁴ amination,⁵ or perfluoroalkylation⁶). In contrast, an oxyamination protocol using simple heteroatom nucleophiles requires an “oxidase” strategy in which the terminal oxidant serves as an electron acceptor and not a functional group donor.⁸ Unfortunately, terminal oxidants that are suitable for non-photochemical reactions can be problematic in photoredox applications. For instance, molecular oxygen,⁹ often identified as an ideal oxidant for organometallic catalysis,¹⁰ is an efficient quencher of the excited states of many photocatalysts¹¹ and reacts rapidly with the organoradical intermediates that are ubiquitous in photoredox chemistry.¹² Similarly, photoredox activation of peroxides affords promiscuously reactive oxygen-centered radicals that can be incompatible with common electron-rich functional groups.¹³ Organic oxidants (e.g., perhaloalkanes, nitroarenes, amine *N*-oxides, or arene diazonium salts)¹⁴ can avoid these problems, but are relatively expensive and produce unattractive stoichiometric byproducts. The identification of a general terminal oxidant that is free from these drawbacks would thus represent a significant advance in photochemical synthesis.

We began by examining Nicewicz's pioneering examples of photocatalytic alkene hydrofunctionalization reactions¹⁵ (Scheme 1). In these studies, photooxidation of an alkene affords an electrophilic radical cation intermediate (**1**⁺). Subsequent trapping of a heteroatomic nucleophile such as an alkyl carbamate results in the generation of a carbon-centered radical intermediate (**2**). Interception by a hydrogen atom donor, often a malononitrile or thiophenol, affords the product of a redox-neutral hydrofunctionalization reaction (**3**). A recent report has also described interception of **2** by catalytic CuCl₃ as an atom-transfer reagent to effect an oxidative aminochlorination.^{4c} We speculated that an appropriate alternative oxidant could react with this radical to produce a formally cationic intermediate; cyclization with facile loss of *tert*-butyl cation would afford oxazolidone products (**4**), effecting a net-oxidative oxyamination reaction.

We began by examining the effect of several photocatalysts and potential terminal oxidants on the oxidative cyclization of carbamate **5a**. Irradiation with a 15 W blue LED for 15 h in the presence of a commercially available pyrylium photocatalyst (TPPT, **7**) and Cu(TFA)₂ as an oxidant affords oxazolidone **6** in 82% yield as a single diastereomer (Table 1, entry 1). Although other strongly oxidizing organic photocatalysts afford similar results, transition metal photoredox catalysts with less positive excited state reduction potentials are ineffective (entries 2 and 3). The addition of a terminal oxidant is required for conversion to oxazolidone (entry 4), and importantly, the identity of the oxidant is critical. Other Cu(II) salts provide diminished yields of the oxyamination product (entries 5 and 6), but an extensive screen of alternate oxidants spanning several different classes revealed that only Cu(II) oxidants are effective. For example, PhI(OAc)₂, TEMPO, MnO₂, and FeCl₃ provide

no conversion to **6**, while air, DDQ, and *t*-BuOOH result in extensive decomposition, which is consistent with the formation of reactive oxygen-centered radical intermediates (entries 7 and 8). Finally, control experiments validated the photocatalytic nature of this reaction; no product is observed in the absence of light or photocatalyst (entries 9 and 10).

The unique suitability of Cu(II) salts in this reaction is consistent with seminal investigations by Kochi, demonstrating that Cu(II) salts are exceptionally rapid and efficient oxidants of carbon-centered radicals.¹⁶ Moreover, copper compounds do not interfere with the photochemistry of common photoredox catalysts and have been utilized as co-catalysts in several recently reported photocatalytic methods.¹⁷ Additionally, Cu(II) salts are attractive, practical stoichiometric oxidants that are readily handled on the bench-top and removed from a reaction mixture by simple extraction or filtration through silica.¹⁸ Copper(II) salts are generally less toxic¹⁹ and less expensive than many common organic terminal oxidants. Although the cost of Cu(TFA)₂ itself is unusually high, on par with common organic oxidants, it can be synthesized from inexpensive CuCO₃ and trifluoroacetic acid, and reactions using either commercial or freshly prepared Cu(TFA)₂ give identical results.

These optimized conditions proved to be effective on a gram-scale batch, affording the oxyamination product in 84% yield (Scheme 2a). A variety of styrenic alkenes undergo facile oxyamination. The arene could be substituted at all positions with common functional groups, including halides, ketones, and aldehydes (Scheme 2b, **8–18**). Heteroaryl styrenes can also undergo oxyamination; while pyridines provide diminished rates, less basic O and S heterocycles were readily tolerated (Scheme 2c, **19–21**). The diastereoselectivity was excellent, except in the cases of very electron-rich styrenes (**11** and **19**).²⁰ Increasing the length of the tether, incorporating heteroatoms, or substituting the carbon chain had little effect on the rate of the cyclization (Scheme 2d, **22–24**). The styrene could be substituted on the alkene moiety as well, both at the α and β positions (Scheme 2e, **25** and **26**). Trisubstituted aliphatic alkenes, however, instead afforded an allylic carbamate (**27**), suggesting that although the excited state photocatalyst could oxidize the substrate, the resulting cationic intermediate undergoes elimination faster than trapping by the carbamoyl oxygen. Finally, we examined intermolecular oxyaminations using simple Boc carbamate as the amine source (Scheme 2f). Both *cis* and *trans* 1,2-disubstituted styrenes (**28** and **29**) react with good yields. Interestingly, methyl cinnamate also underwent oxyamination, despite the presence of the highly electron-withdrawing ester moiety, affording direct access to a highly functionalized oxazolidone scaffold (**31**).

Scheme 3 summarizes several experiments supporting the mechanism proposed in Scheme 4a. First, subjecting hydroamination product **32** to the optimized reaction conditions resulted in the formation of oxazolidone product **33** in 14% yield. Thus, the benzylic radical produced by photooxidation of the arene²¹ can re-enter the catalytic cycle, supporting its role as an intermediate in this process. Second, irradiation of **5a** in the presence of 2 equiv of TPPT, but in the absence of Cu(TFA)₂, resulted in complete decomposition of the substrate without formation of the oxyamination product (eq 2). This result is consistent with the hypothesis that Cu(II) is not merely a terminal oxidant, but is intimately involved in oxidation of the organoradical intermediate, as expected from Kochi's studies. Third, the formation of the final product by loss of *tert*-butyl cation is consistent with the observation

that while Cbz carbamates cyclize effectively, Moc carbamates provide no product, in line with the poor stability of the methyl cation. Finally, because the reaction requires only 1.2 equiv of $\text{Cu}(\text{TFA})_2$, $\text{Cu}(\text{II})$ must be acting as a net two-electron oxidant. It is unclear whether the $\text{Cu}(\text{I})$ intermediate directly turns over the photocatalyst by oxidation of **7** or disproportionates to $\text{Cu}(\text{II})$ and $\text{Cu}(0)$. Both mechanisms would be consistent with the observed precipitation of copper metal from solution during the course of the reaction.

Oxyamination reactions are representative of a broader class of synthetically important oxidative alkene difunctionalizations.²² If this $\text{Cu}(\text{II})$ -mediated photocatalytic strategy could be generalized to the use of alternate nucleophilic reaction partners, it would provide a novel, flexible approach toward the photocatalytic synthesis of a wide range of vicinal heteroatom arrays. Indeed, the use of ureas in place of carbamates affords the products of net alkene diamination (Scheme 4b). The optimal photocatalyst (MesAcrMe^+) and oxidant [$\text{Cu}(\text{OAc})_2$] were slightly different in this case, which demonstrates that these variables can be tuned to achieve optimal reactivity in different transformations. Moreover, the irradiation of dihydronaphthalene with various alcohols in the presence of $\text{Cu}(\text{TFA})_2$ and TPPT results in alkene dioxxygenation (Scheme 4c). Thus, these preliminary results indicate that the identity of each of the heteroatoms introduced across the alkene can be varied and suggest that this reaction design plan might be generalizable to a much wider range of oxidative functionalization reactions.

In summary, copper(II) salts are effective oxidants that enable the oxidative difunctionalization of alkenes using photoredox catalysis. More broadly, these results are exciting because organoradical intermediates are common in photoredox reactions. The ability to divert these intermediates toward cationic reactivity using convenient $\text{Cu}(\text{II})$ oxidants suggests a powerful and conceptually novel approach toward the design of a wide range of new oxidative functionalization reactions. Studies toward this broader goal are ongoing in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This work was funded by the NIH (GM098886). The mass spectroscopy facilities at UW-Madison are funded in part by the NIH (S10 OD020022), as are the NMR facilities (S10 OD012245).

REFERENCES

1. a) For recent reviews of alkene oxyamination, see: O'Brien P "Sharpless Asymmetric Aminohydroxylation: Scope, Limitations, and Use in Synthesis," *Angew. Chem. Int. Ed* 1999, 38, 326–329. b) Nilov D; Reiser O "The Sharpless Asymmetric Aminohydroxylation – Scope and Limitation," *Adv. Synth. Catal* 2002, 344, 1169–1173. c) Bodkin JA; McLeod MD "The Sharpless asymmetric aminohydroxylation," *J. Chem. Soc., Perkin Trans 1* 2002, 2733–2746. d) Donohoe TJ; Callens CKA; Flores A; Lacy AR; Rathi AH "Recent Developments in Methodology for the Direct Oxyamination of Olefins," *Chem. Eur. J* 2011, 17, 58–76. [PubMed: 21207600]
2. a) For selected examples of alkene oxyaminations without preoxidized nitrogen atom donors, see: Alexanian EJ; Lee C; Sorensen EJ "Palladium-Catalyzed Ring-Forming Aminoacetoxylation of

Alkenes,” J. Am. Chem. Soc. 2005, 127, 7690–7691. [PubMed: 15913354] b)Liu G; Stahl SS “Highly Regioselective Pd-Catalyzed Intermolecular Aminoacetoxylation of Alkenes and Evidence for *cis*-Aminopalladation and S_N2 C–O Bond Formation,” J. Am. Chem. Soc. 2006, 128, 7179–7181. [PubMed: 16734468] c)Fuller PH; Kim J-W; Chemler SR “Copper Catalyzed Enantioselective Intramolecular Aminooxygenation of Alkenes,” J. Am. Chem. Soc. 2008, 130, 17638–17639. [PubMed: 19049311] d)Lovick HM; Michael FE “Metal-Free Highly Regioselective Aminotrifluoroacetoxylation of Alkenes,” J. Am. Chem. Soc. 2010, 132, 1249–1251. [PubMed: 20058921] e)Wardrop DJ; Bowen EG; Forslund RE; Sussman AD; Weerasekera SL “Intramolecular Oxamidation of Unsaturated O-Alkyl Hydroxamates: A Remarkably Versatile Entry to Hydroxy Lactams,” J. Am. Chem. Soc. 2010, 132, 1188–1189. [PubMed: 19788297] f)de Haro T; Nevado C “Flexible Gold-Catalyzed Regioselective Oxidative Difunctionalization of Unactivated Alkenes,” Angew. Chem. Int. Ed. 2011, 50, 906–910.g)Farid U; Wirth T “Highly Stereoselective Metal-Free Oxyaminations Using Chiral Hypervalent Iodine Reagents,” Angew. Chem. Int. Ed. 2012, 51, 3462–3465.h)Shen H-C; Wu Y-F; Zhang Y; Fan L-F; Han Z-Y; Gong L-Z “Palladium-Catalyzed Asymmetric Aminohydroxylation of 1,3-Dienes,” Angew. Chem. Int. Ed. 2018, 57, 2372–2376.i)Wu F; Stewart S; Ariyaratna JP; Li W “Aerobic Copper-Catalyzed Alkene Oxyamination for Amino Lactone Synthesis,” ACS Catal. 2018, 8, 1921–1925.

3. a)For reviews, see: Narayanam JMR; Stephenson CRJ “Visible light photoredox catalysis: applications in organic synthesis,” Chem. Soc. Rev 2011, 40, 102–113. [PubMed: 20532341] b)Prior CK; Rankic DA; MacMillan DWC “Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis,” Chem. Rev 2013, 113, 5322–5363. [PubMed: 23509883] c)Romero NA; Nicewicz DA “Organic Photoredox Catalysis,” Chem. Rev 2016, 116, 10075–10166. [PubMed: 27285582] d)Twilton J; Le C; Zhang P; Shaw MH; Evans RW; MacMillan DWC “The merger of transition metal and photocatalysis” Nature Rev. Chem 2017, 1, 0052.e)Zou YQ; Hörmann FM; Bach T “Iminium and enamine catalysis in enantioselective photochemical reactions,” Chem. Soc. Rev 2018, 47, 278–290. [PubMed: 29155908] f)Silvi M; Melchiorre P “Enhancing the potential of enantioselective organocatalysis with light,” Nature 2018, 554, 41–49. [PubMed: 29388950]
4. a)Rueda-Becerril M; Mahé O; Drouin M; Majewski MB; West JF ; Wolf MO; Sammis GM; Paquin JF “Direct C-F bond formation using photoredox catalysis,” J. Am. Chem. Soc 2014, 136, 2637–2641. [PubMed: 24437369] b)Ventre S; Petronijevic FR; MacMillan DWC “Decarboxylative Fluorination of Aliphatic Carboxylic Acids via Photoredox Catalysis,” J. Am. Chem. Soc. 2015, 137, 5654–5657. [PubMed: 25881929] c)Griffin JD; Cavanaugh CL; Nicewicz DA “Reversing the Regioselectivity of Halofunctionalization Reactions through Cooperative Photoredox and Copper Catalysis,” Angew. Chem. Int. Ed 2017, 56, 2097–2100.d)Davies J; Sheikh NS; Leonori D “Photoredox Imino Functionalizations of Olefins,” Angew. Chem. Int. Ed. 2017, 56, 13361–13365.
5. a)Allen LJ; Cabrera PJ; Lee M; Sanford MS “N-Acyloxyphthalimides as Nitrogen Radical Precursors in the Visible Light Photocatalyzed Room Temperature C–H Amination of Arenes and Heteroarenes,” J. Am. Chem. Soc. 2014, 136, 5607–5610. [PubMed: 24702705] b)Greulich TW; Daniliuc CG; Studer A “N-Aminopyridinium Salts as Precursors for N-Centered Radicals – Direct Amidation of Arenes and Heteroarenes,” Org. Lett. 2015, 17, 254–257. [PubMed: 25541887] c)Brachet E; Ghosh T; Ghosh I; König B “Visible light C–H amidation of heteroarenes with benzoyl azides,” Chem. Sci. 2015, 6, 987–992. [PubMed: 29560185] d)Rabet PTG; Fumagalli G; Boyd S; Greaney MF “Benzylic C–H Azidation Using the Zhdankin Reagent and a Copper Photoredox Catalyst,” Org. Lett. 2016, 18, 1646–1649. [PubMed: 27007454]
6. a)Nagib DA; MacMillan DWC “Trifluoromethylation of arenes and heteroarenes by means of photoredox catalysis,” Nature 2011, 480, 224–228. [PubMed: 22158245] b)Iqbal N; Choi S; Kim E; Cho EJ “Trifluoromethylation of Alkenes by Visible Light Photoredox Catalysis,” J. Org. Chem. 2012, 77, 11383–11387. [PubMed: 23167602] c)Ye Y; Sanford MS “Merging Visible-Light Photocatalysis and Transition-Metal Catalysis in the Copper-Catalyzed Trifluoromethylation of Boronic Acids with CF₃I,” J. Am. Chem. Soc. 2012, 134, 9034–9037. [PubMed: 22624669]
7. Stahl SS “Palladium Oxidase Catalysis: Selective Oxidation of Organic Chemicals by Direct Dioxygen-Coupled Turnover,” Angew. Chem. Int. Ed. 2004, 43, 3400–3420.
8. a)For alternate strategies towards photoredox oxyaminations using preoxidized heteroatom donors, see: Davies J; Booth SG; Essafi S; Dryfe RAW; Leonori D “Visible-Light-Mediated Generation of Nitrogen-Centered Radicals: Metal-Free Hydroimination and Iminohydroxylation Cyclization

- Reactions." *Angew. Chemie Int. Ed.* 2015, 54, 14017–14021. b) Hu X-Q; Chen J; Chen J-R; Yan D-M; Xiao W-J "Organophotocatalytic Generation of N- and O-Centred Radicals Enables Aerobic Oxyamination and Dioxygenation of Alkenes," *Chem. Eur. J.* 2016, 22, 14141–14146. [PubMed: 27258972] c) Ren X; Guo Q; Chen J; Xie H; Xu Q.; Lu Z "Visible-Light Promoted Distereodivergent Intramolecular Oxyamidation of Alkenes," *Chem. Eur. J.* 2016, 22, 18695–18699. [PubMed: 27770547] d) Miyazawa K; Koike T; Akita M "Aminohydroxylation of olefins with iminopyridinium ylides by dual Ir photocatalysis and Sc(OTf)₃ catalysis," *Tetrahedron* 2016, 72, 7813–7820.
9. a) For examples of net-oxidative photoredox methods utilizing dioxygen, see: Condie AG; Gonzalez-Gomez JC; Stephenson CRJ "Visible-Light Photoredox Catalysis: Aza-Henry Reactions via C–H Functionalization," *J. Am. Chem. Soc.* 2010, 132, 1464–1465. [PubMed: 20070079] b) Zou Y-Q; Lu L-Q; Fu L; Chang N-J; Rong J; Chen J-R; Xiao W-J "Visible-light-induced oxidation/[3+2] cycloaddition/oxidative aromatization sequence: a photocatalytic strategy to construct pyrrolo[2,1-a]isoquinolines," *Angew. Chem. Int. Ed.* 2011, 50, 7171–7175. c) Zou Y-Q; Chen J-R; Liu X-P; Lu L-Q; Davis RL; Jørgensen KA; Xiao W-J "Highly efficient aerobic oxidative hydroxylation of arylboronic acids: photoredox catalysis using visible light," *Angew. Chem. Int. Ed.* 2012, 51, 784–788. d) Cheng Y; Yang J; Qu Y; Li P "Aerobic visible-light photoredox radical C–H functionalization: catalytic synthesis of 2-substituted benzothiazoles," *Org. Lett.* 2012, 14, 98–101. [PubMed: 22146071] e) Romero NA; Margrey KA; Tay NE; Nicewicz DA "Site-selective arene C–H amination via photoredox catalysis," *Science* 2015, 349, 1326–1330. [PubMed: 26383949] f) McManus JB; Nicewicz DA "Direct C–H Cyanation of Arenes via Organic Photoredox Catalysis," *J. Am. Chem. Soc.* 2017, 139, 2880–2883. [PubMed: 28177237]
10. a) Punniyamurthy T; Velusamy S; Iqbal J "Recent Advances in Transition Metal Catalyzed Oxidation of Organic Substrates with Molecular Oxygen," *Chem. Rev.* 2005, 105, 2329–2364. [PubMed: 15941216] b) Shi Z; Zhang C; Tang C; Jiao N "Recent advances in transition-metal catalyzed reactions using molecular oxygen as the oxidant," *Chem. Soc. Rev.* 2012, 41, 3381–3430. [PubMed: 22358177] c) Campbell AN; Stahl SS "Overcoming the 'Oxidant Problem': Strategies to Use O₂ as the Oxidant in Organometallic C–H Oxidation Reactions Catalyzed by Pd (and Cu)," *Acc. Chem. Res.* 2012, 45, 851–863. [PubMed: 22263575]
11. a) Winterle JS; Kliger DS; Hammond GS "Mechanisms of photochemical reactions in solution. 80. Photochemical oxidation of tris(2,2'-bipyridyl)ruthenium(II) by molecular oxygen," *J. Am. Chem. Soc.* 1976, 98, 3719–3721. b) Srinivasan VS; Podolski D; Westrick NJ; Neckers DC "Photochemical generation of superoxide ion (O₂^{•−}) by Rose Bengal and Ru(bpy)₃²⁺," *J. Am. Chem. Soc.* 1978, 100, 6513–6515. c) Takizawa S; Aboshi R; Murata S "Photooxidation of 1,5-dihydroxynaphthalene with iridium complexes as singlet oxygen sensitizers," *Photochem. Photobiol. Sci.* 2011, 10, 895–903. [PubMed: 21298183]
12. a) Maillard B; Ingold KU; Scaiano JC "Rate constants for the reactions of free radicals with oxygen in solution," *J. Am. Chem. Soc.* 1983, 105, 5095–5099. b) Ohkubo K; Nanjo T; Fukuzumi S "Efficient Photocatalytic Oxygenation of Aromatic Alkene to 1,2-Dioxetane with Oxygen via Electron Transfer," *Org. Lett.* 2005, 7, 4265–4268. [PubMed: 16146403]
13. a) For examples of net-oxidative photoredox methods using peroxides, see: Dai C; Meschini F; Narayanam JMR; Stephenson CRJ "Friedel-Crafts Amidoalkylation via Thermolysis and Oxidative Photocatalysis," *J. Org. Chem.* 2012, 77, 4425–4431. [PubMed: 22458307] b) Blum TR; Zhu Y; Nordeen SA; Yoon TP "Photocatalytic Synthesis of Dihydrobenzofurans by Oxidative [3+2] Cycloaddition of Phenols," *Angew. Chem. Int. Ed.* 2014, 53, 11056–11059. c) Liu Z; Zhang Y; Cai Z; Sun H; Cheng X "Photoredox Removal of *p*-Methoxybenzyl Ether Protecting Group with Hydrogen Peroxide as Terminal Oxidant," *Adv. Synth. Catal.* 2015, 357, 589–593.
14. a) Cano-Yelo H; Deronzier A "Photo-oxidation of some carbinols by the Ru(II) polypyridyl complex-aryl diazonium salt system," *Tetrahedron Lett.* 1984, 25, 5517–5520. b) DiRocco DA; Rovis T "Catalytic asymmetric α -acylation of tertiary amines mediated by a dual catalysis mode: N-heterocyclic carbene and photoredox catalysis," *J. Am. Chem. Soc.* 2012, 134, 8094–8097. [PubMed: 22548244] c) Freeman DB; Furst L; Condie AG; Stephenson CRJ "Functionally Diverse Nucleophilic Trapping of Iminium Intermediates Generated Utilizing Visible Light," *Org. Lett.* 2012, 14, 94–97. [PubMed: 22148974] d) Beatty JW; Douglas JJ; Cole KP; Stephenson CRJ "A scalable and operationally simple radical trifluoromethylation," *Nat. Commun.* 2015, 6, 7919. [PubMed: 26258541] e) Pandey G; Laha R; Singh D "Benzylic C(sp³)-H Functionalization for C–

- N and C–O Bond Formation via Visible Light Photoredox Catalysis” J. Org. Chem 2016, 81, 7161–7171. [PubMed: 27269307]
15. a)Hamilton DS; Nicewicz DA “Direct catalytic anti-Markovnikov hydroetherification of alkenols,” J. Am. Chem. Soc. 2012, 134, 18577–18580. [PubMed: 23113557] b)Nguyen TM; Nicewicz DA “Anti-Markovnikov Hydroamination of Alkenes Catalyzed by an Organic Photoredox System,” J. Am. Chem. Soc. 2013, 135, 9588–9591. [PubMed: 23768239] c)Perkowski AJ; Nicewicz DA “Direct Catalytic Anti-Markovnikov Addition of Carboxylic Acids to Alkenes,” J. Am. Chem. Soc. 2013, 135, 10334–10337. [PubMed: 23808532] d)Nguyen TM; Manohar N; Nicewicz DA “Anti-Markovnikov Hydroamination of Alkenes Catalyzed by a Two-Component Organic Photoredox System: Direct Access to Phenethylamine Derivatives,” Angew. Chem. Int. Ed. 2014, 53, 6198–6201.e)Wilger DJ; Grandjean JMM; Lammert TR; Nicewicz DA “The direct anti-Markovnikov addition of mineral acids to styrenes,” Nat. Chem. 2014, 6, 720–726. [PubMed: 25054943]
16. a)Kochi JK; Bemis A “Carbonium ions from alkyl radicals by electron transfer,” J. Am. Chem. Soc 1968, 90, 4038–4051.b)Kochi JK; Bemis A; Jenkins CL “Mechanism of electron transfer oxidation of alkyl radicals by copper(II) complexes,” J. Am. Chem. Soc 1968, 90, 4616–4625.c)Jenkins CL; Kochi JK “Homolytic and ionic mechanisms in the ligand-transfer oxidation of alkyl radicals by copper(II) halides and pseudohalides,” J. Am. Chem. Soc 1972, 94, 856–865.
17. a)Yoo W-J; Tsukamoto T; Kobayashi S “Visible-Light-Mediated Chan-Lam Coupling Reactions of Aryl Boronic Acids and Aniline Derivatives,” Angew. Chem. Int. Ed. 2015, 54, 6587–6590.b)Perepichka I; Kundu S; Hearne Z; Li C-J “Efficient merging of copper and photoredox catalysis for the asymmetric cross-dehydrogenative-coupling of alkynes and tetrahydroisoquinolines,” Org. Biomol. Chem. 2015, 13, 447–451. [PubMed: 25372475] c)Mao R; Frey A; Balon J; Hu X “Decarboxylative C(sp³)-N cross-coupling via synergetic photoredox and copper catalysis,” Nat. Catal. 2018, 1, 120–126.d)Kautzky JA; Wang T; Evans RW; MacMillan DWC “Decarboxylative Trifluoromethylation of Aliphatic Carboxylic Acids,” J. Am. Chem. Soc. 2018, 140, 6522–6526. [PubMed: 29754491] e)Le C; Chen TQ; Liang T; Zhang P; MacMillan DWC “A radical approach to the copper oxidative addition problem: Trifluoromethylation of bromoarenes,” Science. 2018, 360, 1010–1014. [PubMed: 29853683] f)Liang Y; Zhang X; MacMillan DWC “Decarboxylative sp³ C–N coupling via dual copper and photoredox catalysis,” Nature 2018, 559, 83–88. [PubMed: 29925943]
18. a)Zabawa TP; Kasi D; Chemler SR “Copper (II) Acetate Promoted Intramolecular Diamination of Unactivated Olefins,” J. Am. Chem. Soc. 2005, 127, 11250–11251. [PubMed: 16089447] b)Baran PS; DeMartino MP “Intermolecular Oxidative Enolate Heterocoupling,” Angew. Chem. Int. Ed. 2006, 45, 7083–7086.c)Stuart DR; Fagnou K “The Catalytic Cross-Coupling of Unactivated Arenes,” Science 2007, 316, 1172–1175. [PubMed: 17525334] d)Hyster TK; Rovis T “Rhodium-Catalyzed Oxidative Cycloaddition of Benzamides and Alkynes via C–H/N–H Activation,” J. Am. Chem. Soc. 2010, 132, 10565–10569. [PubMed: 20662529]
19. ICH Q3D. Guideline for Elemental Impurities, http://www.ich.org/fileadmin/Public_Web_Site/ICH_Products/Guidelines/Quality/Q3D/Q3D_Step_4.pdf
20. An experiment resubjecting the major diastereomer of **19** to the reaction conditions did not result in any observable epimerization to the minor isomer. This control experiment suggests that the diastereomer ratios reflect an intrinsic kinetic preference in the C–O bond-forming step.
21. a)Ohkubo K; Mizushima K; Iwata R; Souma K.; Suzuki N; Fukuzumi S “Simultaneous production of *p*-tolualdehyde and hydrogen peroxide in photocatalytic oxygenation of *p*-xylene and reduction of oxygen with 9-mesityl-10-methylacridinium ion derivatives,” Chem. Commun 2010, 46, 601–603.b)Pandey G; Pal S; Laha R “Direct Benzylic C–H Activation for C–O Bond Formation by Photoredox Catalysis” Angew. Chem. Int. Ed 2013, 52, 5146–5149.c)Yi H; Bian C; Hu X; Niu L; Lei A “Visible light mediated efficient oxidative benzylic sp³ C–H to ketone derivatives obtained under mild conditions using O₂,” Chem. Commun 2015, 51, 14046–14049.d)Zhou R; Liu H; Tao H; Yu X; Wu J “Metal-free direct alkylation of unfunctionalized allylic/benzylic sp³ C–H bonds via photoredox induced radical cation deprotonation,” Chem. Sci 2017, 8, 4654–4659. [PubMed: 28970885] e)Mazzarella D; Crisenza GEM; Melchiorre P “Asymmetric Photocatalytic C–H Functionalization of Toluene and Derivatives,” J. Am. Chem. Soc 2018, 140, 8439–8443. [PubMed: 29932655]

22. a) For recent reviews covering oxidative alkene difunctionalization, see: Chemler SR “The Enantioselective Intramolecular Aminative Functionalization of Unactivated Alkenes, Dienes, Allenes and Alkynes for the Synthesis of Chiral Nitrogen Heterocycles”, *Org. Biomol. Chem.* 2009, 7, 3009–3019. [PubMed: 20976023] b) McDonald RI; Liu G; Stahl SS “Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications”, *Chem. Rev.* 2011, 111, 2981–3019. [PubMed: 21428440] c) Yin G; Mu X; Liu G “Palladium(II)-Catalyzed Oxidative Difunctionalization of Alkenes: Bond Forming at a High-Valent Palladium Center”, *Acc. Chem. Res.* 2016, 49, 2413–2423. [PubMed: 27739689] d) Lan X-W; Wang N-X; Xing Y “Recent Advances in Radical Difunctionalization of Simple Alkenes”, *Eur. J. Org. Chem.* 2017, 5821–5851.

■ *Conventional approach: oxyaminations with preoxidized nitrogen atom source*

■ *This work: photocatalytic oxyaminations with external oxidant*

Goals

- low-cost, bench-stable, environmentally benign oxidant
- compatible with photocatalytic reaction conditions
- avoids promiscuous heteroatom-centered radical intermediates

Figure 1.
Oxyamination reactions with and without preoxidized nitrogen atom donors.

Scheme 1.

Diverting photogenerated radical intermediates toward net oxidative functionalization reactions.

Scheme 2. Scope studies for photocatalytic oxyamination reactions^a

^a Unless otherwise noted, all reactions were conducted using 2.5 mol % 7, 1.2 equiv Cu(TFA)₂ in degassed CH₂Cl₂ and irradiated with a 15 W blue LED flood lamp for 15–48 h. Isolated yields are reported. ^b Reaction time 48 h. ^c Reaction time 72 h. ^d Reaction time 7 days.

Scheme 3.

Experiments supporting the proposed mechanism.

Scheme 4.

Working mechanistic hypothesis and extension to alkene difunctionalization

^a Reactions conducted using MesAcrMe⁺BF₄[−] as the photocatalyst and Cu(OAc)₂ as the terminal oxidant. ^b Reactions conducted using 4 equiv of alcohol, TPPT as photocatalyst, and Cu(TFA)₂ as terminal oxidant.

Table 1.Effect of reaction variables on oxyamination.^a

entry	variation from standard conditions	yield
1	None	82%
2	MesAcrMe ⁺ BF ₄ ⁻ instead of TPPT	72%
3	Ir[dF(CF ₃)ppy] ₂ (dtbbpy)PF ₆ instead of TPPT	5%
4	No Cu(TFA) ₂	0% ^b
5	Cu(OTf) ₂ instead of Cu(TFA) ₂	24%
6	Cu(OAc) ₂ instead of Cu(TFA) ₂	7%
7	PhI(OAc) ₂ , TEMPO, MnO ₂ , or FeCl ₃ instead of Cu(TFA) ₂	0% ^b
8	Air, DDQ, or <i>t</i> -BuOOH instead of Cu(TFA) ₂	0% ^c
9	No TPPT	1%
10	No light	0% ^b

^aUnless otherwise noted, all reactions were conducted in degassed CH₂Cl₂ and irradiated with a 15 W blue LED flood lamp for 15 h. Yields determined by ¹H NMR analysis of the unpurified reaction mixtures using phenanthrene as an internal standard.

^bNo conversion.

^cSubstrate decomposition.