Published in final edited form as:

Cogn Behav Pract. 2016 February ; 23(1): 40–50. doi:10.1016/j.cbpra.2014.09.003.

Pilot Open Case Series of Voice over Internet Protocol-delivered Assessment and Behavior Therapy for Chronic Tic Disorders

Emily J. Ricketts¹, Christopher C. Bauer², Dagong Ran³, Michael B. Himle⁴, and Douglas W. Woods²

¹Western Psychiatric Institute and Clinic

²Texas A&M University

³Southern Illinois University

⁴University of Utah

Abstract

Comprehensive Behavioral Intervention for Tics (CBIT) is an efficacious treatment for children with Chronic Tic Disorders (CTDs). Nevertheless, many families of children with CTDs are unable to access CBIT due to a lack of adequately trained treatment providers, time commitment, and travel distance. This study established the interrater reliability between in-person and VoIP administrations of the Yale Global Tic Severity Scale (YGTSS), and examined the preliminary efficacy, feasibility, and acceptability of Voice over Internet Protocol (VoIP)-delivered CBIT for reducing tics in children with CTDs in an open case series. Across in-person and VoIP administrations of the YGTSS, results showed mean agreement of 91%, 96%, and 95% for motor, phonic, and total tic severity subscales. In the pilot feasibility study, four children received 8-weekly sessions of CBIT via VoIP and were assessed at pre- and post-treatment by an independent evaluator. Results showed a 29.44% decrease in clinician-rated tic severity from pre to post-assessment on the YGTSS. Two of the four patients were considered treatment responders at post treatment, using Clinical Global Impressions-Improvement ratings. Therapeutic alliance, parent and child treatment satisfaction and videoconferencing satisfaction ratings were high. CBIT was considered feasible to implement via VoIP, although further testing is recommended.

Keywords

tic disorders; behavior therapy; voice over internet protocol; videoconferencing

Chronic Tic Disorders (CTDs), including Tourette Syndrome (TS) involve sudden, repetitive involuntary motor and/or vocal tics lasting for longer than one year (American Psychiatric Association, 2013). CTDs are associated with impairment in physical, psychological, social,

Direct correspondence concerning this article should be addressed to Douglas W. Woods, PhD, Texas A&M University, Department of Psychology, 4235 TAMU, College Station, TX 77843-4235; phone: (979) 845-2540; fax: (979) 845-4727; dowoods@tamu.edu.

The research reported in this publication was supported by the National Institute of Mental Health of the National Institutes of Health under Award Number F31MH096375. The content is the responsibility of the authors and is not necessarily representative of the views of the National Institutes of Health.

and family functioning and reduced quality of life (Conelea et al., 2011; Cooper, Robertson, & Livingston, 2003; Cutler, Murphy, Gilmour, & Heyman, 2009; Storch et al., 2007). Antipsychotic medications have typically been the first line intervention for CTDs (Gilbert, 2006). Although moderately effective for reducing tics, they are often associated with adverse side effects that limit their clinical use, including: weight gain, sedation, cognitive dulling, depressive symptoms, and neurological side effects (e.g., tardive dyskinesia, dystonia; Scahill et al., 2006). Behavior therapy has emerged as an effective nonpharmacological treatment option for reducing tics (Cook & Blacher, 2007; Himle, Woods, Piacentini, & Walkup, 2006), and has recently been recommended as a first line treatment for tics in European and Canadian best practice guidelines (Steeves et al., 2012; Verdellen, van de Griendt, Hartmann, Murphy, & the ESSTS Guidelines Group, 2011).

Recently, several behavioral techniques including psychoeducation, Habit Reversal Training (HRT), function-based assessment and intervention, self-monitoring, relaxation training, and behavioral rewards (Woods et al., 2008) have been combined to create a multifaceted treatment package called Comprehensive Behavioral Intervention for Tics (CBIT). CBIT was recently compared to supportive psychotherapy and education (PST) in two separate multi-site RCTs in children (Piacentini et al., 2010) and adults (Wilhelm et al., 2012). Acute treatment for both groups involved 8 sessions over 10 weeks. Results of the child study ($n = 126$) demonstrated that 52.5% of the participants in the CBIT group were acute phase treatment responders, compared to 18.5% of the PST group. There were also significantly greater reductions in clinician-rated tic severity, as measured by the Yale Global Tic Severity Scale (YGTSS) in CBIT (30.8%) relative to PST (14.2%); reductions were similar in magnitude to those found in several RCTs of medications (Scahill et al., 2013). Additionally, decreases in the YGTSS impairment scores in the CBIT group (51.2%) were greater compared to PST (29.9%). Responder status was maintained through 6-month follow-up for 87% of the acute-phase CBIT responders.

Despite the efficacy of CBIT, many families of children with CTDs are unable to access the treatment. In a recent national survey examining treatment utilization in children and adults with chronic tic disorders ($n = 672$), treatment-seeking families cited several reasons for not pursuing a behavior therapy option to treat their child's tics; including little time to attend weekly therapy, a lack of treatment providers in their region, and long distances to providers (Woods, Conelea, & Himle, 2010).

One way to increase access to treatment among underserved populations is through the use of videoconferencing (VC) technologies. The traditional VC model, such as that used by Himle et al. (2012), typically involves two clinical or academic sites with specialty VC equipment connected via a dedicated high-speed connection. Although VC has been shown to be efficacious, feasible, and acceptable when used to deliver interventions for a range of psychological disorders (e.g., Capner, 2000; Simpson, 2009), traditional VC has only recently been applied to CTDs. In an initial pilot study, Himle et al. (2010) delivered CBIT via VC to three children with TS and found that all three showed significant reductions in tic severity (Himle, Olufs, Himle, Tucker, & Woods, 2010). In a follow-up study, Himle et al. (2012) randomly assigned 20 children to receive CBIT delivered either via VC or traditional face-to-face delivery. Results showed significant pre to post reductions in clinician-rated tic

severity (YGTSS) in both groups, with mean reductions of 33% and 27% for VC and face-to-face groups, respectively. No significant differences in mean reductions were found between groups. Benchmarking results against findings in the original CBIT study, reductions in tic severity were similar when using VC format. Additionally, both treatment modalities were rated as highly acceptable by parents and children, with no significant differences between groups. There were also no significant differences in parent and child-reported therapeutic alliance between groups (Himle et al., 2012).

Despite its growing popularity, the traditional VC delivery model fails to address several important accessibility barriers. Treatment may be restricted to the locations where the equipment is installed and typically requires patients, and sometimes the therapist, to travel to a dedicated site to use the equipment. Another limitation is that the traditional VC model often requires the support of specially trained personnel at both VC sites to maintain the equipment, assist the therapist and client in connecting and using the equipment, and to provide logistical support to the patient (Hilty, Luo, Morache, & Nesbitt, 2002; Simpson, Bell, Knox, & Mitchell, 2005). Such reliance on support resources not only increases the cost of treatment, it poses logistical barriers that may limit adoption of VC to deliver care.

A newer alternative to traditional VC delivery is the use of Voice over Internet Protocol (VoIP) telephony or web-based VC (e.g., Skype®). This technology allows users to connect over the internet through software using wired or wireless high speed connections and a basic web camera. Delivering CBIT via web-based VC has several potential advantages over traditional VC delivery. First, it benefits patients, as services may be received from their own home computers, reducing costs, and potentially decreasing time missed from work and school to attend appointments. Second, a web camera is a lower cost alternative to the equipment needed for traditional VC. Third, web-based VC benefits clinicians, as they may be able to work directly from their office or be otherwise freed from the constraints posed by the physical stability of traditional VC equipment. Lastly, another advantage is that VoIP allows patients broader, direct access to experts in a particular treatment or field. Although both traditional VC and VoIP delivery provide access to treatment for those who live far away from knowledgeable treatment providers, traditional VC typically requires the patients to drive to the nearest clinic that houses the equipment for treatment.

Despite the potential of VoIP to increase access to treatment, there are potential limitations. Specifically, call quality and reliability are highly dependent on the type and speed of the internet connection (Kazemitabar, Ahmed, Said, & Habsullah, 2010) and may be influenced by certain computer specifications, including processor speed, random access memory, and hard drive disk space (Ramirez, 2011). Additionally, external technical support may not be readily available, particularly for consumers using VoIP in their homes. There is some research assessing the efficacy and feasibility of VoIP-delivered treatment, with promising findings. VoIP treatments for insomnia (Lichstein, Scogin, Thomas, DiNapoli, Dillon, & McFadden, 2013), Social Phobia (Yuen et al., 2013), OCD (Storch et al., 2011), substance abuse (King, Brooner, Peirce, Kolodner, & Kidorf, 2014), and depression in older adults (Choi et al., 2014; Choi, Wilson, Sirrianni, Marinucci, & Hegel, in press) have been associated with significant decreases in target symptoms. Additionally high satisfaction (Choi et al., 2014; King et al., 2014; Storch et al., 2011; Yuen et al., 2013) and a strong

therapeutic alliance (King et al., 2014; Lichstein et al., 2013; Yuen et al., 2013) have been observed. With respect to feasibility, patients have found the modality to be convenient, but technical difficulties (Choi et al., in press; King et al., 2014; Lichstein et al., 2013; Yuen et al., 2013) and challenges reading body language (Lichstein et al., 2013; Storch et al., 2011) are common. Despite preliminary evidence, research on the efficacy and feasibility of VoIP-delivered therapy is still rather limited; and the validity and reliability of administering clinician-rated measures via this modality have yet to be assessed. Therefore two studies were performed as part of research testing the feasibility of using VoIP to assess and treat CTD in children. To gain preliminary data on the feasibility of assessing tic severity over VoIP, Study 1 assessed the interrater agreement between in-person and VoIP-administrations of a primary tic rating scale, the Yale Global Tic Severity Scale (YGTSS; Leckman et al., 1989). In Study 2, the feasibility, acceptability, and preliminary efficacy of VoIP-delivered behavior therapy for CTDs were evaluated.

Study 1

Method

Participants—Participants were 10 former and current patients of a university-based Tic Disorders Specialty Clinic recruited through completed and active studies. Participant ages ranged from 10 to 14 years ($M = 12.39$, $SD = 1.36$). Nine participants were male, and one was female. All participants were identified as Caucasian. Inclusion criteria included (a) residence in the clinic's home state (b) an age between 9 and 17, (c) DSM-IV-TR diagnosis of CTD (Chronic Motor/Vocal Tic Disorder or Tourette Syndrome), and (d) fluency in English. Exclusion criteria included (a) a DSM-IV-TR substance abuse or dependence of conduct disorder diagnosis within the past 3 months, (b) a lifetime DSM-IV-TR diagnosis of pervasive developmental disorder, mania, or psychotic disorder, (c) any serious psychiatric, psychosocial, or neurological condition requiring urgent treatment beyond that provided through the study, and (d) lack of a functional and accessible home computer, and high speed (i.e., cable/DSL) internet connection.

Materials—Skype[®] was used to perform web-based VC. Skype[®], owned by Microsoft Corporation[®], is a peer-to-peer VoIP system, providing free video and voice calling, instant messaging, and file sharing between users (Skype Technologies S. A., 2013). Skype[®] was chosen due to its popularity among consumers, and ease of use (Garfinkel, 2005). Treatment was delivered from a private secure clinic room, using a Dell[®] Optiplex GX 980 desktop computer with a 21.5 in. screen, Logitech[®] c270 web camera, and a high speed (54.0 megabytes per second) wireless local area network internet connection available through the university. The Skype[®] picture-in-picture function was used at all times so the therapist could monitor the participant's movements and body positioning. Participants used their home desktop or laptop computer, high speed internet connection, and a web camera to connect with the therapist. A low-cost Logitech[®] c110 web camera was loaned to any family who did not already own one.

Measures—*Yale Global Tic Severity Scale* (YGTSS; Leckman et al., 1989), a clinician-rated tic severity measure, was administered. It allows for the assessment of tics using five

categories: number, frequency, intensity, complexity, and impairment. Each item is rated on a scale of 0 to 5. The measure has independent subscales for motor and vocal tic severity, each ranging from 0 to 25, an overall impairment scale ranging from 0 to 50, and an overall tic severity score: the sum of the motor and vocal subscales. When delivered face-to-face, the YGTSS has demonstrated acceptable psychometric properties, including good construct validity, internal consistency, and interrater reliability; and acceptable convergent and divergent validity (Leckman et al., 1989; Storch et al., 2005).

Procedure—A brief phone screen was conducted with interested participants to gauge study eligibility. A study evaluator, trained in the administration of the YGTSS, then traveled to the homes of eligible participants to complete the study procedures. During the visits, the evaluator sought parental consent and child assent using Institutional Review Board-approved forms, which highlighted risks of participation including, but not limited to, protecting session content from being overheard by family members and others within the home, and maintaining internet network security and privacy during videoconferencing. The evaluator also assisted family members to download and set-up the Skype® software and to install a web camera for use during the study visit. Families were then connected via Skype® to a primary rater located in a private room in the Clinic. The primary rater, also trained to perform the YGTSS, administered the interview via web-based VC. The evaluator in the home, who was off-camera, served as the secondary rater. They observed the interview and independently rated the child's tics from within the families' homes. Participants were paid \$25 for completion of study procedures.

Results

The interrater agreement between in-person and VoIP administrations of the YGTSS was calculated by dividing lower total score by the higher total score for each subject. Across all 10 participants the average agreement for the YGTSS motor subtotal was 91% (range = 71% - 100%). The YGTSS vocal tic subtotal average was 96% (range = 85% to 100%), and the total score (motor and vocal subtotals) was 96% (range = 88% to 100%).

Study 2

Methods

Participants—Participants were four Caucasian males, between the ages of 10 and 13, diagnosed with TS (see Table 1), who resided within 83.5 miles of the clinic on average. They were selected from the Study 1 sample, as they were the only four participants who had not yet received behavior therapy. Inclusion criteria were (a) residence in the Clinic's home state (b) age between 9 and 17, (c) DSM-IV-TR diagnosis of CTD (Chronic Motor/Vocal Tic Disorder or Tourette Syndrome), (d) a Clinical Global Impressions – Severity score 4 (moderately ill or worse), (e) a YGTSS Score 14 and < 30 for TS or 10 and < 20 for CMTD, (f) medication free or on a stable medication for the treatment of tics, OCD, ADHD, anxiety, and/or depressive disorder for at least 6 weeks, with no planned changes during the study period, and (g) fluency in English. Exclusion criteria were (a) a total YGTSS tic score > 30, considered high severity, selected to remain consistent with other behavior therapy exclusion criteria; for YGTSS scores > 30, a caseness panel discussed

additional treatment options with the family, and determined the appropriateness of the patient's participation in the study), (b) a Wechsler Abbreviated Scale of Intelligence – Vocabulary (The Psychological Corporation, 1999) subtest T-score < 37 (low average range), (c) a DSM-IV-TR diagnosis of substance abuse or dependence, or conduct disorder within the past 3 months, (d) a lifetime DSM-IV-TR diagnosis of pervasive developmental disorder, mania, or psychotic disorder, (e) any serious psychiatric, psychosocial, or neurological condition requiring urgent treatment beyond that provided through the study, (f) four or more previous HRT sessions for TS, (g) lack of a functional and accessible home computer, and/or high speed (i.e., cable/DSL) internet connection, and (h) refusal to sign a release of information form for the child's local primary care physician, mental health professional, or neurologist.

Materials—For a detailed description of computer equipment used by the treatment site and participants during the study period see Table 2.

Measures

Mini International Neuropsychiatric Interview for Schizophrenia and Psychotic Disorders Studies for Children and Adolescents (MINI-Kid; Sheehan et al., 1998). The MINI-Kid is a brief structured clinician-rated diagnostic interview assessing for 27 psychiatric disorders in children and adolescents. The interview has adequate to excellent interrater and test-retest reliability and good to excellent convergent validity for most diagnoses (Sheehan et al., 2010).

Wechsler Abbreviated Scale of Intelligence (WASI; The Psychological Corporation, 1999). The WASI is a psychometrically acceptable (Axelrod, 2002; Canivez, Konold, Collins, & Wilson, 2009; Saklofske, Caravan, & Schwartz, 2000) measure of intellectual functioning for individuals ages 6 to 89 years. The vocabulary subtest was administered during the screening assessment.

YGTSS (Leckman et al., 1989). The YGTSS, described in Study 1, was administered by an independent evaluator (IE) via web-based VC.

Clinical Global Impression-Severity and Improvement Scales (CGI-S & CGI-I; Guy, 1976). The CGI-S and CGI-I are clinician-rated scales assessing global severity (-S) and improvement (-I) in the patient since the baseline assessment. Clinicians rate the patient's severity using the following anchors: 0 (not assessed), 1 (normal, not at all ill), 2 (borderline ill), 3 (mildly ill), 4 (moderately ill), 5 (markedly ill), 6 (severely ill), and 7 (extremely ill). Clinicians assess the patient's improvement using the following ratings, which were slightly modified from the original scale: 0 (not assessed), 1 (very much improved), 2 (much improved), 3 (improved), 4 (minimally improved), 5 (no change), 6 (minimally worse), 7 (much worse), and 8 (very much worse). The scales have good concurrent validity and are sensitive to change (Leon et al., 1993).

Child's Perception of Therapeutic Relationship (Kendall et al., 1997). The Child's Perception of Therapeutic Relationship scale is a 10-item measure, assessing the child's

perceptions of the quality of the therapeutic relationship. Total scores range from 8 to 40, with higher scores reflecting a more positive perception of the therapeutic relationship. The instrument displays adequate internal consistency and reliability (Elvins & Green, 2008).

Treatment Acceptability Questionnaire (TAQ; Hunsley, 1992). The TAQ is a 6-item measure assessing patient and parent reactions to treatment. Items covering acceptability of the treatment, side effects, ethics and effectiveness of the treatment, and knowledge and trustworthiness of the treatment provider were rated by parents on a likert scale of 1 to 7. Total scores on the measure range from 7 to 42, with higher scores reflecting greater acceptance. The questionnaire displays good internal consistency and test-retest reliability, and some indication of concurrent validity (Hunsley, 1992).

Client Satisfaction Questionnaire (CSQ; Larsen, Attkisson, Hargreaves, & Nguyen, 1979). The CSQ is an 8-item measure designed to assess patient satisfaction with healthcare service quality and type, the extent to which the service met patient needs, willingness to recommend the service to others, satisfaction with the degree of help, the extent to which the service helped the patient manage problems more effectively, overall satisfaction, and willingness to return for future services. Individual items are rated on a scale of 1 to 4, with total scores ranging from 8 (low satisfaction) to 32 (high satisfaction). The CSQ possesses high internal consistency (Cox, Brown, Peterson, & Rowe, 1982; Larsen et al., 1979; Roberts & Attkisson, 1983), and excellent concurrent validity (Nguyen, Attkisson, & Steggner, 1983; Larsen et al., 1979).

Videoconferencing Satisfaction Questionnaire. This 12-item parent and child-report measure assesses satisfaction with the VC aspect of treatment, including such domains as the therapeutic relationship and therapist competence, comfort using the equipment, audio and visual quality, the ability of VC to address treatment barriers, and preference for VC relative to in-person treatment. It was adapted from the Telemedicine Satisfaction Questionnaire (Nelson, Barnard, & Cain, 2003). Items are rated on a likert scale of 1 (strongly disagree) to 7 (strongly agree), with total scores ranging from 7 (low VC satisfaction) to 84 (high VC satisfaction).

VoIP Call Quality Ratings. The severity and type of technical difficulties during VC treatment and assessment sessions were tracked during treatment sessions, using questions adapted from Yuen (2010). Ratings were qualitative, and no numerical values were assigned to the descriptors. Severity ratings included: 'None', 'Insignificant' (session quality not affected), 'Minimal' (session quality minimally affected), 'Moderate' (session quality moderately affected), 'Major' (session quality majorly affected), and 'Severe' (could not complete the session). Staff who endorsed the presence of technical difficulties were asked to indicate the type of technical difficulties by selecting from four categories: sound quality (e.g., choppy, echoing, soft, delayed), video quality (e.g., blurry, freezing, choppy, delayed, dropped calls), unable to hear sound, and unable to see video. All relevant types of technical difficulties could be indicated for each session.

Procedure—Participants were recruited through a Tic Disorders Specialty Clinic. Six participants were screened via the phone to gauge eligibility. All were eligible, but two

declined to participate. The first declined due to concerns that receiving treatment via VoIP would not be hands on enough compared to in-person treatment, and the second declined due to a lack of interest in receiving treatment within the context of a research study. Those agreeing to participate were mailed packets including Institutional Review Board-approved consent and assent forms (which addressed home and internet privacy issues like those used in study 1), in addition to standard psychology clinic forms (i.e., consent for treatment, clinic background form, acknowledgement of receipt of privacy practices, email permission), instructions for downloading Skype®, and guidelines for maintaining privacy when using web-based VC. Consent forms were then reviewed and discussed via telephone with interested participants. Also, a web camera was mailed to any participant who did not already own one, for use during the study period. Next, a baseline assessment was conducted by an IE via Skype® using the measures describe above. The IE was a Master's level doctoral student who received training and supervision in administration of the clinician-rated assessments. The purpose of the evaluation was to (a) confirm eligibility, (b) determine initial tic severity, (c) establish the presence of comorbid conditions, and (d) document the type of VC and computer equipment being used by participants for the study. After eligibility was confirmed, a CBIT parent workbook (Woods et al., 2008) was mailed to families.

Treatment was provided by two Master's level therapists each with 2 years' experience performing CBIT. Therapists were trained and supervised according to procedures in the original CBIT trial (Piacentini, 2010). Treatment consisted of 8 weekly sessions, and included several components. Families followed along with the protocol using a parent workbook, listing treatment procedures, homework assignments, and blank homework sheets. Treatment was conducted with both the child and parent together, with flexibility allowed for older adolescents.

In session 1, the therapist provided psychoeducation about tics, created a hierarchical list of tics to treat, discussed tic-related impairment, and formed a behavioral reward program wherein the patient received points towards rewards for participating in treatment. In session 2 function-based assessment and interventions, and HRT were conducted for the first tic on the hierarchy (typically the most bothersome tic). Function-based strategies involved identifying and minimizing any antecedents and consequences to ticcing associated with higher tic severity; HRT involved awareness training, competing response training, and social support. In awareness training, the child was taught to identify when tics or premonitory sensations occurred. This was done by having the child operationally define the tic and any premonitory urges, and instructing the child to practice noticing the tics as they occurred in session. In competing response training, the child selected a behavior that was (a) physically incompatible with the tic, (b) able to be held for 1 minute or until the urge to tic had subsided, and (c) socially discreet. This "competing response" was then done for one minute or until the urge dissipated contingent upon the occurrence of the tic or urge/signal to tic. In the social support component, parents were trained to provide prompts and praise for use of competing responses. As in session 2, the therapist targeted a new tic using function-based interventions and HRT each week in sessions 3 through 8. Sessions 4 and 5 featured instruction in relaxation techniques, including relaxed breathing, and progressive muscle relaxation. Sessions 7 and 8 included a discussion of relapse prevention strategies.

Between sessions, patients and parents completed tic-monitoring homework, wherein 3 to 4 times per week, they independently tallied each occurrence of a targeted tic during a 15–20 minute interval. Self-monitoring homework was reviewed at the beginning of each session by having the parent and child read it to the therapist from the workbook. In between sessions, parents supported their children by reminding them to use the competing response when they noticed the child had forgotten, and praising the child when they saw that the child had correctly used a competing response contingent on tic occurrence.

Following treatment, the IE performed a post-assessment (i.e., YGTSS, CGI scales) via Skype®. Also, participants and parents were prompted via e-mail to complete internet questionnaires (i.e., Child's Perception of Therapeutic Relationship, TAQ, CSQ, and Videoconferencing Satisfaction Questionnaire) on a secure internet surveying website (Qualtrics Labs, Inc. software, 2013). Participants were reimbursed a total of \$100 for completion of both baseline and post-treatment assessments.

Results

Andrew¹—Andrew was a 13 year old male diagnosed with TS and ADHD-Inattentive type. At the time of the study, he was taking several medications for TS and ADHD, including Vyvanse, Sertraline, and Clonidine. His WASI-vocabulary T-score was 73, which fell in the very superior range. At baseline, Andrew received a YGTSS total tic severity score of 32, with tic-related impairment falling in the moderate range, and a CGI-S score of 5 (markedly ill; see Table 3). Andrew entered treatment with several tics, including: eye blinking, squinting, nostril flaring, ear wiggling, toe cracking, coughing, throat clearing, sniffing, repeating words, and emphasizing certain syllables in words. There were several tics he repeated a certain number of times, including making a half smile, knuckle cracking, retracing over his own writing, evening up, hyperextending his elbow, wrist movements, and shoulder shrugs. He experienced tic-related impairment in the form of feelings of embarrassment and stress, and interference in sports. At the final assessment, his total YGTSS score had decreased by 6 points to a score of 26, and he endorsed tic-related impairment in the minimal range. His CGI-S score was 4 (moderately ill), and his CGI-I score was 3 (improved).

During treatment, a few challenges occurred. Andrew's most bothersome tics, which were more complex, did not occur during session. Andrew became irritated by his mother's use of verbal prompts to perform competing responses, as his younger siblings began to mimic her in his presence. This was temporarily resolved when his mother used non-verbal prompts instead. Of note, is that during half of the sessions, the patient's mother was nearby the computer, but did not join the session until the last 10 to 15 minutes, due to the need to care for Andrew's younger siblings. She had reported that she enjoyed joining the session at the end for a recap, as she felt it made her son feel more mature to not have a parent sitting over his shoulder while doing therapy.

Ben—Ben was a 13 year old male with TS. He also met criteria for ADHD-Inattentive type, and his WASI-vocabulary T-Score was 43 (average). His YGTSS total severity score at baseline was 22, his tic-related impairment score fell in the mild range, and his CGI-S score

was 4, indicating moderate symptoms. Four tics were targeted, including sniffing, a neck movement, holding his breath, and eye blinking/darting. Ben's tics were not particularly bothersome to him. His mother attributed this to a lack of awareness of the frequency with which he performed them; however, she perceived Ben's sniffing tic to be annoying, as it was frequent. Ben noted his tics were worse when he was stressed, bored, or tired. Allergies exacerbated his sniffing tic, and it was discovered that his tics occurred more often when they were being discussed. During the final assessment he received a YGTSS total tic score decreased to 14, and endorsed no tic-related impairment. His CGI-S score was 3 (mildly ill) and his CGI-I score was 1 (very much improved).

Due to the mild and infrequent nature of the tics, performing and conveying the importance of function-based assessment and intervention was difficult. During sessions, Ben sat at the computer with his mother or father sitting behind him in the background.

Craig—Craig was an 11-year old male, who resided with his parents, twin sister, and older sister. He met criteria for TS based on the MINI-Kid interview, and reported a history of reading difficulties. His WASI-vocabulary score was 52 (average). His YGTSS total tic severity score was 20, his tic-related impairment fell in the minimal range, and his CGI-S score was 4, indicating moderate illness severity. He reported some tic-related impairment in the form of personal annoyance from having tics and receiving unwanted questions from peers. Craig entered treatment with several tics, including jaw movements, turning his head upwards, skipping then tapping his hip with his palm, shoulder roll, throat clearing, sniffing with a nose scrunch, and spinning his whole body to either side. He indicated that head turning upwards happened more often when outside on sunny days, and the skipping-tapping tic occurred more frequently when he was inside his home, walking around or washing his hands. At the post-assessment his YGTSS total tic score increased to 22, and his tic-related impairment remained in the minimal range throughout the course of treatment. He received a CGI-S score of 4 (moderately ill), and a CGI-I score of 5 (no change).

Several issues occurred in treatment. Homework compliance was fairly poor. Also, Craig often fidgeted and had a playful demeanor in sessions. In the initial sessions he played with the web camera frequently, rotating it so that he and his mother shifted in and out of viewing range, and turning the video feed on and off. This behavior steadily decreased, especially after he was asked to bring a fidget toy to session. He also enjoyed using the Skype® instant messaging feature randomly to chat during sessions which was somewhat distracting. Treatment ended after the fourth session, as it was disrupted due to a virus on the family computer. The problem took several weeks to be properly fixed, then the family was lost to contact for several more weeks.

David—David was a 10 year old male, residing with his parents, twin brother, and older brother. He met criteria for TS, based on the MINI-Kid interview, and his parents also indicated he had mild anxiety. His WASI-vocabulary score was 56 (average). At baseline his YGTSS total tic severity score was 27, and his tic-related impairment was moderate. He received a CGI-S score of 5. His tics, which occurred most often during school hours, included a throat noise, head nod, hard wink, quick breath, and squeak. His head nod caused physical pain and interrupted activities, his vocal tics interfered with speaking, and he

received unwanted attention and questions about his tics from peers. Overall, most tics were reduced from the beginning of treatment, with the exception of the quick breath tic. David also reported less distress and fewer tic hassles. At the post-assessment, his YGTSS total tic score had decreased to 13, and he endorsed no tic-related impairment. His CGI-S score was 3 (mildly ill), and his CGI-I score was 1 (very much improved).

Treatment raised some issues. It was difficult to hear David's vocal tics over VoIP, making awareness training and competing response training challenging. However, his mother helped to identify tic occurrence during awareness training, and prompted and praised David's use of physically incompatible responses during competing response training. Additionally, David's siblings occasionally interrupted sessions, as they were curious about what was happening. During one session the family conducted the session from David's bedroom, which was unusual. In this case, David attended sessions with his mother or both parents.

Therapeutic Alliance and Acceptability—Averaged across all four participants, the mean patient rating for the Child's Perception of Therapeutic Relationship scale was 32.5 out of 40 ($SD = 1.73$; range = 31–35), suggesting a strong therapeutic alliance and was similar to that found among patients receiving VC-delivered CBIT (Himle et al., 2012). The mean parent treatment acceptability (TAQ) rating was high, with an average of 39 out of 42 ($SD = 2.94$; range = 35–42), and was also similar to the mean parent TAQ rating (35.5) found among those receiving VC-delivered CBIT (Himle et al., 2012). Parental treatment satisfaction (CSQ) ratings were strong, with an average of 29.25/32 ($SD = 2.75$; range = 26–32), as were child treatment satisfaction (CSQ) ratings ($M = 30.75/32$; $SD = 1.50$; range = 29–32). In regard to satisfaction with VC, the mean patient and parent ratings were high, with respective scores of 73 ($SD = 5.35$; range = 68–79), and 74.25 ($SD = 5.74$; range = 67–81) out of 84.

VoIP Call Quality—In 71% of the 28 total treatment sessions across participants, no technical difficulties occurred. 'Insignificant' technical difficulties occurred in 17.9% of sessions, 'Minimal' technical difficulties occurred in 7.1% of sessions, and 'Moderate' difficulties occurred in 3.6% of sessions. In regard to type of technical difficulties, 21.4% were related to sound quality (excluding complete inability to hear sound), and 10.7% were related to video quality (excluding complete inability to see video feed). In no treatment sessions were staff completely unable to see video feed or hear sound.

Discussion

In Study 1, the interrater agreement between in-person and VoIP administrations of the YGTSS indicated high reliability between modalities. In Study 2, the preliminary efficacy, feasibility, and acceptability of VoIP-delivered CBIT were assessed. Results showed a 29.44% average decrease in YGTSS tic severity from pre to post, which is similar to the 31% decrease found in the original CBIT trial in children (Piacentini et al., 2010) and in line with the respective ratings of 33% and 27% found in both VC and in-person modalities in the randomized comparison by Himle et al. (2012). Two of the four participants (50%) were classified as treatment responders at post-treatment. This is similar to the response rate

(52.5%) in CBIT delivered face-to-face (Piacentini et al., 2010), however, as this was a case series with a small sample size, comparisons should be made cautiously. The therapeutic alliance and patient and parent satisfaction ratings were high, and the intervention was feasible with respect to technical issues, as 71% of therapy sessions were free of technical difficulties, and 75% of assessment sessions had 'none' or 'insignificant' technical difficulties.

Despite the promising findings, several notable issues emerged during the VoIP delivery of CBIT. First, one participant's family computer was infected with a virus just prior to the fifth treatment session. After taking the computer to be fixed, the family was still unable to use Skype®, and needed to take the computer in a second time. Following this, the family was lost to contact for several weeks, and treatment was not reinitiated. Second, variable sound quality occasionally interfered with treatment delivery, especially during awareness training, competing response training, and relaxation training. During awareness training, it was sometimes challenging to bring certain tics to the child's attention, since they were more difficult to hear. Third, child engagement and sustained attention was occasionally difficult to manage during sessions. If a child is off-task during a session, the clinician may feel less control over the situation over VoIP than in a face-to-face session. Although this could suggest externalizing diagnoses like ADHD may moderate treatment response, of the two participants who had a comorbid diagnosis of ADHD, one was classified as a responder, with a rating of 'very much improved' at post-treatment, and the other was not but still received a rating of 'improved'.

A fourth issue involved the review of homework at each session. Parents and patients read aloud self-monitoring homework to the therapist at the beginning of each session, and the therapist took notes. This is in contrast to face-to-face sessions, in which the therapist is able to view homework first hand. A fifth issue involved the restricted range of view of the patient, due to the equipment set-up (i.e., the computer placed on top of a table, desk, or cart). In contrast, traditional VC formats using dedicated VC systems allow for full view of the body or remote control of movable cameras. In the current VoIP model, therapists only saw the patient's face and top-half of their body. This made addressing tics involving the lower extremities more challenging. A related difficulty involved eye contact. For one therapist, giving the appearance of making eye contact with the patient often required looking into the web camera instead of at the computer screen; however doing so made it challenging for the therapist to see what was on screen.

A sixth issue was reduced parental participation during sessions in some cases. Despite parents being present for all sessions, their participation levels varied. Reasons are unclear, but it is possible that the nature of in-home VC-delivered treatment allows parents to multi-task during sessions. Additionally, receiving treatment in the home can invite interruptions by other family members.

Based on these issues, the following are suggestions for improving treatment quality for home-based VC. For therapists who need to track weekly progress using a self-report symptom questionnaire, it may be helpful to have parents return such documents as email attachments in order to preserve document formatting for easy viewing and scoring. To

enhance awareness training and competing response training, it may be beneficial to utilize parents to identify tics during sessions, so the therapist can call them to the child's attention and remind the child to use the competing response. To address the problem of restricted view of tics, it may be helpful to begin treating a facial tic or other tic involving the upper body first, so the child develops a routine before working on a tic involving the lower extremities. Asking the child to back up from the computer to observe tics or to see if the competing response is being performed correctly can also be helpful. It is also important to remind parents to sit in front of the computer with their child during sessions and avoid multitasking, especially if their child is young. This should also aid children who have difficulty sustaining attention.

Solving eye contact issues is challenging, but it may be helpful if the therapist sits in a chair that allows their eyes to be aligned with or above the web camera so that looking down at the screen will be less noticeable, or sits far enough away from the camera, thus decreasing the ability of the patient to discriminate subtle differences in direction of gaze. If the therapist is unable to have such a set-up then they should focus on viewing the patients on screen, and only occasionally look up into the web camera. Explaining this issue to the patient, prior to treatment, would be beneficial.

The study also raises some ethical considerations. Delivering treatment while patients are in a home setting introduces potential privacy issues with respect to family members interrupting sessions and hearing content. Additionally, maintaining internet network security and privacy during VoIP sessions is a potential concern. Prior to study participation, families were made aware of these issues in the consent process, and received VC guidelines to address these concerns (e.g., conducting sessions in a computer room or room with closeable doors, logging off from Skype[®] following each session, using a firewall or antivirus program, etc.); however, these issues do remain limitations. Furthermore, a general concern regarding distance treatments involves the handling of psychiatric emergencies. In the present study, no psychiatric emergencies occurred, and this issue was addressed preemptively by having families provide contact information and sign a release of information for a local health care provider in the event of an emergency prior to beginning treatment.

Other general design limitations also warrant mention. One major limitation involves the open trial design. A second is that this was a pilot study with four subjects limiting the generality of our conclusions. Another limitation is that one of the six families phone screened for eligibility declined participation because they deemed the treatment modality unacceptable. Also, the need for a computer and high speed internet access may be considered a limitation, as it may be perceived as displaying a bias towards families of higher socioeconomic status. Finally, we did not include a follow-up assessment so we are unable to make conclusions regarding maintenance of gains.

Generally, web-based CBIT was effective and feasible and has the potential to further increase access to behavior therapy for families. Despite the positive outcomes, use of VoIP for therapy purposes may pose some logistical barriers with respect to policy issues for some clinics, depending on the institutional setting. Additionally, use of VoIP-delivered treatment

has not been addressed within most States' licensing laws. Future studies should assess the efficacy of web-based VC delivery of CBIT relative to a waitlist or active control group. It would also be interesting to compare it to traditional VC delivery and face-to-face delivery in terms of efficacy and acceptability.

References

- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th edition). Washington, DC: Author. doi.org/10.1176/appi.books.9780890425596.893619
- Axelrod BN (2002). Validity of the Wechsler Abbreviated Scale of Intelligence and other very short forms of estimating intellectual functioning. *Assessment*, 9, 17–23. doi: 10.1177/1073191102009001003 [PubMed: 11911230]
- Canivez GL, Konold TR, Collins JM, & Wilson G (2009). Construct validity of the Wechsler Abbreviated Scale of Intelligence and Wide Range Intelligence Test: Convergent and structural validity. *School Psychology Quarterly*, 24, 252–265. doi:10.1037/a0018030
- Capner M (2000). Videoconferencing in the provision of psychological services at a distance. *Journal of Telemedicine and Telecare*, 6, 311–319. doi:10.1258/1357633001935969 [PubMed: 11265099]
- Choi NG, Hegel MT, Marti CM, Marinucci ML, Sirrianni L, & Bruce ML (2014). Telehealth problem-solving therapy for depressed low-income homebound older adults *American Journal of Geriatric Psychiatry*. doi:10.1016/j.jagp.2013.01.037
- Choi NG, Wilson NL, Sirrianni L, Marinucci ML, & Hegel MT (in press). Acceptance of home-based telehealth problem-solving therapy for depressed, low-income homebound older adults: Qualitative interviews with the participants and aging-service case managers. *The Gerontologist*.
- Conelea CA, Woods DW, Zinner SH, Budman C, Murphy T, Scahill LD, ...Walkup, T (2011). Exploring the impact of Chronic Tic Disorders on youth: Results from the Tourette Syndrome impact survey. *Child Psychiatry and Human Development*, 42, 219–242. doi:10.1007/s10578-010-0211-4 [PubMed: 21046225]
- Cooper C, Robertson MM, & Livingston G (2003). Psychological morbidity and caregiver burden in parents of children with Tourette's Disorder and psychiatric comorbidity. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42, 1370–1375. doi:10.1097/01.CHI.0000085751.71002.48 [PubMed: 14566175]
- Cook CR, & Blacher J (2007). Evidence-based psychosocial treatments for tic disorders. *Clinical Psychology: Science & Practice*, 14, 252–267. doi:10.1111/j.14682850.2007.00085.x
- Cox JB, Brown TR, Peterson PD, & Rowe MM (1982). A report on a statewide community mental health center outcome study. *Community Mental Health Journal*, 18, 135–150. [PubMed: 7128052]
- Cutler D, Murphy T, Gilmour J, & Heyman I (2009). The quality of life of young people with Tourette syndrome. *Child: Care, Health, and Development*, 35, 496–504. doi:10.1111/j.1365-2214.2009.00983.x
- Elvins R, & Green J (2008). The conceptualization and measurement of therapeutic alliance: An empirical review. *Clinical Psychology Review*, 28, 1167–1187. doi:10.1016/j.cpr.2008.04.002 [PubMed: 18538907]
- Frueh BC, Monnier J, Yim E, Grubaugh AL, Hammer MB, & Knapp RG (2007). A randomized trial of telepsychiatry for post-traumatic stress disorder. *Journal of Telemedicine and Telecare*, 13, 142–147. doi: 10.1258/135763307780677604 [PubMed: 17519056]
- Garfinkel, S. L. (2005). *VoIP and Skype security*. Retrieved February 25, 2011, from <http://skypetips.internetvisitation.org/files/VoIP%20and%20Skype.pdf>
- Gilbert D (2006). Treatment of children and adolescents with tics and Tourette Syndrome. *Journal of Child Neurology*, 21, 690–700. doi:10.1177/08830738060210080401 [PubMed: 16970870]
- Guy W (1976). *ECDEU assessment manual for psychopharmacology* Rev. Rockville, Md: National Institute of Mental Health (DHEW publication no. (ADM) 76–338).

- Hilty DM, Luo JS, Morache C, Marcelo DA, & Nesbitt TS (2002). Telepsychiatry: An overview for psychiatrists. *CNS Drugs*, 16, 527–548. doi:10.2165/00023210-200216080-00003 [PubMed: 12096934]
- Himle MB, Freitag M, Walther M, Franklin SA, Ely L, & Woods DW (2012). A randomized pilot trial comparing videoconference versus face-to-face delivery of behavior therapy for childhood tic disorders. *Behaviour Research and Therapy*, 50, 565–570. doi:10.1016/j.brat.2012.05.009 [PubMed: 22743661]
- Himle MB, Olufs E, Himle J, Tucker BTP, & Woods DW (2010). Behavior therapy for tics via videoconference delivery: An initial pilot test in children. *Cognitive and Behavioral Practice*, 17, 329–337. doi:10.1016/j.cbpra.2010.02.006
- Himle MB, Woods DW, Piacentini J, & Walkup J (2006). A brief review of habit reversal training for Tourette syndrome. *Journal of Child Neurology*, 21, 719–725. doi: 10.1177/08830738060210080101 [PubMed: 16970874]
- Hunsley J (1992). Development of the treatment acceptability questionnaire. *Journal of Psychopathology and Behavioral Assessment*, 14, 55–64. doi:10.1007/BF00960091
- Kazemitabar H, Ahmed S, Nisar K, Said AB, & Habsullah HB (2010). A comprehensive review on VoIP over wireless LAN networks. *Computer Science Letters*, 2
- Kendall PC, Flannery-Schroeder E, Panichelli-Mindel SM, Southam-Gerow M, Henin A, & Warman M (1997). Therapy for youths with anxiety disorders: A second randomized clinical trial. *Journal of Consulting and Clinical Psychology*, 65, 366–380. [PubMed: 9170760]
- King VL, Brooner RK, Peirce JM, Kolodner K, & Kidorf MS (2014). Randomized trial of web-based videoconferencing for substance abuse counseling. *Journal of Substance Abuse Treatment*, 46, 36–42. doi:10.1016/j.jsat.2013.08.009 [PubMed: 24035556]
- Larsen DL, Attkisson C, Hargreaves WA, & Nguyen TD (1979). Assessment of client/patient satisfaction: Development of a general scale. *Evaluation and Program Planning*, 2, 197–207. doi: 10.1016/0149-7189(79)90094-6 [PubMed: 10245370]
- Leckman JF, Riddle MA, Hardin MT, Ort SI, Swartz KL, Stevenson J, & Cohen DJ (1989). The Yale Global Tic Severity Scale: Initial testing of a clinician-rated scale of tic severity. *Journal of the American Academy of Child & Adolescent Psychiatry*, 28, 566–573. doi: 10.1097/00004583-198907000-00015 [PubMed: 2768151]
- Leon AC, Shear MK, Klerman GL, Portera L, Rosenbaum JF, & Goldenberg I (1993). A comparison of symptom determinants of patient and clinician global ratings in patients with panic disorder and depression. *Journal of Clinical Psychopharmacology*, 13, 327–331. doi: 10.1097/00004714-199310000-00005 [PubMed: 8227491]
- Lichstein KL, Scogin F, Thomas J DiNapoli EA, Dillon HR, & McFadden A (2013). Telehealth cognitive behavior therapy for co-occurring insomnia and depression symptoms in older adults. *Journal of Clinical Psychology*, 69, 1056–1065. doi: 10.1002/jclp.22030 [PubMed: 24014056]
- Nelson E, Barnard M, & Cain S (2003). Treating childhood depression over videoconferencing. *Telemedicine Journal and e-Health*, 9, 49–55. doi:10.1089/153056203763317648 [PubMed: 12699607]
- Nguyen TD, Attkisson CC, & Stegner BL (1983). Assessment of patient satisfaction: Development and refinement of a Service Evaluation Questionnaire. *Evaluation and Program Planning*, 6, 299–314. doi:10.1016/0149-7189(83)90010-1 [PubMed: 10267258]
- Piacentini J, Woods DW, Scahill L, Wilhelm S, Peterson AL, Chang S, ... Walkup, JT (2010). Behavior therapy for children with Tourette Disorder: A randomized controlled trial. *The Journal of the American Medical Association*, 303, 1929–1937. doi:10.1001/jama.2010.607 [PubMed: 20483969]
- The Psychological Corporation. (1999). Wechsler Abbreviated Scale of Intelligence. San Antonio, TX: Author.
- Qualtrics Labs, Inc. software (2013). Version 47320 of the Qualtrics Research Suite. Provo, UT: Qualtrics Labs, Inc. <http://www.qualtrics.com>
- Ramirez, M. (2011, April). Communication in “the cloud”: Skype, GoogleTalk, and Google Voice. *Technology*. Retrieved from <http://digitalcommons.calpoly.edu>

- Roberts RE, & Attkisson CC (1983). Assessing client satisfaction among Hispanics. *Evaluation and Program Planning*, 6, 401–403. doi:10.1016/0149-7189(83)90019-8 [PubMed: 10267267]
- Saklofske DH, Caravan G, & Schwartz C (2000). Concurrent validity of the Wechsler Abbreviated Scale of Intelligence (WASI) with a sample of Canadian children. *Canadian Journal of School Psychology*, 16, 87–94. doi:10.1177/082957350001600106
- Scahill L, Erenberg G, Berlin CM, Budman C, Coffey BJ, Jankovic J, ... Tourette Syndrome Association Medical Advisory Board: Practice Committee. (2006). Contemporary assessment and pharmacotherapy of Tourette syndrome. *NeuroRx*, 3, 192–206. doi:10.1016/j.nurx.2006.01.009 [PubMed: 16554257]
- Scahill L, Woods DW, Himle MB, Peterson AL, Wilhelm S, Piacentini J, ... Mink, JW. (2013). Current controversies on the role of behavior therapy in Tourette Syndrome. *Movement Disorders*, 28, 1179–1183. doi:10.1002/mds.25488 [PubMed: 23681719]
- Sheehan DV, Lecubrier Y, Sheehan KH, Amorim P, Janavs J Weiller, E, ... Dunbar, GC. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59 Suppl 20, 22–33.
- Sheehan DV, Sheehan KH, Shytle RD, Janavs J, Bannon Y, Rogers J, ...Wilkinson B (2010). Reliability and validity of the Mini International Neuropsychiatric Interview for Children and Adolescents (MINI-Kid). *The Journal of Clinical Psychiatry*, 71, 313–326. doi:10.4088/JCP.09m05305whi [PubMed: 20331933]
- Simpson S (2009). Psychotherapy via videoconferencing: A review. *British Journal of Guidance & Counseling*, 37, 271–286. doi:10.1080/03069880902957007
- Simpson S, Bell L, Knox J, & Mitchell D (2005). Therapy via videoconferencing: A route to client empowerment? *Clinical Psychology and Psychotherapy*, 12, 156–165. doi:10.1002/cpp.436
- Skype Technologies S. A. (2013). *What is Skype?* Retrieved from <http://about.skype.com/>
- Steeves T, McKinlay BD, Gorman D, Billingham L, Day L, Carroll A, ...Pringsheim, T (2012). Canadian guidelines for the evidence-based treatment of tic disorders: behavioural therapy, deep brain stimulation, and transcranial magnetic stimulation. *The Canadian Journal of Psychiatry*, 57, 144–151. [PubMed: 22398000]
- Storch EA, Caporino NE, Morgan JR, Lewin AB, Rojas A, Brauer L, ...Murphy, TK (2011). Preliminary investigation of web-camera delivered cognitive-behavioral therapy for youth with obsessive-compulsive disorder. *Psychiatry Research*, 189, 407–412. doi:10.1016/j.psychres.2011.05.047 [PubMed: 21684018]
- Storch EA, Merlo LJ, Lack C, Milsom VA, Geffken GR, Goodman WK, & Murphy TK (2007). Quality of life in youth with Tourette's syndrome and chronic tic disorder. *Journal of Clinical Child and Adolescent Psychology*, 36, 217–227. doi:10.1080/15374410701279545 [PubMed: 17484694]
- Storch EA, Murphy TK, Geffken GR, Sajid M, Allen P, Roberti JW, & Goodman WK (2005). Reliability and validity of the Yale Global Tic Severity Scale. *Psychological Assessment*, 17, 486–491. doi:10.1037/1040-3590.17.4.486 [PubMed: 16393016]
- Verdellen C, van de Griendt J, Hartmann A, Murphy T, & the ESSTS Guidelines Group (2011). European clinical guidelines for Tourette Syndrome and other tic disorders. Part III: Behavioural and psychosocial interventions. *European Child & Adolescent Psychiatry*, 20, 197–207. doi: 10.1007/s00787-011-0167-3 [PubMed: 21445725]
- Wilhelm S, Peterson AL, Piacentini J, Woods DW, Deckersbach T, Sukhodolsky DG,...Scahill L (2012). Randomized trial of behavior therapy for adults with Tourette Syndrome. *Archives of General Psychiatry*, 69, 795–803. doi:10.1001/archgenpsychiatry.2011.1528 [PubMed: 22868933]
- Woods DW, Conelea CA, & Himle MB (2010). Behavior therapy for Tourette's disorder: Utilization in a community sample and an emerging area of practice for psychologists. *Professional Psychology: Research and Practice*, 41, 518–525. doi:10.1037/a0021709
- Woods DW, Piacentini J, Chang SW, Deckersbach T, Ginsburg G, Peterson AL, ...Wilhelm S (2008). *Managing Tourette Syndrome: A Behavioral Intervention for Children and Adults*. New York, NY: Oxford University Press Inc.

- Yuen EK (2010). Acceptance-based behavior therapy for social anxiety disorder using videoconferencing (Doctoral dissertation, Drexel University). Retrieved from <http://dspace.library.drexel.edu/>
- Yuen EK, Herbert JD, Forman EM, Goetter EM, Juarascio AS, Rabin S, ...Bouchard, S (2013). Acceptance based behavior therapy for social anxiety disorder through videoconferencing. *Journal of Anxiety Disorders*, 27, 389–397. doi:10.1016/j.janxdis.2013.03.002 [PubMed: 23764124]

Table 1

Patient Characteristics

	Participant 1	Participant 2	Participant 3	Participant 4
Gender	Male	Male	Male	Male
Age	13	13	11	10
Race	Caucasian (non-Hispanic)	Caucasian (non-Hispanic)	Caucasian (non-Hispanic)	Caucasian (non-Hispanic)
WASI-Vocab T-score	73 (very superior)	43 (average)	52 (average)	56 (average)
Mini-Kid Diagnosis	TS; ADHD-inattentive type	TS; ADHD-inattentive type	TS	TS
Medication	Vyvanse, Sertraline, Clonidine	None	None	None

TS = Tourette Syndrome; ADHD = Attention-deficit hyperactivity disorder

Table 2**Computer Equipment and Features**

	Participant 1	Participant 2	Participant 3	Participant 4	Treatment Site
Internet Connection	Wireless	Wireless	Wireless	Wireless	Wi-Fi
Web camera Type	Built-in	Built-in	Separate	Built-in	Separate
Web camera model	N/A	N/A	Logitech c110	N/A	Logitech 270
Additional Equipment	Headset	None	None	None	None
Computer Type	Laptop	Desktop	Desktop	Laptop	Laptop
Computer Location	Basement	Basement	Kitchen	Dining room/Kitchen	Office
Computer Age	1.5 years	7.5 months	7 months	3 weeks	12 months
Computer Make/Model	Gateway NV53	iMac12,2	Lenovo	Hewlett Packard Pavillion G7	Dell Optiplex 980
Operating System	Windows 7	Mac OS 10	Windows 7	Windows 7	Windows XP
Processor	AMD Athlon II	Intel Core i5	Intel R core i5	AMD a6	Intel Core i5
Processor Speed	2.0 GHz	2.7 GHz	2.5 GHz	2.7 GHz	3.2 GHz
Random Access Memory	4 GB	4 GB	6 GB	3.48 GB	3.49 GB
Free Hard Drive Space	199.0 GB	892.9 GB	376.0 GB	575.0 GB	217.0 GB

Table 3

Baseline and Post-treatment Clinical Outcomes

	Baseline			CGI-S	Post				
	YGTSS				YGTSS			CGI-S	CGI-I
	Motor	Vocal	Total		Motor	Vocal	Total		
Participant 1	17	15	32	5	16	10	26	4	3
Participant 2	10	12	22	4	6	8	14	3	1
Participant 3	12	8	20	4	14	8	22	4	5
Participant 4	14	13	27	5	8	5	13	3	1

Note. CGI-Severity scale: 1 = normal, not all ill; 2 = borderline mentally ill; 3 = mildly ill; 4 = moderately ill; 5 = markedly ill; 6 = severely ill; 7 = extremely ill; CGI-Improvement: 1 = very much improved; 2 = much improved; 3 = improved 4 = minimally improved; 5 = no change; 6 = minimally worse; 7 = much worse; 8 = very much worse