Published in final edited form as:

Glia. 2018 December ; 66(12): 2684–2699. doi:10.1002/glia.23521.

NG2 expression in NG2 glia is regulated by binding of SoxE and bHLH transcription factors to a Cspg4 intronic enhancer

Hitoshi Gotoh^{1,4,*}, William M. Wood¹, Kiran D. Patel¹, Daniel C. Factor⁵, Linda L. Boshans¹, Tadashi Nomura⁴, Paul J. Tesar^{5,6}, Katsuhiko Ono⁴, and Akiko Nishiyama^{1,2,3,*}

¹Department of Physiology and Neurobiology, University of Connecticut, Storrs, CT, 06269-3156, USA

²Institute of Systems Genomics, University of Connecticut

³Institute of Brain and Cognitive Science, University of Connecticut

⁴Department of Biology, Kyoto Prefectural University of Medicine, Kyoto, 606-0823, Japan

⁵Department of Genetics and Genome Sciences, Case Western Reserve University School of Medicine, Cleveland OH, 44106, USA

⁶Department of Neurosciences, Case Western Reserve University School of Medicine

Abstract

NG2 is a type 1 integral membrane glycoprotein encoded by the *Cspg4* gene. It is expressed on glial progenitor cells known as NG2 glial cells or oligodendrocyte precursor cells that exist widely throughout the developing and mature central nervous system and vascular mural cells but not on mature oligodendrocytes, astrocytes, microglia, neurons, or neural stem cells. Hence NG2 is widely used as a marker for NG2 glia in the rodent and human. The regulatory elements of the mouse *Cspg4* gene and its flanking sequences have been used successfully to target reporter and Cre recombinase to NG2 glia in transgenic mice when used in a large 200-kilobase bacterial artificial chromosome cassette that contained the 38-kilobase *Cspg4* gene in the center. Despite the tightly regulated cell type- and stage-specific expression of NG2 in the brain and spinal cord, the mechanisms that regulate its transcription have remained unknown. Here, we describe a 1.45-kb intronic enhancer of the mouse *Cspg4* gene that directed transcription of EGFP reporter to NG2 glia but not to pericytes in vitro and in transgenic mice. The 1.45-kb enhancer contained binding sites for SoxE and basic helix-loop-helix transcription factors, and its enhancer activity was augmented cooperatively by these factors, whose respective binding elements were found in close proximity to each other. Mutations in these binding elements abrogated the enhancer activity when tested in the postnatal mouse brain.

*Correspondence should be addressed to: Akiko Nishiyama, tel: 1-860-486-4561; fax: 1-860-486-3303; akiko.nishiyama@uconn.edu or Hitoshi Gotoh, tel: +81-75-703-4941; fax: +81-75-703-4978; hgotoh@koto.kpu-m.ac.jp.

Conflict of interest

None

Keywords

NG2; Cspg4; oligodendrocyte precursor cell; Sox10; Olig2; Ascl1

INTRODUCTION

NG2-expressing glia (NG2 cells) are a major population of glial cells in the adult central nervous system (CNS). They are process-bearing, proliferative cells that continue to generate mature oligodendrocytes throughout life and hence they are also known as oligodendrocyte precursor cells (OPCs) or polydendrocytes (Nishiyama et al., 2009, 2016). NG2 cells are distributed uniformly throughout the CNS, more broadly than oligodendrocytes (Tomassy et al., 2014), and they are highly integrated in the neural networks (Hill and Nishiyama, 2014). The study of this population of glial cells has been greatly facilitated by the identification of NG2 and platelet-derived growth factor receptor alpha (Pdgfra) as the two key cell surface proteins expressed by NG2 cells but not by neurons, astrocytes, mature oligodendrocytes, or microglia (Nishiyama et al., 1996a; Nishiyama, 2012). The NG2 protein, which is encoded by the chondroitin sulfate proteoglycan 4 (Cspg4) gene, is a type 1 integral membrane glycoprotein (Stallcup et al., 1983; Nishiyama et al., 1991), and its human ortholog is also known as the melanoma chondroitin sulfate proteoglycan (Pluschke et al., 1996). NG2 modulates diverse cellular functions including cell adhesion, migration, proliferation, and differentiation (Stallcup and Huang, 2008; Binamé, 2014). NG2 is also expressed by vascular mural cells in the CNS and elsewhere and by tissue-specific progenitor cells of a variety of non-neural lineages.

The onset of NG2 expression in the oligodendrocyte lineage occurs immediately after induction of the SoxE family of high mobility group transcription factors Sox9 and Sox10, which are induced by the basic helix-loop-helix transcription factor Olig2 (Weider and Wegner, 2016). This coincides with oligodendrocyte lineage specification. NG2 is highly expressed on migrating, proliferating progenitor cells and is downregulated upon terminal oligodendrocyte differentiation. NG2 expression is upregulated in a variety of pathological states, such as demyelination (Reynolds et al. 2001), spinal cord injury (Jones et al. 2002), and in neoplasia (Shoshan et al. 1999, Stallcup and Huang, 2008).

Despite the well known changes in NG2 expression under various physiological and pathological conditions and the widespread use of NG2 as a marker for NG2 glia, little is known about the transcriptional regulation of the Cspg4 gene. In transgenic mouse lines in which expression cassettes are inserted into the first exon of the mouse Cspg4 gene placed in the center of a 200-kb bacterial artificial chromosome (BAC), faithful expression of DsRed (Zhu et al., 2008), EGFP (Hughes et al., 2013), and Cre/CreERTM (Zhu et al. 2008, 2011) in NG2-expressing cells has been attained. This indicates that the *cis*-acting regulatory elements that are required for the correct expression of Cspg4 are located within or close to the Cspg4 genomic locus. We have identified a 1.45-kb evolutionarily conserved enhancer in the first intron of the mouse Cspg4 gene. The enhancer targets transcription of a reporter gene to NG2 glia in vivo and in vitro. Both Sox10 and Olig2 bind to their respective binding

elements in the 1.45-kb enhancer, and the enhancer activity is cooperatively augmented by SoxE and basic helix-loop-helix (bHLH) transcription factors.

MATERIALS AND METHODS

Analysis of conserved regions within the *Cspg4* gene

To identify the regions of the mouse *Cspg4* gene that were conserved among species, we analyzed the *Cspg4* gene using VISTA browser in the Vista program (Genomics Division, Lawrence Berkeley National Laboratory and US Department of Energy Joint Genome Institute, <http://genome.lbl.gov/vista/index.shtml>; Visel et al., 2007) and the University of California Santa Cruz (UCSC) Genome Browser. Nucleotide positions are based on the December 2011 *Mus musculus* assembly (Genome Reference Consortium Mouse Build 38 (GRCm38; GCA_000001635.2). We identified the following non-coding regions that were conserved among mammals: 3.0k promoter and 1.5k promoter in the promoter region; intron 1–1 (int1–1), intron 1–2 (int1–2), intron1–3 (int1–3), intron 8 (int8) in intronic sequences; and two regions in the 3' untranslated region (3'-UTR1 and 3'-UTR2) (Figure 1A). The int1–3 region was further subdivided into int1–3a, b, and c subregions, and int1–3b was further subdivided into int1–3bi, 1–3bii, and 1–3biii. The nucleotide positions of these regions are listed in supplemental Table 1. Transcription factor binding sites in int1–3 were searched in the UCSC Genome Browser.

Plasmids

For electroporation into chick embryos and transfecting mixed glial cultures, we used the transposon vector pT2AL200R175-CAGGS-EGFP (kindly provided by Dr. Koichi Kawakami) (Kawakami and Noda 2004; Urasaki et al. 2006) and replaced the chicken beta-actin (CAG) promoter with PCR-amplified promoter/ enhancer regions of the mouse *Cspg4* gene, and inserted it upstream of the SV40 minimal promoter. The SV40 minimal promoter showed little basal activity in chick, but we saw significant background activity in astrocytes when used in mixed glial cultures. Therefore, in subsequent experiments we switched to using the β -globin minimal promoter (Yee and Rigby, 1993). For electroporation into mouse brain, the *Cspg* int1–3b enhancer and β -globin minimal promoter were inserted into the transposon vector upstream of EGFP. For luciferase assays, int1–3bi, int1–3bii, or int1–3biii was PCR-amplified together with the β -globin minimal promoter and inserted into pGL3-Basic vector (Promega) expressing Firefly luciferase. Scrambled mutations were introduced into Sox BE1, Sox BE2, Sox BE3, bHLH BE1, BE3, and bHLH BE4 sequences. For generating transgenic mice, int1–3b enhancer was inserted upstream of *Heat-shock protein 68* (*hsp68*) minimal promoter to drive EGFP transcription. For these experiments, we switched to using the *hsp68* minimal promoter because it had been shown that the *hsp68* minimal promoter gives a low background activity on its own but strong activity when combined with active promoter/enhancer elements in transgenic mice (Kothary et al., 1989). All the primer sequences used for the construction of the plasmids are listed in Supplemental Tables 1 and 2.

Cell culture and transfection

For mixed glial cultures, postnatal day 0–3 (P0–3) mouse forebrains were dissociated into single cells with trypsin and plated onto poly-L-lysine-coated glass coverslips. The cells were then directly transfected or passaged once and transfected. Oli-neu mouse oligodendrocyte precursor cell line (Jung et al. 1995, a gift from Dr. Jacky Trotter, Mainz) was maintained in DMEM/F12 containing 1% horse serum and N2 supplement (Invitrogen, USA). Neuro-2a mouse neuroblastoma cells (ATCC, USA) were maintained in DMEM containing 10% FBS. Plasmids were transfected into these cells using Lipofectamine 2000 (Life Technologies, USA) according to the manufacturer's instructions.

Animals

ICR mice were obtained from SLC (Shizuoka, Japan), FVB mice from Jackson Laboratory (Bar Harbor, Maine), and CD1 mice were from Charles River Laboratories (Wilmington, MA) and were bred in our animal facilities. For perinatal electroporation, mice were anesthetized by hypothermia, approximately 1 µg of the plasmids were injected into the lateral ventricle, and 4 square pulses of 100V and 50 msec each were applied using a NEPA21 electroporator (Nepa Gene, Japan; Bulldog Bio, USA). Transgenic mice were generated in the Gene Transfer and Transgenic Mouse Facility of the University of Connecticut Health Center by injecting linearized plasmids into fertilized CD1 oocytes and transferring them into pseudopregnant mice. Fertilized chick eggs were obtained from a local breeder (Yamagishi, Japan) and were incubated at 37°C. The embryonic stage was determined according to Hamburger and Hamilton (1951). For electroporation, plasmid vectors were injected into the neural tube, and three electric pulses of 30V and 50 msec each were applied. All animal procedures were approved by the animal care committee of University of Connecticut or Kyoto Prefectural University of Medicine.

Immunohistochemistry

Brain, cerebellum, and spinal cords were fixed in 4% paraformaldehyde in phosphate-buffered saline by intracardiac perfusion. Sections were cut using a cryostat or vibratome (Leica, Germany) and subjected to immunohistochemical staining as previously described (Zhu et al., 2011). Primary antibodies used for immunohistochemistry were rabbit anti-GFP (1:200, Invitrogen), chick anti-GFP (1:1000, Abcam and Aves); rabbit anti-NeuN (1:1000, Cell Signaling Technology), rabbit anti-NG2 (1:500, Millipore), rabbit anti-Olig2 (1:500, Millipore); mouse anti-Olig2 (1:1000, Novus Biologicals); mouse anti-GFAP (1:1000, Sigma-Aldrich); and rat anti-CD11 (1:100, Serotec, UK). Stained sections were observed under Zeiss Axiovert M200 fluorescent microscope or confocal microscope (Olympus FV1000; Leica TCS SP2 and SP8).

Chromatin immunoprecipitation (ChIP) analysis

ChIP analysis was performed as described by Carey et al. (2009) and in the manual for SimpleChIP Plus Sonication Kit (Cell Signaling). Approximately 2×10^7 Oli-neu cells grown in 10 ml of medium in a 10-cm dish were cross-linked with 1% methanol-free formaldehyde (Cell Signaling) for 20 minutes followed by quenching with 0.137M glycine. Cross-linked cells were harvested and washed in cold PBS and stored frozen at -80°C . To prepare nuclear

lysate, cells were first lysed in a solution containing 0.5% NP40 and 85 mM KCl in 5 mM Pipes buffer, pH8.0. The nuclear pellet was then lysed in a solution containing 1% SDS and 10 mM EDTA in 50 mM Tris-HCl (pH8.0) buffer and sonicated for 7–10 cycles of 15 seconds on and 45 seconds off at 80% output on a Misonix 705 sonicator using a cup horn to obtain DNA of 0.3–3 kb in size. The chromatin preparation was diluted in a buffer containing 167 mM NaCl, 1.2 mM EDTA, 0.01% SDS, and 1.1% NP-40 in 16.7 mM Tris-HCl buffer pH8.8 and incubated at 4°C overnight with normal rabbit IgG (4 µg/ml, Millipore), rabbit anti-histone H3 (10 µl/ml, Cell Signaling), rabbit anti-Olig2 (4 µg/ml, Millipore), or rabbit anti-Sox10 (1:5,000 dilution; kind gift from Dr. Michael Wegner, Germany). The immune complexes were then incubated in Protein G Magnetic Beads (Cell Signaling) at 4°C for 2 hours. The beads were washed twice with dilution buffer and twice with high salt buffer containing 500 mM NaCl, 1 mM EDTA, and 0.1% SDS in 50 mM Tris-HCl buffer, pH8.0. Bound DNA was eluted in elution buffer containing 10 mM EDTA and 1% SDS and 50 mM Tris-HCl, pH8.0. Cross-links were reversed by incubating the eluted material in proteinase K at 65°C for 2 hours, and DNA was purified using NucleoSpin Gel and PCR Clean-up columns (Macherey-Nagel, USA). The purified DNA was subjected to PCR using int1–3b and intron 2 primers from the mouse *Cspg4* gene (Table 2) and the products were separated on 1.8% agarose gels.

ChIP sequencing datasets and analysis

ChIP-seq datasets from epiblast stem cell derived OPCs were retrieved from GEO (accession # not yet obtained). Adapter and quality trimming was performed using the FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). Data was aligned to the mm9 genome using Bowtie v0.12.9 (Langmead et al. 2009). Multi-aligned reads and PCR duplicates were discarded. WIG tracks were generated using MACS v1.4 (Zhang et al. 2008), then normalized by the median signal and converted to bigwig format using wigToBigWig (<http://hgdownload.cse.ucsc.edu/admin/exe/>). Tracks were visualized using IGV (Thorvaldsdottir et al. 2013).

Luciferase assays

Neuro-2a cells were cotransfected with the *Cspg4* enhancer-Firefly luciferase constructs in pGL3 (Promega) and the Renilla luciferase expression plasmid pGL4.73 (Promega). The details of the luciferase constructs are provided in Supplemental Table 2. Forty-eight hours after transfection, luciferase activity was assayed using the Dual luciferase reporter assay kit (Promega) and a Turner Veritas Microplate Luminometer or a BioTek Synergy HTX Multi-Mode Reader. The relative luciferase activity for each construct was obtained by dividing the Firefly luciferase activity by the Renilla luciferase activity.

Electrophoretic mobility shift assays

The DNA binding regions of mouse Sox10 (amino acid 1–235), full-length Olig2, and Ascl1 were cloned into pT7CFE1-cMyc vector (ThermoFisher, USA). In vitro translation was performed using the TNT Quick Coupled transcription/translation system (Promega, USA). For EMSAs for SoxBE1–3 and bHLH BE1 and 2, 5-prime biotinylated oligonucleotides were obtained through Integrated DNA Technologies (USA) and annealed. For EMSAs for bHLH BE3 and 4, oligonucleotides were 5-prime labeled by [γ -32P]ATP and annealed. The

oligonucleotide probes are listed in supplemental Table 2. Non-labeled oligonucleotides and mutated oligonucleotides in which presumptive transcription binding sites had been substituted with adenine were added to binding solutions to analyze the binding specificity. DNA-protein complexes were separated by electrophoresis using 5.5 % acrylamide gels and analyzed using LightShift Chemiluminescent EMSA Kit (ThermoFisher, USA; for SoxBE1–3 and bHLHBE1, 2) or BAS-5000 Phosphorimager (FujiFilm, Japan) for bHLH BE3 and 4. Mutant competitors were made by introducing scrambled mutation into the core transcription factor binding region.

RESULTS

Identification of a novel intronic enhancer of the mouse *Cspg4* gene

Chick in ovo electroporation—We searched the mouse *Cspg4* gene for regions that were highly conserved among mammals, which could represent potential enhancer regions. The most highly conserved non-coding regions were found in: 1) the promoter, 2) intron 1, 3) intron 8, and 4) the 3'-untranslated region (UTR) (Figure 1A). We first tested whether the conserved 1.5 kb and 3.0 kb promoter fragments could direct transcription in NG2 cells by chick in ovo electroporation assays. Since NG2 is not yet expressed in chick spinal cord at the time of electroporation at embryonic day 3 (E3), the Tol2 transposon system (Kawakami and Noda 2004; Yoshida et al., 2010) was used to allow genomic integration and retention of the expression cassette until NG2-expressing cells emerged around E5. In this binary system, two plasmids were electroporated (Figure 1). The helper plasmid contained the Tol2 transposase cDNA under the control of the cytomegalovirus immediate early enhancer / chicken β -actin promoter (CAG). The donor plasmid contained various elements from the mouse *Cspg4* gene followed by SV40 minimal promoter and EGFP cDNA, which was flanked by the Tol2 element. In the presence of transposase, transposition occurs by excision of the transposon of the donor plasmid and integration of the *Cspg4* enhancer-EGFP cassette into the genome.

We analyzed the phenotype of EGFP+ cells in the electroporated embryos at E15, which is a time point when Olig2+ oligodendrocyte lineage cells become abundantly distributed in the spinal cord parenchyma. The 1.5 kb promoter, which had previously been shown to be active in NG2 cells (Sellers et al. 2009), failed to direct EGFP expression in Olig2+ oligodendrocyte lineage cells (Figure 1B,D). Neither did the 3.0 kb promoter yield Olig2+ EGFP+ cells (not shown). To identify an enhancer region that could direct NG2 expression in NG2 glial cells, we generated additional *Tol2* donor vectors containing other conserved regions of the *Cspg4* genes upstream of EGFP cDNA (Figure 1A) and tested them by in ovo electroporation. EGFP+ cells were detected with intron 1–3 (int1–3, Figure 1C, D) and to a lesser degree with intron 8 (int8) constructs but not with constructs from intron 1–1 (int1–1), intron 1–2 (int1–2), or 3'-untranslated regions 1 or 2 (UTR-1, UTR-2). The large majority of the EGFP+ cells (96.1%) in embryos electroporated with the int1–3 construct expressed Olig2, whereas only 12.2% of the EGFP+ cells in embryos electroporated with the int8 construct were Olig2+ (Figure 1D). These results suggested that int1–3 corresponding to the 3' end of the first intron contained the regulatory elements of the mouse *Cspg4* gene necessary for targeting *Cspg4* expression to NG2 glial cells.

Transfection into mixed glial cells from perinatal mouse brain—To further identify the region in the mouse *Cspg4* gene required for NG2 cell-specific transcription, we transfected EGFP reporter constructs containing different regions of the *Cspg4* gene (Figure 2A, top) into mixed glial cultures from perinatal mouse brains that consisted of astrocytes, NG2 cells, microglia, and vascular cells and compared the ability of the various promoter/enhancer elements to drive EGFP expression in NG2 cells. When we transfected cells with CAG-EGFP, approximately 20% of the EGFP⁺ cells expressed NG2, suggesting adequate delivery of the expression cassette to NG2 cells (not shown). EGFP expression was also detected in cells transfected with EGFP reporter constructs driven by the 1.5 kb promoter (Figure 2A–B, I), 3.0 kb promoter, int1–3, int8, UTR-1 and UTR-2, but not with int1–1 or int1–2 (Figure 2I). However, only the int1–3-EGFP construct resulted in EGFP expression in NG2 cells, with $35.6 \pm 6.1\%$ of EGFP⁺ cells expressing NG2 (Figure 2I). GFAP was detected in approximately 10% of EGFP⁺ cells in cultures transfected with each of the other constructs, including the negative control plasmid driven by the SV40 minimal promoter, suggesting that this level of GFAP expression was due to the nonspecific activity of the SV40 minimal promoter. Other EGFP⁺ cells were flat cells that expressed neither NG2 nor GFAP.

In order to further map the *cis*-element required for the correct expression of *Cspg4*, we subdivided the 4.2-kb int1–3 region into three fragments (int1–3a, 1–3b, and 1–3c), each subregion representing a stretch of conserved sequences (Figure 1A). When we transfected mixed glial cultures with EGFP reporter constructs driven by int1–3a or int1–3c, there was little or no EGFP expression in NG2 cells (Figure 2, C–D, G–H, I). By contrast, the int1–3b - EGFP construct resulted in NG2 expression in $35.4 \pm 7.1\%$ of EGFP⁺ cells with polydendrocyte morphology, similar to that obtained with the entire int1–3-EGFP construct (Figure 2E–F, I). This suggests that the 1.45-kb int1–3b region contained a *cis*-acting element that was necessary for *Cspg4* expression in NG2 cells. Interestingly, this region was the only region of the non-coding sequence of the *Cspg4* gene that was also conserved in the chick.

Cspg4 int1–3b targets EGFP expression to NG2 cells in transgenic mice

To analyze the specificity of the *Cspg4* int1–3b enhancer, we generated transgenic mice by injecting fertilized oocytes with a transgene that contained int1–3b and *hsp68* minimal promoter followed by the EGFP coding sequence (Figure 3A). Two of the three founder lines (*Cspg4*-EGFP1 and *Cspg4*-EGFP2) showed strong EGFP expression in NG2 cells throughout the developing CNS. All subsequent analyses were performed using the *Cspg4*-EGFP1 line. At P21, EGFP was expressed in NG2 cells in the neocortex, corpus callosum, cerebellum, and spinal cord, which also expressed the oligodendrocyte transcription factor Sox10 (Figure 3, B–D, arrows). Labeling for PDGFR α and Olig2 also revealed EGFP expression in Olig2⁺ PDGFR α ⁺ cells (Figure 3E, arrows). EGFP⁺ cells did not express the astrocyte proteins glutamine synthetase (GS; Figure 3F) or glutamate aspartate transporter (GLAST, not shown). Nor did EGFP⁺ cells express the neuronal nuclear antigen NeuN (RbFox3; Figure 3G). In addition to glial cells, NG2 is expressed by vascular pericytes. In *Cspg4*-EGFP1 mice, EGFP was not detected in PDGFR β ⁺ pericytes in the cerebrum, cerebellum, and spinal cord at all ages examined (Figure 3H). Thus, the int1–3b enhancer

specifically targeted transcription in NG2 glial cells but not in vascular pericytes, suggesting differential regulation of *Cspg4* transcription in the two NG2+ cell populations in the CNS.

In P14 *Cspg4*-EGFP1 mice, more than 80% of NG2+ glial cells in the neocortex, corpus callosum, and spinal cord white matter exhibited EGFP fluorescence, while the proportion of NG2 cells that showed EGFP fluorescence in the cerebellum and spinal cord gray matter was slightly lower, between 60 and 75% (Figure 3I). At P60, more than 74% of NG2+ glial cells had EGFP fluorescence, while the proportion was lower in the gray matter ranging from 59% in the spinal cord to 72% in the neocortex (Figure 3J). Overall, there was no statistically significant difference between the proportion of NG2+ cells that were EGFP+ at P14 and P60 in gray or white matter ($p=0.0540$ and 0.9408 , respectively, $n=3$). The slight trend toward a decrease in P60 gray matter likely reflects the decline in NG2 mRNA expression with age (Nishiyama et al., 1996). EGFP fluorescence was sufficiently strong to visualize the cells without immunolabeling for EGFP, and EGFP+ NG2+ cells exhibited the typical polydendrocyte morphology.

We further evaluated the specificity and robustness of EGFP expression within the oligodendrocyte lineage in the neocortex and corpus callosum of *Cspg4*-EGFP1 mice at P14 and P60 (Figure 4). At both ages, the proportion of EGFP+ cells that were NG2+ was greater in the superficial layers of the neocortex than in deeper cortical layers and corpus callosum. At P14, 87% of EGFP+ cells in cortical layers 2/3 were NG2+, whereas 70% and 63% of EGFP+ cells in layers 5/6 and corpus callosum were NG2+, respectively (Figure 4A, D, G, J, M, P). This trend was more pronounced in P60 cerebrum, where 62%, 28%, and 16% of EGFP+ cells in layers 2/3, 5/6, and corpus callosum, respectively, were NG2+. Since mature oligodendrocytes are more abundant in the older brain and exist in a gradient of increasing density from superficial cortical layers to the corpus callosum (Tomassy et al., 2014), we quantified the proportion of EGFP+ cells that were CC1+ oligodendrocytes to determine whether the EGFP+ NG2- cells were oligodendrocytes. At P14, 3%, 24%, and 34% of EGFP+ cells in layers 2/3, 5/6, and corpus callosum, respectively, were labeled by the oligodendrocyte antibody CC1 (Figure 4 B, E, H, K, N, Q). At P60, a greater proportion of EGFP+ cells were oligodendrocytes (38%, 71%, and 88% in layers 2/3, 5/6, and corpus callosum, respectively). NG2+ cells and CC1+ cells accounted for the entire EGFP+ cell population in all three regions at P14 and P60. This was further confirmed in Olig2-labeled sections in which, 99–100% of EGFP+ cells were Olig2+ in all regions at P14 and P60 (Figure 4, C, F, I, L, O, R). Generally, EGFP expression in NG2+ cells was more robust than that in CC1+ cells, and the difference was more apparent when comparing EGFP fluorescence (Figure 3) than immunolabeled EGFP signal (Figure 4) (e.g. compare EGFP+ cells in Figure 3E with arrows (NG2 cells) and those without arrows (oligodendrocytes)). Thus, in the forebrain of *Cspg4*-EGFP1 mice, EGFP expression was highest in NG2 cells and persisted at a lower level in some oligodendrocytes.

Binding of Sox10 and Olig2 to *Cspg4* int1–3b enhancer

The nucleotide sequence of *Cspg4* int1–3 enhancer revealed several potential binding sites for Sox and bHLH transcription factors. To examine whether endogenous Sox and bHLH transcription factors were associated with the int1–3b enhancer of the endogenous *Cspg4*

gene, we performed chromatin immunoprecipitation (ChIP)-PCR on cross-linked extracts from Oli-neu cells, which are an oligodendrocyte precursor cell line that robustly expresses NG2 (Jung et al., 1995). We immunoprecipitated Olig2 and Sox10 and analyzed the associated DNA by PCR using primers from int1–3b. We used primers from intron 2 as negative control (Figure 5A). Int1–3b but not intron 2 sequences were detected in immunoprecipitates formed with antibodies to Olig2 and Sox10 but not with control rabbit IgG. Both int1–3b and intron 2 sequences were detected in total input chromatin and in immunoprecipitates formed with the positive control antibody to histone H3. This suggested that both Olig2 and Sox10 specifically bound to the *Cspg4* int1–3b enhancer.

Active histone posttranslational modifications at *Cspg4* int1–3b

We next examined whether the *Cspg4* int1–3b region was enriched in histone posttranslational modifications that are known to be associated with actively transcribed genes. Mono-methylated histone H3 lysine 4 (H3K4me1) is known to be enriched at enhancer sequences (Heintzman et al. 2009; Heintzman et al. 2007; Wang et al. 2008), and acetylated histone H3 lysine 27 (H3K27ac) marks active enhancer elements (Creyghton et al. 2010; Rada-Iglesias et al. 2011; Wang et al. 2008; Zentner et al. 2011). We examined ChIP-seq datasets for H3K4me1 and H3K27ac in NG2 cells derived from epiblast stem cells. Int1–3b and the surrounding regions were more enriched for both H3K4me1 and H3K27ac than intron 2 (Figure 5B). Thus, the local chromatin state was consistent with enhancer activity at int1–3b but not intron 2 in NG2 cells.

Cooperative activation of *Cspg4* int1–3b enhancer by SoxE and bHLH proteins

The bHLH transcription factors Olig2 and Ascl1 are involved in oligodendrocyte specification, proliferation, and differentiation (Lu et al. 2002; Parras et al. 2007; Takebayashi et al. 2002; Zhou and Anderson 2002; Weider and Wegner, 2016). The SoxE family of transcription factors Sox9 and Sox10 are expressed in committed oligodendrocyte lineage cells (Finzsch et al. 2008; Stolt et al., 2002, 2003; Küspert and Wegner, 2016). To examine whether SoxE and bHLH proteins modulate *Cspg4* int1–3b enhancer activity, we performed luciferase assays in Neuro-2a mouse neuroblastoma cells (Schubert et al. 1969), which do not express NG2. Neuro-2a cells were transfected with *Cspg4*-int1–3b-luciferase construct (int1–3b-luc) in combination with pCAG expression constructs containing SoxE and/or bHLH cDNAs, and luciferase activity was measured after 48 hours (Figure 6A). Values were normalized to luciferase activity in cells cotransfected with int1–3b-luc and the pCAG control vector without any transcription factors.

When added alone, Ascl1 or the SoxE proteins Sox9 and Sox10 slightly increased *Cspg4* int1–3b-luc activity. While Olig2 homodimer is known to function as a transcriptional repressor, Olig2 can activate transcription when it is heterodimerized with an E-box binding bHLH factor such as E47, (Küspert et al., 2011). Transfection of Olig2+E47 did not increase int1–3b-luciferase activity over control or Olig2 alone. However, expression of Sox9 in combination with Olig2+E47 resulted in a 3.6-fold increase in luciferase activity compared to Olig2+E47 and a 1.7-fold increase compared Sox9 alone. The combination of Sox10 and Olig2+E47 led to a 12.2-fold increase over Olig2+E47 and a 2.9-fold increase over Sox10 alone. Similarly, combined expression of Sox10 and Ascl1 resulted in a 4.9-fold increase in

luciferase activity compared with Ascl1 alone and a 2.8-fold increase over Sox10 alone. These observations suggest that there are binding sites for Sox and bHLH proteins within the Cspg4 int1–3b enhancer, and that SoxE and the bHLH proteins act synergistically to increase the enhancer activity.

Identification of Sox10 binding sites

To identify Sox10 binding sites in the Cspg4 enhancer, we divided the 1.45-kb int1–3b segment into three smaller fragments of approximately 500 bp each (int1–3bi, 3bii, and 3biii) and tested them in luciferase assays in Neuro-2a neuroblastoma cells in the presence or absence of Sox10. When the full-length 1.45 kb int1–3b was used to drive luciferase, co-transfection of Sox10 significantly increased the luciferase activity (Figure 6A, top 2 bars). We next performed luciferase assays with a series of deletion mutants that lacked progressively larger segments of int1–3bi, ii, and iii. We identified two 100-bp regions, one in int1–3bi and the other in int1–3biii that were responsive to activation by Sox10 (Figure 6B), suggesting that there were at least two Sox10-binding regions within the int1–3b enhancer.

We searched int1–3bi and int1–3biii sequences for potential Sox-binding motifs that resembled the canonical Sox-binding motif (A/T)(A/T)CAA(A/T)G (Küspert and Wegner, 2016) and found two candidate sites, Sox-BE (binding element) 1 (CACAAAT) and Sox-BE2 (ACACAAAC), in int1–3bi and a third candidate site Sox-BE3 (TCAAAG) in int1–3biii (Supplemental table 1). We generated int1–3bi luciferase constructs with scrambled mutations in Sox-BE1 and Sox-BE2 (Sox-BE1mut or Sox-BE2mut) and an int1–3biii luciferase construct with a scrambled mutation in Sox-BE3 (Sox-BE3mut) and measured luciferase activity after co-transfecting them into Neuro-2a cells with pCAG-Sox10 (Figure 6C). Mutations in Sox-BE1 and Sox-BE2 sites reduced the Sox10-induced luciferase activity to 38.3 and 26.0% of that produced by wild type (wt) int1–3bi-luc, respectively. Furthermore, int1–3bi-luc construct containing mutations in both Sox-BE1 and BE2 sites led to an 85.2% reduction in luciferase activity. This suggests a cooperative function of the two Sox-BE binding sites in int1–3bi. Sox-BE3mut also reduced Sox10-induced luciferase activity to 41.4% of that of wt int1–3biii, although the extent of Sox10 to increase wild type int1–3biii enhancer activity was significantly lower than its ability to increase int1–3bi activity (9.5-fold versus 2-fold increase).

The direct binding of Sox10 to Sox-BE1–3 sequences was analyzed by electrophoretic mobility shift assay (EMSA) (Figure 6D). Incubation of labeled oligonucleotides spanning the Sox-BE1, 2, or 3 sequences with purified Sox10 amino terminal portion containing the DNA binding domain resulted in retardation of the labeled oligonucleotide bands. The observed mobility shift was abolished when we added an excess of unlabeled wild type oligonucleotides as competitors but not mutant oligonucleotides that contained mutations in the Sox binding motifs.

Identification of bHLH binding sites

We took a similar approach to identify bHLH-binding motifs in int1–3b. In Neuro-2a cells, int1–3bi-luc and int1–3biii-luc but not int1–3bii-luc activity was increased in the presence of

Ascl1 (Figure 7A). Consistent with the lack of int1–3b-luciferase response to Olig2+E47 shown in Figure 6A, Olig2+E47 was less effective in increasing transcriptional activity of int1–3bi and int1–3biii. We scanned int1–3bi and int1–3biii subregions for sequences resembling the canonical bHLH binding motif CAXXTG (Chaudhary and Skinner 1999) and found two putative sites in int1–3bi region, bHLH-BE1 (CAATCG) and bHLH-BE2 (CACGCG), and two additional sites in int1–3biii region, bHLH-BE3 CAGCCAG in the sense strand and bHLH-BE4 (CAGAAG) in the antisense strand. When we introduced scrambled mutations into the BE1 site in the int1–3bi luciferase construct, there was a general trend toward higher luciferase activity. This reached a statistically significant level only in the presence of Olig2+E47 but not in control or in the presence of Ascl1 (Figure 7B, left). It is possible that Olig2 binding to this region could repress transcription. By contrast, when we mutated the BE2 site in the int1–3bi luciferase construct, there was a reduction in luciferase activity compared with that obtained with wt int1–3i-luc construct, and this reached statistical significance in the presence of Olig2+E47 as well as with Ascl1 (Figure 7B, left). Mutation of the bHLH BE3 sequence had no effect on int1–3biii luciferase activity in the presence of Olig2+E47 but reduced the activity induced by Ascl1 (Figure 7B, right). Mutations in bHLH BE4 had no effect on int1–3biii luciferase activity.

EMSA analysis showed retardation of labeled bHLH BE1, 3, and 4 by Olig2 and Ascl1, which was diminished in the presence of excess unlabeled wild type but not mutant competitors (Figure 7C), suggesting that bHLH BE1, 3, and 4 were directly recognized by Olig2 and Ascl1. In the case of bHLH BE4, addition of Olig2 or Ascl1 led to the formation of two species of retarded bands, both of which could be competed out by wild type but not mutant bHLH BE4 oligonucleotides, suggesting multimeric binding. Interestingly, Sox and bHLH binding motifs in both int1–3bi and int1–3biii were located within 100 bp of one another. Therefore, the two classes of transcription factors may functionally interact to modulate Cspg4 transcription.

Transcription factor binding sites are necessary for int1–3b enhancer activity in the brain

To examine whether Sox10 and bHLH binding sites were required for Cspg4 enhancer activity *in vivo*, we tested the ability of mutated int1–3b to drive reporter expression after electroporation into the neonatal mouse brain. When we electroporated wild type Tol2-Cspg4-EGFP (Figure 8A) together with Tol2-CAG-mCherry and CAG-Transposase plasmids into P0 brain and analyzed the brains at P14, EGFP+ cells were found in the brain parenchyma, away from the site of electroporation (Figure 8B and D). We next constructed a Tol2-Cspg4mut-EGFP reporter plasmid with scrambled mutations in the six transcription factor binding sites, Sox-BE1, 2, 3 and bHLH-BE1, 3, and 4. When the Tol2-Cspg4mut-EGFP plasmid was electroporated, no EGFP+ cells were observed even though mCherry+ cells were detected along the lateral ventricles in similar numbers to the brains electroporated with wild type Tol2-Cspg4-EGFP plasmid (Figure 8, C and D). These results suggest that the Sox10- and bHLH-binding elements within the Cspg4 int1–3b enhancer were required for Cspg4 transcription in NG2 glial cells *in vivo*.

To confirm that P0 electroporation could target NG2 glia, we verified the efficacy of electroporation by co-electroporating into P0 brain Tol2-CAG-mCherry plasmid encoding

mCherry and the transposase plasmid (CAG-TPase). When the brains were analyzed at P14, there was widespread mCherry expression among GS⁺ astrocytes, Pdgfra⁺ OPCs, CC1⁺ oligodendrocytes, as well as in cells that were negative for these markers (Figure 8E, left), indicating that P0 electroporation with CAG promoter-driven Tol2 constructs targeted a wide range of cell types besides NG2 cells. When we co-electroporated wild type Tol2-Cspg4-EGFP plasmid along with CAG-TPase, all the EGFP⁺ cells were Pdgfra⁺ or NG2⁺ and exhibited polydendrocyte morphology (Figure 8E, right, and 8F). This verified the specificity of the wild type Cspg4 int1–3b enhancer in targeting EGFP expression to NG2 cells in the P0 mouse brain electroporation assay and supported the requirement of one or more of the SoxE and bHLH binding elements in this process.

DISCUSSION

Identification of a 1.45-kb Cspg4 enhancer that directs transcription in NG2 glia

We have identified a 1.45 kb intronic enhancer in the mouse Cspg4 gene (Cspg4-int1–3b) that conferred reporter expression in NG2 glial cells in the mouse and chick CNS but not in NG2-expressing vascular cells. Using int1–3b-luciferase assays in Neuro-2a mouse neuroblastoma cells, we have shown that the enhancer activity was cooperatively increased by SoxE proteins Sox9 or Sox10 in combination with bHLH factors Ascl1 or Olig2+E47. By using mutation analysis, we further demonstrated by luciferase assays and EMSA that the binding sites for SoxE and bHLH factors within the int1–3b enhancer were in close proximity to each other. Using ChIP-PCR of Oli-neu cells extracts, we showed that both Olig2 and Sox10 were enriched at Cspg4 int1–3b enhancer sequence but not at intron 2. The enhancer activity of int1–3b was further independently supported by ChIP-seq data, which showed a greater occupancy of H3K27ac and H3K4me1 around int1–3b region compared with intron 2 in NG2 glial cells derived from mouse epiblast.

A previous study reported that a 1.5-kb mouse Cspg4 promoter sequence targets reporter expression to NG2 cells when used in a retroviral vector injected into injured CNS (Guo et al. 2014; Sellers et al. 2009). However, in our assays using primary mouse mixed glial cultures and chick in ovo electroporation, we did not detect reporter expression in NG2 cells with the 1.5 kb promoter. Moreover, the 1.5 kb promoter resulted in nonspecific reporter expression in neurons and GFAP⁺ astrocytes. In the Sellers study, 15% of the infected cells were reported to be GFAP⁺ astrocytes, and 10% were Iba1⁺ microglia. The long-terminal repeats of the retroviral vector rather than the promoter are likely to have primarily driven the expression of the reporter in the highly proliferative NG2 cells.

The 1.45-kb int1–3b enhancer in the first intron, which was the only non-coding region found to be conserved in chick as well as mammals, directed reporter expression specifically to NG2 glial cells and some oligodendrocytes in vitro and in vivo. While the majority of NG2 cells in Cspg4-EGFP1 transgenic mice expressed EGFP, there were some NG2 cells that did not express EGFP, possibly due to the absence of a critical promoter sequence that was not included in the construct and/or position effect of the integration site. We never detected EGFP expression in astrocytes, which suggests that int1–3b enhancer contains sequences critical for repressing transcription in astrocytes or that astrocytes lack the critical factor that activates enhancer.

In Cspg4-EGFP1 mice, EGFP expression was also detected in some CC1+ oligodendrocytes, although at a lower level than in NG2 cells. A small percentage of the EGFP+ cells that lacked Olig2 or Sox10 were likely to be mature myelinating oligodendrocytes (Muroyama et al., 2005). Since it takes several days for NG2 cells to differentiate into oligodendrocytes, especially in older mice at P60, (Zhu et al., 2011; Hill et al. 2014), which is longer than the half-life of EGFP, the presence of EGFP in oligodendrocytes is likely due to weak activation of EGFP transcription in mature oligodendrocyte rather than residual EGFP protein that had been synthesized in NG2 cells prior to differentiation. The larger proportion of EGFP+ cells that were CC1+ oligodendrocytes in deeper cortical layers and corpus callosum at both P14 and P60 also suggests that the enhancer remains active in new oligodendrocytes. There are likely to be other silencing elements required for rapid downregulation of NG2 transcription during oligodendrocyte differentiation. For example, the 3'-UTR, which was not included in the Cspg4-EGFP1 mice, could contain sequences required for regulation by miRNAs, as shown for downregulation of PDGFR α expression during oligodendrocyte differentiation (Budde et al. 2010; Dugas et al. 2010; Zhao et al. 2010). Notably, >99% of EGFP+ cells were Olig2+ oligodendrocyte lineage cells.

In Cspg4-EGFP1 mice, very faint EGFP expression that was only slightly above background was detected in the subventricular zone and in some cortical neurons. In Cspg4-EGFP2 transgenic line that exhibited a higher level of EGFP, a more widespread distribution of EGFP was detected not only in oligodendrocyte lineage cells but also in neuronal cells (data not shown). By contrast, when the int1–3b enhancer was used to drive EGFP expression in a Tol2 construct by in vivo electroporation in the P0 brain, EGFP was expressed exclusively in NG2 cells (Figure 8). This suggests that cell type specificity of the enhancer could depend on copy number, and at low gene dosage, there is a higher degree of specificity. We did not observe EGFP expression in PDGFR β + pericytes at any of the ages examined, suggesting that NG2 expression in pericytes is regulated through different *cis*-regulatory elements. Thus, the Cspg4 int1–3b sequence will be useful for targeting genes specifically to NG2 glia and not pericytes.

Sox10 and Olig2 binding to the Cspg4 int1–3b enhancer

We have shown that SoxE and bHLH transcription factors activated the Cspg4 int1–3b enhancer, and Olig2 and Sox10 were enriched at int1–3b sequence. Of the SoxE proteins which are Sox8, 9, and 10, Sox9 plays a critical role in the generation of parenchymal NG2 cells in the developing spinal cord. In the absence of Sox9, Olig2+ cells appear normally in the ventricular zone, but the appearance of NG2 cells in the parenchyma and the expression of NG2 and Sox10 are severely affected in the E12.5-E14.5 spinal cord (Stolt et al. 2003). Sox9 is downregulated in oligodendrocyte lineage cells beyond the first postnatal week (Finzsch et al. 2008; Stolt et al. 2003, 2006; Weider and Wegner, 2016). By contrast, Sox10 persists through the lineage and has multiple roles at different stages of differentiation. Thus, Sox10 is likely to play a more prominent role in regulating NG2 expression in postnatal NG2 cells. Our mutation analysis revealed at least three Sox-binding sites in the int1–3b enhancer sequence, each of which partially contributed to activate NG2 transcription. Mutations in both Sox10 BE1 and 2 putative binding sites reduced Sox10-dependent

luciferase activity by 85% suggesting a cooperative role of Sox10. However, the distance between BE1 and 2 are farther than the reported binding sites for Sox10 dimer in other cell types (Peirano and Wegner, 2000), and we did not see a supershifted band in EMSA. ChIP-sequencing of epiblast stem cell-derived OPCs revealed an enrichment of H3K27ac and H3K4me1 at Cspg4 int1–3b, indicative of an open accessible chromatin state consistent with enhancer activity. A recent Sox10 ChIP-sequencing study using mouse spinal cord showed a high degree of Sox10 engagement in the same subregions of int1–3b that contained the Sox-BE sites (Lopez-Anido et al., 2015), further corroborating the accessibility of the Cspg4 int1–3b enhancer to the transcription factors.

Among the bHLH transcription factors, Olig2 and Ascl1 also play a critical role in oligodendrocyte progenitor specification and differentiation (Lu et al. 2002; Parras et al. 2007; Takebayashi et al. 2002; Zhou and Anderson 2002; Yu et al., 2013; Nakatani et al., 2013; Wegener et al., 2015). In Neuro-2a cells, the combination of Sox10 and each of these bHLH factors significantly elevated int1–3b enhancer, suggesting synergistic activity between the two classes of transcription factors. Further mapping of the potential Sox and bHLH binding sites using mutant constructs of int1–3b revealed that the SoxE and bHLH bindings sites were found close to each other within the int1–3bi and int1–3biii subdomains. Enhanced luciferase activity in bHLH BE1 site suggests that Olig2 binding to this site could negatively regulate transcription. An example of where this could occur is when Sox10 and Olig2 expression are elevated as NG2 cells differentiate into mature oligodendrocytes and downregulate NG2 expression. The EMSA assays revealed that Olig2 and Ascl1 share the same three bHLH binding motifs. Thus, the combination of Olig2 and Ascl1 may be involved in a dose-dependent regulation of NG2 transcription, similar to the previously described gene dosage effects of Ascl1 and Olig2 on the specification of PDGFR α ⁺ OPCs in the telencephalon (Parras et al. 2007). Within the two clusters of Sox and bHLH binding motifs in int1–3b, the Sox and bHLH binding motifs were within 100 bp of each other, providing further support for a functional interaction of the two families of transcription factors. Furthermore, the Sox family of proteins have been shown to directly interact with Olig2 (Wissmuller et al. 2006). The inability of the mutated enhancer to direct EGFP transcription after delivery into neonatal mouse brain also supports the importance of these binding sites in the developing brain *in vivo*.

In conclusion, we have identified a novel 1.45-kb Cspg4 intronic enhancer that activates transcription in NG2 glial cells *in vivo* and is coordinately regulated by Sox10 and the bHLH transcription factors Olig2 and Ascl1. The identification and characterization of the enhancer described here not only provides new insight into glial cell-specific transcription of Cspg4, but the enhancer can also be used as a tool for targeting genes specifically to oligodendrocyte lineage cells.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We would like to thank Dr. Koichi Kawakami for providing us the tol2 expression vectors. We thank Dr. Michael Wegner (Erlangen, Germany) for the Sox10 antibody. We also thank Dr. Siu-Pok Yee (University of Connecticut Health Center, USA) for his advice and assistance in generating transgenic mice, Youfen Sun for maintaining the mouse colony, and David Serwanski for tissue processing and immunohistochemistry.

FUNDING

This work was supported by grants from the National Institutes of Health (grant numbers R01 NS074870 to AN, R01 NS073425 to AN); Grants-in-Aid for Scientific Research (grant number 25870618 to HG) from Japan Society for the Promotion of Science and Japan-U.S. Brain Research Cooperative Program.

REFERENCES

- Biname F (2014). Transduction of extracellular cues into cell polarity: the role of the transmembrane proteoglycan NG2. *Mol Neurobiol* 50, 482–493. [PubMed: 24390567]
- Budde H, Schmitt S, Fitzner D, Opitz L, Salinas-Riester G, Simons M. 2010 Control of oligodendroglial cell number by the miR-17–92 cluster. *Development* 137:2127–32. [PubMed: 20504959]
- Chaudhary J, Skinner MK. 1999 Basic helix-loop-helix proteins can act at the E-box within the serum response element of the c-fos promoter to influence hormone-induced promoter activation in Sertoli cells. *Mol Endocrinol* 13:774–86. [PubMed: 10319327]
- Chekenya M, Immervoll H. 2007 NG2/HMP proteoglycan as a cancer therapeutic target. *Methods Mol Biol* 361:93–117. [PubMed: 17172708]
- Dugas JC, Cuellar TL, Scholze A, Ason B, Ibrahim A, Emery B, Zamanian JL, Foo LC, McManus MT, Barres BA. 2010 Dicer1 and miR-219 Are required for normal oligodendrocyte differentiation and myelination. *Neuron* 65:597–611. [PubMed: 20223197]
- Finzsch M, Stolt CC, Lommes P, Wegner M. 2008 Sox9 and Sox10 influence survival and migration of oligodendrocyte precursors in the spinal cord by regulating PDGF receptor alpha expression. *Development* 135:637–46. [PubMed: 18184726]
- Gard AL, Williams WC, 2nd, and Burrell MR (1995). Oligodendroblasts distinguished from O-2A glial progenitors by surface phenotype (O4+GalC-) and response to cytokines using signal transducer LIFR beta. *Dev Biol* 167, 596–608. [PubMed: 7875381]
- Guo Z, Zhang L, Wu Z, Chen Y, Wang F, Chen G. 2014 In vivo direct reprogramming of reactive glial cells into functional neurons after brain injury and in an Alzheimer's disease model. *Cell Stem Cell* 14:188–202. [PubMed: 24360883]
- Hamburger V, Hamilton HL. 1951 A series of normal stages in the development of the chick embryo. *J Morphol* 88:49–92. [PubMed: 24539719]
- Hill RA, and Nishiyama A (2014). NG2 cells (polydendrocytes): listeners to the neural network with diverse properties. *Glia* 62, 1195–1210. [PubMed: 24753030]
- Hill RA, Patel KD, Goncalves CM, Grutzendler J, Nishiyama A. 2014 Modulation of oligodendrocyte generation during a critical temporal window after NG2 cell division. *Nat Neurosci* 17:1518–27. [PubMed: 25262495]
- Hughes EG, Kang SH, Fukaya M, and Bergles DE (2013). Oligodendrocyte progenitors balance growth with self-repulsion to achieve homeostasis in the adult brain. *Nature neuroscience* 16, 668–676. [PubMed: 23624515]
- Jones LL, Yamaguchi Y, Stallcup WB, Tuszynski MH. 2002 NG2 is a major chondroitin sulfate proteoglycan produced after spinal cord injury and is expressed by macrophages and oligodendrocyte progenitors. *J Neurosci* 22:2792–803. [PubMed: 11923444]
- Jung M, Kramer E, Grzenkowski M, Tang K, Blakemore W, Aguzzi A, Khazaie K, Chlichlia K, von Blankenfeld G, Kettenmann H and others. 1995 Lines of murine oligodendroglial precursor cells immortalized by an activated neu tyrosine kinase show distinct degrees of interaction with axons in vitro and in vivo. *Eur J Neurosci* 7:1245–65. [PubMed: 7582098]

- Kang P, Lee HK, Glasgow SM, Finley M, Donti T, Gaber ZB, Graham BH, Foster AE, Novitsch BG, Gronostajski RM, Deneen B. 2012 Sox9 and NFIA coordinate a transcriptional regulatory cascade during the initiation of gliogenesis. *Neuron*. 74(1):79–94. [PubMed: 22500632]
- Kawakami K, Noda T. 2004 Transposition of the Tol2 element, an Ac-like element from the Japanese medaka fish *Oryzias latipes*, in mouse embryonic stem cells. *Genetics* 166:895–9. [PubMed: 15020474]
- Kothary R, Clapoff S, Darling S, Perry MD, Moran LA, Rossant J. 1989 Inducible expression of an hsp68-lacZ hybrid gene in transgenic mice. *Development* 105:707–14. [PubMed: 2557196]
- Küspert M, Hammer A, Bosl MR, Wegner M. 2011 Olig2 regulates Sox10 expression in oligodendrocyte precursors through an evolutionary conserved distal enhancer. *Nucleic Acids Res* 39(4):1280–93. [PubMed: 20959288]
- Küspert M, Wegner M. 2016 Something 2 talk about-transcriptional regulation in embryonic and adult oligodendrocyte precursors. *Brain Res* 1638, 167–82. [PubMed: 26232072]
- Langmead B, Trapnell C, Pop M, Salzberg SL. 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 10:R25. [PubMed: 19261174]
- Lu QR, Sun T, Zhu Z, Ma N, Garcia M, Stiles CD, Rowitch DH. 2002 Common developmental requirement for Olig function indicates a motor neuron/oligodendrocyte connection. *Cell* 109:75–86. [PubMed: 11955448]
- Mabie PC, Mehler MF, Marmur R, Papavasiliou A, Song Q, and Kessler JA (1997). Bone morphogenetic proteins induce astroglial differentiation of oligodendroglial-astroglial progenitor cells. *J Neurosci* 17, 4112–4120. [PubMed: 9151728]
- Muroyama Y, Fujiwara Y, Orkin SH, and Rowitch DH. 2005 Specification of astrocytes by bHLH protein SCL in a restricted region of the neural tube. *Nature* 438:360–363. [PubMed: 16292311]
- Nakatani H, Martin E, Hassani H, Clavairoly A, Maire CL, Viadieu A, Kerninon C, Delmas A, Frahm M, Weber M, Nakafuku M, Zalc B, Thomas JL, Guillemot F, Nait-Oumesmar B, Parras C. 2013 *Ascl1/mash1* promotes brain oligodendrogenesis during myelination and remyelination. *J Neurosci* 33, 9752–9768. [PubMed: 23739972]
- Nishiyama A, Boshans L, Goncalves CM, Wegrzyn J, and Patel KD (2016). Lineage, fate, and fate potential of NG2-glia. *Brain Res* 1638, 116–128. [PubMed: 26301825]
- Nishiyama A, Lin XH, Giese N, Heldin CH, and Stallcup WB (1996). Co-localization of NG2 proteoglycan and PDGF alpha-receptor on O2A progenitor cells in the developing rat brain. *J Neurosci Res* 43, 299–314. [PubMed: 8714519]
- Nishiyama A, Lin XH, Giese N, Heldin CH, and Stallcup WB (1996). Interaction between NG2 proteoglycan and PDGF alpha-receptor on O2A progenitor cells is required for optimal response to PDGF. *J Neurosci Res* 43, 315–330. [PubMed: 8714520]
- Nishiyama A, Dahlin KJ, Prince JT, Johnstone SR, Stallcup WB. 1991 The primary structure of NG2, a novel membrane-spanning proteoglycan. *J Cell Biol* 114:359–71. [PubMed: 1906475]
- Nishiyama A, Komitova M, Suzuki R, Zhu X. 2009 Polydendrocytes (NG2 cells): multifunctional cells with lineage plasticity. *Nat Rev Neurosci* 10:9–22. [PubMed: 19096367]
- Nishiyama A 2012 NG2 cells (Polydendrocytes) chapter 10 in *Neuroglia* 3rd ed., ed Kettenmann H and Ransom B. Oxford University Press.
- Parras CM, Hunt C, Sugimori M, Nakafuku M, Rowitch D, Guillemot F. 2007 The proneural gene *Mash1* specifies an early population of telencephalic oligodendrocytes. *J Neurosci* 27:4233–42. [PubMed: 17442807]
- Peirano RI, Wegner M. 2000 The glial transcription factor *sox10* binds to DNA both as monomer and dimer with different functional consequences. *Nucleic Acids Res* 28, 3047–55 [PubMed: 10931919]
- Pluschke G, Vanek M, Evans A, Dittmar T, Schmid P, Itin P, Filardo EJ, and Reisfeld RA (1996). Molecular cloning of a human melanoma-associated chondroitin sulfate proteoglycan. *Proceedings of the National Academy of Sciences of the United States of America* 93, 9710–9715. [PubMed: 8790396]
- Price MA, Colvin Wanshura LE, Yang J, Carlson J, Xiang B, Li G, Ferrone S, Dudek AZ, Turley EA, McCarthy JB. 2011 CSPG4, a potential therapeutic target, facilitates malignant progression of melanoma. *Pigment Cell Melanoma Res* 24:1148–57. [PubMed: 22004131]

- Reynolds R, Cenci di Bello I, Dawson M, Levine J. 2001 The response of adult oligodendrocyte progenitors to demyelination in EAE. *Prog Brain Res* 132:165–74. [PubMed: 11544985]
- Schubert D, Humphreys S, Baroni C, Cohn M. 1969 In vitro differentiation of a mouse neuroblastoma. *Proc Natl Acad Sci U S A* 64:316–23. [PubMed: 4189500]
- Sellers DL, Maris DO, Horner PJ. 2009 Postinjury niches induce temporal shifts in progenitor fates to direct lesion repair after spinal cord injury. *J Neurosci* 29:6722–33. [PubMed: 19458241]
- Stallcup WB, Arner LS, Levine JM. 1983 An antiserum against the PC12 cell line defines cell surface antigens specific for neurons and Schwann cells. *J Neurosci* 3:53–68. [PubMed: 6337237]
- Stallcup WB, Huang FJ. 2008 A role for the NG2 proteoglycan in glioma progression. *Cell Adh Migr* 2:192–201. [PubMed: 19262111]
- Stolt CC, Lommes P, Sock E, Chaboissier MC, Schedl A, Wegner M. 2003 The Sox9 transcription factor determines glial fate choice in the developing spinal cord. *Genes Dev* 17:1677–89. [PubMed: 12842915]
- Stolt CC, Schlierf A, Lommes P, Hillgartner S, Werner T, Kosian T, Sock E, Kessaris N, Richardson WD, Lefebvre V and others. 2006 Sox9 proteins influence multiple stages of oligodendrocyte development and modulate Sox9 protein function. *Dev Cell* 11:697–709. [PubMed: 17084361]
- Takebayashi H, Nabeshima Y, Yoshida S, Chisaka O, Ikenaka K, Nabeshima Y. 2002 The basic helix-loop-helix factor olig2 is essential for the development of motoneuron and oligodendrocyte lineages. *Curr Biol* 12:1157–63. [PubMed: 12121626]
- Tomassy GS, Berger DR, Chen HH, Kasthuri N, Hayworth KJ, Vercelli A, Seung HS, Lichtman JW, and Arlotta P (2014). Distinct profiles of myelin distribution along single axons of pyramidal neurons in the neocortex. *Science* 344, 319–324. [PubMed: 24744380]
- Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013 Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 14:178–92. [PubMed: 22517427]
- Urasaki A, Morvan G, Kawakami K. 2006 Functional dissection of the Tol2 transposable element identified the minimal cis-sequence and a highly repetitive sequence in the subterminal region essential for transposition. *Genetics* 174:639–49. [PubMed: 16959904]
- Visel A, Minovitsky S, Dubchak I, Pennacchio LA. 2007 VISTA Enhancer Browser--a database of tissue-specific human enhancers. *Nucleic Acids Res* 35:D88–92. [PubMed: 17130149]
- Wegener A, Deboux C, Bachelin C, Frah M, Kerninon C, Seilhean D, Weider M, Wegner M, Nait-Oumesmar B. 2015 Gain of olig2 function in oligodendrocyte progenitors promotes remyelination. *Brain* 138, 120–35. [PubMed: 25564492]
- Weider M and Wegner M. 2016 Sox9 factors: transcriptional regulators of neural differentiation and nervous system development. *Semin Cell Dev Biol*.
- Wissmuller S, Kosian T, Wolf M, Finzsch M, Wegner M. 2006 The high-mobility-group domain of Sox proteins interacts with DNA-binding domains of many transcription factors. *Nucleic Acids Res* 34:1735–44. [PubMed: 16582099]
- Yee SP, Rigby PW. 1993 The regulation of myogenin gene expression during the embryonic development of the mouse. *Genes Dev* 7:1277–89. [PubMed: 8391506]
- Yoshida A, Yamaguchi Y, Nonomura K, Kawakami K, Takahashi Y, Miura M. 2010 Simultaneous expression of different transgenes in neurons and glia by combining in utero electroporation with the tol2 transposon-mediated gene transfer system. *Genes Cells* 15, 501–12. [PubMed: 20384787]
- Yu Y, Chen Y, Kim B, Wang H, Zhao C, He X, Liu L, Liu W, Wu LM, Mao M, Chan JR, Wu J, Lu QR. 2013 Olig2 targets chromatin remodelers to enhancers to initiate oligodendrocyte differentiation. *Cell* 152, 248–61. [PubMed: 23332759]
- Zhang Y, Liu T, Meyer CA, Eickhout J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W and others. 2008 Model-based analysis of ChIP-Seq (MACS). *Genome Biol* 9:R137. [PubMed: 18798982]
- Zhao X, He X, Han X, Yu Y, Ye F, Chen Y, Hoang T, Xu X, Mi QS, Xin M and others. 2010 MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* 65:612–26. [PubMed: 20223198]
- Zhou Q, Anderson DJ. 2002 The bHLH transcription factors OLIG2 and OLIG1 couple neuronal and glial subtype specification. *Cell* 109:61–73. [PubMed: 11955447]

- Zhu X, Bergles DE, Nishiyama A. 2008 NG2 cells generate both oligodendrocytes and gray matter astrocytes. *Development* 135:145–57. [PubMed: 18045844]
- Zhu X, Hill RA, Dietrich D, Komitova M, Suzuki R, and Nishiyama A (2011). Age-dependent fate and lineage restriction of single NG2 cells. *Development* 138, 745–753. [PubMed: 21266410]

Main Points:

A novel 1.45-kb intronic enhancer of the mouse *Cspg4* gene regulates NG2 expression in NG2 glial cells but not pericytes in vivo. SoxE and basic helix-loop-helix transcription factors bind to and activate the enhancer.

Figure 1. Cspg4 enhancer int1–3 drives reporter expression in chick oligodendrocyte lineage cells.

(A) Comparison of Cspg4 genomic regions within mammals using the UCSC genome browser. Genomic regions of mouse Cspg4 gene are shown across the top. Numbered blue boxes represent exons and lines indicate introns. Conserved regions against the mouse sequence are shown as peaks (red: intron; blue: exon). Genomic elements used for promoter / enhancer assays were selected based on conservation among species and are shown as black bars at the bottom.

(B, C) The tol2-Cspg4–1.5kb-promoter-EGFP (B) or tol2-Cspg4-int1–3-EGFP (C) construct was electroporated with the transposase expression plasmid into E3 chick neural tube. At E15, sections were stained for EGFP (green) and Olig2 (red). EGFP+ Olig2+ cells are detected with the int1–3 region but not with the 1.5-kb promoter. Scale bar in B indicates 40 μ m. Crosshairs indicate the positions for orthogonal projections.

(D) Quantification of EGFP+ cells in the chick. The percentages of EGFP+ cells that were Olig2+ (green) or Olig2- (red) with the 1.5-kb promoter (1.5k prom), int1–3 intronic region (int1–3), or intron 8 (int8) are shown. $n=3$ (1.5k prom and int8) or $n=4$ (int1–3b).

Figure 2. Int1-3b enhancer drives reporter expression in mouse NG2 cells in culture.

(A-H) EGFP expression in perinatal mouse mixed glial cultures transfected with EGFP expression constructs containing different *cis*-acting elements from the mouse *Cspg4* gene and immunostained for NG2 (magenta) or GFAP (blue) after 4 days. (A, B) Cells transfected with 1.5kb promoter-EGFP. (C, D) Cells transfected with int1-3a-EGFP. (E, F) Cells transfected with int1-3b-EGFP. Arrows in E and F indicate EGFP+ cells that express NG2 and exhibit the morphology of NG2 cells (G, H) Cells transfected with tol2-int1-3c-EGFP. Scale bars, 50 μ m.

(I) The percentage of EGFP+ cells that were NG2+ (green) or GFAP+ (red) in mixed glial cultures 4 days after transfection with the indicated constructs. n.d. No EGFP+ cells detected. $n=3$

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 3. Characterization of EGFP+ cells in P21 Cspg4-EGFP1 transgenic mice.

(A) A schematic of the transgenic construct. Top shows the 5' portion of the mouse Cspg4 gene on chromosome 9 up to exon 4 (not to scale). Int1–3 is located near the 3' end of intron 1. Bottom shows the key elements in the transgenic mouse line Cspg4-EGFP1. It contains the 1.45 kb int1–3b upstream of the Hsp68 minimal promoter (min prom) followed by the coding sequence of EGFP and polyadenylation signal (pA).

(B–D) Coronal section of the cerebrum (B), sagittal section of the cerebellum (C), and transverse section of the spinal cord (D) from P21 Cspg4-EGFP1 mice stained for Sox10 (red) and NG2 (blue). The green channel is EGFP fluorescence without antibody enhancement. Arrows indicate EGFP+ cells that are also Sox10+ and NG2+. In C, ml =

molecular layer, pcl = Purkinje cell layer, igl = internal granule cell layer, and wm = white matter.

(E) Coronal sections through the neocortex and corpus callosum of P21 transgenic mice immunolabeled for *Pdgfra* (red) and *Olig2* (blue). Green channel represents EGFP fluorescence. Arrows indicate EGFP+ *Pdgfra*+ *Olig2*+ cells.

(F) Coronal section of the neocortex of P21 transgenic mouse stained for GS showing that EGFP+ cells do not express GS.

(G) P21 neocortex stained for the neuronal marker NeuN showing that EGFP fluorescence is not detected in neurons. Crosshairs represent positions for orthogonal projections in panels F-H.

(H) P21 neocortex stained for *Pdgfr β* showing that EGFP fluorescence is not detected in vascular pericytes.

Scale bars in B-H represent 50 μ m.

(I-J) Quantification of the proportion of NG2+ cells that expressed EGFP in the gray and white matter of the cerebrum, cerebellum, and spinal cord at P14 (I) and P60 (J). $n=3$.

Figure 4. Expression EGFP in oligodendrocyte lineage cells in the cerebrum of Cspg4-EGFP mice. Images were taken from 1-μm single confocal z-slices.

(A-F) P14 brain immunolabeling for EGFP and NG2 (A, D), CC1 (B, E), or Olig2 (C, F) in layers 2/3 (A-C) and layers 5/6-corpus callosum junction (D-F).

(G-M) P60 brain immunolabeling for EGFP and NG2 (G, J, M), CC1 (H, K, N), or Olig2 (M, N, O) in layers 2/3 (G-I), layers 5/6 (J-L), and corpus callosum (M-O) of P60 Cspg4-EGFP1 mice.

Scale, 50 μm. Arrows indicate EGFP+ NG2+ cells. Arrowheads indicate EGFP+ CC1+ cells.

P. The proportion of EGFP+ cells that were NG2+ in different regions at P14 and P60.

- Q. The proportion of EGFP⁺ cells that were CC1⁺ in different regions at P14 and P60.
- R. The proportion of EGFP⁺ cells that were Olig2⁺ in different regions at P14 and P60.

A Cspg4 int1-3b in chromatin immunoprecipitates with olig2 and Sox10

B H3K27Ac and H3K4me1 ChIP-seq across int1-3b and int2

Figure 5. ChIP to show binding of Olig2 and Sox10 and active histone marks at int1-3b.

(A) Scheme of the *Cspg4* gene showing the location of int1-3b and intron2 primers at the top. Arrows indicate primers used for ChIP-PCR. The image of a gel below shows PCR detection of int1-3b or intron 2 regions on immunoprecipitates formed from Oli-neu cell nuclear extracts using control rabbit IgG, rabbit anti-histone H3 antibody (positive control), rabbit anti-Olig2 antibody, and rabbit anti-Sox10 antibody. The expected size of int1-3b PCR product is 126 bp and is indicated by an arrowhead on the left margin. The expected size of intron 2 PCR product is 136 bp and is indicated by an arrowhead on the right margin. The positions of 200 bp and 100 bp on the DNA ladder are indicated by arrows. The faster

migrating band across all lanes at the bottom indicated by an asterisk are likely to be primer dimers.

(B) Genome browser tracks at the int1–3b and int2 loci showing ChIP-seq tracks from epiblast stem cell-derived OPCs. H3K4me1 and H3K27ac enhancer-associated histone modifications are enriched at int1–3b, but not at int2.

Figure 6. SoxE binding sites in Cspg4 int1-3b.

(A) Stimulation of Cspg4 int1-3b enhancer activity by SoxE proteins Sox9 and Sox10.

Luciferase activity in Neuro-2a cells co-transfected with Cspg4-int1-3b-luc construct and plasmids encoding the indicated transcription factors. Relative luciferase activity is Firefly luciferase activity normalized against Renilla luciferase activity. All values were normalized to no transcription factor (TF) controls. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. $n = 4$. One-way ANOVA, uncorrected Fisher's LSD test. Asterisks without brackets indicate significant difference compared to no TF control.

(B-D) Sox10-binding sites in Cspg4 int1-3b

(B) Luciferase assays in Neuro-2a cells comparing the enhancer activity of wt int1–3b or its various deletion mutants in the presence or absence of Sox10. Left: A schematic of the deletion mutants. Gray bars represent enhancer segments, lines represent deleted sequences, black bars are β -globin minimal promoter (mp), and white bars are Firefly luciferase coding sequence (luc). Right: Each of these constructs was transfected into Neuro-2a cells with (dark bars) or without (light bars) Sox10 expression construct. Values indicate Firefly luciferase activity normalized to Renilla luciferase activity, showing that Sox10-dependent increase in luciferase activity was detected with int1–3bi or int1–3biii but not with int1–3bii sequences the enhancer.

(C) Luciferase assay with int1–3b constructs mutated in the putative Sox binding motifs. There were two presumptive Sox binding motifs within 3bi and one in 3biii. Top: a schematic showing the positions (black bars) and sequences of SoxBE1 and SoxBE2 in int1–3bi and SoxBE3 in int1–3biii. Bottom: graph showing fold increase of int1–3bi-luc or int1–3biii-luc activity induced by Sox10 overexpression relative to control without Sox10. Values are relative Luciferase activity normalized to wt int1–3bi control without TF. $n=4$, two-way ANOVA, uncorrected Fisher's LSD test, * $p < 0.05$, **** $p < 0.0001$

(D) EMSA using Sox-binding motifs within the Cspg4 int1–3b enhancer. Biotinylated oligonucleotides were incubated with Sox10 DNA binding domain, separated by acrylamide gel electrophoresis, and detected by chemiluminescence. Shifted bands are indicated by arrowheads. Arrows indicate free probe. The intensity of the shifted bands was reduced by adding unlabeled wild type competitors (W) but not by mutant competitors (M) containing mutated sequences within the core binding motifs.

A Luciferase assay: int1-3b deletion mutants and bHLH proteins

B Luciferase assay: int1-3b bHLH BE1-4 mutants

C EMSA: Olig2 and Ascl1 binding to int1-3b bHLH BE1-4

Figure 7. bHLH binding sites in Cspg4 int1-3b

(A) Luciferase assays in Neuro-2a cells comparing the enhancer activity of wt 1.4 kb int1–3b or its various deletion mutants in the presence or absence of bHLH proteins. Left: A schematic of the deletion mutants. Gray bars represent enhancer segments, lines represent deleted sequences, black bars are β -globin minimal promoter (mp), and white bars are Firefly luciferase coding sequence (luc). Right: Each region was inserted upstream of luciferase and cotransfected into Neuro-2a cells with Olig2+E47 or Ascl1. Upregulation of luciferase activity was observed in the presence of Ascl1 with int1–3bi-luciferase and 1–3biii-luciferase constructs.

(B) Luciferase assay with int1–3b constructs mutated in the putative bHLH binding motifs. Top: a schematic showing the positions (black bars) and sequences of bHLH BE1 and BE2 in int1–3bi and bHLH BE3 and BE4 in int1–3biii. Bottom: Graphs showing changes in relative luciferase activity driven by bHLH BE1 and BE2 mutants in int1–3bi (left) and changes in the relative luciferase activity driven by bHLH BE3 and BE4 mutants in int1–3biii (right) induced by the bHLH Olig2+E47 (gray bar) or Ascl1 (black bars). Values were normalized to wt int1–3bi luciferase activity (left) or wt int1–3biii luciferase activity (right). $n=4$, two-way ANOVA, uncorrected Fisher's LSD test, ** $p < 0.01$, *** $p < 0.01$, and **** $p < 0.0001$).

(C) EMSA using oligonucleotides containing the bHLH binding motifs. [γ - ^{32}P]ATP-labeled or biotinylated oligonucleotides were incubated with in vitro-translated Olig2 or Ascl1 and separated by gel electrophoresis followed by chemiluminescence detection or autoradiography. Arrowheads indicated shifted bands. Free probe is seen at the bottom. The intensity of the shifted bands is reduced in the presence of unlabeled competitors (W) but not in the presence of unlabeled mutated competitors that have mutations within the core binding motifs (M).

Figure 8. Requirement of int1–3b for correct reporter expression

(A) Schema of the electroporated plasmids

(B-D) Number and distribution of EGFP+ and mCherry+ cells two weeks after electroporation of wt Tol2-Cspg4-EGFP (B) or mutated Tol2-Cspg4mut-EGFP (C) plasmids into P0 brains along with Tol2-CAG-mCherry and CAG-TPase plasmids. EGFP+ cells (arrowheads in B) are detected only after electroporation with wt Tol2-Cspg4-EGFP but not with mutated Tol2-Cspg4mut-EGFP. (D) Total number of mCherry+ cells and EGFP+ cells detected in each electroporated animal.

Scale bar in C indicates 50 μm.

(E) Left: Phenotype of mCherry+ cells two weeks after electroporation of in Tol2-CAG-mCherry and CAG-TPase plasmids into P0 brains. Right: Phenotype of EGFP+ cells two weeks after electroporation of Tol2-Cspg4-EGFP containing wt int1–3b enhancer and CAG-TPase plasmids into P0 brains.

(F) A section of a P14 mouse electroporated with Tol2-Cspg4-EGFP at P0 and stained for NG2 (magenta) and CC1 (cyan), showing an EGFP+ cell that is also NG2+. Scale bar indicates 20 μ m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript