Published in final edited form as:

Am J Cardiol. 2019 January 15; 123(2): 206–211. doi:10.1016/j.amjcard.2018.10.002.

Decade Long Trends (2001-2011) in the Incidence Rates of Initial Acute Myocardial Infarction

Robert J. Goldberg, PhD^a, Mayra Tisminetzky, MD, PhD^{a,b}, Hoang V. Tran, MD, MPH, PhD^c, Jorge Yarzebski, MD, MPH^a, Darleen Lessard, MS^a, and Joel M. Gore, MD^a

^aDepartment of Quantitative Health Sciences, University of Massachusetts Medical School, Worcester, MA, 01655

^bMeyers Primary Care Institute and the Division of Geriatric Medicine, University of Massachusetts Medical School, Worcester, MA

^cDepartment of Medicine, Bridgeport Hospital, Yale New Haven Health, Bridgeport, CT

Abstract

Despite the magnitude and impact of acute coronary disease, there are limited population-based data in the U.S. describing relatively recent trends in the incidence rates of acute myocardial infarction (AMI). The objectives of this study were to describe decade long (2001-2011) trends in the incidence rates of initial hospitalized episodes of AMI, with further stratification of these rates by age, sex, and type of AMI, among residents of central Massachusetts hospitalized at 11 area medical centers. The study population consisted of 3,737 adults hospitalized with a first AMI at 11 medical centers in central Massachusetts on a biennial basis between 2001 and 2011. The median age of this study population was 70 years, 57% were men, and 90% were white. Patients hospitalized during the most recent study years (2009/11) were younger, more likely to be men, have more comorbidities, and less in-hospital complications as compared with those in the earliest study years (2001/03). The overall age-adjusted hospital incidence rates (per 100,000 persons) of initial AMI declined (from 319 to 163), for men (from 422 to 219), women (from 232 to 120), for patients with a STEMI (129 to 56), and for those with an NSTEMI (190 to 107) between 2001 and 2011, respectively. In conclusion, the incidence rates of initial AMI declined appreciably in residents of central Massachusetts who were hospitalized with AMI during the years under study.

Keywords

acute myocardial infarction; population-based investigation; incidence rates

Address editorial correspondence to: **Robert J. Goldberg**, PhD, Division of Epidemiology of Chronic Diseases and Vulnerable Populations, Department of Quantitative Health Sciences, University of Massachusetts Medical School, 368 Plantation St, Worcester, MA 01605, Tel: 508-856-3991; Fax: (508) 856-8993, robert.goldberg@umassmed.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

Despite the magnitude and impact of coronary heart disease, a limited number of population-based studies in the United States have examined trends in the incidence rates of acute coronary disease (1–8). The population-based and longitudinal clinical/epidemiological studies that have monitored trends in the incidence rates of acute myocardial infarction (AMI) have noted relatively stable rates in the 1980's and 1990's with some evidence of declines thereafter (1–8). A limited number of studies have furthered our understanding of the descriptive epidemiology of AMI according to its major phenotypic manifestations of ST segment elevation AMI (STEMI) and non-ST segment elevation AMI (NSTEMI) (7,9–12).

The objectives of this community-wide observational study among residents of central Massachusetts hospitalized at all metropolitan Worcester medical centers with an independently validated AMI were to describe decade long trends (2001–2011) in the incidence rates of initial hospitalized episodes of AMI, with further stratification of these rates by age, sex, and type of AMI.

METHODS

The Worcester Heart Attack Study is a population-based investigation that is examining long-term trends in the incidence, hospital, and post-discharge case-fatality rates of AMI among residents of the Worcester metropolitan area hospitalized at all 11 greater Worcester medical centers in central Massachusetts on an approximate biennial basis over the past several decades (2,13–16).

Details of this study have been described previously (2,13–16). In brief, the medical records of residents of the Worcester metropolitan area hospitalized for possible AMI at all central Massachusetts medical centers on a biennial basis between 2001 and 2011 were individually reviewed and a diagnosis of AMI was validated according to predefined criteria. Because we were interested in examining trends in the magnitude of incident (initial) cases of AMI, patients with a history of previously diagnosed AMI (n=2,049) were excluded. This study was approved by the Institutional Review Board at the University of Massachusetts Medical School.

Data on patient's demographic characteristics, medical history, clinical, and in-hospital treatment data were abstracted from the hospital medical records of metropolitan Worcester residents with confirmed AMI by trained study physicians and nurses. Information was collected about patient's age, sex, comorbidities, type of AMI (STEMI vs. NSTEMI) (9), and hospital cardiac medications and diagnostic/interventional procedures. Information was also collected about the occurrence of clinically significant in-hospital complications including stroke (17), atrial fibrillation (18), heart failure (19), cardiogenic shock (20), and death.

Data analysis

Differences in the distribution of demographic and clinical characteristics between patients hospitalized for an initial AMI in the respective time periods were examined by using chi-square tests and ANOVA for categorical and continuous variables, respectively. We pooled our 6 individual study years into 3 two year time periods for purposes of these analyses and for ease of data interpretation. The annual incidence rates of AMI were calculated in a standard manner using available census data for residents of central Massachusetts. Age-adjusted incidence rates of AMI were computed by means of the direct method of standardization, using population census estimates for the greater Worcester population in 2010.

RESULTS

A total of 3,737 residents of central Massachusetts were hospitalized with an incident (initial) independently validated AMI at all 11 greater Worcester medical centers during the six study years between 2001 and 2011. The median age of this patient population was 70 years, 57% were men, and 90% were white.

Patients hospitalized during the most recent study years (2009/11) were significantly younger and included an increasingly greater proportion of men and individuals with a history of anemia, chronic kidney disease, hypertension, and hyperlipidemia but a lower proportion of those with previously diagnosed heart failure (Table 1). There was a significant increase in the proportion of individuals who presented with multiple previously diagnosed comorbidities over time. The risk of developing several major adverse events during the patient's acute hospitalization decreased during the years under study (Table 1).

In examining changing trends in the hospital incidence rates of AMI, the unadjusted incidence rates of initial AMI declined appreciably between our initial study year of 2001 (275 per 100,000) and our most recent study year of 2011 (165 per 100,000) (Figure 1). The age-adjusted incidence rates also decreased appreciably between 2001 (319 per 100,000) and 2011 (163 per 100,000) (Figure 2).

The age-adjusted incidence rates of initial AMI decreased by nearly one half between 2001 (422 per 100,000) and 2011 (219 per 100,000) for both men (Figure 3) and for women (232 per 100,000 in 2001) and (120 per 100,000 in 2011) (Figure 3).

The age-specific incidence rates of initial AMI declined among those less than 65 years old between 2001 (144 per 100,000) and 2011 (83 per 100,000), among individuals aged 65-74 years old between 2001 (745 per 100,000) and 2011 (310 per 100,000), and in individuals aged 75 years and older during similar study years (1,265 to 613 per 100,000, respectively) (Figure 4).

The age-adjusted incidence rates of an initial STEMI decreased between 2001 (129 per 100,000) and 2011 (56 per 100,000) as did the age-adjusted incidence rates of an initial NSTEMI (190 per 100,000 in 2001) (107 per 100,000 in 2011) (Figure 5).

DISCUSSION

The results of this clinical/epidemiological study suggest encouraging declines in the incidence rates of AMI during the years under study as well as changing demographic and clinical characteristics of those hospitalized with AMI.

Patients hospitalized with an initial AMI in this primarily Caucasian population tended to be younger, have a greater proportion of most of the previously diagnosed comorbidities we examined, and experienced favorable declines in hospital death rates and several clinical factors of prognostic importance during the most recent as compared with earliest study years.

In examining changes over time (1987-2006) in the characteristics of patients hospitalized with an incident AMI in Olmsted County, Minnesota, there were declines in the proportion of women, in those diagnosed in a higher Killip class, and in patients who developed a STEMI (8). In an analysis of more than 3.4 million patients who had a first listed diagnosis of AMI between 1979 and 2005 in the National Hospital Discharge Survey (6), there was an increase in the mean age of patients included during this period with an increasing proportion of women and of those with selected comorbidities. These changing trends in the descriptive epidemiology of AMI underscore the public health and clinical importance of monitoring contemporary trends in patient's demographic and clinical characteristics for purposes of carrying out surveillance of a major chronic disease in the U.S.

We observed encouraging declines in the incidence rates of AMI during the years under study overall and in patients further categorized according to their age, sex, and type of AMI. Several other studies have found similar findings. In a study of 3,810 incident AMI's that occurred among residents of Olmsted County, MN between 1995 and 2012, the age and sex adjusted incidence rates of AMI declined over time, with the most notable declines observed during the most recent years under study (21). Using data from more than 125,000 hospitalizations for AMI among adults enrolled in 13 U.S. health plans in the Cardiovascular Research Network between 2000 and 2008, the age and sex adjusted incidence rates of AMI decreased by an average of nearly 4% annually (22); the incidence rates of STEMI decreased by nearly 9% annually, whereas the incidence rates of NSTEMI increased and subsequently decreased over time.

Interpretation of trends in the incidence rates of AMI is difficult due to the characteristics of the populations studied, lack of information about the lifestyle and treatment practices of the population from which cases of AMI occurred, years under study, diagnostic criteria for AMI utilized, and whether hospital records were independently reviewed to validate the diagnosis of AMI.

The declines in the incidence rates of AMI that were observed in the present study as well as in prior investigations were most likely due to improvements in primary prevention efforts, including both lifestyle alterations and pharmacological interventions, though we lacked data on the major coronary predisposing factors and preventive practices among residents of central Massachusetts.

We found that the incidence rates of AMI declined for patients hospitalized with an STEMI or NSTEMI, with a larger decrease observed for patients with a STEMI. Other studies using unvalidated diagnostic codes also found similar trends with a larger decline observed in the occurrence rates of STEMI compared with NSTEMI during the periods under study (10). In a study of more than 125,000 AMI's identified by discharge codes in the Cardiovascular Research Network, the incidence rates of STEMI decreased by more than 50% between 2000 and 2008, whereas the incidence rates of NSTEMI slightly decreased during this period (22). Since the increased use of laboratory biomarker assays over time may have artifactually affected the observed trends in these incidence rates, it is possible that this effect is greater in patients who develop an NSTEMI where the electrocardiogram is less indicative of an AMI and the diagnosis of AMI is more often based on elevated cardiac biomarker findings.

We found similar relative declines of approximately 45% in the hospital incidence rates of initial AMI in both men and women residents of central Massachusetts between 2001 and 2011. In the Atherosclerosis Risk in Communities (ARIC) Study, the annual incidence rates of AMI declined by 4.3% among white men, 3.8% among white women, 2.9% among black women, and 1.5% among black men between 1987 and 2008 (23). Data from the U.S. and other countries have also shown a decrease in the incidence rates of AMI in both men and women (4,9,24,25).

We found that the incidence rates of initial AMI declined over time among patients in all age strata, with lesser absolute and relative declines observed in younger as compared with patients 65 years and older. In the Cardiovascular Research Network, the incidence rates of initial AMI declined by more than 4% annually among participants aged 55 years or older which was considerably greater than the annual decline of approximately 1% among younger HMO enrollees (7).

In the present era of changing patient characteristics, working definitions of AMI, and use of more aggressive lifestyle and treatment approaches to targeted high risk individuals, our findings highlight the importance of surveillance of acute coronary disease at the population level with further in-depth studies needed to attempt to disentangle the reasons for observed trends in AMI incidence rates in the U.S. (26).

Acknowledgements:

We wish to express our appreciation to all persons involved in the review of data for this project during the years under study. This study was made possible through the cooperation of the administration, medical records, and cardiology departments of participating Worcester metropolitan area hospitals.

Funding Sources: Grant support for this project was provided by the National Heart, Lung, and Blood Institute (RO1 HL35434). The efforts of Drs. Gore, Goldberg, and Yarzebski were supported in part by funding from the National Heart, Lung and Blood Institute U01HL105268.

Declaration of Interest: Grant support for this project was provided by the National Heart, Lung, and Blood Institute (RO1 HL35434). Drs. Gore, Goldberg, and Yarzebski's effort was supported in part by funding from the National Heart, Lung and Blood Institute U01HL105268.

REFERENCES

1. Rosamond WD, Chambless LE, Folsom AR, Cooper LS, Conwill DE, Glegg L, Wang CH, Heiss G. Trends in the incidence of myocardial infarction and in mortality due to coronary heart disease, 1987 to 1994. *N Engl J Med* 1998;339:861–867. [PubMed: 9744969]
2. Goldberg RJ, Yarzebski J, Lessard D, Gore J. A two-decade (1975 to 1995) long experience in the incidence, in-hospital and long-term case-fatality rates of acute myocardial infarction: a communitywide perspective. *J Am Coll Cardiol* 1999;33:1533–1539. [PubMed: 10334419]
3. Roger VL, Jacobsen SJ, Weston SA, Goraya T, Killian BB, Reeder GS, Kottke TE, Yawn BP, Frye RL. Trends in the incidence and survival of patients with hospitalized myocardial infarction, Olmsted County, Minnesota, 1979 to 1994. *Ann Intern Med* 2002;136:341–348. [PubMed: 11874305]
4. McGovern PG, Pankow JS, Shahar E, Doliszny KM, Folsom A, Blackburn H, Luepker R. Recent trends in acute coronary heart disease — mortality, morbidity, medical care, and risk factors. *N Engl J Med* 1996;334:884–890. [PubMed: 8596571]
5. Chen J, Normand ST, Wang Y, Drye EE, Schreiner BS, Krumholz HM. Recent declines in hospitalizations for acute myocardial infarction for Medicare fee-for-service beneficiaries: Progress and continuing challenges. *Circulation* 2010;121:1322–1328. [PubMed: 20212281]
6. Fang J, Alderman MH, Keenan NL, Ayala C. Acute myocardial infarction hospitalization in the United States, 1979 to 2005. *Am J Med* 2010;123:259–266. [PubMed: 20193835]
7. Yeh RW, Sidney S, Chandra M, Sorel M, Selby JV, Go AS. Population trends in the incidence and outcomes of acute myocardial infarction. *N Engl J Med* 2010;362:2155–2165. [PubMed: 20558366]
8. Roger VL, Weston SA, Gerber Y, Killian BB, Dunlay SM, Jaffe AS, Bell MR, Kors J, Yawn BP, Jacobsen SJ. Trends in incidence, severity, and outcome of hospitalized myocardial infarction. *Circulation* 2010;121:863–869. [PubMed: 20142444]
9. McManus DD, Gore JM, Yarzebski J, Spencer F, Lessard D, Goldberg RJ. Recent trends in the incidence, treatment, and Outcomes of patients with STEMI and NSTEMI. *Am J Med* 2011; 124:40–47. [PubMed: 21187184]
10. Montalescot G, Dallongeville J, Van Belle E, Rouanet S, Baulac C, Degrandart A, Vicaute E. STEMI and NSTEMI: are they so different? 1 year outcomes in acute myocardial infarction as defined by the ESC/ACC definition (the OPERA registry). *Eur Heart J* 2007;12:1409–1417.
11. Boyer NM, Laskey WK, Cox M, Hernandez AF, Peterson ED, Bhatt DL, Cannon CP, Fonarow GC. Trends in clinical, demographic, and biochemical characteristics of patients with acute myocardial infarction from 2003 to 2008: a report from the American Heart Association Get With The Guidelines Coronary Artery Disease. *JAMA* 2012;1:e001206. [PubMed: 23130159]
12. Roe MT, Messenger JC, Weintraub WS, Cannon CP, Fonarow GC, Dai D, Chen AY, Klein LW, Masoudi FA, McKay C, Hewitt K, Brindis RG, Peterson ED, Rumsfeld JS. Treatments, trends, and outcomes of acute myocardial infarction and percutaneous coronary intervention. *J Am Coll Cardiol* 2010;56:254–263. [PubMed: 20633817]
13. Goldberg RJ, Gorak EJ, Yarzebski Y, Hosmer DW, Jr, Dalen P, Gore JM, Alpert JS, Dalen JE. A communitywide perspective of sex differences and temporal trends in the incidence and survival rates after acute myocardial infarction and out-of-hospital deaths caused by coronary heart disease. *Circulation* 1993;87:1947–1953. [PubMed: 8504508]
14. Floyd KC, Yarzebski J, Spencer FA, Lessard D, Dalen JE, Alpert JS, Gore JM, Goldberg RJ. A 30 year perspective (1975–2005) into the changing landscape of patients hospitalized with initial acute myocardial infarction: Worcester Heart Attack Study. *Circ Cardiovasc Qual Outcomes* 2009;2:88–95. [PubMed: 20031820]
15. Tisminetzky M, McManus DD, Gore JM, Yarzebski J, Coles A, Lessard D, Goldberg RJ. Thirty Year Trends in Patient Characteristics, Treatment Practices, and Long-Term Outcomes of Adults Aged 35 to 54 Years Hospitalized With Acute Myocardial Infarction. *Am J Cardiol* 2014;113:1137–1141. [PubMed: 24507173]
16. Chen H-Y, Gore JM, Lapane KL, Yarzebski J, Person SD, Gurwitz JH, Kiefe CI, Goldberg RJ. A 35-year perspective (1975–2009) into the long-term prognosis and hospital management of patients

discharged from the hospital after a first acute myocardial infarction. *Am J Cardiol* 2015;116:24–29. [PubMed: 25933734]

17. Hariri E, Tisminetzky M, Lessard D, Yarzebski J, Gore J, Goldberg RJ. Twenty-five year (1986-2011) trends in the incidence and death rates of stroke complicating acute myocardial infarction. *Am J Med* 2018; (In Press).
18. Saczynski JS, McManus D, Zhou Z, Spencer F, Yarzebski J, Lessard D, Gore JM, Goldberg RJ. Trends in atrial fibrillation complicating acute myocardial infarction. *Am J Cardiol* 2009;104:169–174. [PubMed: 19576341]
19. Spencer FA, Meyer TE, Goldberg RJ, Yarzebski J, Hatton M, Lessard D, Gore JM. Twenty year trends (1975–1995) in the incidence, in-hospital and long-term death rates associated with heart failure complicating acute myocardial infarction: a community-wide perspective. *J Am Coll Cardiol* 1999;34:1378–1387. [PubMed: 10551682]
20. Goldberg RJ, Makam RC, Yarzebski J, McManus DD, Lessard D, Gore JM. Decade long trends (2001-2011) in the incidence and hospital death rates associated with the in-hospital development of cardiogenic shock after acute myocardial infarction. *Circ Cardiovasc Qual Outcomes* 2016;9:117–125. [PubMed: 26884615]
21. Gerber Y, Weston SA, Jiang R, Roger VL. The changing epidemiology of myocardial infarction in Olmsted County, Minnesota, 1995-2012. *Am J Med* 2015;128:144–151. [PubMed: 25261010]
22. Reynolds K, Go AS, Leong TK, Boudreau DM, Cassidy-Bushrow AE, Fortmann SP, Goldberg RJ, Gurwitz JH, Magid D, Margolis KL, McNeal CJ, Newton KM, Novotny R, Quesenberry CP, Jr, Rosamond WD, Smith DH, VanWormer JJ, Vupputuri S, Waring SC, Williams MS, Sidney S. Trends in incidence of hospitalized acute myocardial infarction in the cardiovascular research network (CVRN). *Am J Med* 2017;130:317–327. [PubMed: 27751900]
23. Rosamond WD, Chambless LE, Heiss G, Mosley TH, Coresh J, Whitsel E, Wagenknecht L, Ni H, Folsom AR. Twenty-two year trends in incidence of myocardial infarction, CHD mortality, and case-fatality in four US communities, 1987 to 2008. *Circulation* 2012;125:1848–1857. [PubMed: 22420957]
24. Briffa T, Nedkoff L, Peeters A, Tonkin A, Hung J, Ridout SC, Knuiman M. Discordant age and sex-specific trends in the incidence of a first coronary heart disease event in Western Australia from 1996 to 2007. *Heart* 2011 ;97:400–404. [PubMed: 21212133]
25. Schmidt M, Jacobsen JB, Lash TL, Bøtker HE, Sørensen HT. 25 year trends in first time hospitalisation for acute myocardial infarction, subsequent short and long term mortality, and the prognostic impact of sex and comorbidity: a Danish nationwide cohort study. *BMJ* 2012;344:e356. [PubMed: 22279115]
26. Huffman MD. Maturing methods for cardiovascular disease and stroke surveillance in the United States. *JAMA Cardiology* 2018;3:390 [PubMed: 29641822]

Figure 1:
Trends in the hospital incidence rates of initial acute myocardial infarction: Worcester Heart Attack Study

Figure 2:
Trends in the age-adjusted hospital incidence rates of initial acute myocardial infarction:
Worcester Heart Attack Study

Figure 3:
Trends in the age-adjusted hospital incidence rates of initial acute myocardial infarction by sex: Worcester Heart Attack Study

Figure 4:
Trends in the age-adjusted hospital incidence rates of initial acute myocardial infarction by age group: Worcester Heart Attack Study

Figure 5:
Trends in the age-adjusted hospital incidence rates of initial acute myocardial infarction (AMI) by type of AMI: Worcester Heart Attack Study

Table 1:

Characteristics of patients hospitalized with an initial acute myocardial infarction according to time period

Characteristic	2001/03 (n=1,547) (%)	2005/07 (n=1,172) (%)	2009/11 (n=1,018) (%)
Age (years)			
<65	35.3	35.9	43.3
65-74	20.9	17.8	19.8
75	43.8	46.3	36.8
Male	56.9	54.9	59.1
White race	87.9	90.6	91.9**
NSTEMI	61.5	64.6	65.5
Length of stay (days, median)	4	4	3
Medical history			
Anemia	4.5	11.8	10.0**
Atrial fibrillation	11.8	12.2	11.0
Chronic kidney disease	11.8	17.5	18.0**
Chronic obstructive pulmonary disease	17.3	13.6	13.3*
Depression	14.5	15.8	16.2
Diabetes	28.0	28.8	31.7
Heart failure	15.0	16.5	12.2*
Hyperlipidemia	44.2	55.5	65.6**
Hypertension	64.8	70.6	69.7*
Stroke	9.3	9.2	7.5
Number of comorbidities			**
0-1	33.9	29.4	27.9
Any 2	28.1	26.1	24.5
Any 3	20.2	20.4	22.4
4 or more	17.7	24.2	25.3
Complications during hospitalization			
Atrial fibrillation	21.7	21.5	15.1**
Cardiogenic shock	4.9	6.1	4.6
Heart failure	34.4	34.0	28.2*
Stroke	1.7	0.6	1.8*
Death	9.6	9.8	6.5*

*
P<.05**
P<.01***
P<.001