EPA Public Access

Author manuscript

Environ Sci Technol. Author manuscript; available in PMC 2019 September 04.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Environ Sci Technol. 2018 September 04; 52(17): 10067–10077. doi:10.1021/acs.est.8b02169.

Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) Transcriptome Profile Changes Induced by Diesel Emissions Generated with CeO₂ Nanoparticle Fuel Borne Catalyst

Jay R. Reichman^{†,*}, Paul T. Rygielwicz[†], Mark G. Johnson[†], Michael A. Bollman[†], Bonnie M. Smith[†], Q. Todd Krantz[‡], Charly J. King[‡], Kasey D. Kovalcik[§], and Christian P. Andersen[†]

[†]Western Ecology Division, National Health and Environmental Effects Laboratory, US Environmental Protection Agency, Corvallis, Oregon, 97333, USA

[‡]Environmental Public Health Division, National Health and Environmental Effects Laboratory, US Environmental Protection Agency Research Triangle Park, North Carolina, 27711, USA

[§]Exposure Methods and Measurements Division, National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina 27711, United States

Abstract

It is important to understand molecular effects on plants exposed to compounds released from use of products containing engineered nanomaterials. Here, we present mRNA sequencing data on transcriptome impacts to Douglas-fir following 2 weeks of sublethal exposure to 30:1 diluted airborne emissions released from combustion of diesel fuel containing engineered CeO₂ nanoparticle catalysts (DECe). Our hypothesis was that chamber exposure to DECe would induce distinct transcriptome changes in seedling needles compared with responses to conventional diesel exhaust (DE) or filtered DECe Gas Phase. Significantly increased uptake/binding of Ce in needles of DECe treated seedlings was 2.7X above background levels and was associated with altered gene expression patterns. All 225 Blast2GO gene ontologies (GOs) enriched by up-regulated DECe transcripts were nested within GOs for DE, however, 29 of 31 enriched GOs for down-regulated DECe transcripts were unique. MapMan analysis also identified 3 pathways enriched with DECe

*Correspondence: Jay R. Reichman, Western Ecology Division, National Health and Environmental Effects Laboratory, US Environmental Protection Agency, Corvallis, Oregon, 97333, USA. Tel: 541-754-4643. reichman.jay@epa.gov.

Supporting Information

Detailed Materials and Methods. Data quality indicators for inorganic ions by ICP-MS. Generalized schematic of exposure configuration. Distribution of BLASTX Hits, B2GO mapping & annotations for psme_transcriptome v203. Distribution of Mercator assignments for psme_transcriptome v203. Sets of genes that were significantly regulated by DE, DECe and DECe Gas Phase treatments. Blast2GO gene ontologies enriched for genes that were significantly regulated by DE, DECe and DECe Gas Phase treatments. MapMan pathway bins enriched for genes that were significantly down-regulated by DE, DECe and DECe Gas Phase treatments. Unique Blast2GO gene ontologies enriched for genes that were significantly down-regulated by DECe treatment. Unique MapMan pathway bins enriched for genes that were significantly down-regulated by DECe treatment. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Data availability

The RNA-sequencing data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus^{81–82} and are accessible through GEO Series accession number GSE107902: (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE107902>).

Disclaimer

This manuscript has been subjected to Agency review and has been approved for publication. The views expressed in the present study are those of the author(s) and do not necessarily reflect the views or policies of the US Environmental Protection Agency. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

down-regulated transcripts. There was prominent representation of genes with attenuated expression in transferase, transporter, RNA regulation and protein degradation GOs and pathways. CeO₂ nanoparticle additive decreased and shifted molecular impact of diesel emissions. Widespread use of such products and chronic environmental exposure to DECe may adversely affect plant physiology and development.

Graphical Abstract

Keywords

plant; RNA-sequencing; metal oxide engineered nanomaterials; EMN-enabled products

Introduction

The vast majority of effects research conducted on engineered nanomaterials (ENMs) involves direct exposure to pristine particles. These investigations are important for revealing physiochemical characteristics of individual ENMs, and can identify specific surface properties or other characteristics that can lead to toxicity. While these studies can demonstrate important ENM properties, they provide little insight into the actual risk these materials pose to environmental health and safety since most are incorporated into products that likely alter both the toxicity and distribution of the nanoparticles once they reach the environment. There is a need for empirical data on the impacts to organisms persistently exposed to compounds that are released into the environment from the use, degradation or weathering of products containing ENMs, which themselves may become transformed in the process.¹

Metal oxide nanoparticles are widely used in surface coatings and other materials, including additives to diesel as fuel borne catalysts (FBCs) to improve fuel economy and increase oxidation of particulates.² When cerium dioxide nanoparticles (nCe) are added to diesel fuel, burning efficiency increases, producing emissions with characteristics that differ from conventional diesel exhaust (DE).²⁻⁵ A key difference of diesel exhaust with nCe (DECe) is that it contains potentially reactive forms of Ce, which are released into the environment in aerosols that can be inhaled and settle on various surfaces including soils and plants. Previous analyses showed that laboratory-generated DECe samples contained nano CeO₂

agglomerated within the soot matrix⁶ and mixtures that also include individual CeO₂ nanoparticles.⁷ A recent report detailed the transformations of nCe within the diesel fuel additive Envirox™ (Energenics Europe Ltd., Kidlington, Oxfordshire, U.K.) during combustion from a small engine.⁸ Surfactant coated aggregates of the original FBC were melted and recrystallized following combustion. The resulting 50–300 nm, near-spherical, crystals released within the DECe had superstructures containing Ce(III), and approximately 30% of those had a new hydrocarbon coating.

There is now widespread on-road use of Envirox™ and similar products in Canada, Europe and Asia Pacific, which increases the likelihood of near-road exposure of organisms to nCe.^{6, 9–10} For example, a 5-fold increase of environmental deposition of cerium from FBC (~0.3 ng/m³) above ambient was measured near a bus depot in Newcastle-on-Tyne, UK, where the only likely source of extra cerium was from a single bus fleet using Envirox™.^{9, 11} Since 2011, nCe FBCs have only been registered for off-road use in the United States. Because off-road vehicles include diesel farm equipment, there is currently potential in the US for crop plants and other vegetation to be exposed to nCe FBCs. Modelling simulations also indicate that if nCe FBCs were used in all on-road and off-road diesel vehicles in the Central and Eastern US, there would be an increase of surface-level atmospheric cerium concentrations from an estimated average of 0.5 ng/m³ up to 22 ng/m³, especially near highways and in large cities.¹⁰

Given the prolonged exposures to DECe that are currently taking place in several countries, it is important to understand how exposure to these emissions affect various organisms.¹² It has previously been shown that rats exposed to DECe had more adverse pulmonary effects than from DE alone. That DECe exposure also resulted in Ce accumulation in lung tissue and systemic translocation of Ce to the liver.¹³ Recently, exposure of human alveolar epithelial cells to diesel particulate matter derived from FBCs-doped fuels showed decreased viability and altered gene expression patterns within a broad range of biochemical pathways, which increases human health concerns.¹⁴ Effects of DECe on plants have not been reported to date.

Here, we focused on molecular-level impacts of DECe to Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco), which is one of the most ecologically and economically important tree species in the world.^{15–16} We tested the hypothesis that 2 weeks of controlled exposure to DECe will induce certain distinct transcriptome changes in Douglas-fir needles compared to responses to DE or filtered DECe Gas Phase. Specific objectives of these experiments were to: 1) Characterize the physical and chemical properties of diesel emissions (both gas- and particle-phase) generated by combustion of diesel fuels ± nano-CeO₂ FBC; 2) Quantify Ce deposition on needles and that which was also bound to or taken up by needles; 3) Compare transcriptome changes in Douglas-fir seedlings exposed to DECe, DE and DECe Gas Phase to Air Controls via mRNA-sequencing; and 4) Identify unique enrichment of gene ontologies and pathways with significantly regulated transcripts. The results demonstrate important effects of exposure to compounds released from use of an EMN-enabled consumer product.

Materials and Methods

Generation and characterization of DE, DECe and DECe gas phase emissions

Emissions for plant exposure experiments were generated using two identical Yanmar L70 diesel engines. The base-fuel run in one engine was ultralow sulfur diesel (<16 ppm.), while the nCe doped fuel used in the second engine was prepared by adding Envirox™ at 10 ppm concentration.

Four Hazelton 1000 (984 L) exposure chambers (one each for DE, DECe, DECe Gas Phase, and Air Control) were each outfitted with two LED lights to supply photosynthetically active radiation to plants during the exposures. The DE and DECe chambers were supplied with whole exhaust that was diluted 30:1 with HEPA filtered air resulting in the target particle concentration of ~1000 µg/m³. The DECe gas phase chamber utilized the same diluted emissions but the particles were filtered out using a HEPA filter. The air control chamber received charcoal / Purafil HEPA filtered air. A schematic of the exposure configuration is shown in Supporting Information Figure S1.

All treatment chamber concentrations (exhaust dilutions) were controlled by periodic adjustments of the dilution air based on continuous particulate matter mass concentrations. Other chamber parameters that were measured included particle size, concentrations of gasses (O₂, CO, NO, NO₂ and NO_X) as described in ¹⁷, as well as temperature and relative humidity. The overall exposure regime was 6 hours/day × 5 days/week × 2 weeks. Filtered air was pumped through all chambers during the times between the exposures. Statistical differences between mean chamber parameters were determined by ANOVA with post-hoc Tukey HSD multiple comparison procedures, which were performed in R v3.1.1.¹⁸ Adjusted *P*-values < 0.05 were considered significant.

Tree Seedling Source and Growth Conditions

Douglas-fir seedlings were obtained from a commercial nursery (Weyerhaeuser, Aurora, OR), and half-sibling plants were used to minimize genetic variation among individuals. Seedlings were individually planted in potting media amended with a slow-release fertilizer (14–14–14 nitrogen-phosphate-potash) with minor elements. In each chamber, the LED lights were set on a 16-hour light:8-hour dark cycle and plant pots were placed in a shared tray. Seedlings were bottom-watered daily with deionized water, which was added directly to the trays. There were 7 seedlings per chamber and plants were re-distributed within the exposure chambers mid-way through the exposure period to minimize potential edge effects. Chamber experiments were run during the month of December prior to bud burst and needle growth and the associated transcriptional shifts that typically begin in early spring in the northern hemisphere.¹⁹

Inductively Coupled Plasma Mass Spectrometry (ICP-MS)

ICP-MS was used to quantify the total Ce deposited on needles during exposure and that which was still bound to the plant tissue after washes. Following the exposure period, all needles were collected from seedlings in each treatment. To separate the amount of Ce adhering to needle surfaces from that which was embedded or taken up, needles were

washed in 100 mL of pH 4.5 simulated rainwater²⁰ and dried at 60°C. The wash was retained.

Needle wash solutions and ~5 g of biomass representing 100s of ground needles/sample were digested in preparation for ICP-MS analysis following US EPA Method 3010A.²¹ An Element 2™ Inductively Coupled Plasma Mass Spectrometry Instrument (Thermo Scientific, Inc., Waltham, MA) was used to analyse the sample digests to quantify Ce and other metals. Data quality standards for the ICP-MS analysis are shown in Supporting Information Table S1. Amounts of Ce were reported as ng of Ce/cm² of Douglas-fir needle surface area based on a subset of needles from 5 randomly chosen plants from each control and treatment group analyzed. A conversion factor of 42.8 cm²/g was derived from the total dry needle area and mass measured in triplicate for these seedlings. The area to biomass ratio was then used to estimate needle area for the specific mass of needles analyzed per sample. The total Ce deposited was estimated from the sum of the Ce in the wash plus the Ce taken up by the needles. Statistical differences between mean Ce levels in samples were also determined by ANOVA with post-hoc Tukey HSD multiple comparison procedures, as described above.

RNA-Sequencing

Samples, library preps and data collection—Immediately after the exposure ended, ~30 needles (~200 mg) grown in the previous year were collected from each seedling and flash-frozen in liquid nitrogen. Total RNA was extracted from needles using PureLink® Plant RNA Reagent per the manufacturer's protocol (Ambion® by Life Technologies™, Inc. Austin, TX). The integrity of RNA samples was quantified with the NanoDrop 1000 (Thermo Scientific, Inc.). RNA concentrations were determined with a Qubit Fluorometer (Thermo Scientific, Inc.) and stored at –80°C until further processing.

Messenger RNA was purified from total RNA samples with FastTrack® MAG mRNA Isolation Kits (Ambion, Inc.). Then, RNA-sequencing libraries were prepared with PrepX RNA-Seq for Illumina Library Kits using an Apollo 324™ automated system (Wafergen Biosystems, Inc, Freemont, CA). Clusters were amplified from pooled libraries with TruSeq PE Cluster Kit v3-cBot-HS kits on a cBot System (Illumina Inc., San Diego, CA). RNA sequencing data were collected on a HiScan™ SQ System and base calls were made with Illumina Casava version 1.8.2 (Illumina Inc.).

Templated assemblies—Illumina reads for each of our seedlings were auto trimmed and aligned to a fasta file of 173,882 tree-derived sequences from the Douglas-fir psme_transcriptome_v203 (GenBank: GFFY000000000.1)^{15, 19, 22} using SeqMan NGen v13.1 (DNASTAR, Inc., Madison, WI).

Gene expression analyses—Gene expression levels were quantified using QSeq/ArrayStar v13.1 (DNASTAR, Inc.) with Reads Assigned per Kilobase of target per Million mapped reads (RPKM) Normalization. Sets of regulated transcripts in treated samples were identified as those with 2-fold change (Log₂) and 95% confidence across biological replicates compared to Air Controls. The confidence was based on Student's t-test ($P < 0.05$) with Benjamini & Hochberg Multiple Testing Correction (FDR < 0.05).

Transcript identification, mapping, annotation and enrichment analyses—The identities of psme_transcriptome_v203 template sequences were updated by blast-2.3.0+ BLASTX²³ against the non-redundant (nr) database downloaded from NCBI on 1/14/2016. The BLASTX output was imported to Blast2Go v3.3 (BioBam, Inc. Valencia, Spain) for mapping, annotation and gene ontology (GO) enrichment analyses. Statistically significant enrichment of GOs was determined by Fisher's exact test ($P \leq 0.05$) with Benjamini & Hochberg Multiple Testing Correction (FDR ≤ 0.05).

Also, pathway enrichment analyses of regulated transcripts were performed with MapMan v3.5.1R2.²⁴ To produce the mapping file needed for MapMan analyses, psme_transcriptome_v203 sequences were uploaded to the Mercator Pipeline for Automated Sequence Annotation at <http://mapman.gabipd.org/web/guest/app/mercator>.²⁵ Enrichment of MapMan pathway bins was determined by Wilcoxon Ranked Sum tests ($P \leq 0.05$) with Benjamini & Hochberg multiple testing correction (FDR ≤ 0.05).

Detailed Materials and Methods are available in the Supporting Information.

Results & Discussion

Characterization of Exposures and Ce Uptake

The average particle concentrations for the DE and DECe exposures were 1157.8 ± 187.0 (S.E.) and $1002.6 \pm 7.6 \mu\text{g}/\text{m}^3$ respectively. These values were not significantly different from each other and both were near the target concentration of $1000 \mu\text{g}/\text{m}^3$ (Table 1). The average particle volume measured for DECe was 15% higher than that for DE, however, the difference was not significant. It is likely that the DECe had a smaller average particle size than the DE at the exit of the engine, as would be expected. The particle size measurements were taken downstream after dilution and transport that may have caused minor volumetric increase or agglomeration in the DECe particulates. Overall, the particle concentration and volumes for DE and DECe were more like each other than those within the DECe Gas Phase or Air Controls, both of which were filtered. The few counts of larger particles at low concentrations within the DECe Gas Phase and Air Control chambers were most likely condensed vapors including water that passed through the filter. Concentrations of CO, NO, NO₂, NO_x and Relative Humidity were higher for DECe than DE, and HEPA filtration to produce the DECe Gas Phase did not appreciably change these parameters. Thus as planned, the chamber gasses for DECe Gas Phase closely resembled those of the DECe, allowing direct comparison between treatments.

Analyses of ICP-MS data demonstrated that Ce amounts in and on the Douglas-fir needles from the DECe exposure were significantly greater than that in all other treatments (Figure 1). These sums included Ce taken up by needles and forms of Ce loosely adhering to the exterior of needles from the emissions plus the background Ce. The mean for total Ce present in DECe samples was $25 \pm 1.1 \mu\text{g}/\text{cm}^2$. The total Ce in and on the DECe Gas Phase needles was 84.7 % less than that for DECe and was not significantly different from the Air Control (Figure 1A). Simulated rainwater washes removed over 99% of the excess Ce from DECe exposed needle exteriors and essentially all from the DECe Gas Phase samples. Nevertheless, there were still $11.6 \pm 0.8 \text{ ng}/\text{cm}^2$ of Ce that were either taken up by the DECe

needles or securely bound to the needle surfaces. This remaining quantity was significantly higher than the amount in any other treatment (Figure 1B).

RNA-Sequencing Assemblies & Comparison of Regulated Transcripts

Table 2 shows SeqMan NGen statistics for the 26 Douglas-fir mRNA-Seq assemblies in this project. A sample number of 7 was planned for each experimental condition, however, sequence data were not available for 1 DECe Gas Phase sample and 1 Air Control due to insufficient RNA yield from these samples. The mean number of trimmed Illumina reads assembled per sample ranged from $3,080,314 \pm 1,347,303$ for DE up to $10,776,209 \pm 1,732,380$ for Air Controls. In all cases, reads that were assembled had Seq Scores of 90–100% match to the psme_transcriptome_v203 template.

Under all three treatments, there were substantially more down-regulated than up-regulated transcripts compared to Air Controls (Figure 2A-C). The percentages of down-regulated genes were 97.7%, 94.2% and 98.3% for DE, DECe and DECe Gas Phase, respectively. The number of regulated genes (2-fold change and 95% confidence in expression level) and R^2 (linear correlation) for each comparison were as follows: A) DE, 39,372, 0.58; B) DECe, 4,873, 0.87; C) DECe Gas Phase, 1944, 0.91. Figure 2D is a heatmap generated by hierarchical clustering of expression patterns for the combined set of 39,910 regulated genes across the treatments (see values in compartments of Figure 3). DE contained an abundance of strongly regulated transcripts as is evident from the heatmap.

Numerically, the DECe and DECe Gas Phase sets of differentially expressed genes had more in common with those of DE than with each other (Figure 3). There were 4554 regulated genes shared between DECe and DE and 1666 in common between DECe Gas Phase and DE. On the other hand, only 938 such genes were shared between DECe and DECe Gas Phase. Excluding shared genes, there were 34,031, 260 and 219 genes that were uniquely regulated under DE, DECe and DECe Gas Phase treatments respectively (Supporting Information Table S2). It is surprising that the simple addition of the CeO₂ nanoparticle fuel borne catalyst so effectively reduced the number of regulated genes in seedlings exposed to DECe vs. DE. A detailed analysis of DE induced transcriptome changes is beyond the scope of this paper, but clearly use of the FBC in our experiments decreased and shifted the molecular impact of the emissions in Douglas-fir needles.

While the DECe set had the greatest proportion of up-regulated genes (281 out of 4873), only 7.5% of these were unique to this exposure (Figure 3). By contrast respectively, 51.5% and 70.9% of genes that were up-regulated under DECe Gas Phase and DE were unique to those treatments. Two hundred and sixty DECe up-regulated genes were shared with the other treatments, and 92% were in common with DE. The majority (91.9%) of transcripts that were significantly regulated uniquely in DECe treated seedlings were down-regulated.

Transcript identification, mapping, annotation and bin assignments

To evaluate the importance of differential expression patterns of mRNAs, it is essential to determine the identity of as many regulated transcripts as possible, what proteins they code for and what physiological roles those proteins play. There were 105,696 tree-derived nuclear, chloroplast and mitochondrial sequences within the psme_transcriptome_v203

template that met the thresholds for BLASTX hits against the NCBI nr database. Of these identified transcripts, 68,574 had successful Blast2GO gene ontology mapping, and 32,122 were additionally annotated with enzyme codes. A group of 68,186 template sequences were Blasted without hits, and another 25,741 were excluded without analysis because they were previously identified as rRNA or originating from bacteria or fungi (Supporting Information Figure S2).

Mercator assigned 30,489 psme_transcriptome_v203 sequences to MapMan bins in the process of generating a mapping file for pathway analyses. There were 143,393 transcripts not assigned. The same set excluded from Blast2GO mapping and annotation was also omitted from Mercator assignments (Supporting Information Figure S3).

Enrichment of Gene Ontologies and Pathways Unique to DECe Exposure

The three sets of genes significantly regulated by DE, DECe and DECe Gas Phase were each split into up- and down-regulated subsets for separate GO enrichment analyses. Blast2GO gene ontologies enriched by significantly up-regulated DECe transcripts were nested within those for DE, however, 93.5% of enriched GOs for significantly down-regulated DECe transcripts were unique (Supporting Information Table S3). Figure 4 shows the rank order number of DECe down-regulated transcripts in the test sets statistically associated with each of these enriched Molecular Function, Biological Process, and Cellular Component GOs. DECe attenuated expression of genes that affect functions of transferases, kinases, transmembrane transporters, transcription factors, diester hydrolases and RNA polymerase III; processes of protein phosphorylation, responses to stimuli, hormones and chemicals, carbohydrate transport, RNA polymerase III transcription, and cellular anion homeostasis; plus, components of the plasma membrane.

The same subsets of up- and down-regulated genes were also utilized for pathway enrichment analyses. There were no MapMan bins that were enriched with DECe up-regulated transcripts at the corrected P -value of 0.05. However, MapMan identified RNA regulation of transcription, protein degradation, and lipid metabolism pathways that were uniquely enriched with DECe down-regulated transcripts (Supporting Information Table S4). Figure 5 presents the rank order number of DECe down-regulated transcripts for these pathways. The prominent overlaps between the GO and pathway results are effects on proteins and control of transcription.

Although the majority of Douglas-fir transcripts significantly regulated by DECe exposure were similarly affected by DE exposure, DECe induced several transcriptome profile changes that were distinct, supporting our central hypothesis (Figure 3; Supporting Information Table S3). Even though the DE treatment resulted in more total transcriptome changes overall when compared to DECe, suggesting greater overall impact of the conventional diesel emissions, there were still important responses in DECe exposed seedlings that could result in altered health. The particular systems affected by DECe may point to potential molecular initiating events (MIEs) that can be explored to determine if they lead to adverse outcome pathways (AOPs) in plants.²⁶

Important DECe Transcriptome Changes That May Affect Physiology & Development

Here we discuss various biological roles of proteins that are coded for by the most frequently represented transcripts within DECe enriched GOs and pathways. DECe driven reduction of gene expression (and protein levels) in the 4 categories identified below may adversely affect plant physiology and development.

Transferase Activities—Our results demonstrate that there was substantial impact on the Douglas-fir seedlings from DECe exposure on the molecular function of *Transferase Activities, Transferring Phosphorus-Containing Groups*. This GO was enriched by 177 down-regulated transcripts (Figure 4; Supporting Information Table S5). Transcripts enriching the *Kinase Activity* and *Protein Kinase Activity* GOs were co-assigned to this broader group. Affected transcripts include calcium-dependent protein kinases (CDPKs) that play essential roles in signaling for pathogen-associated defense, responses to hormones and growth.^{27–32} Calcineurin B-like protein (CBL)-interacting protein kinases (CIPKs) are involved in decoding cellular influx signatures of Ca^{2+} that are early signaling events resulting from environmental stressors.^{33–36} Cysteine-rich receptor kinases (CRKs) control aspects of plant development and oxidative stress adaptation. Group V CRKs are involved in stomatal closing following reactive oxygen species production by NADPH oxidase that occurs after intracellular Ca^{2+} signaling.³⁷ G-type lectin S-receptor-like serine threonine-kinases (GsSRKs) are regulated by Ca^{2+} /CaM and mediate plant tolerance to abiotic salt stress.^{38–40} Leucine-rich repeat (LRR) receptor kinases (LRR-RKs) such as flagellin sensing 2 (FLS2) transduce variable signals from the environment, into changes in nuclear gene expression and thereby work to coordinate growth-defense trade-offs.^{41–42} L-type lectin-domain containing receptor kinases (LecRKs) are important to both plant developmental and adaptive processes. LecRKs contain a partially conserved hydrophobic-binding site that may recognize small hydrophobic ligands, such as plant hormones or microbial associated molecular patterns.⁴³ Hormone-sensitive lipase 1 (HSL1) is one of several downstream receptor kinases involved in response to endoplasmic reticulum stress needed for cell survival.⁴⁴ Without adequate expression of transferase genes, several critical plant biochemical systems would fail.

Transmembrane Transporter Activity—Another prominently affected molecular function GO was that of *Transmembrane Transporter Activity*, which was enriched by 102 down-regulated transcripts (Figure 4; Supporting Information Table S5). Among Douglas-fir subtypes, ATP-binding cassette (ABC) transporter proteins are localized in multiple membrane types such as plasma membranes, tonoplasts, chloroplasts, mitochondria and peroxisomes.⁴⁵ ABC transporters are involved in auxin transport, cuticle formation, lipid catabolism, xenobiotic detoxification, disease resistance, stomatal guard cell function, and male reproductive tissue development.^{46–48} Certain ABC transporters have been shown to have calcium-dependent signal transduction, modulate calcium homeostasis, and be sensitive to calcium channel inhibition by lanthanide series metal cations.^{45–46}

Aquaporin membrane channels are particularly important in plant functions since they enable transport of water and small solutes and dissolved gasses across biological membranes.⁴⁹ In plants, aquaporins play key roles in hydraulic regulation and nutrient

transport in roots and leaves. They contribute to plant growth, development and responses to stress. The gating of plasma membrane intrinsic proteins (PIP) aquaporins, can be post-translationally regulated by phosphorylation, Ca^{2+} and protons (H^+).⁵⁰ NOD26-like intrinsic protein (NIP) aquaporins localize to the plasma membrane or the endoplasmic reticulum. Many NIPs have a CDPK phosphorylation site at the carboxyl terminus.⁴⁹

The calcium transporters affected by DECE in Douglas-fir include mitochondrial calcium uniporters (MCUs) and plasma membrane calcium-transporting ATPases (PMCA). MCUs complex with other macromolecules that function together to control Ca^{2+} up-take across the inner membrane of mitochondria. MCUs are important in the regulation of aerobic metabolism and cell survival.^{51–53} PMCA are critical ion pumps that expel Ca^{2+} from the cytosol to restore intracellular Ca^{2+} resting concentration after transitory elevation during Ca^{2+} signaling.⁵⁴ Movement of many macromolecules, ions and dissolved gasses across membranes is dependent on transporters.

RNA Regulation of Transcription—The *RNA Regulation of Transcription* pathway bin was enriched by 202 DECE down-regulated transcripts, which was the largest set identified by MapMan analysis (Figure 5; Supporting Information Table S6). Here, DECE exposure reduced expression of genes that code for members of several transcription factor/binding protein families. These polypeptides themselves target the promoters and control expression of hundreds of downstream regulon genes. APETALA2/Ethylene-responsive element binding proteins (AP2/EREBP; AP2/ERF) are a large family of plant specific transcription factors that includes four major subfamilies; AP2, ERF, dehydration-responsive elements-binding proteins (DREBs) and related to ABI3/VP1 proteins (RAVs). AP2/EREBPs variously regulate genes needed to protect cells and sustain growth in response to abiotic stressors such as cold, dehydration, heat shock and salinity.^{55–56} Nineteen Douglas-fir DECE down-regulated AP2/EREBP transcripts coded for DREB A-5s, ERF B-3s, B-4s, B-5s and RAV2. These results are in contrast to recently reported up-regulation of AP2/EREBP *DREB1B/CBF1*, *DREB1C/CBF2*, *DREB1D/CBF4* and *DREB2A* genes in *Arabidopsis thaliana* leaves following exposure to pristine CeO_2 ENMs.⁵⁷ Although down-regulation of these genes differed from earlier results using native CeO_2 particles, the sensitivity of these gene complexes to both CeO_2 nanoparticles and DECE indicates they may alter the plants ability to deal with naturally occurring stresses. While these differences may be related to morphological and physiological distinctions between the species, they could be driven by the altered reactivity of DECE cerium components vs. pristine CeO_2 ENMs and routes of exposure in the experiments.

Basic-helix-loop-helix (bHLH) transcription factors are important regulators of diverse systems in plants including anthocyanin biosynthesis, phytochrome signaling, globulin expression, fruit dehiscence, carpel and epidermal development.⁵⁸ They are also involved in a wide range of developmental processes and in response to biotic and abiotic stresses.⁵⁹

Douglas-fir had several regulated zinc finger binding protein transcription factors including DNA-binding with one finger proteins (DOFs), such as C2C2 DOFs (high cambial activity 2 (HCA2) & cycling DOF factor 3 (CDF3). DOFs are associated with several plant-specific phenomena including light, phytohormone and defense responses, seed development and

germination.⁶⁰ GATA factors contain a class IV zinc finger motif. In plants, GATA DNA motifs have been implicated in light-dependent and nitrate-dependent control of transcription.⁶¹ GATA-9 is under circadian regulation and is involved in control of light-responsive, tissue-specific transcription.⁶²

Homeobox transcription factors with a homeodomain associated to a leucine zipper (HD-ZIPs) are unique to plants. Class I HD-ZIPs are regulated by external factors such as drought, extreme temperatures, osmotic stresses and illumination conditions. Expression is specific to different tissues and organs of plants. They have roles in development and responses to environmental conditions, particularly abiotic stresses. Class III HD-ZIPs function as developmental regulators of apical meristem, vascular bundles, adaxial domains of lateral organs and auxin transport.⁶³

MYB (myeloblastosis) domain transcription factors are key in regulatory networks controlling development, metabolism and responses to abiotic and biotic stresses.⁶⁴ Among Douglas-fir regulated transcripts, 21 MYBs code for pentatricopeptide repeat (PPR)-like proteins that play essential roles in mitochondria and chloroplasts.⁶⁵ PPRs are involved in many post-transcriptional processes such as splicing, editing, processing and translation of chloroplast and mitochondrial RNAs.^{66–67}

NAC (NAM, ATAF1,2, CUC2) domain transcription factors function in plant embryonic, floral and vegetative development. NACs are also involved in lateral root formation, auxin signaling, defense against pathogens and abiotic stress responses.⁶⁸

The DECE treatment down-regulated transcripts for Douglas-fir homologs of WRKY domain transcription factors 6, 7, 21, 31, 35, 41, 42, 47, 53 and 72. These WRKYs mediate responses to cold, drought, heat stress, herbivory and salt. They are also important in adapting to phosphorus deprivation, ROS signaling, senescence processes and sugar starvation signaling. See references within reviews.^{69–72} As noted above, our results contrasted with responses in *Arabidopsis* exposed to pristine CeO₂ nanoparticles, where the only *WRKY* gene expression change reported was up-regulation of 38.⁵⁷ WRKY38 tunes plant basal defense responses⁷³ and affects seed germination.⁶⁹ Perturbations of transcription factor levels can have cascading effects on down-stream gene regulon networks.

Protein Degradation—As a final illustration, 53 of the 93 Douglas-fir transcripts that enriched the *Protein Degradation* pathway code for RING/U-box zinc finger E3 ubiquitin ligases (Figure 5; Supporting Information Table S6). Members of this large family of ligases recognize specific target substrates for ubiquitination and mediating transfer of ubiquitin. These modifications can lead to proteolysis, modulation of protein function, structure, assembly, and/or localization.⁷⁴ Previous results suggest that a RING E3 ubiquitin ligase is important to induction of cell death and regulation of defense response to microbial pathogens.⁷⁵ There is also evidence that many RING E3 ligases interact with either calcium binding proteins or proteins related to calcium mediated pathways.⁷⁶ Clearly, there was an abundance of molecular impacts from DECE exposure that could affect the physiology and development of Douglas-fir trees.

Further Comparisons and Research Directions

In this report, we present data on the molecular impacts to plants from sublethal exposure to diesel emissions generated with CeO₂ nanoparticles as a fuel borne catalyst. Our results indicate that divergent DECe treatment effects were associated with significantly elevated needle uptake of cerium, rather than chamber particulate matter concentration (DECe $\approx$ DE) or gas composition (DECe $\approx$ DECe Gas Phase). In a recent report, *Chlamydomonas reinhardtii* internalized PVP-coated nCe, and when this freshwater algal species was exposed to supra-environmental concentration (10,000 μ g/L), 34 GOs were enriched with down-regulated transcripts. Most of these gene ontologies were related to chloroplast, photosynthesis and stress responses.⁷⁷ As described above, DECe also repressed expression of some transcripts with similar roles in Douglas-fir.

There may be other components of the DECe particulate matter that significantly contribute to transcriptome changes in Douglas-fir, though this remains to be determined. Interestingly, human alveolar epithelial cells (A549) exposed to DECe particulate matter also had altered expression of genes involved in transcription, cell proliferation/cell cycle/cell aging, metabolism, response to stimulus, cell adhesion, cell signal transduction, response to organic substances/drugs and transport.¹⁴ As in our DECe treatment, human genes related to control of transcription were also down-regulated. In that study, the authors pointed to several constituents of the particulate matter as potentially affecting gene expression, including; polycyclic aromatic hydrocarbons, fractions of organic carbon and physical properties of metallic nanoparticles. Additional research will be needed to identify any other drivers of the observed gene expression changes in Douglas-fir. It is also important to test whether these changes are reflective of conserved MIEs in response to DECe exposure that are evident in other species, potentially leading to predictable downstream effects at larger scales.²⁶

One possible type of initiating event that warrants additional research involves interference with calcium related processes, which were affected by exposure to DECe. As discussed above, several of the down-regulated transcripts that enriched DECe GOs and pathways code for proteins that involve Ca²⁺ binding, mediated signaling and/or transport. Although calcium related processes are ubiquitous in plant biochemical reactions, there is a possible mechanism by which Ce from DECe exposure reduced expression of these genes. The CeO₂ nanoparticles within EnviroxTM have very low solubility in aqueous solutions, but, combustion can increase this, especially when solutions contain organic matter.⁷⁸ Under physiological conditions inside the Douglas-fir needles, some fraction of the cerium taken up from DECe exposure may have been exchangeable Ce(III). Certain lanthanide series metal cations (including Ce³⁺) can occupy Ca²⁺ binding sites on various proteins, sometimes activating but often inhibiting the functions of many enzymes and transporters. Ce³⁺ has a similar ionic radius yet higher charge to volume ratio compared to Ca²⁺.⁷⁹ Thus, Ce³⁺ may competitively displace and replace Ca ions in biological systems, which can lead to toxic effects.⁸⁰

The wide range of transcriptome responses of Douglas-fir needles to DECe exposure suggests that there may be several other important MIEs that could be explored for their roles in altering plant growth and development. As noted above, these responses may involve interactions between combustion derived hydrocarbon surface coatings on nCe and plant cell

walls and membranes. Other MIEs may occur due to the physiochemical reactivity of combustion derived CeO₂ superstructures that do not involve release of Ce³⁺ ions.

While this study was of relatively short duration, it highlighted changes in gene expression that may lead to downstream effects at larger scales involving plant physiology and development, especially with chronic exposure to DECe. The results also suggest that studies employing native particles may not completely capture the dynamic range of responses to ENMs when they are released from product use. Future studies on effects of DECe exposure should include a diverse assortment of plant species to determine whether these responses are conserved across taxonomic groups, and also to determine if other metal ENMs elicit similar responses resulting from shared molecular initiating events. In addition, phenotypic changes that are linked to specific gene expression shifts need to be identified.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We appreciate the ICP-MS analysis performed by Dr. Kasey Kovalcik at the US EPA, National Exposure Research Laboratory. We thank the Genomic Research Core at the US EPA, National Health and Environmental Effects Research Laboratory for collecting the RNA-Seq data. The psme_transcriptome_v203 fasta was kindly shared by Dr. Richard Cronn (USDA Forest Service) and Dr. Sanjuro Jogdeo (Oregon State University). Funding for all authors on this project was provided by the US Environmental Protection Agency, Office of Research and Development.

References

- (1). Boyes WK; Thornton BLM; Al-Abed SR; Andersen CP; Bouchard DC; Burgess RM; Hubal EAC; Ho KT; Hughes MF; Kitchin K; Reichman JR; Rogers KR; Ross JA; Rygielwicz PT; Scheckel KG; Thai S-F; Zepp RG; Zucker RM A comprehensive framework for evaluating the environmental health and safety implications of engineered nanomaterials. *Critical Reviews in Toxicology* 2017, 1–44, DOI: 10.1080/10408444.2017.1328400.
- (2). HEI Health Effects Institute Communication 9: Evaluation of Human Health Risk from Cerium Added to Diesel Fuel; Health Effects Institute: Boston, MA, USA, 2001.
- (3). Skillas G; Qian Z; Baltensperger U; Matter U; Burtscher H The influence of additives on the size distribution and composition of particles produced by diesel engines. *Combustion Science and Technology* 2000, 154 (1), 259–273.
- (4). Jung H; Kittelson DB; Zachariah MR The influence of a cerium additive on ultrafine diesel particle emissions and kinetics of oxidation. *Combustion and Flame* 2005, 142 (3), 276–288.
- (5). Zhang J; Nazarenko Y; Zhang L; Calderon L; Lee K-B; Garfunkel E; Schwander S; Tetley TD; Chung KF; Porter AE Impacts of a nanosized ceria additive on diesel engine emissions of particulate and gaseous pollutants. *Environmental Science & Technology* 2013, 47 (22), 13077–13085. [PubMed: 24144266]
- (6). Batley GE; Halliburton B; Kirby JK; Doolette CL; Navarro D; McLaughlin MJ; Veitch C Characterization and ecological risk assessment of nanoparticulate CeO₂ as a diesel fuel catalyst. *Environmental Toxicology and Chemistry* 2013, 32 (8), 1896–1905. [PubMed: 23595783]
- (7). Gantt B; Hoque S; Fahey KM; Willis RD; Delgado-Saborit JM; Harrison RM; Zhang KM; Jefferson DA; Kalberer M; Bunker KL; Conny JM; Bhawe PV; Weinstein JP; Pye HOT Factors Affecting the Ambient Physicochemical Properties of Cerium-Containing Particles Generated by Nanoparticle Diesel Fuel Additive Use. *Aerosol Science and Technology* 2015, 49 (6), 371–380, DOI: 10.1080/02786826.2015.1027809.

- (8). Dale JG; Cox SS; Vance ME; Marr LC; Hochella MF Transformation of Cerium Oxide Nanoparticles from a Diesel Fuel Additive during Combustion in a Diesel Engine. *Environmental Science & Technology* 2017, 51 (4), 1973–1980, DOI: 10.1021/acs.est.6b03173. [PubMed: 28112928]
- (9). Gantt B; Hoque S; Willis RD; Fahey KM; Delgado-Saborit JM; Harrison RM; Erdakos GB; Bhavé PV; Zhang KM; Kovalcik K; Pye HOT Near-Road Modeling and Measurement of Cerium-Containing Particles Generated by Nanoparticle Diesel Fuel Additive Use. *Environmental Science & Technology* 2014, 48 (18), 10607–10613, DOI: 10.1021/es502169p. [PubMed: 25144365]
- (10). Erdakos GB; Bhavé PV; Pouliot GA; Simon H; Mathur R Predicting the Effects of Nanoscale Cerium Additives in Diesel Fuel on Regional-Scale Air Quality. *Environmental Science & Technology* 2014, 48 (21), 12775–12782, DOI: 10.1021/es504050g. [PubMed: 25271762]
- (11). Johnson AC; Park B Predicting contamination by the fuel additive cerium oxide engineered nanoparticles within the United Kingdom and the associated risks. *Environmental Toxicology and Chemistry* 2012, 31 (11), 2582–2587, DOI: 10.1002/etc.1983. [PubMed: 22893546]
- (12). Cassee FR; van Balen EC; Singh C; Green D; Muijsers H; Weinstein J; Dreher K Exposure, Health and Ecological Effects Review of Engineered Nanoscale Cerium and Cerium Oxide Associated with its Use as a Fuel Additive. *Critical Reviews in Toxicology* 2011, 0 (0), 1–16, DOI: doi:10.3109/10408444.2010.529105.
- (13). Snow SJ; McGee J; Miller DB; Bass V; Schladweiler MC; Thomas RF; Krantz T; King C; Ledbetter AD; Richards J; Weinstein JP; Conner T; Willis R; Linak WP; Nash D; Wood CE; Elmore SA; Morrison JP; Johnson CL; Gilmour MI; Kodavanti UP Inhaled Diesel Emissions Generated with Cerium Oxide Nanoparticle Fuel Additive Induce Adverse Pulmonary and Systemic Effects. *Toxicological Sciences* 2014, 142 (2), 403–417, DOI: 10.1093/toxsci/kfu187. [PubMed: 25239632]
- (14). Zhang Z-H; Balasubramanian R Effects of Cerium Oxide and Ferrocene Nanoparticles Addition As Fuel-Borne Catalysts on Diesel Engine Particulate Emissions: Environmental and Health Implications. *Environmental Science & Technology* 2017, 51 (8), 4248–4258, DOI: 10.1021/acs.est.7b00920. [PubMed: 28346827]
- (15). Howe GT; Yu J; Knaus B; Cronn R; Kolpak S; Dolan P; Lorenz WW; Dean JF A SNP resource for Douglas-fir: de novo transcriptome assembly and SNP detection and validation. *BMC Genomics* 2013, 14 (1), 137, DOI: 10.1186/1471-2164-14-137. [PubMed: 23445355]
- (16). Hermann RK; Lavender DP Douglas-fir planted forests. *New Forests* 1999, 17 (1), 53–70, DOI: 10.1023/a:1006581028080.
- (17). Gordon CJ; Schladweiler MC; Krantz T; King C; Kodavanti UP Cardiovascular and thermoregulatory responses of unrestrained rats exposed to filtered or unfiltered diesel exhaust. *Inhalation Toxicology* 2012, 24 (5), 296–309. [PubMed: 22486347]
- (18). R Core Team R: A language and environment for statistical computing, R Foundation for Statistical Computing: Vienna, 2013.
- (19). Cronn R; Dolan PC; Jogdeo S; Wegrzyn JL; Neale DB; St. Clair JB; Denver DR Transcription through the eye of a needle: daily and annual cyclic gene expression variation in Douglas-fir needles. *BMC Genomics* 2017, 18 (1), 558, DOI: 10.1186/s12864-017-3916-y. [PubMed: 28738815]
- (20). Uchimiya M; Cantrell KB; Hunt PG; Novak JM; Chang S Retention of heavy metals in a Typic Kandiodult amended with different manure-based biochars. *Journal of Environmental Quality* 2012, 41 (4), 1138–1149. [PubMed: 22751056]
- (21). EPA SW-846 Test Method 3010A: Acid Digestion of Aqueous Samples and Extracts for Total Metals for Analysis by Flame Atomic Absorption Spectroscopy (FLAA) or Inductively Coupled Plasma Spectroscopy (ICP). <https://www.epa.gov/hw-sw846/sw-846-test-method-3010a-acid-digestion-aqueous-samples-and-extracts-total-metals-analysis> (accessed December 4, 2017).
- (22). Dendrome P.menziesii (Douglas Fir) Transcriptome v2.0: 19.57 kbp from contig1:1..19,567. http://dendrome.ucdavis.edu/fgb2/psme_transcriptome2/.
- (23). Camacho C; Madden T; Coulouris G; Avagyan V; Ma N; Tao T; Agarwala R BLAST command line applications user manual BLAST® Help. Bethesda, MD: National Center for Biotechnology Information (US) 2008.

- (24). Usadel B; Poree F; Nagel A; Lohse M; Czedik-Eysenberg A; Stitt M A guide to using MapMan to visualize and compare Omics data in plants: a case study in the crop species, Maize. *Plant, Cell & Environment* 2009, 32 (9), 1211–1229, DOI: 10.1111/j.1365-3040.2009.01978.x.
- (25). Lohse M; Nagel A; Herter T; May P; Schroda M; Zrenner R; Tohge T; Fernie AR; Stitt M; Usadel B Mercator: a fast and simple web server for genome scale functional annotation of plant sequence data. *Plant, Cell & Environment* 2014, 37 (5), 1250–1258, DOI: 10.1111/pce.12231.
- (26). Ankley GT; Bennett RS; Erickson RJ; Hoff DJ; Hornung MW; Johnson RD; Mount DR; Nichols JW; Russom CL; Schmieder PK; Serrano JA; Tietge JE; Villeneuve DL Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk assessment. *Environmental Toxicology and Chemistry* 2010, 29 (3), 730–741, DOI: 10.1002/etc.34. [PubMed: 20821501]
- (27). Cheng S-H; Willmann MR; Chen H-C; Sheen J Calcium signaling through protein kinases. The Arabidopsis calcium-dependent protein kinase gene family. *Plant Physiology* 2002, 129 (2), 469–485. [PubMed: 12068094]
- (28). Dubiella U; Seybold H; Durian G; Komander E; Lassig R; Witte C-P; Schulze WX; Romeis T Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. *Proceedings of the National Academy of Sciences* 2013, 110 (21), 8744–8749, DOI: 10.1073/pnas.1221294110.
- (29). Matschi S; Hake K; Herde M; Hause B; Romeis T The calcium-dependent protein kinase CPK28 regulates development by inducing growth phase-specific, spatially restricted alterations in jasmonic acid levels independent of defense responses in Arabidopsis. *The Plant Cell* 2015, 27 (3), 591–606. [PubMed: 25736059]
- (30). Romeis T; Ludwig AA; Martin R; Jones JD Calcium-dependent protein kinases play an essential role in a plant defence response. *The EMBO Journal* 2001, 20 (20), 5556–5567. [PubMed: 11597999]
- (31). Mori IC; Murata Y; Yang Y; Munemasa S; Wang Y-F; Andreoli S; Tiriack H; Alonso JM; Harper JF; Ecker JR CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca²⁺-permeable channels and stomatal closure. *PLoS Biology* 2006, 4 (10), e327. [PubMed: 17032064]
- (32). Boudsocq M; Willmann MR; McCormack M; Lee H; Shan L; He P; Bush J; Cheng S-H; Sheen J Differential innate immune signalling via Ca²⁺ sensor protein kinases. *Nature* 2010, 464 (7287), 418–422, DOI: 10.1038/nature08794. [PubMed: 20164835]
- (33). Kolukisaoglu Ü; Weinl S; Blazevic D; Batistic O; Kudla J Calcium Sensors and Their Interacting Protein Kinases: Genomics of the Arabidopsis and Rice CBL-CIPK Signaling Networks. *Plant Physiology* 2004, 134 (1), 43–58, DOI: 10.1104/pp.103.033068. [PubMed: 14730064]
- (34). Luan S The CBL–CIPK network in plant calcium signaling. *Trends in Plant Science* 2009, 14 (1), 37–42, DOI: 10.1016/j.tplants.2008.10.005. [PubMed: 19054707]
- (35). Weinl S; Kudla J The CBL–CIPK Ca²⁺-decoding signaling network: function and perspectives. *New Phytologist* 2009, 184 (3), 517–528, DOI: 10.1111/j.1469-8137.2009.02938.x. [PubMed: 19860013]
- (36). Sanyal SK; Rao S; Mishra LK; Sharma M; Pandey GK Plant Stress Responses Mediated by CBL–CIPK Phosphorylation Network. *The Enzymes* 2016, 40, 31–64, DOI: 10.1016/bs.enz.2016.08.002. [PubMed: 27776782]
- (37). Bourdais G; Burdiak P; Gauthier A; Nitsch L; Salojärvi J; Rayapuram C; Idänheimo N; Hunter K; Kimura S; Merilo E; Vaattovaara A; Oracz K; Kaufholdt D; Pallon A; Anggoro DT; Glów D; Lowe J; Zhou J; Mohammadi O; Puukko T; Albert A; Lang H; Ernst D; Kollist H; Brosché M; Durner J; Borst JW; Collinge DB; Karpinski S; Lyngkjær MF; Robatzek S; Wrzaczek M; Kangasjärvi J; on behalf of the, C. R. K. C. Large-Scale Phenomics Identifies Primary and Fine-Tuning Roles for CRKs in Responses Related to Oxidative Stress. *PLOS Genetics* 2015, 11 (7), e1005373, DOI: 10.1371/journal.pgen.1005373. [PubMed: 26197346]
- (38). Sun X-L; Yu Q-Y; Tang L-L; Ji W; Bai X; Cai H; Liu X-F; Ding X-D; Zhu Y-M GsSRK, a G-type lectin S-receptor-like serine/threonine protein kinase, is a positive regulator of plant tolerance to salt stress. *Journal of Plant Physiology* 2013, 170 (5), 505–515, DOI: 10.1016/j.jplph.2012.11.017. [PubMed: 23276523]

- (39). Yang L; Ji W; Zhu Y; Gao P; Li Y; Cai H; Bai X; Guo D GsCBRLK, a calcium/calmodulin-binding receptor-like kinase, is a positive regulator of plant tolerance to salt and ABA stress. *Journal of Experimental Botany* 2010, 61 (9), 2519–2533. [PubMed: 20400529]
- (40). Yang T; Chaudhuri S; Yang L; Chen Y; Poovaiah B Calcium/calmodulin up-regulates a cytoplasmic receptor-like kinase in plants. *Journal of Biological Chemistry* 2004, 279 (41), 42552–42559. [PubMed: 15292241]
- (41). Belkhadir Y; Yang L; Hetzel J; Dangel JL; Chory J The growth–defense pivot: crisis management in plants mediated by LRR-RK surface receptors. *Trends in Biochemical Sciences* 2014, 39 (10), 447–456. [PubMed: 25089011]
- (42). Robatzek S; Wirthmueller L Mapping FLS2 function to structure: LRRs, kinase and its working bits. *Protoplasma* 2013, 250 (3), 671–681, DOI: 10.1007/s00709-012-0459-6. [PubMed: 23053766]
- (43). Bouwmeester K; Govers F Arabidopsis L-type lectin receptor kinases: phylogeny, classification, and expression profiles. *Journal of Experimental Botany* 2009, 60 (15), 4383–4396, DOI: 10.1093/jxb/erp277. [PubMed: 19773388]
- (44). Bonilla M; Cunningham KW Mitogen-activated Protein Kinase Stimulation of Ca²⁺ Signaling Is Required for Survival of Endoplasmic Reticulum Stress in Yeast. *Molecular Biology of the Cell* 2003, 14 (10), 4296–4305, DOI: 10.1091/mbc.E03-02-0113. [PubMed: 14517337]
- (45). Kang J; Park J; Choi H; Burla B; Kretschmar T; Lee Y; Martinoia E Plant ABC Transporters. *The Arabidopsis Book* 2011, e0153, DOI: 10.1199/tab.0153. [PubMed: 22303277]
- (46). Rea PA Plant ATP-binding cassette transporters. *Annu. Rev. Plant Biol* 2007, 58, 347–375. [PubMed: 17263663]
- (47). Garroum I; Bidzinski P; Daraspe J; Mucciolo A; Humbel BM; Morel J-B; Nawrath C Cuticular Defects in *Oryza sativa* ATP-binding Cassette Transporter G31 Mutant Plants Cause Dwarfism, Elevated Defense Responses and Pathogen Resistance. *Plant and Cell Physiology* 2016, 57 (6), 1179–1188, DOI: 10.1093/pcp/pcw066. [PubMed: 27121976]
- (48). Zhao G; Shi J; Liang W; Zhang D ATP binding cassette G transporters and plant male reproduction. *Plant Signaling & Behavior* 2016, 11 (3), e1136764, DOI: 10.1080/15592324.2015.1136764. [PubMed: 26906115]
- (49). Kaldenhoff R; Fischer M Aquaporins in plants. *Acta Physiologica* 2006, 187 (1–2), 169–176, DOI: 10.1111/j.1748-1716.2006.01563.x. [PubMed: 16734753]
- (50). Li G; Santoni V; Maurel C Plant aquaporins: Roles in plant physiology. *Biochimica et Biophysica Acta (BBA) - General Subjects* 2014, 1840 (5), 1574–1582, DOI: 10.1016/j.bbagen.2013.11.004. [PubMed: 24246957]
- (51). De Stefani D; Raffaello A; Teardo E; Szabo I; Rizzuto R A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature* 2011, 476 (7360), 336–340, DOI: <http://www.nature.com/nature/journal/v476/n7360/abs/nature10230.html#supplementary-information>. [PubMed: 21685888]
- (52). Stefani DD; Rizzuto R; Pozzan T Enjoy the Trip: Calcium in Mitochondria Back and Forth. *Annual Review of Biochemistry* 2016, 85 (1), 161–192, DOI: 10.1146/annurev-biochem-060614-034216.
- (53). Wagner S; De Bortoli S; Schwarzländer M; Szabò I Regulation of mitochondrial calcium in plants versus animals. *Journal of Experimental Botany* 2016, 67 (13), 3809–3829, DOI: 10.1093/jxb/erw100. [PubMed: 27001920]
- (54). Strehler EE Plasma membrane calcium ATPases: From generic Ca²⁺ sump pumps to versatile systems for fine-tuning cellular Ca²⁺. *Biochemical and Biophysical Research Communications* 2015, 460 (1), 26–33, DOI: 10.1016/j.bbrc.2015.01.121. [PubMed: 25998731]
- (55). Mizoi J; Shinozaki K; Yamaguchi-Shinozaki K AP2/ERF family transcription factors in plant abiotic stress responses. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* 2012, 1819 (2), 86–96, DOI: 10.1016/j.bbagrm.2011.08.004. [PubMed: 21867785]
- (56). Sakuma Y; Liu Q; Dubouzet JG; Abe H; Shinozaki K; Yamaguchi-Shinozaki K DNA-Binding Specificity of the ERF/AP2 Domain of Arabidopsis DREBs, Transcription Factors Involved in Dehydration- and Cold-Inducible Gene Expression. *Biochemical and Biophysical Research Communications* 2002, 290 (3), 998–1009, DOI: 10.1006/bbrc.2001.6299. [PubMed: 11798174]

- (57). Tumburu L; Andersen CP; Rygiewicz PT; Reichman JR Molecular and physiological responses to titanium dioxide and cerium oxide nanoparticles in *Arabidopsis*. *Environmental Toxicology and Chemistry* 2017, 36 (1), 71–82, DOI: 10.1002/etc.3500. [PubMed: 27212052]
- (58). Buck MJ; Atchley WR Phylogenetic Analysis of Plant Basic Helix-Loop-Helix Proteins. *Journal of Molecular Evolution* 2003, 56 (6), 742–750, DOI: 10.1007/s00239-002-2449-3. [PubMed: 12911037]
- (59). Le Hir R; Castelain M; Chakraborti D; Moritz T; Dinant S; Bellini C AtbHLH68 transcription factor contributes to the regulation of ABA homeostasis and drought stress tolerance in *Arabidopsis thaliana*. *Physiologia Plantarum* 2017, 160 (3), 312–327, DOI: 10.1111/pp.12549. [PubMed: 28369972]
- (60). Yanagisawa S The Dof family of plant transcription factors. *Trends in Plant Science* 2002, 7 (12), 555–560, DOI: 10.1016/S1360-1385(02)02362-2. [PubMed: 12475498]
- (61). Reyes JC; Muro-Pastor MI; Florencio FJ The GATA Family of Transcription Factors in *Arabidopsis* and Rice. *Plant Physiology* 2004, 134 (4), 1718–1732, DOI: 10.1104/pp.103.037788. [PubMed: 15084732]
- (62). Manfield IW; Devlin PF; Jen C-H; Westhead DR; Gilmartin PM Conservation, Convergence, and Divergence of Light-Responsive, Circadian-Regulated, and Tissue-Specific Expression Patterns during Evolution of the *Arabidopsis* GATA Gene Family. *Plant Physiology* 2007, 143 (2), 941–958, DOI: 10.1104/pp.106.090761. [PubMed: 17208962]
- (63). Ariel FD; Manavella PA; Dezar CA; Chan RL The true story of the HD-Zip family. *Trends in Plant Science* 2007, 12 (9), 419–426, DOI: 10.1016/j.tplants.2007.08.003. [PubMed: 17698401]
- (64). Dubos C; Stracke R; Grotewold E; Weisshaar B; Martin C; Lepiniec L MYB transcription factors in *Arabidopsis*. *Trends in Plant Science* 2010, 15 (10), 573–581, DOI: 10.1016/j.tplants.2010.06.005. [PubMed: 20674465]
- (65). Lurin C; Andrés C; Aubourg S; Bellaoui M; Bitton F; Bruyère C; Caboche M; Debast C; Gualberto J; Hoffmann B; Lecharny A; Le Ret M; Martin-Magniette M-L; Mireau H; Peeters N; Renou J-P; Szurek B; Taconnat L; Small I Genome-Wide Analysis of *Arabidopsis* Pentatricopeptide Repeat Proteins Reveals Their Essential Role in Organelle Biogenesis. *The Plant Cell* 2004, 16 (8), 2089–2103, DOI: 10.1105/tpc.104.022236. [PubMed: 15269332]
- (66). Delannoy E; Stanley WA; Bond CS; Small ID Pentatricopeptide repeat (PPR) proteins as sequence-specificity factors in post-transcriptional processes in organelles. *Biochemical Society Transactions* 2007, 35 (6), 1643–1647, DOI: 10.1042/bst0351643. [PubMed: 18031283]
- (67). Schmitz-Linneweber C; Small I Pentatricopeptide repeat proteins: a socket set for organelle gene expression. *Trends in Plant Science* 2008, 13 (12), 663–670, DOI: 10.1016/j.tplants.2008.10.001. [PubMed: 19004664]
- (68). Olsen AN; Ernst HA; Leggio LL; Skriver K NAC transcription factors: structurally distinct, functionally diverse. *Trends in Plant Science* 2005, 10 (2), 79–87, DOI: 10.1016/j.tplants.2004.12.010. [PubMed: 15708345]
- (69). Bakshi M; Oelmüller R WRKY transcription factors: Jack of many trades in plants. *Plant Signaling & Behavior* 2014, 9 (2), e27700–1–18, DOI: 10.4161/psb.27700. [PubMed: 24492469]
- (70). Chen L; Song Y; Li S; Zhang L; Zou C; Yu D The role of WRKY transcription factors in plant abiotic stresses. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* 2012, 1819 (2), 120–128, DOI: 10.1016/j.bbagr.2011.09.002. [PubMed: 21964328]
- (71). Phukan UJ; Jeena GS; Shukla RK WRKY Transcription Factors: Molecular Regulation and Stress Responses in Plants. *Frontiers in Plant Science* 2016, 7, 1–14, DOI: 10.3389/fpls.2016.00760. [PubMed: 26858731]
- (72). Rushton PJ; Somssich IE; Ringler P; Shen QJ WRKY transcription factors. *Trends in Plant Science* 2010, 15 (5), 247–258, DOI: 10.1016/j.tplants.2010.02.006. [PubMed: 20304701]
- (73). Kim K-C; Lai Z; Fan B; Chen Z *Arabidopsis* WRKY38 and WRKY62 Transcription Factors Interact with Histone Deacetylase 19 in Basal Defense. *The Plant Cell Online* 2008, 20 (9), 2357–2371, DOI: 10.1105/tpc.107.055566.
- (74). Deshaies RJ; Joazeiro CAP RING Domain E3 Ubiquitin Ligases. *Annual Review of Biochemistry* 2009, 78 (1), 399–434, DOI: 10.1146/annurev.biochem.78.101807.093809.

- (75). Lee DH; Choi HW; Hwang BK The Pepper E3 Ubiquitin Ligase RING1 Gene, *CaRING1*, Is Required for Cell Death and the Salicylic Acid-Dependent Defense Response. *Plant Physiology* 2011, 156 (4), 2011–2025, DOI: 10.1104/pp.111.177568. [PubMed: 21628629]
- (76). Mukherjee R; Das A; Chakrabarti S; Chakrabarti O Calcium dependent regulation of protein ubiquitination – Interplay between E3 ligases and calcium binding proteins. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 2017, 1864 (7), 1227–1235, DOI: 10.1016/j.bbamcr.2017.03.001. [PubMed: 28285986]
- (77). Taylor NS; Merrifield R; Williams TD; Chipman JK; Lead JR; Viant MR Molecular toxicity of cerium oxide nanoparticles to the freshwater alga *Chlamydomonas reinhardtii* is associated with supra-environmental exposure concentrations. *Nanotoxicology* 2016, 10 (1), 32–41, DOI: 10.3109/17435390.2014.1002868. [PubMed: 25740379]
- (78). Park B; Donaldson K; Duffin R; Tran L; Kelly F; Mudway I; Morin J-P; Guest R; Jenkinson P; Samaras Z; Giannouli M; Kouridis H; Martin P Hazard and Risk Assessment of a Nanoparticulate Cerium Oxide-Based Diesel Fuel Additive—A Case Study. *Inhalation Toxicology* 2008, 20 (6), 547–566, DOI: 10.1080/08958370801915309. [PubMed: 18444008]
- (79). Evans CH Interesting and useful biochemical properties of lanthanides. *Trends in Biochemical Sciences* 1983, 8 (12), 445–449.
- (80). Paoli L; Fiorini E; Munzi S; Sorbo S; Basile A; Loppi S Uptake and acute toxicity of cerium in the lichen *Xanthoria parietina*. *Ecotoxicology and Environmental Safety* 2014, 104, 379–385, DOI: 10.1016/j.ecoenv.2014.02.028. [PubMed: 24747927]
- (81). Edgar R; Domrachev M; Lash AE Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Research* 2002, 30 (1), 207–210, DOI: 10.1093/nar/30.1.207. [PubMed: 11752295]
- (82). Barrett T; Wilhite SE; Ledoux P; Evangelista C; Kim IF; Tomashevsky M; Marshall KA; Phillippy KH; Sherman PM; Holko M; Yefanov A; Lee H; Zhang N; Robertson CL; Serova N; Davis S; Soboleva A NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Research* 2013, 41 (D1), D991–D995, DOI: 10.1093/nar/gks1193. [PubMed: 23193258]

Figure 1. ICP-MS quantification of total Ce for Douglas-fir needles A) after experimental treatments and B) still bound in needle tissue after simulated rainwater wash. Significant differences among means are designated by lowercase letters and are based on adjusted P -values ≤ 0.05 .

Figure 2. RNA-Seq scatter plots (A-C) and heatmap (D) of Log₂ expression levels of Douglas-fir transcriptomes under DE, DECe, and DECe Gas Phase treatments relative to Air Controls. In the scatter plots, each point is the mean expression for a gene across all biological replicates for a treatment vs. those for Air Controls. Warmer (orange) and cooler (blue) colors indicate up and down-regulation respectively. Selected genes shown as points in white have 2-fold change and 95% confidence in expression levels. Rows of horizontal bars in the hierarchical clustering heatmap correspond to the combined set of 39,910 genes meeting the selection criteria. The color scale applies to expression levels within the heatmap.

Figure 3.

RNA-Seq Venn diagram of genes sets from DE, DECe and DECe Gas Phase treatments with 2-fold changes and 95% confidence in expression levels compared to Air Controls. Numbers of transcripts are within parentheses. ↑ and ↓ follow counts of up and down-regulated transcripts within each set respectively. Circle areas are not to scale.

Figure 4. Blast2GO identified A) Molecular Function, B) Biological Process, and C) Cellular Component enriched gene ontologies (GOs) that are unique to Douglas-fir transcripts that were down-regulated by DECE.

Figure 5.
Enrichment of MapMan pathway bins that are unique to Douglas-fir transcripts down-regulated by DECe.

Table 1.

Chamber Operating Conditions during 2-Week Exposure of Douglas-fir Seedlings to DE, DECe, DECe Gas Phase and Air Control Lower case letters indicate significant/non-significant differences across rows using Tukey HSD (adjusted $p = 0.05$)

Chamber Parameter	N	DE	DECe	DECe Gas Phase	Air Control
Particle Concentration (ug/m ³)	7	1157.8 ± 187.0 a	1002.6 ± 7.6 a	32.2 ± 1.7 b	0.8 ± <0.1 b
Total Particle Number (#/day)	7	5,723,333.3 ± 176,548.9 a	1,848,333.3 ± 88,266.4 b	10,423.7 ± 6,306.7 c	1,226.0 ± 714.3 c
Particle Volume (nm ³)	7	86.6 ± 4.6 b	99.3 ± 3.4 b	227.4 ± 23.5 a	256.7 ± 29.5 a
O ₂ (%)	11	20.6 ± 0.1 b	20.0 ± 0.1 c	20.0 ± 0.1 c	20.9 ± 0.1 a
CO (ppm)	11	19.0 ± 0.9 b	38.6 ± 3.1 a	44.2 ± 3.0 a	0.6 ± 0.1 c
NO (ppm)	11	14.6 ± 1.1 b	31.0 ± 1.7 a	34.3 ± 2.0 a	0.8 ± 0.1 c
NO ₂ (ppm)	11	2.1 ± 0.1 b	4.8 ± 0.4 a	4.8 ± 0.4 a	1.2 ± 0.2 b
NO _x (ppm)	11	16.7 ± 1.0 b	35.8 ± 1.9 a	39.0 ± 2.3 a	2.0 ± 0.2 c
Relative Humidity (%)	11	41.6 ± 0.3 c	60.4 ± 2.3 a	54.1 ± 2.5 ab	49.2 ± 2.6 bc
Temperature (°C)	11	23.8 ± 0.1 b	23.4 ± 0.1 c	25.7 ± <0.1 a	22.2 ± <0.1 d

Each value represents the average ± standard error of the mean daily values for each exposure day.

Abbreviations: DE, diesel exhaust; DECe, diesel exhaust from fuel containing cerium oxide nanoparticles

Table 2.

Douglas-fir RNA-Seq Assembly Statistics

Treatment	N	Trimmed Reads Assembled	Seq Score < 100% 90%	Seq Score = 100%
DE	7	3,005,892 ± 1,232,163	1,961,141 ± 604,844	1,044,750 ± 639,241
DECe	7	6,377,217 ± 2,664,219	3,610,828 ± 1,294,510	2,766,389 ± 1,381,977
DECe Gas Phase	6	6,232,900 ± 1,732,194	3,421,737 ± 786,960	2,811,164 ± 961,163
Air Control	6	10,776,209 ± 473,252	5,679,812 ± 300,553	5,096,396 ± 395,883

Each value represents the mean ± standard error

Abbreviations: DE, diesel exhaust; DECe, diesel exhaust from fuel containing cerium oxide nanoparticles