Published in final edited form as:

Org Lett. 2018 September 21; 20(18): 5550–5553. doi:10.1021/acs.orglett.8b02133.

An Atypical Mode of [3+2]-Cycloaddition: Pseudo-1,3-Dipole Behavior in Reactions of Electron-Deficient Thioamides with Benzyne

Juntian Zhang[#], Annika C. S. Page[#], Vignesh Palani, Junhua Chen, and Thomas R. Hoye^{*}
Department of Chemistry, 207 Pleasant St. SE, University of Minnesota, Minneapolis, MN 55455

[#] These authors contributed equally to this work.

Abstract

Thioamides bearing electron-withdrawing groups on the thiocarbonyl carbon atom react with benzyne [generated by the hexadehydro-Diels–Alder (HDDA) cycloisomerization] in an unprecedented fashion. Namely, the dihydrobenzothiazole products are consistent with a pathway involving initial formation of a stabilized ammonium ylide by a rare type of [3+2]-cycloaddition reaction. The fate of this species depends upon the nature of the R group(s) attached to the ylide nitrogen atom. The demonstration of new modes of reactivity represents the major advance arising from this study.

Graphical Abstract

We recently reported reactions between benzyne such as **II** (thermally generated from **I**) and aromatic thioamides such as **III** to produce dihydrobenzothiazines such as **V** (Figure 1a, structures designated by Roman **I–V** refer to species that appeared in our previous report).¹ We rationalized this process as proceeding through the series of intermediates **IVa–IVc**. With this in mind, we have been surprised to now see an alternative type of product (and mode of reaction) to arise upon heating **1a** in the presence of electron deficient thioamides such as **3a**. This produced, instead, the dihydrobenzothiazole **5aa**.² We propose that this transformation proceeds by a rare type of net [3+2]-cycloaddition (yellow highlight) to produce the ammonium ylide^{3,4} **4aa**, followed by ethylene elimination. We are unaware of

^{*}Corresponding Author hoye@umn.edu.

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website. Experimental details for the preparation of new compounds; spectroscopic data for their characterization, including copies of ¹H and ¹³C NMR spectra; and computed (DFT) geometries and energies of benzyne intermediates and transition states (PDF).

The authors declare no competing financial interests.

any similar reactions between a thioamide and a 2π (alkene or alkyne) component, although related transformations of trithiocarbonate and dithiocarbamate⁵ functional groups have been reported.⁶

Although the reaction between **2a** and **3a** may evoke the notion of a 1,3-dipolar cycloaddition,⁷ it differs in important ways. More specifically, each of the non-sacrificial resonance contributors of classical 1,3-dipoles has the same overall bond order, regardless of whether it is of the allyl or propargyl type (Figure 1b). Each is a zwitterionic species with a formal positive charge on the central atom and negative charge on one of its terminal atoms.⁸ The [3+2]-cycloaddition products that arise from engagement of a 1,3-dipole with the π -bond of a dipolarophile are, formally, uncharged at all atoms in the principal resonance contributor of the newly created five-membered ring. The nature of **3a** as well as its conversion to the ylide **4aa** are just the opposite—that is, **3a** does not meet the definition of a 1,3-dipole, and now, the resulting adduct **4aa** is zwitterionic.

We turned to DFT calculations to help further assess this novel type of [3+2]-cycloaddition (Figure 2). The computed reaction profile (with minima represented by **6–8** and transition structures by **TS1** and **TS2**) for the model addition of *N,N*-diethyl trifluorothioacetamide (**3a**) to *o*-benzyne (**6**) is shown in Figure 2. The computed exergonicity for the formation of the ammonium ylide **7** as well as for its further conversion to **8** and ethylene (ca. 20 and 50 kcal•mol⁻¹, respectively) are notable. We identified a transition structure (**TS1**) corresponding to a low barrier process for a concerted cycloaddition of **3a** to **6**. The geometry of that TS suggests that the addition proceeds with the nucleophilic portion of the thioamide leading the way—that is, the carbon-sulfur bond formation is more advanced than that of the carbon-nitrogen (cf. indicated bond lengths in **TS1** vis-à-vis those in **7**). The 1,2-zwitterion character in **7** sets up a facile intramolecular elimination reaction via **TS2** to produce **8**, the simplified analog of **5aa**, and ethylene. The low activation barrier for this step (ca. 15 kcal•mol⁻¹) is also notable.⁹

To gain direct experimental evidence for the ejection of an alkene from an ylide such as **4aa** (or **7**), we studied the reaction of triyne **1a** and *N,N*-dioctyltrifluorothioacetamide (**3b**, Figure 3). A 2.2:1 ratio of the isomeric *N*-octylbenzothiazolines **5ab** and **5ab'** was formed.¹⁰ In addition, a very nearly equimolar ratio of 1-octene was observed in the crude reaction mixture (¹H NMR and GC-MS). This result is consistent with the eliminative fragmentation shown in species **4ab**.

Another example of this new type of cycloaddition (Figure 4) was revealed when the *N*-allyl trifluorothioacetamide **3c** (3 equiv) was heated (85 °C) with triyne **1a** (Figure 1a). The dihydrobenzothiazole **5ac**^{10b} results from a [3,2]-sigmatropic migration of the allyl group, which again implicates the intermediacy of the ammonium ylide **4ac**, formed through capture of **2a** by the polarized thioamide **3c**.

We then explored a series of reactions (Table 1) to probe additional aspects of the reaction. First, would other electron withdrawing groups in the thioamide also promote this new type of cycloaddition reaction? The reactions using thioamides **3d–f**, containing –CF₃, –CN, and –CO₂Me groups, respectively, all gave rise to products (**5ad**,^{10b} **5ae**,^{10b} and **5af**,^{10b}

respectively) that demonstrate just that. Second, is the reaction unique to benzyne **2a**? No, as evidenced by the conversions of **1b** and **1c** to products **5bb**^{10b} and **5cc**,^{10b} respectively, via eliminative or [3,2]-sigmatropic processes within the ammonium ylides **4bb** or **4cc**, respectively.

Trapping with the pyrrolidine-containing thioamide **3g** revealed an additional new mode of reaction (Figure 5) beyond the products of Stevens rearrangement (**5ag**) and elimination (**5ag'**).¹¹ Here, we also isolated the novel, dihydrobenzo-1,4-thiazine derivative **5ag''**.^{10b,12} We suggest that this formally ring expanded compound arises from a C–H insertion event within the intermediate, isomeric carbene **9**, arising from dissociation of the nitrogen atom from the ammonium ylide carbon. This overall process reveals yet another facet of the reactivity of the ammonium ylides encountered in this study. The reverse reaction, generation of an ammonium ylide from a carbene, is known.¹³

A comment about the observed regioselectivities leading to the major constitutional isomer in these reactions is warranted. In most cases the major product is consistent with the well-established preference for benzyne **2a** to add nucleophilic species¹⁴ at the sp-hybridized carbon with the larger internal bond angle^{15,16} (see red triangle in **2a** in Fig. 1a). In contrast, the benzyne derived from **1b** and **1c** generally show a lower degree of regioselectivity for the addition of nucleophiles.¹⁴ The rr for the product **5cc** and its isomer is only 1.5:1. However, **5bb** was formed as the sole dihydrobenzothiazole. This substrate (and intermediate benzyne), uniquely among all studied here, contains a bulky aryl substituent adjacent to one of the benzyne carbons and thus sterically guides the nucleophilic sulfur atom to add to the distal benzyne carbon. This demonstrates that steric differences encountered during benzyne trapping reactions can override (at least small) inherent electronic preferences.¹⁷

In conclusion, this study of reactions of thioamides bearing an electron withdrawing substituent on the thiocarbonyl carbon (cf. **3**) with benzyne generated by the HDDA reaction of tri- and tetrayne precursors (cf. **2**) has led to the identification of a new type of [3+2]-cycloaddition reaction. The sulfur and nitrogen atoms of the thioamide engage the benzyne sp-carbon atoms to produce an intermediate ammonium ylide structure (cf. **4**). This species shows a variety of different paths for reaction, including several heretofore unseen, as deduced from the products that are produced. This adds to the overall body of understanding of these relatively rarely encountered species. Overall, this work represents another instance¹⁴ in which the HDDA cycloisomerization reaction has allowed a fundamentally new type of transformation to be discovered.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Support for this research was provided by the National Institutes of General Medical Sciences of the U.S. Department of Health and Human Services (R01 GM65597, then R35 GM127097). A.C.S.P. and V.P. appreciate the support of a Gleysteen-Heisig fellowship for undergraduate students. Some of the NMR data were obtained with an instrument acquired with funds from the NIH Shared Instrumentation Grant program (S10OD011952)

REFERENCES

1. Palani V; Chen J; Hoye TR Reactions of HDDA-derived benzyne with thioamides: Synthesis of dihydrobenzothiazino-heterocyclics. *Org. Lett.* 2016, 18, 6312–6315. [PubMed: 27978648]
2. The structures of products 5 (and intermediate ammonium ylides 4) presented in this manuscript carry two letter descriptors that reflect, first, the HDDA substrate (1a-c) and, second, the electron deficient thioamide (3a-f) from which they were generated. Also, see ref. 10 for a discussion of the fact that isomers arising from two regioisomeric modes of addition, indicated by “rr”, are typically observed and for the basis of assignment of the constitution of 5aa.
3. For reviews of ammonium ylide chemistry, see: Clark JS In *Nitrogen, Oxygen and Sulfur Ylide Chemistry*, Ed., Oxford University Press, Oxford, 2002; Wittig G *Achievements and problems in ylid chemistry* *J. Organomet. Chem.* 1975, 100, 279–287.
4. For examples of rearrangement reactions of N–C ylides, see: Stevens TS; Creighton EM; Gordon AB; MacNicol M *CCCCXXIII.—Degradation of quaternary ammonium salts. Part I.* *J. Chem. Soc.* 1928, 3193–3197. Sommelet M On a particular mode of intramolecular rearrangement. *Hebd. C. R. Seances Acad. Sci.* 1937, 205, 56. Vanecko JA; Wan H; West FG Recent advances in the Stevens rearrangement of ammonium ylides: application to the synthesis of alkaloid natural products. *Tetrahedron*, 2006, 62, 1043–1062. Clayden J; Donnard M; Lefranc J; Tetlow DJ Quaternary centres bearing nitrogen (α -tertiary amines) as products of molecular rearrangements. *Chem. Commun.* 2011, 47, 4624–4639.
- 5 (a). Nakayama J; Kimata A; Tanikuchi H; F. Takahashi Reactions of benzyne with 1,3-benzodithiole-2-thione and related compounds: formation of novel tetracyclic sulfonium salts and their reactions leading to dibenzo-1,3,6-trithiocin derivatives. *Bull. Chem. Soc. Jpn* 1996, 2349–2354; (b) Nakayama J; Kimata A; Tanikuchi H; Takahashi F Reactions of 1,3-benzodithiole-2-thione and ethylene trithiocarbonate with benzyne generated from 2-carboxybenzenediazonium chloride: preparation of novel bicyclic sulfonium salts by trapping 1,3-dipolar cycloaddition intermediates. *Chem. Comm.* 1996, 205–206.
6. Hwu JR; Hsu YC Stereospecific benzyne-induced olefination from β -amino alcohols and its application to the total synthesis of (–)-1-deoxy-D-fructose *Chem. Eur. J.* 2011, 4727–4731.
7. For recent studies of 1,3-dipolar cycloaddition reactions in which arynes were the primary dipolarophiles, see Ikawa T; Masuda S; Takagi A; Akai S 1,3- and 1,4-Benzdiyne equivalents for regioselective synthesis of polycyclic heterocycles. *Chem. Sci.* 2016, 7, 5206–5211; [PubMed: 30155171] Ikawa T; Kaneko H; Masuda S; Ishitsubo E; Tokiwa H; Akai S Trifluoromethanesulfonyloxy-group-directed regioselective (3 + 2) cycloadditions of benzyne for the synthesis of functionalized benzo-fused heterocycles. *Org. Biomol. Chem.* 2015, 13, 520–526; [PubMed: 25378144] Li P; Zhao J; Wu C; Larock RC; Shi F Synthesis of 3-substituted indazoles from arynes and N-tosylhydrazones. *Org. Lett.* 2011, 13, 3340–3343; [PubMed: 21630698] Spiteri C; Keeling S; Moses JE New synthesis of 1-substituted-1H-indazoles via 1,3-dipolar cycloaddition of in situ generated nitrile imines and benzyne. *Org. Lett.* 2010, 12, 3368–3371. [PubMed: 20608661] For a recent review that includes a subsection compiling 1,3-dipolar cycloaddition reactions, see Wu, C.; Shi, F. A closer look at aryne chemistry: Details that remain mysterious. *Asian J. Org. Chem* 2013, 2, 116–125.
8. Huisgen R 1,3-Dipolar Cycloadditions—Introduction, Survey, Mechanism In *1,3-Dipolar Cycloaddition Chemistry*; Padwa A Ed.; Wiley: New York, 1984; Vol. 1, pp 3–6.
9. We also computed the reaction energy for the transformation of the analog of 3a in which the CF₃ substituent was replaced with a CH₃ group to give the CH₃ analog of 7. That step is less exergonic (5 kcal•mol^{–1}, see Supporting Information) than that of 3a to 7. Whereas it was a trivial matter to locate TS1, we have been unable to identify a bound TS for the case of the methyl analog. The presence of the electron-withdrawing substituent in the model thioamide 3a appears to play an important role.
10. (a) In many of the reactions reported here, two constitutional isomers were formed, arising from competitive modes of addition of the thioamide to the various unsymmetrical benzyne. The assignment of constitution to the major and minor isomers 5ab and 5ab' rests on the nOe's shown in blue in Figure 3. (b) For many of the additional isomeric pairs (ratios indicated as rr), only the major isomer is shown in the manuscript [see Supporting Information (SI) for additional details].

The constitution of the major product is assigned on the basis of the very high field chemical shift of the aromatic H10 proton for the N-arylated products (5ac, 5ad, and 5cc), an nOe analogous to that for 5ab for the N-alkylated products (5aa, 5ae, 5af, 5ag'', and 5bb), and an abnormally high-field resonance for the N-CH₂ because of shielding by the nearby p-carbomethoxyphenyl group in 5bb.

11. We also observed a small amount (6% yield) of what we judged to be the analogous elimination product, the vinyl ether 5af', accompanying product 5af (Table 1).

12. This compound was obtained as a single diastereomer by the NCHCHS vicinal coupling constant (4.0 Hz) did not allow for a the confident assignment of its relative configuration.
13. Bamford WR; Stevens TS; Wright JW 829. Degradation of quaternary ammonium salts. Part IX. Attempts to prepare presumed intermediates having co-ordinately linked carbon. J. Chem. Soc. 1952, 4334–4338.
14. Diamond OJ; Marder TB Methodology and applications of the hexadehydro-Diels–Alder (HDDA) reaction. Org. Chem. Front. 2017, 4, 891–910.
15. Marell DJ; Furan LR; Woods BP; Lei X; Bendelsmith AJ; Cramer CJ; Hoyer TR; Kuwata KT Mechanism of the Intramolecular Hexadehydro-Diels-Alder Reaction. J. Org. Chem. 2015, 80, 11744–11754. [PubMed: 26270857]
16. Cheong PHY; Paton RS; Bronner SM; Im G-YJ; Garg NK; Houk KN Indolyne and Aryne Distortions and Nucleophilic Regioselectivities. J. Am. Chem. Soc. 2010, 132, 1267–1269 [PubMed: 20058924]
17. Medina JM; Mackey JL; Garg NK; Houk KN The Role of Aryne Distortions, Steric Effects, and Charges in Regioselectivities of Aryne Reactions. J. Am. Chem. Soc. 2014, 136, 15798–15805. [PubMed: 25303232]

a previous (III to V) vs. these (3a to 5aa) reactions of benzynes + thioamides

b classical 1,3-dipolar cycloadditions

Figure 1. Background.

(a) Previous and current work reporting divergent (left. vs. right, respectively) behavior in the trapping reactions of benzynes with thioamides. (b) Classical 1,3-dipolar cycloaddition reactions.

^a See ref. 2 for description of the two letter structure designations.

Figure 2. DFT [SMD(benzene)//Mo6-2X/6-311+G(d,p)] calculations for the [3+2]-cycloaddition and subsequent ethylene elimination for the hypothetical reaction between *o*-benzyne (**6**) and the *N,N*-diethyl thioamide **3a**.

Figure 3.

Use of the *N,N*-dioctylthioamide **3b** allowed identification of 1-octene, the elimination product from **4ab**, and led to the structural assignment of the major and minor products **5ab** and **5ab'**, respectively, by the indicated nOe 's.

Figure 4. Product **5ac** from [3,2]-sigmatropic rearrangement also support the intermediacy of ylide **4ac**.

Figure 5.

Pyrrolidino thioamide **3g** gave rise to three distinct types of products.

Table 1.

Products 5 arising from several different electron-deficient thioamides 3 and several different benzyne precursors 1 via the mechanistic events suggested in the intermediates ammonium ylides 4.

substrate 1	thioamide 3	product 5	[fate of N-ylide 4]
<p>1a</p>	<p>3d</p>	<p>5ad (88%, 11.6:1 rr)</p>	<p>4ad</p>
<p>1a +</p> <p>3e</p>	<p>5ae (98%, 8.5:1 rr)</p>	<p>4ae</p>	
<p>1a +</p> <p>3f</p>	<p>5af (70%, 3.1:1 rr)</p>	<p>4af</p>
<p>1b</p>	<p>3b</p>	<p>5bb (57%)</p>	<p>4bb</p>
<p>1c</p>	<p>3c</p>	<p>5cc (53%, 1.5:1 rr)</p>	<p>4cc</p>