Published in final edited form as:

Drug Alcohol Depend. 2019 January 01; 194: 288–295. doi:10.1016/j.drugalcdep.2018.10.008.

Influence of comorbid drug use disorder on receipt of evidence-based treatment for alcohol use disorder among VA patients with alcohol use disorder and Hepatitis C and/or HIV

Madeline C. Frost^a, Theresa E. Matson^{a,b}, Judith I. Tsui^c, and Emily C. Williams^{a,d}

^aHealth Services Research and Development (HSR&D) Center of Innovation for Veteran-Centered and Value-Driven Care, Veterans Affairs (VA) Puget Sound Health Care System, 1660 South Columbian Way, Seattle WA 98108; Tessa.E.Matson@kp.org; Madeline.Frost@va.gov; Emily.Williams3@va.gov

^bKaiser Permanente Washington Health Research Institute, 1730 Minor Ave, Seattle, WA 98101; Tessa.E.Matson@kp.org

^cDepartment of Medicine, University of Washington School of Medicine, 1959 NE Pacific St, Seattle, WA 98195; tsuij@uw.edu

^dDepartment of Health Services, University of Washington School of Public Health, 1959 NE Pacific St, Seattle, WA 98195; Emily.Williams3@va.gov

Abstract

Background: Alcohol use is risky for patients with hepatitis C virus (HCV) and/or human immunodeficiency virus (HIV) infection, but alcohol use disorder (AUD) treatment is underutilized in these populations. Comorbid drug use disorders (DUD) are common, but their influence on AUD treatment receipt is understudied. We evaluated the association between DUD and AUD treatment receipt in two national samples of patients with AUD, those with HIV and those with HCV, in the U.S. Veterans Health Administration.

Methods: Samples included patients with AUD and HCV and/or HIV among positive alcohol screens (AUDIT-C 5) documented 10/01/09-5/30/13 in the national electronic health record. Poisson regression models estimated incidence rate ratios for receiving specialty treatment (stop codes) and pharmacotherapy (filled prescription for naltrexone, disulfiram, acamprosate, or topiramate) within 365 days of positive alcohol screening for patients with DUD versus those without. Models were clustered on patient and adjusted for potential confounders.

Correspondence Madeline C. Frost, MPH, VA Puget Sound Health Care System, 1660 South Columbian Way, Seattle, WA 98108, (206) 764-2822, Madeline.Frost@va.gov.

Contributors

Ms. Frost led data analysis, interpretation, and manuscript writing. Dr. Williams led the creation of the secondary dataset and provided mentorship for data analysis, interpretation and revision of the manuscript. Ms. Matson and Dr. Tsui contributed to interpretation of findings and revision of the manuscript. All authors contributed to and have approved the final manuscript.

Conflict of Interest

The authors declare no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Results: Among 22,039 patients with HCV/AUD, 45.2% (N=9,964) had DUD, which was associated with receiving specialty treatment [adjusted incidence rate ratio: 1.89 (95% confidence interval 1.82-1.96)] and pharmacotherapy [aIRR: 1.50 (1.37-1.65)]. Among 1,834 patients with HIV/AUD, 56.9% (N=1,043) had DUD, which was associated with receiving specialty treatment [aIRR: 1.94 (1.68-2.24)], but not pharmacotherapy.

Conclusions: Rates of AUD treatment receipt among patients with AUD and HCV and/or HIV were low overall, but likelihood of treatment receipt was generally higher among those with comorbid DUD. Future research should investigate mechanisms underlying these associations, such as enhanced readiness for treatment or differential provider prescribing or referral practices.

Keywords

Alcohol; Alcohol Use Disorder; Treatment; Comorbidity; HCV; HIV

1. Introduction

Unhealthy alcohol use and alcohol use disorders are highly prevalent among people with hepatitis C virus (HCV) and among people with human immunodeficiency virus (HIV) (Armstrong et al., 2006; CDC, 2016; Galvan et al., 2002; Taylor et al., 2016; Tsui et al., 2016; Williams et al., 2016b), and alcohol use has unique harmful effects for persons with each of these infectious diseases. Alcohol and HCV have synergistic impacts on health. Among people with HCV, those who drink heavily are at increased risk for HCV-related liver diseases and mortality compared with those who do not (Donato et al., 2002; Fuster et al., 2014; Hutchinson et al., 2005; Safdar and Schiff, 2004; Szabo et al., 2006; Szabo et al., 2010; Younossi et al., 2013), and among people with alcohol use disorder, those with HCV experience more healthcare utilization and mortality compared to those without HCV (Fuster et al., 2015; Tsui et al., 2006). Unhealthy alcohol use may also result in decreased receipt of antiviral treatment for HCV (Lin et al., 2017) and accelerated HCV viral replication (McCartney and Beard, 2010). Among those with HIV, alcohol use negatively influences HIV care engagement and adherence and multiple HIV-related outcomes, including mortality risk (Edelman et al., 2018; Vagenas et al., 2015; Williams et al., 2016a; Williams et al., 2018). Further, research suggests that people with HIV may be more susceptible to alcohol-related harm compared to those without HIV using alcohol at similar levels (Justice et al., 2016; McGinnis et al., 2016). Thus, people with HCV and/or HIV are particularly important target populations for the reduction of unhealthy alcohol use.

Alcohol use disorder (AUD) is the most severe form of unhealthy alcohol use (Saitz, 2005). There are effective behavioral and pharmacological treatments for AUD (Anton et al., 2006; Dawson et al., 2006; Jonas et al., 2014; Lingford-Hughes et al., 2012; National Institute on Alcohol Abuse and Alcoholism, 2007; Pettinati et al., 2005; Weisner et al., 2003), and specialty addictions treatment and/or prescription of pharmacotherapy are recommended for treatment of AUD (Department of Veterans Affairs, 2015; Jonas et al., 2014; National Institute for Health and Care Excellence, 2015; NIAAA, 2007). Pharmacological options include three FDA-approved medications (acamprosate, disulfiram, and oral or injectable naltrexone) (Jonas et al., 2014; NIAAA, 2007), and a fourth, topiramate, was found to have strong evidence for treatment of AUD in a meta-analysis (Jonas et al., 2014). Despite strong

evidence for these treatments, they are underused among patients with AUD (Cohen et al., 2007; Harris et al., 2012; Rubinsky et al., 2015), and multiple barriers to treatment receipt and implementation have been documented (Alanis-Hirsch et al., 2016; Finlay et al., 2017; Hagedorn et al., 2016; Harris et al., 2013; Knudsen and Roman, 2014; Ober et al., 2017; Oliva et al., 2011; Williams et al., 2017a). People with infectious diseases (HCV and HIV) may be even less likely to receive evidence-based treatment for AUD compared to people without these conditions (Owens et al., 2018; Williams et al., 2017d).

Drug use is also prevalent among people with HCV and people with HIV (Alter, 2007; Armstrong et al., 2006; Bing et al., 2001; CDC, 2016), and comorbid drug use disorder (DUD) is common among people with AUD (Grant et al., 2015; Grant and Pickering, 1996; Grant et al., 2016; Stinson et al., 2005). Previous analyses of nationally-representative samples of U.S. residents have found that co-occurring DUD increases the likelihood of alcohol treatment utilization among people with AUD in the general population (Cohen et al., 2007; Dawson et al., 2012; Grant and Pickering, 1996; Ilgen et al., 2011; Stinson et al., 2005). In contrast, one previous study among patients from the national Veterans Health Administration (VA) found that DUD was not a significant predictor of receipt of specialty treatment for any substance use disorder (drugs or alcohol) among people with elevated levels of alcohol consumption (Glass et al., 2010). Specific to populations with infectious diseases, one analysis of data from the Women's Interagency HIV Study found that among women living with HIV who reported hazardous drinking, illicit drug use was associated with increased alcohol treatment utilization (Hu et al., 2016). Relatedly, another study among people with HIV found that while alcohol dependence was negatively associated with substance use treatment, drug dependence was positively associated with treatment (Orwat et al., 2011). To our knowledge, no studies have examined the association between comorbid DUD and receipt of AUD treatment among people with HCV specifically.

Further examination of the impact of comorbid DUD on receipt of treatment for AUD among people with HCV and HIV is warranted, particularly in a real-world care setting. The Veterans Health Administration (VA) is the largest provider of HCV and HIV care in the U.S. (Backus et al., 2015; Department of Veterans Affairs, 2017). Understanding the impact of comorbid DUD diagnosis on AUD treatment receipt among patient populations with HCV and HIV who have AUD and receive care in VA may have important implications for provider practice, as these populations have a high prevalence of comorbid DUD and are particularly susceptible to the harms of alcohol use. Thus, the present study aimed to evaluate the association between comorbid DUD and receipt of evidence-based AUD treatments among patients with HCV and patients with HIV. Results may inform future efforts to increase the provision of evidence-based care for AUD in these patient populations, both within and outside the VA healthcare system.

2. Methods

2.1 Setting, data source, and samples

The present study was conducted in the national VA health care system, comprised over 1,200 health care facilities serving over 9 million patients across the U.S. (Department of Veterans Affairs, 2018), and reflects a secondary analysis of data collected for a large study

that assessed receipt of alcohol-related care across HIV status among VA patients with unhealthy alcohol use as determined by annual alcohol screening (Williams et al., 2017d). For the parent study, secondary data were extracted from the VA Corporate Data Warehouse (CDW), which is housed within VA's Informatics and Computing Infrastructure (VINCI), a national repository of clinical, pharmacy, administrative, enrollment, financial and utilization data for VA patients. Data were extracted for all patients with an outpatient appointment between 10/1/09 and 5/30/13 who had at least one documented positive alcohol screen during the study period. A positive alcohol screen was defined as an Alcohol Use Disorders Identification Test Consumption (AUDIT-C) score ≥ 5 , consistent with the national performance measure for brief intervention used across the VA system (Lapham et al., 2012; Williams et al., 2017c). To examine outcomes, each screen was followed up to one year until 5/30/14. Multiple positive screens from the same patient could be included in the dataset during the study period if screens were at least nine months apart, consistent with VA's requirements for annual alcohol screening and the past-year timeframe of the screening instrument. Study procedures, including a waiver of HIPAA authorization and a waiver of written consent, were approved by the Institutional Review Board at VA Puget Sound Health Care System.

Two samples of patients were included in the present secondary analysis: 1) patients with HCV and AUD, and 2) patients with HIV and AUD. These samples were identified based on diagnostic codes from the International Classification of Disease, Ninth Revision, Clinical Modification (ICD-9-CM) in the 0-365 days prior to a positive screen and are not mutually exclusive (i.e., persons with either HIV or HCV may also have been diagnosed with the other condition). AUD was defined as having a documented diagnosis for alcohol abuse or dependence, excluding in remission (309.9-303.92, 305.0-305.02). HCV was defined as having one of the following diagnoses: chronic hepatitis C with (070.44) or without (070.54) hepatic coma, acute hepatitis C with (070.41) or without (070.51) hepatic coma, unspecified viral hepatitis C with (070.71) or without (070.70) hepatic coma, or hepatitis C carrier (02.62). HIV was defined as having two outpatient or one inpatient diagnostic code(s) for AIDS (042) and/or HIV infection (V08), consistent with prior research (Fultz et al., 2006).

2.2 Measures

2.2.1 Primary independent variable.—Comorbid DUD diagnosis was based on ICD-9-CM diagnostic codes in the 0-365 days prior to a positive AUDIT-C screen. Patients with abuse or dependence diagnoses for stimulants (including amphetamines and cocaine; 304.2-304.22, 305.6-305.62, 304.4-304.42, 305.7-305.72), opioids (304.0-304.32, 304.7-304.72, 305.5-305.52), cannabis (304.3-304.32, 305.2-305.22), hallucinogens (304.5-304.52, 305.3-305.32), or sedatives (304.1-304.12, 305.4-305.42) were classified as having comorbid DUD (excluding diagnoses for patients in remission).

2.2.2 Outcomes.—The two outcomes of interest were receipt of any specialty AUD treatment and receipt of any AUD pharmacotherapy within 365 days following a positive AUDIT-C screen. Receipt of specialty AUD treatment was measured based on visit codes for inpatient or outpatient addiction treatment with an accompanying AUD diagnosis at the time of the visit code (Williams et al., 2014a). This definition includes outpatient codes for

individual, home visit and group substance use disorder treatment, and inpatient codes for residential rehabilitation, compensated work therapy, high intensity treatment, psychiatric and domiciliary care (a complete list of VA stop codes and bed sections included in this definition can be found in Williams et al., 2017d). Receipt of AUD pharmacotherapy was measured as at least one filled prescription for acamprosate, disulfiram, topiramate, or oral or injectable naltrexone, based on the date the medication was dispensed for inpatients and the date the medication was filled for outpatients (Harris et al., 2010).

2.2.3 Covariates.—Covariates included demographic characteristics and mental health conditions, based on their associations with both drug use disorders and receipt of alcohol-related care (Bensley et al., 2017; Chen et al., 2018; Cohen et al., 2007; Dawson et al., 2012; Grant et al., 2016; Ilgen et al., 2011; Stinson et al., 2005; Williams et al., 2017b; Williams et al., 2012; Williams et al., 2017c). Demographic characteristics included sex (male/female), age group (18-29, 30-44, 45-64, 65+) categorized consistent with previous studies (Williams et al., 2017d; Williams et al., 2014b), race/ethnicity (Asian American/ Pacific Islander, Black, Hispanic/Latino, Native American, White), marital status (divorced/separated, married, never married/single, widowed) and eligibility for VA co-payment status (full VA coverage, service connection <50%, no coverage) included as a proxy for socioeconomic status (Williams et al., 2012; Young et al., 2003). Mental health conditions were measured based on ICD-9-CM diagnostic codes in the 0-365 days prior to a positive AUDIT-C screen, and included major depression, anxiety disorder, post-traumatic stress disorder, other mood disorder, and serious mental illness (schizophrenia, bipolar disorder, psychosis). Additionally, fiscal year of positive AUDIT-C screen (2010, 2011, 2012, 2013) was included as a covariate to adjust for an increase in rates of alcohol-related care provided in VA nationally over time (Lapham et al., 2012).

2.3 Analyses

Patient characteristics collected at the time of first positive AUDIT-C screen were described at the patient level among patients in each sample: those with HCV and AUD and those with HIV and AUD. Chi-square tests of independence were used to assess differences across comorbid DUD status in each sample.

Consistent with previous studies (Owens et al., 2018; Williams et al., 2017d), modified Poisson regression models (Zou, 2004) were used to estimate relative rate and 95% confidence intervals of the two outcomes across comorbid DUD status, with positive AUDIT-C screens as the unit of analysis. Poisson regression was chosen to estimate relative rates (rather than odds ratios estimated through logistic regression) because outcomes were expected to be common (Greenland, 1995; Zou, 2004). Analyses were clustered at the patient level and standard errors were calculated using the robust sandwich estimator to account for correlation between AUDIT-C screens of the same patient, as well as for misspecification of the variance structure in the Poisson model (Greenland, 1995; Liang and Zeger, 1986; Zou, 2004). To examine the magnitude of effects, marginal predictions were calculated to describe predicted prevalence and 95% confidence intervals of the two outcomes in each group. Models were first fit unadjusted to identify the existing association and prevalence in the samples, then subsequently adjusted for factors that may confound

associations, including demographic characteristics, fiscal year of AUDIT-C screen and mental health diagnoses. Secondary analyses examined the association between specific DUDs (opioid, stimulant, and other) and all outcomes, and examined outcomes among patients diagnosed with both HIV and HCV. Analyses were conducted using Stata 15 software (StataCorp, 2017).

3. Results

3.1 Sample descriptions

Between 10/1/09 and 5/30/13, 830,825 patients (10.3% of patients with outpatient appointments eligible for AUDIT-C screening) screened positive for unhealthy alcohol use (AUDIT-C ≥ 5) with 1,172,606 total positive screens identified. Of these patients, 22,039 patients (2.7%; 30,765 screens) had diagnosed HCV and AUD, and 1,834 patients (0.2%; 2,401 screens) had diagnosed HIV and AUD. Among patients with HCV and AUD (N=22,039), 9,964 (45.2%) had a comorbid DUD, and among patients with HIV and AUD (N=1,834), 1,043 (56.9%) had a comorbid DUD. Characteristics of patients in both samples are presented in Tables 1 (HCV) and 2 (HIV) overall and across comorbid DUD status. For patients with HCV and AUD, most characteristics differed significantly across comorbid DUD status (p-values <0.05). Those with comorbid DUD were more likely to be female, younger, Black/African American, never married/single, to have full VA coverage or service connection $<50\%$, and to be diagnosed with major depression, other mood disorder, PTSD, anxiety disorder, or serious mental illness (Table 1). For patients with HIV and AUD, some characteristics differed significantly across comorbid DUD status (p-values <0.05). Specifically, those with comorbid DUD were more likely to be age 45-64 and Black/African American, and to be diagnosed with major depression, other mood disorder, PTSD, anxiety disorder, and serious mental illness (Table 2).

3.2 Association between comorbid DUD and receipt of AUD treatment

Among patients with HCV and AUD, 8,160 (37.0%) had documented AUD specialty addictions treatment and 1,693 (7.7%) had documented AUD pharmacotherapy within 365 days of positive AUDIT-C screen. Among patients with HIV and AUD, 741 (40.4%) had documented AUD specialty addictions treatment and 108 (5.9%) had documented AUD pharmacotherapy within 365 days of positive AUDIT-C screen.

Results of observation-level regression models for both samples are presented in Table 3. Among those with AUD and HCV, comorbid DUD was associated with increased likelihood of receiving both specialty addictions treatment and AUD pharmacotherapy in both unadjusted and adjusted analyses (Table 3). After adjustment, the predicted prevalence of receiving specialty addictions treatment was 52.8% (51.9-53.7) for those with DUD and 22.4% (21.8-23.1) for those without DUD ($p<0.001$), and the predicted prevalence of receiving AUD pharmacotherapy was 11.3% (10.8-11.9) and 5.6% (5.2-6.0) for those with and without DUD, respectively ($p<0.001$; Table 3). Although both associations were significant, the absolute difference in predicted prevalence for pharmacotherapy (difference of 5.7%) was smaller than for specialty treatment (difference of 30.4%).

Among patients with AUD and HIV, comorbid DUD was associated with increased likelihood of receiving AUD specialty treatment in both unadjusted and adjusted models, but was not significantly associated with receiving pharmacotherapy (Table 3). After adjustment, the predicted prevalence of receiving specialty addictions care was 52.7% (49.9-55.5) for those with DUD and 22.7% (20.0-25.4) for those without DUD ($p<0.001$), and the predicted prevalence of receiving AUD pharmacotherapy was 7.2% (5.7-8.8) and 5.1% (3.7-6.5) for those with and without DUD, respectively ($p=0.345$; Table 3).

Secondary analyses examining the association between specific DUDs (opioid, stimulant, and other) and all outcomes are presented in Supplementary Tables 1a-1c. Results were similar to the primary analyses among all subgroups, except that among patients with HIV, opioid use disorder was significantly associated with increased receipt of pharmacotherapy (Supplementary Table 1a). Results of secondary analyses among patients diagnosed with both HIV and HCV mirrored results among patients with only HIV (data not presented).

4. Discussion

This analysis of two large, non-mutually exclusive national samples of VA patients with AUD—those with HCV, and those with HIV—found that rates of AUD specialty addictions treatment and pharmacotherapy were low overall, particularly for pharmacotherapy (7.7% for HCV and 5.9% for HIV). However, having a comorbid DUD was associated with increased likelihood of receiving specialty addictions treatment for both groups and of receiving pharmacotherapy for those with HCV.

Findings from the present national study of VA patients with AUD and HCV and/or HIV are consistent with those of epidemiologic surveys of the general population (Cohen et al., 2007; Dawson et al., 2012; Grant and Pickering, 1996; Ilgen et al., 2011; Stinson et al., 2005), and studies of people with HIV (Hu et al., 2016; Orwat et al., 2011). Additionally, findings are similar to a previous VA study, which found that posttraumatic stress disorder was associated with increased receipt of AUD specialty addictions treatment and AUD pharmacotherapy (Chen et al., 2018), suggesting that other comorbidities may also increase receipt of treatment for AUD. However, one previous study of VA patients found that among patients with severe unhealthy alcohol use identified by alcohol screening, DUD was not a significant predictor of receiving specialty treatment for any substance use disorder, including drugs or alcohol (Glass et al., 2010). It is possible that differences between the previous study and the present study are related to differing samples and outcomes. In the present study, potentially eligible patients first screened positive for unhealthy alcohol use and then were included in analyses if they also had an AUD diagnosis, and the outcome was specialty treatment for AUD; in the Glass et al. study (2010), patients were included if they screened positive for unhealthy alcohol use, and the outcome was specialty treatment for any alcohol and/or drug use disorder. Differences may also relate to increased attention to provision of alcohol-related care in VA since the time of the previous study (Department of Veterans Affairs, 2015; Lapham et al., 2012). Finally, differences may relate to the specific focus in the present study on patients with infectious diseases in addition to AUD. Given the compounding influences of both AUD and DUD on both HCV and HIV, determinants of

AUD treatment may differ in this sample from those in the general population of VA patients with severe unhealthy alcohol use and comorbid DUD.

The reasons underlying the association between comorbid DUD and receipt of treatment for AUD in these samples are unknown, but may be due to both provider and patient-level factors. For instance, providers who offer substance use treatment may have increased contact with persons with comorbid DUD, thus offering increased opportunities for treatment referral or provision. Studies have found that other mental health comorbidities (Chen et al., 2018; Glass et al., 2010) and greater alcohol use severity (Arndt et al., 2002; Burman et al., 2004; Kaner et al., 2001; Richmond et al., 1996; Volk et al., 1996) are associated with increased likelihood of receiving substance use disorder treatment and brief alcohol counseling. It is possible that multiple diagnoses and/or greater severity of problems may increase perceptions of risk, which could increase patient readiness to seek treatment and/or provider advice to seek treatment. Providers may also be less likely to prescribe pharmacotherapy or refer patients to AUD treatment if they do not have comorbid DUD, if they view AUD as less harmful than DUD. Alternatively, or in addition, patients with AUD and HCV and/or HIV who have comorbid DUD may be more likely to take up AUD treatment once it is prescribed or referred given multiple compounding health risks. This national study was conducted using secondary EHR data and thus was unable to examine the respective roles of provider practices and patient uptake of treatment. However, future research is needed to assess the relative contributions of these two factors and how they are impacted by comorbid DUD diagnosis among patients with HCV and/or HIV, which may inform strategies to increase receipt of AUD treatment among these patients with and without comorbid DUD.

Overall, rates of AUD treatment were low across all groups, but especially among persons without comorbid DUD. These findings are concerning given the substantial health risks AUD poses to people with HIV and/or HCV, and given that sizeable proportions in each of these groups had diagnosed AUD only, without comorbid DUD (54.8% among HCV and 43.1% among HIV). Future work is needed to increase linkage to treatment, and especially to increase use of pharmacological treatments, among patients with infectious diseases who have diagnosed AUD but no comorbid DUD. It should be noted that some medications to treat AUD are metabolized through the liver (e.g., disulfiram, naltrexone), and therefore may raise concerns among providers who treat patients with HCV and other liver conditions. However, studies have demonstrated that these drugs are safe for persons with HCV and/or risk of hepatotoxicity with regular monitoring of liver function (Mitchell et al., 2012; Saxon et al., 1998; Tetrault et al., 2012).

There are several limitations to this study. Extracting EHR data enabled a large secondary analysis, but also introduced several limitations related to internal validity. Firstly, although the AUD specialty addictions treatment outcome was defined as having an AUD diagnosis accompanying the visit code, there may have been some misclassification such that this measure captured treatment for other substance use disorders. Similarly, the AUD pharmacotherapy outcome may both over- and under-estimate medication treatment for AUD. Specifically, our measure included medications that are prescribed for indications other than AUD (naltrexone and topiramate) but did not differentiate between indications,

and therefore may have resulted in overestimation of rates of AUD pharmacotherapy. On the other hand, we did not measure other medications that are increasingly being prescribed to treat AUD, such as gabapentin, and thus may have underestimated medication treatment. Unmeasured confounders that were not captured with EHR data may have resulted in residual confounding. Additionally, we are unable to interpret the association between DUD and receipt of AUD treatment as being evidence of causality in this analysis.

There are also limits to external validity. While VA is an ideal setting in which to conduct this study due to its ability to capture alcohol use and alcohol-related care in EHR data and large populations of patients receiving care for HCV and HIV, it may differ from other healthcare systems in several ways. As the VA is an integrated healthcare system that has implemented routine alcohol screening and in which addictions treatment is recommended in clinical guidelines (Department of Veterans Affairs, 2015), rates of treatment observed here may be higher than in other healthcare systems. Also, the VA patient population is largely male and older than other patient populations. Finally, because we identified patients with AUD among those with positive alcohol screening, results may not be generalizable to patients with AUD who do not receive alcohol screening and screen positive for unhealthy alcohol use.

Despite these limitations, this study demonstrated that among VA patients with AUD and HCV and those with AUD and HIV, rates of AUD specialty addictions treatment and pharmacotherapy were low overall, but significantly increased with presence of a comorbid DUD diagnosis (with the exception of pharmacotherapy among patients with HIV, which was non-significant). Findings suggest that having a comorbid DUD does not pose a barrier to receiving AUD treatment, but instead generally increases the likelihood of receiving treatment for AUD in these populations. Future research is needed to assess how comorbid DUD impacts provider practices and patient uptake of treatment, which may inform strategies to increase receipt of AUD treatment among the larger population of patients who do not have comorbid DUD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the National Institute on Alcohol Abuse and Alcoholism (R21 AA022866). Dr. Williams is supported by a Career Development Award from VA HSR&D (CDA 12-276).

Role of the Funding Source

The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

References

- Alanis-Hirsch K, Croff R, Ford JH, Johnson K, 2nd, Chalk M, Schmidt L, McCarty D, 2016 Extended-release naltrexone: A qualitative analysis of barriers to routine use. *J. Subst. Abuse Treat.* 62, 68–73. [PubMed: 26654934]

- Alter MJ, 2007 Epidemiology of hepatitis C virus infection. *World J. Gastroenterol.* 13, 2436–2441. [PubMed: 17552026]
- Anton RF, O'Malley SS, Ciraulo DA, Cisler RA, Couper D, Donovan DM, Gastfriend DR, Hosking JD, Johnson BA, LoCastro JS, Longabaugh R, Mason BJ, Mattson ME, Miller WR, Pettinati HM, Randall CL, Swift R, Weiss RD, Williams LD, Zweben A, 2006 Combined pharmacotherapies and behavioral interventions for alcohol dependence: The COMBINE study: A randomized controlled trial. *JAMA* 295, 2003–2017. [PubMed: 16670409]
- Armstrong GL, Wasley A, Simard EP, McQuillan GM, Kuhnert WL, Alter MJ, 2006 The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann. Intern. Med.* 144, 705–714. [PubMed: 16702586]
- Arndt S, Schultz SK, Turvey C, Petersen A, 2002 Screening for alcoholism in the primary care setting: Are we talking to the right people? *J. Fam. Pract.* 51, 41–46. [PubMed: 11927063]
- Backus L, Czarnogorski M, Yip G, Thomas BP, Torres M, Bell T, Ross D, 2015 HIV care continuum applied to the US Department of Veterans Affairs: HIV virologic outcomes in an integrated health care system. *J. Acquir. Immune Defic. Syndr.* 69, 474–480. [PubMed: 25835603]
- Bensley KM, Harris AH, Gupta S, Rubinsky AD, Jones-Webb R, Glass JE, Williams EC, 2017 Racial/ethnic differences in initiation of and engagement with addictions treatment among patients with alcohol use disorders in the veterans health administration. *J. Subst. Abuse Treat.* 73, 27–34. [PubMed: 28017181]
- Bing EG, Burnam MA, Longshore D, Fleishman JA, Sherbourne CD, London AS, Turner BJ, Eggan F, Beckman R, Vitiello B, Morton SC, Orlando M, Bozzette SA, Ortiz-Barron L, Shapiro M, 2001 Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Arch. Gen. Psychiatry* 58, 721–728. [PubMed: 11483137]
- Burman ML, Kivlahan DR, Buchbinder MB, Broglio K, Zhou XH, Merrill JO, McDonnell MB, Fihn SD, Bradley KA, 2004 Alcohol-related advice for VA primary care patients: Who gets it, who gives it? *J. Stud. Alcohol* 65, 621–630. [PubMed: 15536772]
- Centers for Disease Control and Prevention (CDC), 2016 Behavioral and clinical characteristics of persons receiving medical care for HIV Infection: Medical Monitoring Project, United States 2013 Cycle (June 2013–May 2014). Retrieved from: <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-hssr-mmp-2013.pdf>.
- Chen JA, Owens MD, Browne KC, Williams EC, 2018 Alcohol-related and mental health care for patients with unhealthy alcohol use and posttraumatic stress disorder in a National Veterans Affairs cohort. *J. Subst. Abuse Treat.* 85, 1–9. [PubMed: 29291765]
- Cohen E, Feinn R, Arias A, Kranzler HR, 2007 Alcohol treatment utilization: Findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Drug Alcohol Depend.* 86, 214–221. [PubMed: 16919401]
- Dawson DA, Goldstein RB, Grant BF, 2012 Factors associated with first utilization of different types of care for alcohol problems. *J. Stud. Alcohol Drugs* 73, 647–656. [PubMed: 22630803]
- Dawson DA, Grant BF, Stinson FS, Chou PS, 2006 Estimating the effect of help-seeking on achieving recovery from alcohol dependence. *Addiction* 101, 824–834. [PubMed: 16696626]
- Department of Veterans Affairs, 2015 VA/DoD Clinical Practice Guideline for the Management of Substance Use Disorders. <http://www.healthquality.va.gov/guidelines/MH/sud/VADoDSUDCPGRevised22216.pdf>.
- Department of Veterans Affairs, 2017 Hepatitis C: Information for Providers. <https://www.hepatitis.va.gov/pdf/Hepatitis-C-Factsheet-Providers.pdf>.
- Department of Veterans Affairs, 2018 Veterans Health Administration - About VHA. Accessed October 4, 2018. <https://www.va.gov/health/aboutVHA.asp>.
- Donato F, Tagger A, Gelatti U, Parrinello G, Boffetta P, Albertini A, Decarli A, Trevisi P, Ribero ML, Martelli C, Porru S, Nardi G, 2002 Alcohol and hepatocellular carcinoma: The effect of lifetime intake and hepatitis virus infections in men and women. *Am. J. Epidemiol.* 155, 323–331. [PubMed: 11836196]
- Edelman EJ, Williams EC, Marshall BDL, 2018 Addressing unhealthy alcohol use among people living with HIV: Recent advances and research directions. *Curr. Opin. Infect. Dis.* 31, 1–7. [PubMed: 29176446]

- Finlay AK, Ellerbe LS, Wong JJ, Timko C, Rubinsky AD, Gupta S, Bowe TR, Burden JL, Harris AH, 2017 Barriers to and facilitators of pharmacotherapy for alcohol use disorder in VA residential treatment programs. *J. Subst. Abuse Treat.* 77, 38–43. [PubMed: 28476269]
- Fultz SL, Skanderson M, Mole LA, Gandhi N, Bryant K, Crystal S, Justice AC, 2006 Development and verification of a "virtual" cohort using the National VA Health Information System. *Med. Care* 44, S25–S30. [PubMed: 16849965]
- Fuster D, Cheng DM, Quinn EK, Nunes D, Saitz R, Samet JH, Tsui JI, 2014 Chronic hepatitis C virus infection is associated with all-cause and liver-related mortality in a cohort of HIV-infected patients with alcohol problems. *Addiction* 109, 62–70. [PubMed: 24112091]
- Fuster D, Sanvisens A, Bolao F, Serra I, Rivas I, Tor J, Muga R, 2015 Impact of hepatitis C virus infection on the risk of death of alcohol-dependent patients. *J. Viral Hepat.* 22, 18–24. [PubMed: 25131721]
- Galvan FH, Bing EG, Fleishman JA, London AS, Caetano R, Burnman MA, Longshore D, Morton SC, Orlando M, Shapiro M, 2002 The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: Results from the HIV cost and services utilization study. *J. Stud. Alcohol* 63, 179–186. [PubMed: 12033694]
- Glass JE, Perron BE, Ilgen M, Chermack ST, Ratliff S, Zivin K, 2010 Prevalence and correlates of specialty substance use disorder treatment for Department of Veterans Affairs Healthcare System patients with high alcohol consumption. *Drug Alcohol Depend.* 112, 150–155. [PubMed: 20656425]
- Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, Pickering RP, Ruan WJ, Smith SM, Huang B, Hasin DS, 2015 Epidemiology of DSM-5 alcohol use disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry* 72, 757–766. [PubMed: 26039070]
- Grant BF, Pickering RP, 1996 Comorbidity between DSM-IV alcohol and drug use disorders: Results from the National Longitudinal Alcohol Epidemiologic Survey. *Alcohol Health Res. World* 20, 67–72.
- Grant BF, Saha TD, Ruan WJ, Goldstein RB, Chou SP, Jung J, Zhang H, Smith SM, Pickering RP, Huang B, Hasin DS, 2016 Epidemiology of DSM-5 drug use disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *JAMA Psychiatry* 73, 39–47. [PubMed: 26580136]
- Greenland S, 1995 Avoiding power loss associated with categorization and ordinal scores in dose-response and trend analysis. *Epidemiology* 6, 450–454. [PubMed: 7548361]
- Hagedorn HJ, Brown R, Dawes M, Dieperink E, Myrick DH, Oliva EM, Wagner TH, Wisdom JP, Harris AH, 2016 Enhancing access to alcohol use disorder pharmacotherapy and treatment in primary care settings: ADaPT-PC. *Implement. Sci.* 11, 64. [PubMed: 27164835]
- Harris AH, Ellerbe L, Reeder RN, Bowe T, Gordon AJ, Hagedorn H, Oliva E, Lembke A, Kivlahan D, Trafton JA, 2013 Pharmacotherapy for alcohol dependence: Perceived treatment barriers and action strategies among Veterans Health Administration service providers. *Psychol. Serv.* 10, 410–419. [PubMed: 23356858]
- Harris AH, Kivlahan DR, Bowe T, Humphreys KN, 2010 Pharmacotherapy of alcohol use disorders in the Veterans Health Administration. *Psychiatr. Serv.* 61, 392–398. [PubMed: 20360279]
- Harris AHS, Oliva E, Bowe T, Humphreys KH, Kivlahan D, Trafton JA, 2012 Pharmacotherapy of alcohol use disorders in the Veterans Health Administration: Patterns of receipt and persistence. *Psychiatr. Serv.* 63, 679–685. [PubMed: 22549276]
- Hu X, Harman J, Winterstein AG, Zhong Y, Wheeler AL, Taylor TN, Plankey M, Rubtsova A, Cropsey K, Cohen MH, Adimora AA, Milam J, Adedmeji A, Cook RL, 2016 Utilization of alcohol treatment among HIV-positive women with hazardous Drinking. *J. Subst. Abuse Treat.* 64, 55–61. [PubMed: 26961420]
- Hutchinson SJ, Bird SM, Goldberg DJ, 2005 Influence of alcohol on the progression of hepatitis C virus infection: A meta-analysis. *Clin. Gastroenterol. Hepatol.* 3, 1150–1159. [PubMed: 16271348]

- Ilgen MA, Price AM, Burnett-Zeigler I, Perron B, Islam K, Bohnert AS, Zivin K, 2011 Longitudinal predictors of addictions treatment utilization in treatment-naïve adults with alcohol use disorders. *Drug Alcohol Depend.* 113, 215–221. [PubMed: 20828944]
- Jonas DE, Amick HR, Feltner C, Bobashev G, Thomas K, Wines R, Kim MM, Shanahan E, Gass CE, Rowe CJ, Garbutt JC, 2014 Pharmacotherapy for adults with alcohol use disorders in outpatient settings: A systematic review and meta-analysis. *JAMA* 311, 1889–1900. [PubMed: 24825644]
- Justice AC, McGinnis KA, Tate JP, Braithwaite RS, Bryant KJ, Cook RL, Edelman EJ, Fiellin LE, Freiberg MS, Gordon AJ, Kraemer KL, Marshall BDL, Williams EC, Fiellin DA, 2016 Risk of mortality and physiologic injury evident with lower alcohol exposure among HIV infected compared with uninfected men. *Drug Alcohol Depend.* 161, 95–103. [PubMed: 26861883]
- Kaner EF, Heather N, Brodie J, Lock CA, McAvoy BR, 2001 Patient and practitioner characteristics predict brief alcohol intervention in primary care. *Br. J. Gen. Pract.* 51, 822–827. [PubMed: 11677706]
- Knudsen HK, Roman PM, 2014 Dissemination, adoption, and implementation of acamprosate for treating alcohol use disorders. *J. Stud. Alcohol Drugs* 75, 467–475. [PubMed: 24766759]
- Lapham GT, Achtmeyer CE, Williams EC, Hawkins EJ, Kivlahan DR, Bradley KA, 2012 Increased documented brief alcohol interventions with a performance measure and electronic decision support. *Med. Care* 50, 179–187. [PubMed: 20881876]
- Liang KY, Zeger SL, 1986 Longitudinal data analysis using generalized linear models. *Biometrika* 73, 13–22.
- Lin M, Kramer J, White D, Cao Y, Tavakoli-Tabasi S, Madu S, Smith D, Asch SM, El-Serag HB, Kanwal F, 2017 Barriers to hepatitis C treatment in the era of direct-acting anti-viral agents. *Aliment. Pharmacol. Ther.* 46, 992–1000. [PubMed: 28949020]
- Lingford-Hughes AR, Welch S, Peters L, Nutt DJ, 2012 BAP updated guidelines: evidence-based guidelines for the pharmacological management of substance abuse, harmful use, addiction and comorbidity: Recommendations from BAP. *J. Psychopharmacol.* 26, 899–952. [PubMed: 22628390]
- McCartney EM, Beard MR, 2010 Impact of alcohol on hepatitis C virus replication and interferon signaling. *World J. Gastroenterol.* 16, 1337–1343. [PubMed: 20238400]
- McGinnis KA, Fiellin DA, Tate JP, Cook RL, Braithwaite RS, Bryant KJ, Edelman EJ, Gordon AJ, Kraemer KL, Maisto SA, Justice AC, For the Veterans Aging Cohort Study, 2016. Number of drinks to "feel a buzz" by HIV status and viral load in men. *AIDS Behav.* 20, 504–511. [PubMed: 26936030]
- Mitchell MC, Memisoglu A, Silverman BL, 2012 Hepatic safety of injectable extended-release naltrexone in patients with chronic hepatitis C and HIV infection. *J. Stud. Alcohol Drugs* 73, 991–997. [PubMed: 23036218]
- National Institute for Health and Care Excellence, 2015 Alcohol use disorders: Diagnosis, assessment and management of harmful drinking and alcohol dependence National Institute for Health and Care Excellence, Manchester, England.
- National Institute on Alcohol Abuse and Alcoholism (NIAAA), 2007 Helping Patients Who Drink Too Much: A Clinician's Guide (Updated 2005 Edition). National Institutes of Health, U.S. Department of Health and Human Services, Washington, D.C..
- Ober AJ, Watkins KE, Hunter SB, Ewing B, Lamp K, Lind M, Becker K, Heinzerling K, Osilla KC, Diamant AL, Setodji CM, 2017 Assessing and improving organizational readiness to implement substance use disorder treatment in primary care: Findings from the SUMMIT study. *BMC Fam. Pract.* 18, 107. [PubMed: 29268702]
- Oliva EM, Maisel NC, Gordon AJ, Harris AH, 2011 Barriers to use of pharmacotherapy for addiction disorders and how to overcome them. *Curr. Psychiatry. Rep.* 13, 374–381. [PubMed: 21773951]
- Orwat J, Saitz R, Tompkins CP, Cheng DM, Dentato MP, Samet JH, 2011 Substance abuse treatment utilization among adults living with HIV/AIDS and alcohol or drug problems. *J. Subst. Abuse Treat.* 41, 233–242. [PubMed: 21700412]
- Owens MD, Ioannou GN, Tsui JI, Edelman JE, Greene PA, Williams EC, 2018 Receipt of alcohol-related care among patients with HCV and unhealthy alcohol use. *Drug Alcohol Depend.* 188, 79–85. [PubMed: 29754030]

- Pettinati HM, Weiss RD, Dundon W, Miller WR, Donovan D, Ernst DB, Rounsaville BJ, 2005 A structured approach to medical management: A psychosocial intervention to support pharmacotherapy in the treatment of alcohol dependence. *J. Stud. Alcohol Suppl* 170–178. [PubMed: 16223068]
- Richmond R, Kehoe L, Heather N, Wodak A, Webster I, 1996 General practitioners' promotion of healthy life styles: What patients think. *Aust. N. Z. J. Public Health* 20, 195–200. [PubMed: 8799096]
- Rubinsky AD, Chen C, Batki SL, Williams EC, Harris AH, 2015 Comparative utilization of pharmacotherapy for alcohol use disorder and other psychiatric disorders among U.S. Veterans Health Administration patients with dual diagnoses. *J. Psychiatr. Res.* 69, 150–157. [PubMed: 26343607]
- Safdar K, Schiff ER, 2004 Alcohol and hepatitis C. *Semin. Liver Dis.* 24, 305–315. [PubMed: 15349807]
- Saitz R, 2005 Clinical practice. Unhealthy alcohol use. *N. Engl. J. Med.* 352, 596–607. [PubMed: 15703424]
- Saxon AJ, Sloan KL, Reoux J, Haver VM, 1998 Disulfiram use in patients with abnormal liver function test results. *J. Clin. Psychiatry* 59, 313–316. [PubMed: 9671344]
- StataCorp, 2017 Stata Statistical Software: Release 15. StataCorp LLC, College Station, TX.
- Stinson FS, Grant BF, Dawson DA, Ruan WJ, Huang B, Saha T, 2005 Comorbidity between DSM-IV alcohol and specific drug use disorders in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Drug Alcohol Depend.* 80, 105–116. [PubMed: 16157233]
- Szabo G, Aloman C, Polyak SJ, Weinman SA, Wands J, Zakhari S, 2006 Hepatitis C infection and alcohol use: A dangerous mix for the liver and antiviral immunity. *Alcohol. Clin. Exp. Res.* 30, 709–719. [PubMed: 16573590]
- Szabo G, Wands JR, Eken A, Osna NA, Weinman SA, Machida K, Joe Wang H, 2010 Alcohol and hepatitis C virus: Interactions in immune dysfunctions and liver damage. *Alcohol. Clin. Exp. Res.* 34, 1675–1686. [PubMed: 20608905]
- Taylor AL, Denniston MM, Kleven RM, McKnight-Eily LR, Jiles RB, 2016 Association of Hepatitis C Virus with alcohol use among U.S. adults: NHANES 2003–2010. *Am. J. Prev. Med* 51, 206–215. [PubMed: 27178884]
- Tetrault JM, Tate JP, McGinnis KA, Goulet JL, Sullivan LE, Bryant K, Justice AC, Fiellin DA, 2012 Hepatic safety and antiretroviral effectiveness in HIV-infected patients receiving naltrexone. *Alcohol. Clin. Exp. Res.* 36, 318–324. [PubMed: 21797892]
- Tsui JJ, Pletcher MJ, Vittinghoff E, Seal K, Gonzales R, 2006 Hepatitis C and hospital outcomes in patients admitted with alcohol-related problems. *J. Hepatol.* 44, 262–266. [PubMed: 16226823]
- Tsui JJ, Williams EC, Green PK, Berry K, Su F, Ioannou GN, 2016 Alcohol use and hepatitis C virus treatment outcomes among patients receiving direct antiviral agents. *Drug Alcohol Depend.* 169, 101–109. [PubMed: 27810652]
- Vagenas P, Azar MM, Copenhaver MM, Springer SA, Molina PE, Altice FL, 2015 The impact of alcohol use and related disorders on the HIV continuum of care: A systematic review: Alcohol and the HIV continuum of care. *Curr. HIV/AIDS Rep* 12, 421–436. [PubMed: 26412084]
- Volk RJ, Steinbauer JR, Cantor SB, 1996 Patient factors influencing variation in the use of preventive interventions for alcohol abuse by primary care physicians. *J. Stud. Alcohol* 57, 203–209. [PubMed: 8683970]
- Weisner C, Matzger H, Kaskutas LA, 2003 How important is treatment? One-year outcomes of treated and untreated alcohol-dependent individuals. *Addiction* 98, 901–911. [PubMed: 12814496]
- Williams EC, Achtmeyer CE, Young JP, Berger D, Curran G, Bradley KA, Richards J, Siegel MB, Ludman EJ, Lapham GT, Forehand M, Harris AHS, 2017a Barriers to and facilitators of alcohol use disorder pharmacotherapy in primary care: A qualitative study in five VA clinics. *J. Gen. Intern. Med.* 33, 258–267. [PubMed: 29086341]
- Williams EC, Gupta S, Rubinsky AD, Glass JE, Jones-Webb R, Bensley KM, Harris AHS, 2017b Variation in receipt of pharmacotherapy for alcohol use disorders across racial/ethnic groups: A

- national study in the U.S. Veterans Health Administration. *Drug Alcohol Depend.* 178, 527–533. [PubMed: 28728114]
- Williams EC, Hahn JA, Saitz R, Bryant K, Lira MC, Samet JH, 2016a Alcohol use and Human Immunodeficiency Virus (HIV) Infection: Current knowledge, implications, and future directions. *Alcohol. Clin. Exp. Res.* 4, 2056–2072.
- Williams EC, Joo YS, Lipira L, Glass JE, 2016b Psychosocial stressors and alcohol use, severity, and treatment receipt across human immunodeficiency virus (HIV) status in a nationally representative sample of US residents. *Subst. Abuse* 38, 1–9.
- Williams EC, Lapham GT, Hawkins EJ, Rubinsky AD, Morales LS, Young BA, Bradley KA, 2012 Variation in documented care for unhealthy alcohol consumption across race/ethnicity in the Department of Veterans Affairs Healthcare System. *Alcohol. Clin. Exp. Res.* 36, 1614–1622. [PubMed: 22404130]
- Williams EC, Lapham GT, Rubinsky AD, Chavez LJ, Berger D, Bradley KA, 2017c Influence of a targeted performance measure for brief intervention on gender differences in receipt of brief intervention among patients with unhealthy alcohol use in the Veterans Health Administration. *J. Subst. Abuse Treat.* 81, 11–16. [PubMed: 28847450]
- Williams EC, Lapham GT, Shortreed SM, Rubinsky AD, Bobb JF, Bensley KM, Catz SL, Richards JE, Bradley KA, 2017d Among patients with unhealthy alcohol use, those with HIV are less likely than those without to receive evidence-based alcohol-related care: A national VA study. *Drug Alcohol Depend.* 174, 113–120. [PubMed: 28324813]
- Williams EC, McGinnis KA, Edelman EJ, Matson TE, Gordon A, Marshall BD, Bryant KJ, Rubinsky AD, Lapham GT, Satre DD, Richards JE, Catz SL, Fiellin DA, Justice AC, Bradley KA, 2018 Level of alcohol use associated with HIV care continuum targets in a national U.S. sample of persons living with HIV receiving healthcare. *AIDS Behav.* [Epub ahead of print].
- Williams EC, Rubinsky AD, Chavez LJ, Lapham GT, Rittmueller SE, Achtmeyer CE, Bradley KA, 2014a An early evaluation of implementation of brief intervention for unhealthy alcohol use in the US Veterans Health Administration. *Addiction* 109, 1472–1481. [PubMed: 24773590]
- Williams EC, Rubinsky AD, Lapham GT, Chavez LJ, Rittmueller SE, Hawkins EJ, Grossbard JR, Kivlahan DR, Bradley KA, 2014b Prevalence of clinically recognized alcohol and other substance use disorders among VA outpatients with unhealthy alcohol use identified by routine alcohol screening. *Drug Alcohol Depend.* 135, 95–103. [PubMed: 24360928]
- Young BA, Maynard C, Boyko EJ, 2003 Racial differences in diabetic nephropathy, cardiovascular disease, and mortality in a national population of veterans. *Diabetes Care* 26, 2392–2399. [PubMed: 12882868]
- Younossi ZM, Zheng L, Stepanova M, Venkatesan C, Mir HM, 2013 Moderate, excessive or heavy alcohol consumption: Each is significantly associated with increased mortality in patients with chronic hepatitis C. *Aliment. Pharmacol. Ther.* 37, 703–709. [PubMed: 23432436]
- Zou G, 2004 A modified poisson regression approach to prospective studies with binary data. *Am. J. Epidemiol.* 159, 702–706. [PubMed: 15033648]

Highlights

- Among VA patients with alcohol use disorder (AUD) and HCV and/or HIV, about half had drug use disorder (DUD)
- DUD was associated with increased receipt of specialty addictions treatment for AUD
- DUD was associated with receipt of pharmacotherapy in persons with HCV, but not HIV

Table 1.

Characteristics of VA outpatients with HCV and AUD: Overall and across comorbid drug use disorder status

	DUD (N=9,964)		No DUD (N=12,075)		Chi-Square value	p-value	Total (N=22,039)	
	N	(%)	N	(%)			N	(%)
Female	208	(2.1)	207	(1.7)	4.12	0.042	415	(1.9)
Age					215.79	<0.001		
18-29	95	(1.0)	32	(0.3)			127	(0.6)
30-44	376	(3.8)	239	(2.0)			615	(2.8)
45-64	9,144	(91.8)	10,992	(91.0)			20,136	(91.4)
65+	349	(3.5)	812	(6.7)			1,161	(5.3)
Race/ethnicity					953.66	<0.001		
American Indian/Alaska Native	145	(1.5)	174	(1.5)			319	(1.5)
Asian American/Pacific Islander	99	(1.0)	137	(1.2)			236	(1.1)
Black/African American	4,465	(45.5)	2,984	(25.7)			7,449	(34.8)
Hispanic/Latino	661	(6.7)	858	(7.4)			1,519	(7.1)
White	4,435	(45.2)	7,465	(64.3)			11,900	(55.6)
Marital status					381.07	<0.001		
Divorced/Separated	4,459	(45.0)	5,583	(46.5)			10,042	(45.8)
Married	1,473	(14.9)	2,816	(23.5)			4,289	(19.6)
Never married/Single	3,588	(36.2)	3,158	(26.3)			6,746	(30.8)
Widowed	397	(4.0)	454	(3.8)			851	(3.9)
Opioid use disorder	3,308	(33.2)	0	(0.0)	4.7e+3	<0.001	3,308	(15.0)
Stimulant use disorder	7,090	(71.2)	0	(0.0)	1.3e+4	<0.001	7,090	(32.2)
Other drug use disorder*	4,968	(49.9)	0	(0.0)	7.8e+3	<0.001	4,968	(22.5)
VA eligibility status					12.04	0.002		
Full VA coverage	1,730	(17.4)	1,897	(15.7)			3,627	(16.5)
Service connection <50%	1,791	(18.0)	2,146	(17.8)			3,937	(17.9)
Non-service connected	6,443	(64.7)	8,032	(66.5)			14,475	(65.7)
Fiscal year of first AUDIT-C					3.36	0.339		
2010	4,347	(43.6)	5,128	(42.5)			9,475	(43.0)
2011	2,731	(27.4)	3,343	(27.7)			6,074	(27.6)
2012	1,970	(19.8)	2,450	(20.3)			4,420	(20.1)
2013	916	(9.2)	1,154	(9.6)			2,070	(9.4)
Major depression	2,387	(24.0)	1,237	(10.2)	747.05	<0.001	3,624	(16.4)
Anxiety disorder	2,085	(20.9)	1,464	(12.1)	313.00	<0.001	3,549	(16.1)
Post-Traumatic Stress Disorder (PTSD)	2,851	(28.6)	2,057	(17.0)	422.74	<0.001	4,908	(22.3)
Other mood disorder	5,974	(60.0)	4,128	(34.2)	1.5e+3	<0.001	10,102	(45.8)
Serious mental illness	2,747	(27.6)	1,186	(9.8)	1.2e+3	<0.001	3,933	(17.9)

* Includes cannabis, hallucinogen, and/or sedative

Table 2.

Characteristics of VA outpatients with HIV and AUD: Overall and across comorbid drug use disorder status

	DUD (N=1,043)		No DUD (N=791)		Chi-Square value	p-value	Total (N=1,834)	
	N	(%)	N	(%)			N	(%)
Female	25	(2.4)	13	(1.6)	1.26	0.262	38	(2.1)
Age					45.46	<0.001		
18-29	14	(1.3)	16	(2.0)			30	(1.6)
30-44	129	(12.4)	106	(13.4)			235	(12.8)
45-64	876	(84.0)	597	(75.5)			1,473	(80.3)
65+	24	(2.3)	72	(9.1)			96	(5.2)
Race/ethnicity					142.80	<0.001		
American Indian/Alaska Native	12	(1.2)	8	(1.0)			20	(1.1)
Asian American/Pacific Islander	7	(0.7)	6	(0.8)			13	(0.7)
Black/African American	770	(74.4)	373	(48.4)			1,143	(63.3)
Hispanic/Latino	73	(7.1)	71	(9.2)			144	(8.0)
White	173	(16.7)	312	(40.5)			485	(26.9)
Marital status					6.75	0.080		
Divorced/Separated	333	(32.2)	289	(37.1)			622	(34.3)
Married	109	(10.6)	90	(11.5)			199	(11.0)
Never married/Single	553	(53.5)	370	(47.4)			923	(50.9)
Widowed	38	(3.7)	31	(4.0)			69	(3.8)
Opioid use disorder	259	(24.8)	0	(0.0)	228.72	<0.001	259	(14.1)
Stimulant use disorder	896	(85.9)	0	(0.0)	1.3e+3	<0.001	896	(48.9)
Other drug use disorder*	446	(42.8)	0	(0.0)	446.93	<0.001	446	(24.3)
VA eligibility status					2.90	0.235		
Full VA coverage	197	(18.9)	133	(16.8)			330	(18.0)
Service connection <50%	185	(17.7)	162	(20.5)			347	(18.9)
Non-service connected	661	(63.4)	496	(62.7)			1,157	(63.1)
Fiscal year of first AUDIT-C					1.38	0.711		
2010	400	(38.4)	283	(35.8)			683	(37.2)
2011	289	(27.7)	233	(29.5)			522	(28.5)
2012	239	(22.9)	185	(23.4)			424	(23.1)
2013	115	(11.0)	90	(11.4)			205	(11.2)
Major depression	285	(27.3)	115	(14.5)	43.13	<0.001	400	(21.8)
Anxiety disorder	183	(17.6)	110	(13.9)	4.44	0.035	293	(16.0)
Post-Traumatic Stress Disorder (PTSD)	233	(22.3)	109	(13.8)	21.72	<0.001	342	(18.7)
Other mood disorder	653	(62.6)	307	(38.8)	102.12	<0.001	960	(52.3)
Serious mental illness	324	(31.1)	94	(11.9)	94.05	<0.001	418	(22.8)

* Includes cannabis, hallucinogen, and/or sedative

Table 3.

Receipt of AUD treatment among VA patients with and without comorbid drug use disorder among those with HCV/HIV and AUD

	DUD		No DUD		IRR ^a	95% CI	P-value ^b
	%	95% CI	%	95% CI			
Patients with HCV and AUD (N=22,039)							
Receipt of any AUD specialty addictions treatment within 365 days following positive screen (AUDIT-C 5) ^c							
Unadjusted	52.7	(51.8-53.6)	22.1	(21.4-22.7)	2.39	(2.31-2.47)	<0.001
Adjusted ^c	52.8	(51.9-53.7)	22.4	(21.8-23.1)	1.89	(1.82-1.96)	<0.001
Receipt of any AUD pharmacotherapy within 365 days following positive screen (AUDIT-C 5) ^d							
Unadjusted	11.2	(10.7-11.8)	5.5	(5.1-5.8)	2.05	(1.89-2.23)	<0.001
Adjusted ^c	11.3	(10.8-11.9)	5.6	(5.2-6.0)	1.50	(1.37-1.65)	<0.001
Patients with HIV and AUD (N=1,834)							
Receipt of any AUD specialty addictions treatment within 365 days following positive screen (AUDIT-C 5)							
Unadjusted	52.9	(50.1-55.8)	22.5	(19.7-25.2)	2.35	(2.06-2.69)	<0.001
Adjusted ^c	52.7	(50.0-55.5)	22.7	(20.0-25.4)	1.94	(1.68-2.24)	<0.001
Receipt of any AUD pharmacotherapy within 365 days following positive screen (AUDIT-C 5) ^d							
Unadjusted	7.1	(5.6-8.6)	5.2	(3.7-6.6)	1.37	(0.97-1.93)	0.071
Adjusted ^c	7.2	(5.7-8.8)	5.1	(3.7-6.5)	1.21	(0.81-1.81)	0.345

^aIncident Rate Ratio.

^bp-value from test to evaluate if IRR is equal to one.

^cAdjusted for gender, age, race/ethnicity, marital status, VA eligibility status, fiscal year in which positive AUDIT-C screen occurred, and mental health conditions.

^dIncludes acamprosate, disulfiram, topiramate, oral or injectable naltrexone.