Identification of potential drug targets and inhibitor of the pathogenic bacteria *Shigella flexneri* 2a through the subtractive genomic approach

Arafat Rahman Oany¹ · Mamun Mia¹ · Tahmina Pervin² · Md. Nazmul Hasan³ · Akinori Hirashima⁴

Received: 27 February 2018 / Accepted: 28 May 2018 / Published online: 4 June 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Shigella flexneri 2a is one of the most pathogenic bacteria among the *Shigella* spp., which is responsible for dysentery and causes masses of deaths throughout the world per year. A proper identification of the potential drug targets and inhibitors is crucial for the treatment of the shigellosis due to their emerging multidrug resistance (MDR) patterns. In this study, a systematic subtractive approach was implemented for the identification of novel therapeutic targets of *S. flexneri* 2a (301) through genome-wide metabolic pathway analysis of the essential genes and proteins. Ligand-based virtual screening and ADMET analyses were also made for the identification of potential inhibitors as well. Initially, we found 70 essential unique proteins as novel targets. After subsequent prioritization, finally we got six unique targets as the potential therapeutic targets and their three-dimensional models were built thereafter. Aspartate- β -semialdehyde dehydrogenase (ASD), was the most potent target among them and used for docking analysis through ligand-based virtual screening. The compound 3 (PubChem CID: 11319750) suited well as the best inhibitor of the ASD through ADMET and enzyme inhibition capacity analysis. To end, we hope that our proposed therapeutic targets and its inhibitors might give some breakthrough to treat shigellosis efficiently in in vitro.

Keywords ADMET · Molecular docking · ASD · MDR · Dysentery

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s40203-018-0048-2>) contains supplementary material, which is available to authorized users.

✉ Arafat Rahman Oany
arafatr@outlook.com

✉ Akinori Hirashima
ahirasim@agr.kyushu-u.ac.jp

¹ Department of Biotechnology and Genetic Engineering, Life Science Faculty, Mawlana Bhashani Science and Technology University, Tangail 1902, Bangladesh

² Biotechnology and Genetic Engineering Discipline, Life Science School, Khulna University, Khulna, Bangladesh

³ Department of Genetic Engineering and Biotechnology, Jessore University of Science and Technology, Jessore, Bangladesh

⁴ Laboratory of Pesticide Chemistry, Division of Molecular Biosciences, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812-8581, Japan

Introduction

Bacillary syndrome or shigellosis is mostly caused by *Shigella* species, a member of Enterobacteriaceae family, and identified as the etiologic agent since 1897 (Garza-Mayers et al. 2015; Jennison and Verma 2004; Jin et al. 2002). *Shigella* is a gram negative, facultative anaerobe, inordinately skillful enteric bacterial pathogens, mainly induced diarrhea by occupying and disseminating into the follicle linked membranous cells, through using the type III secretion system, which necessitates forking out virulence factor into infected cells (Garza-Mayers et al. 2015; Perdomo et al. 1994). Most of the common features and genetic relatedness found to the member of the genus *Escherichia* apparently indicate a subtype of *E. coli* (Jin et al. 2002). Contaminated water is the most common source of *Shigella* and a fecal–oral route is the ordinary path of contagion which is increased via unhygienic conditions (Hale and Keusch 1996; Jennison and Verma 2004). Estimation resulted that about 164.7 million people develop shigellosis per year throughout the world and approximately 1.1 million fallen into the deaths. In underdeveloped countries, approximately 163.2

million personnel are affected by shigellosis or bacillary dysentery per annum, of which most of them are children under the age of five estimating 69% (Kotloff et al. 1999).

Shigella flexneri, *S. boydii*, *S. sonnei* and *S. dysenteriae* are the four species of *Shigella* where higher death rate caused by *S. flexneri* in the majority of developing countries (Bennish and Wojtyniak 1991; Jennison and Verma 2004).

Each of the species also divided into several serotypes relies on the variations of biochemical test and O-antigen. *S. flexneri* possess—13 serotypes, *S. dysenteriae*—12, *S. boydii*—18 and *S. Sonnei*—1 serotype, respectively (Hale and Keusch 1996; Jennison and Verma 2004). *S. flexneri* has been the most commonly isolated species globally and accounts for 60% of cases in the unindustrialized countries (Kotloff et al. 1999). *S. flexneri* has the power to up-regulate acid resistance genes in the presence of low acidity which allows the low infectious dose to host. As a result, *S. flexneri*, especially subspecies *S. flexneri* 2a, is being more infectious than the other species where a few cells are able to induce disease in adult personnel. The estimated dose of *S. flexneri* to cause disease is 100 cells on an average (DuPont et al. 1989; Small et al. 1994).

When *Shigella* start to penetrate, replicate and spread into the mucosal epithelial cells after reaching the colon, the host innate immune system subsequently raises up an inflammatory response against that behavior, which leads to the destruction of the colonic epithelial layer. Shigellosis is the clinical feature of this phenomenon (Philpott et al. 2000). About 18.4% *S. flexneri* of the total *Shigella* isolates had been reported to CDC during the time period in between 1989 and 2002 (Gupta et al. 2004). Generally, 1b, 2a, 3a, 4a and 6 are the most common serotypes of *S. flexneri* in developing countries, whereas several enterotoxins producing 2a strains are prevalent serotype in industrialized countries (DuPont et al. 1969; Fasano et al. 1995; Kotloff et al. 1999). For both sporadic and endemic shigellosis, *S. flexneri* 2a is the primary felon in the US. According to the report of the National Communicable Disease Center, there is one-fourth of all *Shigella* isolates accounted for *S. flexneri* 2a from 1963 to 1966, whereas 11,405 isolates found in 1967 (US Department of Health, Education, and Welfare, Public Health Service, National Communicable Disease Center 1968; Eichner et al. 1968). In China, the principal causal agent of dysentery is *S. flexneri* serotype 2a, where 50–70% cases accounted for 2a of about 10 million affected peoples in every year (Mei et al. 1989). The genome of *S. flexneri* 2a (301) strain was sequenced in 2002. A chromosome of 4,607,203 bp and a virulence plasmid named pCP301 of 221,618 bp make up the whole genome of *S. flexneri* 2a (301) strain. About a sequence of ~3.9 Mb found similar to *E. coli* K12 and O157 strain. The genome contains 314 IS elements and 64 S-islands, where seven *ipaH* genes located in five large S-islands. For better understanding about its

pathogenicity, some of the other virulence genes and their functions have been identified which are related to infections (Jin et al. 2002).

The treatment of Shigellosis largely depends on conventional antibiotic therapy, although the risk of shigellosis can be minimized by washing hands before handling foods. Conventional antibiotic treatment is allowed when infections reach to acute or chronic conditions. But in most cases, antibiotic treatment cannot satisfy the thrust of the therapeutic implications due to rising antibiotic resistance of *Shigella* species. It has been reported that *Shigella* spp. acclimatize resistance capability to mostly used antibiotics like ampicillin, tetracycline, streptomycin, nalidixic acid and sulfamethoxazole-trimethoprim (Dutta et al. 2002). Ciprofloxacin, a third generation antibiotic is the no longer useful treatment for shigellosis in South Asian countries including Bangladesh (Talukder et al. 2004, 2006). As there is no vaccine available that can provide adequate protection against this infectious agent and the current preventive treatment schemes are found to be inadequate, it strongly suggests that there is an urgent and continuing need to find new drugs and vaccine candidates to undertake this baneful agent.

A lot of novel scheme for the identification of bacterial drug targets has been pioneered by the available information of the bacterial genome sequences (Allsop 1998; Chan et al. 2010). As many drawbacks like time-consuming, costly are found in traditional drug discovery methods, advanced genome-based technology has opened an attractive choice to identify drug targets to the researchers. At the genome level, drug targets for any harmful infectious agent are currently easier due to the accessibility of each host and pathogen genome sequence information (Allsop 1998; Stumm et al. 2002). Nowadays, for the development of potential drug and vaccine targets against various infective pathogens, the most effective choice is computational based techniques (Amineni et al. 2010; Barh and Kumar 2009; Damte et al. 2013; Hasan et al. 2016; Oany et al. 2014b, 2017). The essential protein set for pathogen survival, however not in the host, can be identified by using subtractive and comparative genomics approach combined with metabolic pathway (Rahman et al. 2014). Identification of non-homologous essential genes for pathogens by subtraction from both host and pathogen metabolic pathway assures no interaction of drugs with host targets. On the other hand, the selection of conserved proteins can be emphasized by comparative genomics method among several species as the most prosperous targets (Galperin and Koonin 1999; Kobayashi et al. 2003; Koonin et al. 1998; Tatusov et al. 1997). Drug targets for many of the infectious agents can be designed by using advanced bioinformatics tools with the help of genomics, proteomics, and metabolomics.

In the present study, we have used the subtractive genomics approach to identify the potential therapeutic targets and

its inhibitors through the analysis of the metabolic pathways of the pathogen *S. flexneri* 2a, for the identification of better treatment of shigellosis.

Materials and methods

The entire procedure for the identification of appropriate drug targets and prediction of inhibitor is shown in Fig. 1 through the schematic diagram.

Pathway analysis of the host and pathogen

The Kyoto Encyclopedia of Genes and Genomes (KEGG) database (Kanehisa et al. 2006, 2009) was explored for

metabolic pathways for both human genomes and *Shigella flexneri* 2a (301). The identification numbers of all pathways from both organisms were listed. Thereafter, a manual comparison was made between the host and pathogen metabolic pathways. Those pathways that did not appear in the human genomes, but were present in the pathogen, according to the KEGG database annotations, were selected as unique to *Shigella flexneri* 2a. The remaining pathways were listed as common. Finally, amino acid sequences of proteins from common and unique pathways were retrieved from UniProtKB (Magrane 2011) database and stored as FASTA format.

Fig. 1 Schematic diagram of the whole study

Identification of non-homologous proteins from the pathogen

The proteins identified in unique and common pathways of the pathogen were subjected to Protein Basic Local Alignment Search Tool (BLAST) analysis against the human proteome by NCBI-BLASTp program (Johnson et al. 2008). The hits output was filtered on the basis of expectation value (e-value) inclusion threshold being set to 0.003, and a minimum bit score of 100. Proteins, that did not have hits below the e-value inclusion threshold of 0.003, were selected as non-homologous proteins. The non-redundant and Swiss-Prot databases were used for the analysis (Oany et al. 2014a).

Identification of essential proteins and their localization

For the identification of essential proteins of the pathogen, we used STRING (<http://string-db.org>), (Franceschini et al. 2012) a database of known and predicted protein interactions that work through physical and functional associations derived from genomic context, high-throughput experiments, co-expression and previous knowledge. We used STRING as an alternative to the Database of Essential Gene (DEG) due to unavailability of our target in the database. The subcellular localizations of all the essential proteins were predicted by CELLO v2.5 (Yu et al. 2004) and PSORTb (Gardy et al. 2004) server for cross-validation.

Novel target identification

Proteins located in the cytoplasm are the prioritized targets for the drug. We tried to find out novel targets from the essential proteins those are sorted initially, for better target identification. Therefore, the DrugBank (Wishart et al. 2007) database was used to identify the unique target. The BLASTp analysis with default parameters of the DrugBank database was used to identify the druggability as well as the unique targets. The hits were denoted as the common target and the non-hits were denoted as unique targets and used for prioritizing.

Drug target prioritization

The drug target prioritization of the novel targets was made through sequential molecular and structural criteria for evaluation of the suitability of each target for being potential drug targets (Agüero et al. 2008). This prioritization involved the calculation of molecular weight (MW) using computational tools available at Swiss-Prot database (Bairoch and Apweiler 2000). Physiological and chemical properties of the proteins are essential for considering protein as a suitable drug target, including; theoretical isoelectric point

(pI), instability index, aliphatic index, and hydrophobicity (Oany et al. 2014c). All these assessments were done with ProtParam tool of ExPASy (Gasteiger et al. 2005).

Determination of *N*-glycosylation sites in the protein is another important criterion for prioritizing drug target and linked to resistance to degradation, inferring that drug target proteins have an extended lifetime in vivo (Peter-Katalinic 2005). The *N*-glycosylation is assessed by NetNGlyc v1.0 Server (<http://www.cbs.dtu.dk/services/NetNGlyc/>). ModBase (Pieper et al. 2006) and Protein Data Bank (PDB) (Deshpande et al. 2005) were also assessed for the resolved three-dimensional structure of the targeted proteins for the further study. Finally, Supertarget (<http://insilico.charite.de/supertarget/>) and Potential Drug Target Database (Gao et al. 2008) were assessed for securing the novel drug targets.

Homology modelling and model validation

Homology model of the top predicted targets were done by MODELLER v9 (Šali et al. 1995) and top scoring templates were selected for the model construction. Then the predicted models were employed for the validation through the PROCHECK (Laskowski et al. 1996) tools of the SWISS-MODEL Workspace (Arnold et al. 2006). Superposition between the template and the structure was also analyzed.

Electrostatic potential analysis and energy minimization

The electrostatic potential of the predicted model was built by APBS (Unni et al. 2011) server through transforming. pdb to.pqr file and the visualization was done by PyMOL molecular graphics system (Schrödinger 2015). For further securing the validation of the predicted model, the energy minimization was also done. The GROMOS 96 (Scott et al. 1999) force field was used for the minimization of the structure and validated by “Ramachandran Plot” statistic by using SWISS-MODEL Workspace (Arnold et al. 2006).

Active site analysis and virtual screening

The active site of the final proposed model was constructed by the computed atlas of surface topography of proteins (CASTp) server. This server builds a model through the pocket algorithm of the alpha shape theory (Dundas et al. 2006).

The virtual screening was done by a series of combinatorial approaches initiated with BindingDB (Liu et al. 2006) for getting insight about the potential inhibitors. Thereafter, Zinc (Irwin and Shoichet 2005) database which contains over 4.6 million compounds in ready-to-dock in 3D format was used for the screening purpose. Finally, PubChem (Bolton et al. 2008) database was also assessed for the screening

compounds for further confirmation with bioactivities. The Autodock Vina (Trott and Olson 2010) was used for the screening based on the mother compound with a validated inhibition. All the ligands selected for the docking studies were converted into the .pdbqt format using Open Babel toolbox (O'Boyle et al. 2011). The grid generated for molecular docking at the center X: -10.382 , Y: 58.6496 and Z: -29.0307 . The dimensions (Angstrom) used for the docking analysis were X: 25.00 , Y: 25.00 and Z: 25.00 .

Ligand properties and ADMET analysis

Ligand structural and behavioral properties were analyzed for the cross-validation of the ligands as a suitable therapeutic candidate. Wide ranges of properties were analyzed including Lipinski's Rule (Lipinski 2004), drug-likeness, enzymatic inhibition, molecular weight, and bioavailability. These parameters were assessed through various online server and tools including, Molinspiration (<http://www.molinspiration.com>), SEARCh (<http://sea.bkslab.org/search/>), Drug-likeness tool (Oprea 2000) and OSIRIS Property Explorer (Sander 2001).

The absorption, distribution, metabolism, excretion, and toxicity (ADMET) were also assessed for the selected compounds. The ADME SARfari (Davies et al. 2015), admet-SAR (Cheng et al. 2012), Swiss database (Wirth et al. 2012) and QikProp (Schrödinger 2012) were used for these assessments.

Results

Identification of unique and common pathways

After the initial analysis from the KEGG database, we found 27 common pathways of the host–pathogen and 39 unique pathways of the pathogen only. A total of 234 proteins from the host and a total of 170 proteins from the pathogen common pathways' were retrieved from the UniProt database. On the other hand, a total of 234 proteins were retrieved from the pathogen unique pathways' for further analysis. All these common and unique pathways with appropriate entry IDs are described in Table 1. The genes for the pathogen unique pathways were also retrieved and described in the Table S1. The complete pathway distribution of the pathogen is depicted in the Figure S1.

Non-homologous essential protein identification and their localization

All pathogen pathway specific proteins were checked for the BLASTp analysis to identify the non-homologous proteins against the host. A total of 18 proteins from the common

pathway and a total of 124 proteins of the unique pathways of the pathogen were detected as non-homologous.

STRING database retrieved the essential proteins from all the non-homologous proteins of the pathogen based on the highest confidence score of 0.9. All the experimental, co-expression and neighborhood data were selected for screening out the essential proteins. Well-isolated networks with lower seed were removed from the analysis for securing the best prediction. Finally, we got 13 essential proteins from the common pathways and 83 from the unique pathways. The interactive network is depicted in the Figure S2.

Both CELLO and PSORTb predicted the subcellular localizations of all the essential proteins including all the unique pathogen proteins (Fig. 2). The cytoplasmic region was the top most hit among them and finally, we found 7 membrane and 89 cytoplasmic essential proteins from both common and unique pathways.

Novel target identification and prioritization

The DrugBank database retrieved the novel targets through the BLASTp analysis with the default parameters. Finally, we found 70 unique proteins as a novel targets and went for further analysis. Distributions of these essential proteins in different metabolic pathways are visualized in Fig. 3.

In prioritization scheme, we employed several criteria to sort-out the best possible targets. These include several cutoff values comprising, molecular weight < 100 KDa; $pI < 7.2$; instability index < 40 ; aliphatic index in between 95 and 104; hydrophobicity > -0.142 . These assessment results are described in the Table S2. Through the physico-chemical properties analysis, we found 32 sorted proteins for the *N*-glycosylation analysis (Table S3). The parameter for the *N*-glycosylation analysis was > 2 *N*-glycosylated amino acids (Galperin and Koonin 1999; Oany et al. 2014c). Before the assessment of the super-target and potential therapeutic target databases, we secured 12 potential novel targets and to end with six targets (Table 2).

Homology modelling and validation

Homology models of the final six targets were constructed through MODELLER and are depicted in Fig. 4. These models are visualized by PyMOL molecular graphics system. Validations of these models were checked by constructing the "Ramachandran Plot" and its statistics (Table S4) with appropriate templates' information and identity. Finally, we selected the protein P0A9R1 for the further study based on the best target selection. The superposition between the template 1t4b_A and the modeled protein P0A9R1 was also analyzed as a part of the validation process with RMSD of 0.146 Å (Fig. 5).

Table 1 Host-pathogen common and pathogen unique pathways from KEGG database

KEGG entry name	Pathway ID	Pathway name
Host-pathogen common pathway		
M00001	00010	Glycolysis (Embden–Meyerhof pathway)
M00002	00010	Glycolysis, core module involving three-carbon compounds
M00003	00010	Gluconeogenesis
M00307	00620	Pyruvate oxidation
M00009	00020	Citrate cycle (TCA cycle, Krebs cycle)
M00010	00020	Citrate cycle, first carbon oxidation
M00011	00020	Citrate cycle, second carbon oxidation
M00004	00030	Pentose phosphate pathway (pentose phosphate cycle)
M00006	00030	Pentose phosphate pathway, oxidative phase
M00007	00030	Pentose phosphate pathway, non-oxidative phase
M00005	00230	PRPP biosynthesis
M00632	00052	Galactose degradation, Leloir pathway
M00082	00061	Fatty acid biosynthesis, initiation
M00086	00071	Beta-oxidation, acyl-CoA synthesis
M00087	00071	Beta-oxidation
M00565	00520	Nucleotide sugar biosynthesis, glucose ► UDP-glucose
M00554	00052	Nucleotide sugar biosynthesis, galactose ► UDP-galactose
M00048	00230	Inosine monophosphate biosynthesis
M00049	00230	Adenine ribonucleotide biosynthesis
M00050	00230	Guanine ribonucleotide biosynthesis
M00051	00240	Uridine monophosphate biosynthesis
M00052	00240	Pyrimidine ribonucleotide biosynthesis
M00020	00260	Serine biosynthesis
M00015	00330	Proline biosynthesis
M00120	00770	Coenzyme A biosynthesis
M00133	00330	Polyamine biosynthesis
Pathogen unique pathway		
M00168	00710	Crassulacean acid metabolism (CAM), dark
M00579	00720	Phosphate acetyltransferase-acetate kinase pathway
M00176	00920	Assimilatory sulfate reduction
M00008	00030	Entner–Doudoroff pathway
M00012	00630	Glyoxylate cycle
M00631	00030	D-Galacturonate degradation
M00061	00030	D-Glucuronate degradation
M00565	00500	Trehalose biosynthesis
M00093	00564	Phosphatidylethanolamine (PE) biosynthesis
M00060	00540	Lipopolysaccharide biosynthesis
M00063	00540	CMP-KDO biosynthesis
M00064	00540	ADP-L-glycero-D-manno-heptose biosynthesis
M00096	00900	C5 isoprenoid biosynthesis
M00364	00900	C10–C20 isoprenoid biosynthesis
M00053	00240	Pyrimidine deoxyribonucleotide biosynthesis
M00018	00260	Threonine biosynthesis
M00021	00270	Cysteine biosynthesis
M00019	00290	Valine/isoleucine biosynthesis
M00570	00290	Isoleucine biosynthesis
M00432	00290	Leucine biosynthesis
M00016	00300	Lysine biosynthesis
M00028	00220	Ornithine biosynthesis
M00026	00340	Histidine biosynthesis

Table 1 (continued)

KEGG entry name	Pathway ID	Pathway name
M00022	00400	Shikimate pathway
M00023	00400	Tryptophan biosynthesis
M00024	00400	Phenylalanine biosynthesis
M00025	00400	Tyrosine biosynthesis
M00127	00730	Thiamine biosynthesis
M00125	00740	Riboflavin biosynthesis
M00124	00750	Pyridoxal biosynthesis
M00119	00770	Pantothenate biosynthesis
M00572	00780	Pimeloyl-ACP biosynthesis
M00123	00780	Biotin biosynthesis
M00126	00670	Tetrahydrofolate biosynthesis
M00121	00860	Heme biosynthesis
M00550	00053	Ascorbate degradation
M00117	00130	Ubiquinone biosynthesis
M00116	00130	Menaquinone biosynthesis
M00136	00330	GABA biosynthesis

Fig. 2 Subcellular localization of the unique pathogen proteins of different pathways **Fig. 3** Distribution of all the essential proteins of the pathogen with percentage

Electrostatic potentiality and energy minimization

The electrostatic potentiality of the template 1t4b_A and the modeled protein was analyzed for the identification of the energy distribution of the protein and showed in Fig. 6. The comparative energy distribution revealed that the distribution is similar to the target and the model. For better model construction, we applied energy minimization process onto the modeled protein through the GROMOS 96 force field. The force-field energies of the whole structure before and after minimization were -6404.763 and $-16,370.318$ kJ/mol respectively.

Active site exploration and virtual screening

CASTp server predicted different active sites of our desired protein with different volume score and we picked the best large volume as the final active site (Fig. 7). The molecular surface area of the active site was $1940.291 \text{ \AA}^2$.

For the virtual screening, we docked 26 compounds based on the initially targeted compound screened from the BindingBD (Table S5). The top six molecules of best binding energies along with the standard were investigated further for the identification of potential drug candidate (Table 3; Fig. 8). Docking poses with 0.0 RMSD were selected as the final binding energies and interacting amino acid residues are depicted in Fig. 9.

Ligand properties and ADMET data analysis

Different properties of the predicted small molecules (ligand) including drug-likeness, Lipinski's rules, and

Table 2 Six novel therapeutic targets and their basic properties

Sl no.	Uniprot ID	Gene name	Length	Molecular weight (Da)	Metabolic pathway	Subcellular location	PDB ID
1.	Q83MD3	dxr	398	43346.9	C5 isoprenoid biosynthesis	Cytoplasmic	No
2.	P0A6L3	dapA	292	31269.9	Lysine biosynthesis	Cytoplasmic	No
3.	P0A9R1	asd	367	40017.8	Lysine biosynthesis	Cytoplasmic	No
4.	Q83IN8	lysC	449	48517.7	Lysine biosynthesis	Cytoplasmic	No
5.	P0A9M9	pta	714	77172.0	Phosphate acetyltransferase-acetate kinase pathway	Cytoplasmic	No
6.	Q83MH9	thrB	310	33579.5	Threonine biosynthesis	Cytoplasmic	No

Fig. 4 Three-dimensional model of all the targeted proteins. Here, cyan, magenta, blue, yellow, orange and purple color denote the protein, P0A6L3, P0A9M9, Q83MD3, Q83IN8 Q83MH9 and P0A9R1, respectively

enzyme inhibitory activities were assessed for the identification of the best therapeutic candidates (Table 4). All the ADMET properties were also assessed for the selected small molecules and cross-check with different server and tools for the validity of the results. These properties are described in Table 5.

Discussion

This is the first comprehensive subtractive genomics analysis of different metabolic pathways of *Shigella flexneri* 2a for the identification of prospective drug targets through sequential approaches described schematically in Fig. 1. Metabolic pathways are crucial for the survival of the bacteria and hence it serves as a prime target for the development of new therapeutics (Duffield et al. 2010). In this study, we have

reinforced to find out new therapeutic targets that should be novel and potential for getting more striking outputs. As a part of this process, we have tried to figure out unique targets which solely exist in the pathogen only for avoiding any potential side effects.

Initial analysis revealed that the pathogen consists of a wide array of metabolic pathways which are unique to pathogen only and non-homolog as well. Interactive network analyses among the unique pathways' proteins are used for the novel targets' identification and cytoplasmic proteins are the topmost priority among them due to easier purification and assay (Guruprasad et al. 1990; Oany et al. 2014c).

Based on the physiological and chemical properties we prioritized drug targets initially. The molecular weight of the drug targets should be < 100 KDa for securing the appropriate interactions with the therapeutic candidates and we considered this criterion for our selection. The lower theoretical

Fig. 5 Superposition between the template and model. In this figure, template is shown as yellow and the predicted model is shown as blue color. The RMSD of the superposition is 0.146 Å

isoelectric point (pI) and instability index are associated with the stability of the protein in in vitro condition. This is well applicable to the therapeutic target also and we had carefully chosen the targets according to these parameters (Bakheet and Doig 2009). Hydrophobicity of the therapeutic target is higher than the non-therapeutic candidates and that cause the lower polarity of the target (Lange et al. 2000). For this reason, those targets with hydrophobicity > -0.142 were selected for the therapeutic targets. Finally, targets sorting with > 2 *N*-glycosylated amino acids further strengthen our prioritization through securing appropriate folding and assembly of the targets and cause less prone to degradation.

Later in the prioritization steps we found six therapeutic targets. Among them, proteins P0A6L3, P0A9R1 and Q83IN8 encoded by the genes *dapA*, *asd* and *lysC*, respectively, are associated with lysine biosynthesis pathway. Protein Q83MD3 encoded by the gene *dxr* is associated with C5 isoprenoid biosynthesis and Q83MH9 (*thrB*) is responsible for threonine biosynthesis. All these targets are very much vital for the pathogen survival. We further performed an extended literature review and structural feature analysis for gathering insights about the feasibility of these targets and finally the proteins P0A9R1 and Q83MD3 were selected.

Enzyme 1-deoxy-D-xylulose 5-phosphate reductoisomerase (DXR) is the second enzyme of the methylerythritol phosphate (MEP) to isoprenoid biosynthesis pathway. The

enzyme DXR converts 1-deoxy-D-xylulose 5-phosphate (DXP) into the branched compound 2-C-methyl-D-erythritol 4-phosphate (MEP) which requires isomerization and subsequently an NADPH-dependent reduction. This MEP pathway is unique in most of the pathogenic bacteria and apicomplexan parasite like *Plasmodium falciparum* and absent in human, therefore, becoming an attractive target for the development of novel antimalarial and antibacterial compounds (Sangari et al. 2010). Mevalonate (MVA) pathway, an alternative to the MEP, is usually used in many bacteria for the synthesis of isoprenoid and bacteria show a significant plasticity for this pathway through bypassing. As a consequence, resistance to antibiotics is rising (Sakamoto et al. 2003; Sangari et al. 2010).

Aspartate- β -semialdehyde dehydrogenase (ASD) was the second target of our analysis and another potential therapeutic target for the development of new drugs against the aspartate pathway. This enzyme is responsible for the production of many important amino acids as well as very much crucial for the growth and development of the organism (Gerdes et al. 2003; Kobayashi et al. 2003). The absence in the human and wide distribution in many bacteria and plants cause an extended focus for the researcher on this validated drug-target enzyme. For these reasons, we had selected this protein for the further investigation.

Homology model, based on the template 1t4b_A with the high sequence identity of 99.728% secured the model validity and furthermore, superposition RMSD of 0.146 Å is an indicator of the highly valid model. Energy minimization steps also prepare the model ready to dock organization on its large active site pocket.

Petrosamine B, a pyridoacridine alkaloid, isolated from the marine sponge has been used as an inhibitor of the enzyme ASD and works as a weak inhibitor (Carroll et al. 2005). Through focusing this point we had explored the virtual screening for the similar compounds to find-out more potential therapeutic candidates.

The compound 3 (PubChem CID: 11319750) with a binding energy of -9.9 kcal/mol was found to be the most potent inhibitor of the ASD in comparison with the Petrosamine B and two other potent compounds 1 (PubChem CID: 24882646) and 2 (PubChem CID: 23425964) with a binding energy of -7.6 , -10.1 and -10.0 kcal/mol, respectively as the compound 3 possesses the highest enzyme inhibitory and drug-likeness score of 0.72 and 0.2, respectively. There is no violation of the Lipinski's rules for all the compounds and retain the bioavailability of 0.55 as well.

The ADMET profile analysis of all the targeted compounds likewise revealed that the compound 3 was also a potent therapeutic candidate in comparison with others. The intestinal absorption of the compound 3 was 100% and highly water soluble. Blood-brain barrier (BBB) permeability of the compound 3 was 0.2 and the highest permeability

Fig. 6 Electrostatic potential of the proposed target POA9R1 (top structure) and the template 1T4B_A (bottom structure). In the energy distribution, blue color indicates higher energy and red indicates lower energy potentiality. White color indicates energy neutral state

Fig. 7 Active site determination of the targeted protein with energy distribution view

showed by the compound 5 of 0.413. The fraction of the unbound drug is higher for the compound 4 of 0.002 while the compound 3 had 0.143, indicating the high efficiency of the compounds. Total clearance was higher for the

compound 3 of 0.708 and highest of 0.916 for the compound 5 and also showed no AMES toxicity and carcinogenicity.

Through the comparative analysis, finally, we concluded that the compound 3 would be the best possible candidate

Table 3 Docking analysis of the screened compounds

Com- pound no.	ID	RMSD	Binding energy (docking) kcal/mol
1.	PubChem CID: 24882646	0.0	− 10.1
2.	PubChem CID: 23425964	0.0	− 10.0
3.	PubChem CID: 11319750	0.0	− 9.9
4.	PubChem CID: 25251877	0.0	− 9.6
5.	PubChem CID: 188755	0.0	− 9.6
6.	PubChem CID: 11044976	0.0	− 9.2
7.	Petrosamine-B (PubChem CID:21778132)	0.0	− 7.6

as an inhibitor of the ASD because of its relative higher ADMET profile as well as highest enzyme inhibitory and drug-likeness scores.

Conclusion

Shigellosis now has become a global concern for the developed as well as poor countries throughout the world and causes millions of deaths every year. Through the ever-rising resistance patterns of the pathogen, *Shigella* has become a major threat to us. So, the development of better targets and

Fig. 8 Selected small molecules (ligands) for drug designing. The identities (PubChem CID) of these small molecules are 24882646, 23425964, 11319750, 25251877, 188755, 11044976 and 21778132 (Petrosamine-B), respectively, according to the figure numbers

Fig. 9 Molecular docking interaction analysis. **a** All the ligands are docked into the active site pocket of the protein and different ligands are shown by different colors. Ligands are colored in red, green, blue, yellow, magentas, cyan and orange accordance to the order of the

ligands in **b**. **b** Interacting amino acid residues of the protein with different ligands. The interaction descriptions are shown graphically in the bottom of **b**

Table 4 Ligand properties analysis of the selected small molecules

Ligand properties	PubChem CID: 24882646	PubChem CID: 23425964	PubChem CID: 11319750	PubChem CID: 25251877	PubChem CID: 188755	PubChem CID: 11044976	Petrosamine-B (PubChem CID:21778132)
Molecular formula	$C_{21}H_{16}BrN_3O_2$	$C_{21}H_{17}BrN_3O_2$	$C_{21}H_{17}BrN_3O_2^+$	$C_{19}H_{11}NO_2$	$C_{21}H_{19}BrN_3O^+$	$C_{17}H_{14}N_4O_2$	$C_{21}H_{18}BrN_3O_2^{+2}$
Molecular weight (g/mol)	422.27464	423.28258	423.28258	285.29614	409.29906	306.31866	424.29052
LogP	2.53	2.46	1.99	4.86	2.39	1.49	2.00
Hydrogen bond donor	0	1	0	0	0	0	1
Hydrogen bond acceptor	4	4	4	3	3	6	3
Rotatable bond	0	0	0	0	0	2	0
Total polar surface area (TPSA)	57.95	55.12	51.97	43.10	34.90	68.22	53.21
Lipinski's rule violations	0	0	0	0	0	0	0
Enzyme inhibitor	0.52	0.52	0.72	0.26	0.19	0.28	0.72
Aqueous solubility LogS	-3.97	-3.97	-4.11	-7.67	-3.99	-5.64	-4.11
Drug-likeness model score	-0.36	-0.23	0.20	-1.10	0.04	-0.55	-0.06
Bioavailability score	0.55	0.55	0.55	0.55	0.55	0.55	0.55
Binding energy (docking) kcal/mol	-10.1	-10.0	-9.9	-9.6	-9.6	-9.2	-7.6

Table 5 ADME/T properties analysis of the selected small molecules

Ligand properties	PubChem CID: 24882646	PubChem CID: 23425964	PubChem CID: 11319750	PubChem CID: 25251877	PubChem CID: 188755	PubChem CID: 11044976	Petrosamine B (PubChem CID:21778132)
Absorption							
Blood brain barrier	0.8809	0.7620	0.9545	0.9956	0.9246	0.9656	0.9382
Water solubility	– 4.988	– 5.339	– 5.176	– 4.869	– 4.051	– 3.431	– 4.424
Caco ₂ permeability	1.268	1.248	1.264	1.4	1.241	1.023	1.268
Intestinal absorption (human)	100	95.274	100	97.825	100	100	100
Skin permeability	– 3.277	– 3.377	– 3.294	– 2.735	– 3.322	– 3.333	– 3.197
P-Glycoprotein substrate	Yes	Yes	Yes	Yes	Yes	Yes	Yes
P-Glycoprotein I inhibitor	Yes	Yes	Yes	No	Yes	No	Yes
Distribution							
BBB permeability	0.148	0.183	0.2	0.317	0.413	– 0.405	0.2
CNS permeability	– 1.697	– 1.532	– 1.704	– 0.862	– 2.096	– 2.259	– 1.704
Fraction unbound (human)	0.152	0.139	0.143	0.002	0.273	0.317	0.238
Metabolism							
CYP450 1A2 inhibitor	Yes	Yes	Yes	Yes	Yes	Yes	Yes
CYP2C19 inhibitor	No	No	No	Yes	No	No	No
CYP3A4 substrate	Yes	Yes	Yes	Yes	Yes	Yes	No
Excretion							
Total clearance	0.675	0.7	0.708	0.727	0.916	0.414	0.887
Renal OCT2 substrate	No	No	No	No	No	No	No
Toxicity							
AMES toxicity	Yes	Yes	No	No	No	Yes	Yes
Max. tolerated dose (human)	0.218	0.205	0.223	1.049	0.119	0.331	0.141
Carcinogens	No	No	No	No	No	No	No
Acute oral toxicity	0.6124	0.6220	0.6357	0.7867	0.6176	0.7066	0.6736

therapeutic candidates are inevitable for the upcoming centuries. Present comprehensive whole genomics and proteomics-based study of the metabolic pathways' of the pathogenic strain *S. flexneri* 2a gives us a pedal stands to think about its potential therapeutics. Finally, we strongly believe that the identified therapeutic targets and its potential inhibitors would contribute to treating shigellosis in a proficient manner in in vitro and suggesting further study about its potency.

Acknowledgements Authors' declares no acknowledgements for this work.

Funding No funding was received

Compliance with ethical standards

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- Agüero F, Al-Lazikani B, Aslett M, Berriman M, Buckner FS, Campbell RK, Carmona S, Carruthers IM, Chan AE, Chen F (2008) Genomic-scale prioritization of drug targets: the TDR targets database. *Nat Rev Drug Discov* 7:900–907
- Allsop AE (1998) Bacterial genome sequencing and drug discovery. *Curr Opin Biotechnol* 9:637–642
- Amineni U, Pradhan D, Marisetty H (2010) In silico identification of common putative drug targets in *Leptospira interrogans*. *J Chem Biol* 3:165–173
- Arnold K, Bordoli L, Kopp J, Schwede T (2006) The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics* 2:195–201
- Bairoch A, Apweiler R (2000) The SWISS-PROT protein sequence database and its supplement TrEMBL in 2000. *Nucleic Acids Res* 28:45–48
- Bakheet TM, Doig AJ (2009) Properties and identification of human protein drug targets. *Bioinformatics* 25:451–457
- Barh D, Kumar A (2009) In silico identification of candidate drug and vaccine targets from various pathways in *Neisseria gonorrhoeae*. *In Silico Biol* 9:225–231

- Bennish ML, Wojtyniak BJ (1991) Mortality due to shigellosis: community and hospital data. *Rev Infect Dis* 13:245–251
- Bolton EE, Wang Y, Thiessen PA, Bryant SH (2008) PubChem: integrated platform of small molecules and biological activities. *Annu Rep Comput Chem* 4:217–241
- Carroll AR, Ngo A, Quinn RJ, Redburn J, Hooper JN (2005) Petrosamine B, an inhibitor of the helicobacter pylori enzyme aspartyl semialdehyde dehydrogenase from the Australian Sponge *Oceanapia* sp. *J Nat Prod* 68:804–806
- Chan JN, Nislow C, Emili A (2010) Recent advances and method development for drug target identification. *Trends Pharmacol Sci* 31:82–88
- Cheng F, Li W, Zhou Y, Shen J, Wu Z, Liu G, Lee PW, Tang Y (2012) admetSAR: a comprehensive source and free tool for assessment of chemical ADMET properties. ACS Publications, New York, pp 3099–3105
- Damte D, Suh JW, Lee SJ, Yohannes SB, Hossain MA, Park SC (2013) Putative drug and vaccine target protein identification using comparative genomic analysis of KEGG annotated metabolic pathways of *Mycoplasma hyopneumoniae*. *Genomics* 102:47–56
- Davies M, Dedman N, Hersey A, Papadatos G, Hall MD, Cucurull-Sanchez L, Jeffrey P, Hasan S, Eddershaw PJ, Overington JP (2015) ADME SARfari: comparative genomics of drug metabolizing systems. *Bioinformatics* 31:1695–1697
- Deshpande N, Address KJ, Bluhm WF, Merino-Ott JC, Townsend-Merino W, Zhang Q, Knezevich C, Xie L, Chen L, Feng Z (2005) The RCSB Protein Data Bank: a redesigned query system and relational database based on the mmCIF schema. *Nucleic Acids Res* 33:D233–D237
- Duffield M, Cooper I, McAlister E, Bayliss M, Ford D, Oyston P (2010) Predicting conserved essential genes in bacteria: in silico identification of putative drug targets. *Mol BioSyst* 6:2482–2489
- Dundas J, Ouyang Z, Tseng J, Binkowski A, Turpaz Y, Liang J (2006) CASTp: computed atlas of surface topography of proteins with structural and topographical mapping of functionally annotated residues. *Nucleic Acids Res* 34:W116–W118
- DuPont HL, Hornick RB, Dawkins AT, Snyder MJ, Formal SB (1969) The response of man to virulent *Shigella flexneri* 2a. *J Infect Dis* 119:296–299
- DuPont DL, Levine MM, Hornick RB, Formal SB (1989) Inoculum size in shigellosis and implications for expected mode of transmission. *J Infect Dis* 159:1126–1128
- Dutta S, Rajendran K, Roy S, Chatterjee A, Dutta P, Nair G, Bhat-tacharya S, Yoshida S (2002) Shifting serotypes, plasmid profile analysis and antimicrobial resistance pattern of shigellae strains isolated from Kolkata, India during 1995–2000. *Epidemiol Infect* 129:235–243
- Eichner ER, Gangarosa EJ, Goldsby JB (1968) The current status of shigellosis in the United States. *Am J Public Health Nations Health* 58:753–763
- Fasano A, Noriega FR, Maneval D Jr, Chanasongcram S, Russell R, Guandalini S, Levine MM (1995) *Shigella* enterotoxin 1: an enterotoxin of *Shigella flexneri* 2a active in rabbit small intestine in vivo and in vitro. *J Clin Invest* 95:2853
- Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P, Von Mering C (2012) STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 41:D808–D815
- Galperin MY, Koonin EV (1999) Searching for drug targets in microbial genomes. *Curr Opin Biotechnol* 10:571–578
- Gao Z, Li H, Zhang H, Liu X, Kang L, Luo X, Zhu W, Chen K, Wang X, Jiang H (2008) PDTD: a web-accessible protein database for drug target identification. *BMC Bioinform* 9:104
- Gardy JL, Laird MR, Chen F, Rey S, Walsh C, Ester M, Brinkman FS (2004) PSORTb v. 2.0: expanded prediction of bacterial protein subcellular localization and insights gained from comparative proteome analysis. *Bioinformatics* 21:617–623
- Garza-Mayers AC, Miller KA, Russo BC, Nagda DV, Goldberg MB (2015) *Shigella flexneri* regulation of ARF6 activation during bacterial entry via an IpgD-mediated positive feedback loop. *MBio* 6:e02584–e02614
- Gasteiger E, Hoogland C, Gattiker A, Duvaud SE, Wilkins MR, Appel RD, Bairoch A (2005) Protein identification and analysis tools on the ExPASy server. Humana Press, New York, pp 571–607
- Gerdes S, Scholle M, Campbell J, Balazsi G, Ravasz E, Daugherty M, Somera A, Kyrpides N, Anderson I, Gelfand M (2003) Experimental determination and system level analysis of essential genes in *Escherichia coli* MG1655. *J Bacteriol* 185:5673–5684
- Gupta A, Polyak CS, Bishop RD, Sobel J, Mintz ED (2004) Laboratory-confirmed shigellosis in the United States, 1989–2002: epidemiologic trends and patterns. *Clin Infect Dis* 38:1372–1377
- Guruprasad K, Reddy BB, Pandit MW (1990) Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence. *Protein Eng Des Sel* 4:155–161
- Hale TL, Keusch GT (1996) *Shigella*
- Hasan MN, Oany AR, Hirashima A (2016) Comparative structure analysis of tyramine- β -hydroxylase from fruit fly and ADME/T-based profiling of 1-arylimidazole-2 (3H)-thiones as potential inhibitors. *Int J Anal Bio-Sci* 4:37–45
- Irwin BK, Shoichet BK (2005) ZINC—a free database of commercially available compounds for virtual screening. *J Chem Inf Model* 45:177–182
- Jennison AV, Verma NK (2004) *Shigella flexneri* infection: pathogenesis and vaccine development. *FEMS Microbiol Rev* 28:43–58
- Jin Q, Yuan Z, Xu J, Wang Y, Shen Y, Lu W, Wang J, Liu H, Yang J, Yang F (2002) Genome sequence of *Shigella flexneri* 2a: insights into pathogenicity through comparison with genomes of *Escherichia coli* K12 and O157. *Nucleic Acids Res* 30:4432–4441
- Johnson M, Zaretskaya I, Raytselis Y, Merezukh Y, McGinnis S, Madden TL (2008) NCBI BLAST: a better web interface. *Nucleic Acids Res* 36:W5–W9
- Kanehisa M, Goto S, Hattori M, Aoki-Kinoshita KF, Itoh M, Kawashima S, Katayama T, Araki M, Hirakawa M (2006) From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Res* 34:D354–D357
- Kanehisa M, Goto S, Furumichi M, Tanabe M, Hirakawa M (2009) KEGG for representation and analysis of molecular networks involving diseases and drugs. *Nucleic Acids Res* 38:D355–D360
- Kobayashi K, Ehrlich SD, Albertini A, Amati G, Andersen K, Arnaud M, Asai K, Ashikaga S, Aymerich S, Bessieres P (2003) Essential *Bacillus subtilis* genes. *Proc Natl Acad Sci* 100:4678–4683
- Koonin EV, Tatusov RL, Galperin MY (1998) Beyond complete genomes: from sequence to structure and function. *Curr Opin Struct Biol* 8:355–363
- Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, Swerdlow DL, Sansonetti PJ, Adak G, Levine M (1999) Global burden of *Shigella* infections: implications for vaccine development and implementation of control strategies. *Bull World Health Organ* 77:651
- Lange BM, Rujan T, Martin W, Croteau R (2000) Isoprenoid biosynthesis: the evolution of two ancient and distinct pathways across genomes. *Proc Natl Acad Sci* 97:13172–13177
- Laskowski RA, Rullmann JAC, MacArthur MW, Kaptein R, Thornton JM (1996) AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. *J Biomol NMR* 8:477–486
- Lipinski CA (2004) Lead-and drug-like compounds: the rule-of-five revolution. *Drug Discov Today Technol* 1:337–341
- Liu T, Lin Y, Wen X, Jorissen RN, Gilson MK (2006) BindingDB: a web-accessible database of experimentally determined protein–ligand binding affinities. *Nucleic Acids Res* 35:D198–D201

- Magrane M (2011) UniProt Knowledgebase: a hub of integrated protein data. Database
- Mei Y, Liu H, Xu J (1989) Cloning and application of genus specific DNA probes for *Shigella*. Chin J Epidemiol 10:167–170
- Oany SAI, Ahmad SAI, Kibria KK, Hossain MU, Jyoti TP (2014a) A hypothetical protein of *Alteromonas macleodii* AltDE1 (amad1_06475) predicted to be a cold-shock protein with RNA chaperone activity. Gene Regul Syst Biol 8:141–147
- Oany AR, Emran AA, Jyoti TP (2014b) Design of an epitope-based peptide vaccine against spike protein of human coronavirus: an in silico approach. Drug Des Dev Ther 8:1139–1149
- Oany AR, Jyoti TP, Ahmad SAI (2014c) An in silico approach for characterization of an aminoglycoside antibiotic-resistant methyltransferase protein from *Pyrococcus furiosus* (DSM 3638). Bioinform Biol Insights 8:65–72
- Oany AR, Pervin T, Mia M, Hossain M, Shahnaiz M, Mahmud S, Kibria K (2017) Vaccinomics approach for designing potential peptide vaccine by targeting *Shigella* spp. serine protease autotransporter subfamily protein SigA. J Immunol Res
- O'Boyle NM, Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR (2011) Open Babel: an open chemical toolbox. J Cheminf 3:33
- Oprea TI (2000) Property distribution of drug-related chemical databases. J Comput Aided Mol Des 14:251–264
- Perdomo O, Cavaillon J, Huerre M, Ohayon H, Gounon P, Sansonetti P (1994) Acute inflammation causes epithelial invasion and mucosal destruction in experimental shigellosis. J Exp Med 180:1307–1319
- Peter-Katalinić J (2005) Methods in enzymology: *O*-glycosylation of proteins. Methods Enzymol 405:139–171
- Philpott DJ, Edgeworth JD, Sansonetti PJ (2000) The pathogenesis of *Shigella flexneri* infection: lessons from in vitro and in vivo studies. Philos Trans R Soc Lond B Biol Sci 355:575–586
- Pieper U, Eswar N, Davis FP, Braberg H, Madhusudhan MS, Rossi A, Marti-Renom M, Karchin R, Webb BM, Eramian D (2006) MODBASE: a database of annotated comparative protein structure models and associated resources. Nucleic Acids Res 34:D291–D295
- Rahman MA, Noore MS, Hasan MA, Ullah MR, Rahman MH, Hossain MA, Ali Y, Islam MS (2014) Identification of potential drug targets by subtractive genome analysis of *Bacillus anthracis* A0248: an in silico approach. Comput Biol Chem 52:66–72
- Sakamoto Y, Furukawa S, Ogihara H, Yamasaki M (2003) Fosmidomycin resistance in adenylate cyclase deficient (*cya*) mutants of *Escherichia coli*. Biosci Biotechnol Biochem 67:2030–2033
- Šali A, Potterton L, Yuan F, van Vlijmen H, Karplus M (1995) Evaluation of comparative protein modeling by MODELLER. Proteins Struct Funct Bioinform 23:318–326
- Sander T (2001) OSIRIS property explorer. Org Chem Portal
- Sangari FJ, Pérez-Gil J, Carretero-Paulet L, García-Lobo JM, Rodríguez-Concepción M (2010) A new family of enzymes catalyzing the first committed step of the methylerythritol 4-phosphate (MEP) pathway for isoprenoid biosynthesis in bacteria. Proc Natl Acad Sci 107:14081–14086
- Schrödinger L (2012) QikProp, version 3.5. New York, NY
- Schrödinger LLC (2015) The PyMOL Molecular Graphics System, version 1.8
- Scott WR, Hünenberger PH, Tironi IG, Mark AE, Billeter SR, Fennen J, Torda AE, Huber T, Krüger P, van Gunsteren WF (1999) The GROMOS biomolecular simulation program package. J Phys Chem A 103:3596–3607
- Small P, Blankenhorn D, Welty D, Zinser E, Slonczewski JL (1994) Acid and base resistance in *Escherichia coli* and *Shigella flexneri*: role of *rpoS* and growth pH. J Bacteriol 176:1729–1737
- Stumm G, Russ A, Nehls M (2002) Deductive genomics. Am J Pharmacogenom 2:263–271
- Talukder KA, Khajanchi BK, Islam MA, Dutta DK, Islam Z, Safa A, Khan G, Alam K, Hossain M, Malla S (2004) Genetic relatedness of ciprofloxacin-resistant *Shigella dysenteriae* type 1 strains isolated in south Asia. J Antimicrob Chemother 54:730–734
- Talukder KA, Khajanchi BK, Islam MA, Islam Z, Dutta DK, Rahman M, Watanabe H, Nair GB, Sack DA (2006) Fluoroquinolone resistance linked to both *gyrA* and *parC* mutations in the quinolone resistance-determining region of *Shigella dysenteriae* type 1. Curr Microbiol 52:108–111
- Tatusov RL, Koonin EV, Lipman DJ (1997) A genomic perspective on protein families. Science 278:631–637
- Trott O, Olson AJ (2010) AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. J Comput Chem 31:455–461
- Unni S, Huang Y, Hanson RM, Tobias M, Krishnan S, Li WW, Nielsen JE, Baker NA (2011) Web servers and services for electrostatics calculations with APBS and PDB2PQR. J Comput Chem 32:1488–1491
- US Department of Health, Education, and Welfare, Public Health Service, National Communicable Disease Center (1968) *Shigella* surveillance report no. 15
- Wirth M, Zoete V, Michielin O, Sauer WH (2012) SwissBioisostere: a database of molecular replacements for ligand design. Nucleic Acids Res 41:D1137–D1143
- Wishart DS, Knox C, Guo AC, Cheng D, Shrivastava S, Tzur D, Gautam B, Hassanali M (2007) DrugBank: a knowledgebase for drugs, drug actions and drug targets. Nucleic Acids Res 36:D901–D906
- Yu CS, Lin CJ, Hwang JK (2004) Predicting subcellular localization of proteins for Gram-negative bacteria by support vector machines based on *n*-peptide compositions. Protein Sci 13:1402–1406

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.