Published in final edited form as:

Connect Tissue Res. 2018 September ; 59(5): 437–446. doi:10.1080/03008207.2018.1485666.

Partial-width injuries of the rat rotator cuff heal with fibrosis

Elisabeth A. Lemmon^a, Ryan C. Locke^b, Adrianna K. Szostek^a, Elahe Ganji^c, and Megan L. Killian^b

^aDepartment of Animal and Food Sciences and Biomedical Engineering, University of Delaware, Newark, DE, USA

^bDepartment of Biomedical Engineering, University of Delaware, Newark, DE, USA

^cDepartment of Mechanical Engineering and Biomedical Engineering, University of Delaware, Newark, DE, USA

Abstract

Purpose: Identify the healing outcomes following a partial-width, full-thickness injury to the rotator cuff tendon-bone attachment and establish if the adult attachment can regenerate the morphology of the healthy attachment.

Hypothesis: We hypothesized that a partial-width injury to the attachment would heal via fibrosis and bone remodeling, resulting in increased cellularity and extra-cellular matrix deposition, reduced bone volume (BV), osteoclast presence, and decreased collagen organization compared to shams.

Materials and Methods: A partial-width injury was made using a biopsy punch at the center one-third of the rat infraspinatus attachment. Contralateral limbs underwent a sham operation. Rats were sacrificed at 3 and 8 weeks after injury for analyses. Analyses performed at each time point included cellularity (Hematoxylin & Eosin), ECM deposition (Masson's Trichrome), BV (micro-computed tomography; microCT), osteoclast activity (Tartrate Resistant Acid Phosphatase; TRAP), and collagen fibril organization (Picrosirius Red). Injured and sham shoulders were compared at both 3 and 8 weeks using paired, two-way ANOVAs with repeated measures (Sidak's correction for multiple comparisons).

Results: Cellularity and ECM deposition increased at both 3 and 8 weeks compared to sham contralateral attachments. BV decreased and osteoclast presence increased at both 3 and 8 weeks compared to sham contralateral limbs. Collagen fibril organization was reduced at 3 weeks after injury compared to 3-week sham attachments.

Conclusions: These findings suggest that a partial-width injury to the rotator cuff attachment does not fully regenerate the native structure of the healthy attachment. The injury model healed via scar-like fibrosis and did not propagate into a full-width tear after 8 weeks of healing.

Correspondence to: Megan L. Killian.

Disclosure statement

No potential conflict of interest was reported by the authors.

Keywords

Collagen organization; osteoclast; rotator cuff healing; tendon-bone attachment; tendon injury

Introduction

Rotator cuff tears are a common orthopedic injury, and rates of rotator cuff injuries and repairs have risen drastically in the United States over the past decade (1,2). Depending on the location and severity of the tear, clinical recommendations for surgery and physical therapy vary substantially (3,4). For partial-width tears, conservative treatments (e.g., physical therapy) are encouraged before surgical intervention (4–6). The success of rotator cuff repair depends on the health of the cuff at time of repair (7–11) as well as the restoration of the morphology and strength of the attachment (12–17). Following full-width tears, repair of the torn attachment rarely results in full regeneration of the native morphology and structure of the attachment (7,15,18–20). In small animal models of cuff healing, a fibrous scar tissue, characterized by reduced collagen organization, increased vascularization, and increased cell density, are often observed (15,19,20). Loss of the structural integrity of the attachment is considered one of the primary causes of high rates of re-injury after rotator cuff repair (7,20). However, animal models used to study rotator cuff healing have primarily relied on full-width injuries that require surgical reattachment of cuff tendons to their bony footprint for structural reintegration. Healing of cuff tendons during early stages of tear propagation, such as in the case of partial-width injuries, has been investigated *in vivo* using animal models (20,23). However, the healing of partial-width injuries focused at the attachment, without surgical repair and augmentation, has only recently been explored (22).

In the present study, we aimed to develop and validate a new model of rotator cuff injury to investigate the natural healing process of the attachment. Using a rat model of partial-width, full-thickness injury at the attachment, we hypothesized that, although the attachment would remain partially intact, it would not heal with the same structural quality as an intact, uninjured attachment. We tested this hypothesis by assessing the fibrotic response, extra-cellular matrix (ECM) deposition, bone remodeling, and collagen organization of the healing attachment.

Materials and methods

Animal model

Adult Sprague Dawley rats ($N=8$ females, $N=8$ males for *in vivo* healing; ~200–250 g) and adult Long Evans rats ($N=20$ female dams, for *ex vivo* validation at time zero) were used in accordance with the University of Delaware Institutional Animal Care and Use Committee approval. Rats underwent a surgical procedure under anesthesia (isoflurane carried by 1% oxygen) to model a partial-width rotator cuff injury at the center of the infraspinatus (IS) attachment (Figure 1A–C) (24). The IS tendon was exposed and the forearm was internally rotated. A 0.3 mm diameter biopsy punch (Robbins, Chatham, NJ, USA) was placed in the center 1/3 of the tendon width spanning the attachment/bone, and the punch permeated the

fibrocartilage, tendon, and cortical bone (Supplemental Figure 1A–D). The injured shoulder was randomized between rats, and the contralateral shoulder underwent a sham operation to mimic the procedure without the biopsy punch permeation. The surgical site was closed using 5–0 Vicryl suture (Ethicon Inc., Somerville, New Jersey), and rats were given bupivacaine hydrochloride (0.05 mg/kg) as analgesia. Rats were separated into two groups: 3-week healing ($N = 8$; 4 females and 4 males) and 8-week healing ($N = 8$, 4 females and 4 males). Rats were sacrificed with carbon dioxide asphyxiation and thoracotomy. The 3 and 8 week time points were chosen to evaluate the proliferation (3 weeks) and remodeling (8 weeks) phases of tendon-bone healing.

Biomechanics

Validation of the model was evaluated *ex vivo* using biomechanical testing ($N = 18$). Adult, female Long Evans rats, used as breeders in an unrelated study (Department of Psychology and Brain Sciences), were sacrificed after weaning of their first-born litters. Dams were not treated with any pharmacological drugs in previous studies. Superficial (through soft tissue/fibro-cartilage) and deep (permeating the cortical bone) injury defects ($N = 8–10$ shoulders each) were made evaluating the biomechanical properties of the defect at time zero and establish whether a superficial or deep defect result in comparable biomechanical outcomes. Rats were randomized as to which limb received the defect, and the contralateral limb underwent a sham operation to mimic the procedure, similar to *in vivo* approaches. After performing the punch, the IS tendon-to-bone attachments were dissected from the surrounding musculature and bone. The IS muscle belly was carefully removed from the IS tendon using a #11 scalpel blade. The IS tendon attachment at the proximal humerus was left intact for mechanical testing. Uniaxial tensile tests were performed (Instron 5943, Norwood, MA) and the cross-sectional area of the attachments were measured using microCT. To prevent failure at the growth plate during testing, a hole was drilled in the humeral diaphysis and then steel wire was passed through the hole and wrapped posterior to anterior around the humeral head. Tendons were tested in a PBS bath using custom fixtures to ensure uniaxial loading (Supplemental Figure 2A–C). A 0.01N tare load was applied and the test configuration was imaged at the start of the test to measure gauge length of the sample, followed by five preconditioning cycles to 5% strain at a rate of 0.2 %/s. Following preconditioning, samples were held for 30 s at initial gauge length and then loaded to failure at 0.2 %/s. Load/displacement data were recorded throughout the test (preconditioning and load-to-failure) and data were converted to stress/strain data based on the initial cross-sectional area from micro-computed tomography (microCT) (stress) and the gauge length from the start of the test (strain). Stiffness (N/mm) and ultimate load (N) were calculated from load/displacement curves, and Young's modulus (MPa) and ultimate stress (MPa) were calculated from stress/strain curves, with stiffness and Young's modulus as the slope of the respective curves in the linear region and ultimate load/stress as the maximum load/stress of the respective curves. Data were calculated using MATLAB (MathWorks, Natick, MA, USA).

Micro-computed tomography

We validated our injury model via microCT *ex vivo* to ensure the correct location of the punch at the attachment at time of injury (Supplemental Figure 3A–F). The shoulder

complexes of female Long Evans rats ($N=2$ for each group; previously sacrificed) were used within 2 hours of sacrifice and unilateral shoulder complexes were exposed to perform partial-width injuries with the biopsy punch at superficial (through soft tissue/fibrocartilage) and deep (permeating the cortical bone) intervals. For both validation and *in vivo* healing studies at 3- and 8-week healing, the shoulders were carefully removed to isolate the IS tendons and corresponding muscles. Humeri were cut at the mid diaphysis distal to the deltoid tuberosity using bone shears. Shoulders were fixed in their anatomical position in 4% paraformaldehyde ($N=2$ each for control, superficial, and deep injuries for validation studies; $N=8$ each for 3- and 8-week healing time points). Shoulders were scanned in air using microCT (Scanco μ CT35; 20- μ m voxel size, 45 kV, cone beam, 177 μ A, 800-ms integration time). The tendon-bone attachments were analyzed for structural quality based on three-dimensional reconstructions produced in OsiriX DICOM Viewer (v8.0.2., Pixmeo, Switzerland) and quantitatively assessed using Scanco software. Bone morphometric properties were quantified for the (1) entire humeral head and (2) the injury region using total volume (TV), bone volume (BV), and bone volume/total volume ratio (BV/TV) (Supplemental Figure 4A–B). The humeral head measurements included the humeral head proximal to the growth plate, trabecular bone, and cortical bone. The injury region measurements comprised the IS tendon-bone region proximal to the growth plate, including both the trabecular and cortical bone (Scanco, Switzerland).

Histology

After microCT, shoulders were decalcified in 14% EDTA (Sigma-Aldrich, St. Louis, MO, USA) and processed for paraffin sectioning. Attachments were sectioned at 6- μ m thickness and stained using Hematoxylin & Eosin (H&E) for cellular density, Picrosirius Red for collagen organization, Masson's Trichrome for fibrosis and ECM localization, and Tartrate Resistant Acid Phosphatase (TRAP) for osteoclast staining. Stained sections were imaged using an epi-fluorescent microscope (Axio.Observer. Z1, Carl Zeiss, Thornwood, NY). Cell density within the attachment was measured using a custom MATLAB code on a similarly sized region of interest between samples (MATLAB, Supplemental Figure 5A–B') (25). Sections stained with Picrosirius Red were imaged using circular polarized light microscopy, and the deviation of aligned collagen fibrils was evaluated using a custom MATLAB code (Supplemental Figure 6A–B) (26). TRAP-stained sections were analyzed using Osteomeasure software (Ostometrics, Decatur, GA, USA) to quantify osteoclast surface area at the injury site (Oc.S.) relative to the total bone surface area (B.S.).

Statistical analyses

All statistical comparisons were performed using Prism (v7.0, Graphpad, La Jolla, California, USA). To determine if sex differences influenced our quantitative microCT and histological outcomes, an ordinary three-way ANOVA was performed (injury, sex, and time point) with a Tukey multiple comparisons test (Supplemental Figure 7A–G). If sex had very little impact we consolidated the data for comparisons using a two-way ANOVA. Within animal comparisons for humeral head BV/TV, injury region BV/TV, cell density, Picrosirius Red circular standard deviations, and Oc.S./B.S were analyzed using two-way ANOVAs (3 vs. 8 weeks) with repeated measures (sham vs. injured) and Sidak's correction for multiple comparisons. All quantitative data are presented as mean $\pm$ 95% confidence intervals (CI).

Results

Previous studies have illustrated impaired mechanical properties of partial and full-width rotator cuff tears after injury as well as after injury with repair (19,20). Based on validation biomechanical results, we performed the deep injury *in vivo* instead of the superficial injury, as the deep injury resulted in significant decreases in stiffness, Young's modulus, ultimate load, and ultimate stress compared to the uninjured control (Supplemental Figure 1). The superficial injury did not lead to significant differences in biomechanical properties compared to uninjured controls (Supplemental Figure 1). In this study, we did not find significant differences in sex within groups for the outcomes assessed, and therefore we consolidated the data for comparisons between injury groups and time points using a two-way ANOVA.

Gross observations

At the time of dissection for both the 3- and 8-week time points, the injured attachments were intact and none of the injuries had developed into full-width tears. No obvious morphometric differences were observed between the injured and sham muscle bellies or humeral heads. Gross examination of the healed injury site indicated increased fibrosis of the fascia surrounding IS tendon-bone attachments.

Micro-computed tomography

Similar to our validation results, which showed that the deep injury permeated the cortical bone at time zero, the *in vivo* deep injury permeated the cortical and trabecular bone at the IS attachment, which did not return to its normal morphology by 8-week post-injury (Figure 2A-F). As visualized in microCT reconstructions, 3-week injured attachments showed reduced amount of bone at the injury site; by 8-week post-injury, the defect began to fill in with mineral, although this remodeling was incomplete (Figure 2C, F). Quantitatively, humeral head BV/TV was significantly lower at both 3 and 8 weeks for the injured groups compared to the sham groups (combined male and female groups, Figure 2G). In addition, BV/TV increased with age of the rat for both male and female rats, regardless of injury group (Figure 2G).

The injury region had significantly reduced BV/TV at 3-week post-injury compared to the 8-week injured group (Figure 2H). There were no significant differences in injury region BV/TV between the 3- and 8-week sham (Figure 2H). Measurements for both the humeral head TV of the attachments as well as injury region TV were not significantly different across all groups (Supplemental Figure 2C–D).

Histology

Qualitatively, the injured attachments showed increased ECM deposition and decreased collagen organization, as seen using histological imaging. At 3-week post-injury, increased ECM deposition (shown in red + staining) was apparent compared to sham attachments (Figure 3B-B'). At 8 weeks, the injured attachments had increased evidence of fibrovascular scar tissue in addition to fatty infiltration of the tendon compared to sham attachments

(Figure 3D–D'). Collagen production/remodeling was apparent, indicated by the blue + stain, at both 3 and 8 weeks after injury (Figure 3B' and 3D').

At 3-week post-injury, the attachments appeared unorganized and/or had non-existent organized collagen at the injury site (Figure 3A'). Sham attachments at both 3 and 8 weeks had organized collagen (yellow) at the tendon-bone attachment, with a clear transitioned in organization between tendon and bone (Figures 3A–C). At 8-week post-injury, attachments demonstrated evidence of reorganization of collagen at the injury site compared to the 3-week injured attachments (Figure 3C'). The 3-week injured attachments had, quantitatively, decreased collagen organization compared to sham attachments (Figure 3E).

TRAP-positive osteoclasts were prevalent at the injured attachment, indicating an increase in bone remodeling, at both 3 and 8 weeks (Figure 4A–D'). Injured attachments had increased Oc.S./B.S. at 3 and 8 weeks compared to the contralateral sham attachments (Figure 4F). Oc.S./B.S. was significantly higher in the injured 3-week attachments compared to the injured 8-week attachments (Figure 4F). There was no significant difference in Oc.S./B.S. for 3-week sham attachments compared to 8-wk injured attachments. The overall cell density was significantly higher for both time points at the injured tendon-bone attachments compared to the sham attachments (Figure 4E).

Discussion

Overview

The structure of the healthy tendon-bone attachment is ideal for its function to transmit forces generated by muscle to bone (24,26–28). Although the attachment resists failure, injuries commonly occur near the attachment within rotator cuff tendons (7,18,29–31). Rotator cuff tears may necessitate surgical repair of the tendon back to bone, however conservative treatments can also be effective in re-establishing cuff strength and health (3,32–36). In the present study, we investigated cuff healing using a new model of partial-width injury in the rat.

Here, we used this newly established injury model of a partial-width rotator cuff injury to quantitatively and qualitatively investigate the morphological properties of the healing attachment at 3 and 8 weeks post-injury.

Study significance

In this study, we showed that partial-width, full-thickness injuries led to substantial deficits in cellular and structural composition, including impaired collagen alignment, increased osteoclast activity, and decreased BV compared to sham-operated attachments. These data support that healing of the mature tendon-bone attachment is scar-mediated, coinciding with previous work (22). In our study, short-term following injury (3 weeks) led to decreased collagen organization compared to uninjured groups, yet the attachment was able to somewhat overcome this deficit in the longer term (i.e., by 8-week post-injury). This suggests that the attachment continues to remodel and re-organize the injury site. Notably, the biopsy punch used to create the injury permeated the cortical bone, which may have resulted in increased osteoclast activity at the injury site, indicative of a bone marrow-

derived healing response (22,37). The observed increase in bone resorption, driven by osteoclast activity, likely led to reduced BV/TV in the injured groups at 3 and 8 weeks post-injury. Additionally, the injury may have led to reduced loading, due to pain or discomfort, resulting in a reduction in muscle forces transmitted to bone, which could have also contributed to the decreased BV/TV that we observed (9). Reduced muscle loading of the tendon-bone attachment is detrimental to tendon-bone healing (8,9), yet it remains unknown if a partial-width injury to the tendon-bone attachment influences the muscle loading of the attachment in our model. Per our *ex vivo* validation studies, the injured attachments likely have decreased biomechanical properties at time of injury due to loss of structural integrity and removal of attachment tissue. Interestingly, none of the partial-width injuries resulted in a full-width injury, supporting recent evidence that the attachment may resist failure even when damaged, albeit less so than a healthy attachment (24). Therefore, although full regeneration of the structure of the tendon-bone attachment was not observed, the scar tissue formed at the injury site may be mechanically sufficient to resist tear propagation.

Research significance

Although tear propagation was not observed in this study, rotator cuff tears in the clinic may propagate into full-width injuries that require surgical repair (38,40). Therefore, partial injuries to rotator cuff tendons are sometimes surgically repaired to prevent tear propagation (18,29). Hence, many studies have aimed to understand tear propagation (24,3942) and various scenarios of rotator cuff repair (7,18,29,31,34,4347). Additionally, studies have aimed to develop partial-width tear models in rotator cuff tendons as well as in patellar and Achilles tendon groups (48,49). Despite the plethora of research in the field of rotator cuff repair, repairs often fail due to poor re-integration of tendon to bone (7,50). As such, studies of attachment healing have focused on tissue-engineering approaches for improving attachment repair and establishing the mechanical and biological factors that are involved in repair healing (8,9,15,19,20,22,23,3234,37,51–65). Few studies have investigated the innate healing response of the attachment (i.e., attachment healing without repair). One such study identifies a role of Gli1+ cells, key regulators of attachment development, in attachment regeneration via direct punch-like injury to neonatal mouse attachments (22). Although these cells mediate neonate regeneration, Gli1+ cells were scarce during scar-mediated healing in the mature attachment (22). Interestingly, scar-mediated healing has been shown to occur both in the mature mouse and in our rat model of healing, indicating that aging may limit the capabilities for tissue regeneration, regardless of species. The results of our study are consistent with other animal models of rotator cuff injury (8,9,11,19,20,5,61), elucidating that the composition of the attachment varies at different time points after an acute injury. In one study of acute rotator cuff tendon injury and repair, attachments exhibited a wound healing response with a lack of restoration achievable compared to uninjured shoulders (15). Only a few studies, however, have focused on the healing composition of partial-width injuries without repair or augmentation (20). In this study, we established that a model of partial-width rotator cuff injury in the rat shoulder leads to achievable restoration, although scar-like and fibrotic, at the attachment without surgical repair.

Clinical significance

An increased understanding of the healing properties of partial-width rotator cuff injuries are necessary to establish optimal treatment plans for patients with these injuries. The small animal injury model presented here will be beneficial in determining the innate healing properties of the attachment, which can contribute to pharmacological interventions or treatment protocols for partial-width tears.

Limitations

The clinical relevancy of this model is somewhat limited, as most partial-width tears reside on the bursal side of the attachment and include the lateral attachments of the tendon to bone (19,33,34). However, this model is useful to evaluate the injury and healing characteristics of the attachment, directly. Additionally, in this model, the punch biopsy permeated the cortical bone of the humeral head and removed a portion of bone, which could be contribute to the loss of BV/TV in the injured attachments compared to the sham-operated attachments. The permeation of the cortical bone could have also facilitated the interaction of marrow components with the injury site, potentially altering the natural healing response. Furthermore, the differences in time points and sex among the rats may contribute to slight variations in the healing of the attachments. Although this study did not detect significant differences in bone morphometry or quantitative histological outcomes between male and female groups, it is possible that sex differences in healing outcomes exist. Our estimates of sample size using power analyses for detecting sex differences suggest that we would require a sample size of at least 15 per group. Although potentially relevant to healing outcomes of rotator cuff injuries, comparisons of sex differences were not the primary focus for this study. Future work should focus on the healing effects of rotator cuff injuries dependent on sex. The discrepancy between the BV/TV for the entire humeral head and injury region may be due to the sensitivity of the threshold parameters of the microCT software, resulting in undeveloped or less dense bone not being distinguished. Lastly, biomechanical testing was not performed for this injury *in vivo*. Thus, the strength of the integrity of the attachment is unknown, despite decreased structural quality at the injury site of the attachment after healing.

Conclusions

A partial-width, full-thickness injury to the IS attachment of the rotator cuff compromises the quality of the attachment. This study established the structural quality of the attachment following healing in a new rodent model. Proliferation and remodeling of the attachment during healing resulted in attachments that were structurally inferior compared to an uninjured attachment even after 8 weeks of healing. This study contributes a novel *in vivo* model of a partial-width, full-thickness rotator cuff injury that can be used as the basis for further research to evaluate healing properties and tear propagation of the attachment with moderate to vigorous exercise, immobilization, or pharmacological interventions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Special thanks to Terry Kokas, Histology Specialist III (Nemours-A.I. duPont Hospital for Children Histochemistry and Tissue Processing Core; supported by NIH-NIGMS: P20 GM103446 and the state of Delaware) and Crystal Idleburg, HT (ASCP) (Washington University Musculoskeletal Research Center, NIH NIAMS: P30 AR057235) for assistance with histology. The injury illustration (Figure 1A) was created by Katelyn McDonald, MA, CMI (Certified Medical Illustrator).

Funding

This work was supported by the Delaware Space Grant Consortium [DESGC NNX15AI19H]; National Center for Medical Rehabilitation Research [K12HD073945]; National Institute of General Medical Sciences [8 P20 GM103446-16]; University of Delaware Research Foundation [UDRF 16A01396]; Washington University Musculoskeletal Research Center (NIH P30 AR057235); Delaware Rehabilitation Institute (P30GM103333).

References

1. Vitale MA, Vitale MG, Zivin JG, Braman JP, Bigliani LU, Flatow EL, Rotator cuff repair: an analysis of utility scores and cost-effectiveness. *Journal of Shoulder and Elbow Surgery*. 2007 16:2181–187.
2. Colvin AC, Egorova N, Harrison AK, Moskowitz A, Flatow EL, National trends in rotator cuff repair. *J Bone Joint Surg Am*. 2012 94:3227–233.
3. Dunn WR, Kuhn JE, Sanders R, An Q, Baumgarten KM, Bishop JY, et al. 2013 Neer Award: predictors of failure of nonoperative treatment of chronic, symptomatic, full-thickness rotator cuff tears. *J Shoulder Elbow Surg*. 2016 25(8):1303–1311. [PubMed: 27422460]
4. Li S, Sun H, Luo X, Wang K, Wu G, Zhou J, et al. The clinical effect of rehabilitation following arthroscopic rotator cuff repair: a meta-analysis of early versus delayed passive motion. *Medicine*. 2018 97(2):e9625. [PubMed: 29480870]
5. Kong BY, Cho M, Lee HR, Choi YE, Kim SH, Structural evolution of nonoperatively treated high-grade partial-thickness tears of the supraspinatus tendon. *American Journal of Sports Medicine*. 2018 46 (1):79–86. [PubMed: 28949249]
6. Frisch KE, Marcu D, Baer GS, Thelen DG, Vanderby R, The influence of partial and full thickness tears on infraspinatus tendon strain patterns. *Journal of Biomechanical Engineering*. 2014 136:5051004.
7. Galatz LM, Ball CM, Teefey SA, Middleton WD, Yamaguchi K, The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *The Journal of Bone and Joint Surgery, American volume*. 2004 86-A(2):219–224. [PubMed: 14960664]
8. Galatz LM, Charlton N, Das R, Kim HM, Havlioglu N, Thomopoulos S, Complete removal of load is detrimental to rotator cuff healing. *Journal of Shoulder and Elbow Surgery*. 2009 18(5):669–675. [PubMed: 19427237]
9. Killian ML, Cavinatto L, Shah SA, Sato EJ, Ward SR, Havlioglu N, et al. The effects of chronic unloading and gap formation on tendon-to-bone healing in a rat model of massive rotator cuff tears. *Journal of Orthopaedic Research*. 2014 32(3):439–447. [PubMed: 24243733]
10. Sato E, Killian M, Choi A, Lin E, Choo A, Rodriguez-Soto A, et al. Architectural and biochemical adaptations in skeletal muscle and bone following rotator cuff injury in a rat model. *Journal of Bone and Joint Surgery*. 2015 97(7):565–573. [PubMed: 25834081]
11. Cavinatto L, Malavolta EA, Pereira CAM, Miranda-Rodrigues M, Silva LCM, Gouveia CH, et al. Early versus late repair of rotator cuff tears in rats. *Journal of Shoulder and Elbow Surgery*. 2018 27(4):606–613. [PubMed: 29274903]
12. Mellado JM, Calmet J, Olona M, Esteve C, Camins A, Pérez Del Palomar L, et al. Surgically repaired massive rotator cuff tears: MRI of tendon integrity, muscle fatty degeneration, and muscle atrophy correlated with intraoperative and clinical findings. *American Journal of Roentgenology*. 2005 184(5):1456–1463. [PubMed: 15855096]
13. Chung SW, Oh JH, Gong HS, Kim JY, Kim SH, Factors affecting rotator cuff healing after arthroscopic repair. *The American Journal of Sports Medicine*. 2011 39:102099–2107.

14. Kim Y- K, Jung K- H, Kim J- W, Kim U- S, Hwang D- H, Factors affecting rotator cuff integrity after arthroscopic repair for medium-sized or larger cuff tears: a retrospective cohort study. *Journal of shoulder and elbow surgery*. 2018 27(6):1012–1020. [PubMed: 29290609]
15. Galatz LM, Sandell LJ, Rothermich SY, Das R, Mastny A, Havlioglu N, et al. Characteristics of the rat supraspinatus tendon during tendon-to-bone healing after acute injury. *Journal of Orthopaedic Research*. 2006 24(3):541–550. [PubMed: 16456829]
16. Park JS, Park HJ, Kim SH, Oh JH, Prognostic factors affecting rotator cuff healing after arthroscopic repair in small to medium-sized tears. *Am J Sports Med*. 2015 43:102386–2392.
17. Kim KC, Shin HD, Lee WY, Repair integrity and functional outcomes after arthroscopic suture-bridge rotator cuff repair. *J Bone Joint Surg Am*. 2012 94:8e48.
18. Boileau P, Brassart N, Watkinson DJ, Carles M, Hatzidakis AM, Krishnan SG, Arthroscopic repair of full-thickness tears of the supraspinatus: does the tendon really heal? *The Journal of Bone and Joint Surgery, American Volume*. 2005 87:61229–1240.
19. Thomopoulos S, Williams GR, Soslowsky LJ, Tendon to bone healing: differences in biomechanical, structural, and compositional properties due to a range of activity levels. *Journal of Biomechanical Engineering*. 2003 125(1):106–113. [PubMed: 12661203]
20. Carpenter JE, Thomopoulos S, Flanagan CL, DeBano CM, Soslowsky LJ, Rotator cuff defect healing: a biomechanical and histologic analysis in an animal model. *Journal of Shoulder and Elbow Surgery*. 1998 7:6599–605.
21. Derwin KA, Baker AR, Codsi MJ, Iannotti JP, et al. Assessment of the canine model of rotator cuff injury and repair. *Journal of shoulder and elbow surgery/American Shoulder and Elbow Surgeons*. 2007 16(5 Suppl)S140–S8.
22. Schwartz AG, Galatz LM, Thomopoulos S, Enthesis regeneration: a role for Gli1+ progenitor cells. *Development (Cambridge, England)*. 2017 144(7):1159–1164.
23. Derwin KA, Codsi MJ, Milks RA, Baker AR, McCarron JA, Iannotti JP, Rotator cuff repair augmentation in a canine model with use of a woven poly-L-lactide device. *The Journal of Bone and Joint Surgery, American Volume*. 2009 91(5):1159–1171. [PubMed: 19411465]
24. Locke RC, Peloquin JM, Lemmon EA, Szostek A, Elliott DM, Killian ML, Strain distribution of intact rat rotator cuff tendon-to-bone attachments and attachments with defects. *Journal of Biomechanical Engineering*. 2017 139(11):doi:10.1115/1.4038111.
25. David MA, Smith MK, Pilachowski RN, White AT, Locke RC, Price C, Early, focal changes in cartilage cellularity and structure following surgically induced meniscal destabilization in the mouse. *Journal of Orthopaedic Research*. 2017 35(3):537–547. [PubMed: 27664978]
26. Thomopoulos S, Marquez JP, Weinberger B, Birman V, Genin GM, Collagen fiber orientation at the tendon to bone insertion and its influence on stress concentrations. *J Biomech*. 2006 39(10):1842–1851. [PubMed: 16024026]
27. Rossetti L, Kuntz LA, Kunold E, Schock J, Muller KW, Grabmayr H, et al. The microstructure and micromechanics of the tendon-bone insertion. *Nat Mater*. 2017 16(6):664–670. [PubMed: 28250445]
28. Kanazawa T, Gotoh M, Ohta K, Shiba N, Nakamura K, Novel characteristics of normal supraspinatus insertion in rats: an ultrastructural analysis using three-dimensional reconstruction using focused ion beam/scanning electron microscope tomography. *Muscles Ligaments Tendons J*. 2014 4(2):182–187. [PubMed: 25332933]
29. Lo IKY, Burkhart SS, Double-row arthroscopic rotator cuff repair: re-establishing the footprint of the rotator cuff. *Arthroscopy: The Journal of Arthroscopic & Related Surgery: Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*. 2003 19 (9):1035–1042.
30. Tashjian RZ, Hollins AM, Kim HM, Teefey SA, Middleton WD, Steger-May K, et al. Factors affecting healing rates after arthroscopic double-row rotator cuff repair. *The American Journal of Sports Medicine*. 2010 38(12):2435–2442. [PubMed: 21030564]
31. Thomopoulos S, Parks WC, Rifkin DB, Derwin KA, Mechanisms of tendon injury and repair. *Journal of Orthopaedic Research*. 2015 33(6):832–839. [PubMed: 25641114]

32. Ji X, Chen Q, Thoreson AR, Qu J, An K- N, Amadio PC, et al. Rotator cuff repair with a tendon-fibrocartilage-bone composite bridging patch. *Clinical Biomechanics*. 2015 30(9):976–980. [PubMed: 26190097]
33. Lewington MR, Ferguson DP, Smith TD, Burks R, Coady C, Wong IH- B, Graft utilization in the bridging reconstruction of irreparable rotator cuff tears: a systematic review. *The American Journal of Sports Medicine*. 2017 45(13):3149–3157. [PubMed: 28345960]
34. Ostrander RV, Klauser JM, Menon S, Hackel JG, Ultrasound and functional assessment of transtendinous repairs of partial-thickness articular-sided rotator cuff tears. *Orthopaedic Journal of Sports Medicine*. 2017 5(3):2325967117697375. [PubMed: 28451606]
35. Goldberg BA, Nowinski RJ, Matsen FA, 3rd., Outcome of nonoperative management of full-thickness rotator cuff tears. *Clin Orthop Relat Res*. 2001382:99–107.
36. Zingg PO, Jost B, Sukthankar A, Buhler M, Pfirrmann CW, Gerber C, Clinical and structural outcomes of nonoperative management of massive rotator cuff tears. *J Bone Joint Surg Am*. 2007 89(9):1928–1934. [PubMed: 17768188]
37. Nakagawa H, Morihara T, Fujiwara H, Kabuto Y, Sukenari T, Kida Y, et al. Effect of footprint preparation on tendon-to-bone healing: a histologic and biomechanical study in a rat rotator cuff repair model. *Arthroscopy: The Journal of Arthroscopic & Related Surgery: Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*. 2017 33(8):1482–1492.
38. Reilly P, Amis AA, Wallace AL, Emery RJH, Mechanical factors in the initiation and propagation of tears of the rotator cuff. Quantification of strains of the supraspinatus tendon in vitro. *The Journal of Bone and Joint Surgery British Volume*. 2003 85(4):594–599. [PubMed: 12793570]
39. Reilly P, Amis AA, Wallace AL, Emery RJH, 2003 Supraspinatus tears: propagation and strain alteration. *Journal of Shoulder and Elbow Surgery*. 12:2134–138.
40. Miller RM, Fujimaki Y, Araki D, Musahl V, Debski RE, Strain distribution due to propagation of tears in the anterior supraspinatus tendon. *Journal of Orthopaedic Research*. 2014 32:1283–1289. [PubMed: 24985532]
41. Andarawis-Puri N, Kuntz AF, Kim S- Y, Soslowsky LJ, Effect of anterior supraspinatus tendon partial-thickness tears on infraspinatus tendon strain through a range of joint rotation angles. *Journal of Shoulder and Elbow Surgery*. 2010 19(4):617–623. [PubMed: 20080051]
42. Andarawis-Puri N, Ricchetti ET, Soslowsky LJ, Rotator cuff tendon strain correlates with tear propagation. *J Biomech*. 2009 42(2):158–163. [PubMed: 19064267]
43. Derwin KA, Baker AR, Spragg RK, Leigh DR, Iannotti JP, Commercial extracellular matrix scaffolds for rotator cuff tendon repair. Biomechanical, biochemical, and cellular properties. *The Journal of Bone and Joint Surgery, American Volume*. 2006 88(12):2665–2672. [PubMed: 17142417]
44. Derwin KA, Badylak SF, Steinmann SP, Iannotti JP, Extracellular matrix scaffold devices for rotator cuff repair. *Journal of Shoulder and Elbow Surgery*. 2010 19(3):467–476. [PubMed: 20189415]
45. Barber FA, Burns JP, Deutsch A, Labbé MR, Litchfield RB, A prospective, randomized evaluation of acellular human dermal matrix augmentation for arthroscopic rotator cuff repair. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2012 28(1):8–15. [PubMed: 21978432]
46. Yang G, Rothrauff BB, Tuan RS, Tendon and ligament regeneration and repair: clinical relevance and developmental paradigm. *Birth Defects Research Part C, Embryo Today: Reviews*. 2013 99(3):203–222.
47. Thangarajah T, Pendegrass CJ, Shahbazi S, Lambert S, Alexander S, Blunn GW, Augmentation of rotator cuff repair with soft tissue scaffolds. *Orthopaedic Journal of Sports Medicine*. 2015 3(6): 2325967115587495. [PubMed: 26665095]
48. Beason DP, Kuntz AF, Hsu JE, Miller KS, Soslowsky LJ, Development and evaluation of multiple tendon injury models in the mouse. *J Biomech*. 2012 45 (8):1550–1553. [PubMed: 22405494]
49. Yoshida R, Alaei F, Dyrna F, Kronenberg MS, Maye P, Kalajzic I, et al. Murine supraspinatus tendon injury model to identify the cellular origins of rotator cuff healing. *Connect Tissue Res*. 2016 57(6):507–515. [PubMed: 27184388]

50. Kim HM, Dahiya N, Teefey SA, Middleton WD, Stobbs G, Steger-May K, et al. Location and initiation of degenerative rotator cuff tears: an analysis of three hundred and sixty shoulders. *The Journal of Bone and Joint Surgery, American Volume*. 2010 92 (5):1088–1096. [PubMed: 20439653]
51. Li X, Xie J, Lipner J, Yuan X, Thomopoulos S, Xia Y, Nanofiber scaffolds with gradations in mineral content for mimicking the tendon-to-bone insertion site. *Nano Letters*. 2009 9(7):2763–2768. [PubMed: 19537737]
52. Gulotta LV, Kovacevic D, Packer JD, Deng XH, Rodeo SA, Bone marrow-derived mesenchymal stem cells transduced with scleraxis improve rotator cuff healing in a rat model. *The American Journal of Sports Medicine*. 2011 39(6):1282–1289. [PubMed: 21335341]
53. Gulotta LV, Kovacevic D, Packer JD, Ehteshami JR, Rodeo SA, Adenoviral-mediated gene transfer of human bone morphogenetic protein-13 does not improve rotator cuff healing in a rat model. *The American Journal of Sports Medicine*. 2011 39 (1):180–187. [PubMed: 20956264]
54. Zhao S, Zhao J, Dong S, Huangfu X, Li B, Yang H, et al. Biological augmentation of rotator cuff repair using bFGF-loaded electrospun poly(lactide-co-glycolide) fibrous membranes. *International Journal of Nanomedicine*. 2014 9:2373–2385. [PubMed: 24868155]
55. Peterson DR, Ohashi KL, Aberman HM, Piza PA, Crockett HC, Fernandez JJ, et al. 2015 Evaluation of a collagen-coated, resorbable fiber scaffold loaded with a peptide basic fibroblast growth factor mimetic in a sheep model of rotator cuff repair. *Journal of Shoulder and Elbow Surgery*. 24(11):1764–1773. [PubMed: 26251198]
56. Degen RM, Carbone A, Carballo C, Zong J, Chen T, Lebaschi A, et al. The effect of purified human bone marrow-derived mesenchymal stem cells on rotator cuff tendon healing in an athymic rat. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2016 32(12):2435–2443. [PubMed: 27282111]
57. Kanazawa T, Gotoh M, Ohta K, Honda H, Ohzono H, Shimokobe H, et al. Histomorphometric and ultra-structural analysis of the tendon-bone interface after rotator cuff repair in a rat model. *Scientific Reports*. 2016 6:33800. [PubMed: 27647121]
58. Omi R, Gingery A, Steinmann SP, Amadio PC, An K-N ZC, Rotator cuff repair augmentation in a rat model that combines a multilayer xenograft tendon scaffold with bone marrow stromal cells. *Journal of Shoulder and Elbow Surgery*. 2016 25(3):469–477. [PubMed: 26387915]
59. Ye C, Zhang W, Wang S, Jiang S, Yu Y, Chen E, et al. Icaritin promotes tendon-bone healing during repair of rotator cuff tears: a biomechanical and histological study. *International Journal of Molecular Sciences*. 2016 17:11.
60. Shah SA, Korpakakis I, Havlioglu N, Ominsky MS, Galatz LM, Thomopoulos S, Sclerostin Antibody treatment enhances rotator cuff tendon-to-bone healing in an animal model. *The Journal of Bone and Joint Surgery American Volume*. 2017 99(10):855–864. [PubMed: 28509826]
61. Killian ML, Cavinatto LM, Ward SR, Havlioglu N, Thomopoulos S, Galatz LM, Chronic degeneration leads to poor healing of repaired massive rotator cuff tears in rats. *The American Journal of Sports Medicine*. 2015 43(10):2401–2410. [PubMed: 26297522]
62. Sato E, Killian M, Choi A, Lin E, Esparza M, Galatz L, et al. Skeletal muscle fibrosis and stiffness increase after rotator cuff tendon injury and neuromuscular compromise in a rat model. *Journal of Orthopaedic Research*. 2014 32(9):1111–1116. [PubMed: 24838823]
63. Wildemann B, Klatte F, Biological aspects of rotator cuff healing. *Muscles, Ligaments and Tendons Journal*. 2011 1(4):161–168.
64. Thomopoulos S, Hattersley G, Rosen V, Mertens M, Galatz L, Williams GR, et al. The localized expression of extracellular matrix components in healing tendon insertion sites: an in situ hybridization study. *Journal of Orthopaedic Research*. 2002 20(3):454–463. [PubMed: 12038618]
65. Soslowsky LJ, Carpenter JE, DeBano CM, Banerji I, Moalli MR, Development and use of an animal model for investigations on rotator cuff disease. *Journal of Shoulder and Elbow Surgery*. 1996 5(5):383–392. [PubMed: 8933461]

Figure 1.

Rotator cuff infraspinatus injury model and microCT reconstructions of sham and injured groups. (A) Illustration of the human rotator cuff muscle groups. In our animal model, a 0.3 mm circular injury was made at the center of the infraspinatus attachment. MicroCT renders of (B) sham and (C) injured humeral heads. The orange region is the injury location.

Figure 2.

Humeral head BV/TV was significantly reduced in injured attachments compared to the sham forearms for both time points. (A–F) MicroCT reconstructions of sham and injured forearms at 3 and 8 weeks. (A–C) Frontal plane views of microCT reconstructions showing the IS ridge of the humeral head for (A) 3 wk sham, (B) 3 wk injured, and (C) 8 wk injured attachments. (D–F) Transverse cut-plane (D) 3 wk sham, (E) 3 wk injured, and (F) 8 wk injured. White arrowheads: Cortical bone injury in the injury region. (G) Humeral head BV/TV and (H) Injury region BV/TV for 3 wk control, 3 wk injured, 8 wk control, and 8 wk injured attachments. Bars indicate significant differences between groups ($p < 0.05$, mean $\pm$ 95% CI).

Figure 3.

Organization at the IS attachment was impaired in 3 wk injured attachments compared to 3 wk sham attachments. (A–C') Transverse histology stained with Picrosirius Red of (A) 3 wk sham, (A') 3 wk injured, (C) 8 wk sham, and (C') 8 wk injured. (B–D') Transverse histology stained with Masson's Trichrome of (B) 3 wk sham, (B') 3 wk injured, (D) 8 wk sham, and (D') 8 wk injured. White arrowheads in A & A' highlight regions of organized collagen, with (A) sham attachments having noticeably more organized collagen compared to (A') injured attachments. White arrowheads in B' highlight ECM deposition at the injured IS attachment, as well as (D') fatty accumulation in the tendon. Black arrowhead in (D') highlights fibrosis at the injured attachment. (E) Circular Standard Deviation (quantification of collagen fibril alignment) was significantly increased for 3 wk injured attachments compared to sham. Bars indicate significant differences between groups ($p < 0.05$, mean $\pm$ 95% CI). Images taken at 5 $\times$ magnification, scale bar: 200 μ m.

Figure 4.

Cell Density and Oc.S./B.S. ratio were significantly greater in the injured attachments compared to the sham-operated attachments for both time points. (A–D) Transverse histological images at 5× magnification of (A) 3 wk sham, (B) 3 wk injured, (C) 8 wk sham, and (D) 8 wk injured. Scale bar: 200 μm. (B'–D') 20× magnification of (B') 3 wk injured and (D') 8 wk injured. Scale bar: 30 μm. (E) Cell density and (F) Osteoclast surface (Oc.S.) to bone surface (B.S.) ratio. Bars indicate significant differences between groups ($p < 0.05$, mean $\pm$ 95% CI).