CASE REPORT

Starvation-induced diplopia and weakness: a case of beriberi and Wernicke's encephalopathy

Timothy Xin Zhong Tan,¹ Kheng Choon Lim,² Charles Chan Chung,³ Than Aung³¹Department of Emergency Medicine, Singapore General Hospital, Singapore²Department of Radiology, Singapore General Hospital, Singapore³Department of Internal Medicine, Singapore General Hospital, Singapore**Correspondence to**Timothy Xin Zhong Tan,
timothy.tbj13@gmail.com

Accepted 12 December 2018

SUMMARY

A 56-year-old teetotaler man with hypertension and gout presented with a week duration of painless worsening diplopia on a background of loss of weight and appetite, generalised lethargy and weakness for 1 year. On examination, he was noted to be hypothermic and tachycardic with generalised muscle wasting. Proximal myopathy, lower limb fasciculations and areflexia, restricted bilateral eye abduction and nystagmus were observed. Blood investigations demonstrated compensated lactic acidosis, acute kidney injury and leucocytosis. A nerve conduction study showed severe length-dependent axonal sensorimotor polyneuropathy. This was a diagnostic dilemma until an MRI brain revealed symmetrical signal abnormality and enhancement in the periaqueductal area indicative of Wernicke's encephalopathy, caused by thiamine deficiency from poor nutrition. Beriberi, also caused by thiamine deficiency, accounted for his tachycardia, polyneuropathy, areflexia, hypothermia and biochemical abnormalities. Both beriberi and Wernicke's encephalopathy are medical emergencies, which were treated with intravenous thiamine to good effect.

BACKGROUND

Wernicke's encephalopathy (WE) and beriberi are uncommon life-threatening pathologies that are challenging yet critical to diagnose. They can present with a variety of non-specific constitutional symptoms as in this case and may require extensive investigations before definitive diagnosis is made. A high index of suspicion and astute radiological experience is therefore crucial for early diagnosis, allowing for early treatment and improved chances of a complete recovery. It is also unusual for a cognitively intact patient with adequate social support to have suffered for so long before presentation, resulting in him developing extensive clinical signs and symptoms. Although there are reports of WE in literature, reports of beriberi are rare in developed countries where a balanced diet and medical care are easily accessible. There is therefore little published information with regard to the symptoms, complications and management of a patient with beriberi.

CASE PRESENTATION

A 56-year-old Malay man retired waste collector who has a medical history of hypertension and gout, non-compliant to medications presented with generalised lethargy and weakness in his lower limbs

and progressing to his upper limbs, associated with loss of appetite and loss of weight for 1 year duration. This started after a flare of gout caused him to be completely bedbound and assisted in basic activities of daily living. During this period, his daily diet only comprised of an apple or orange, a small packet of dry biscuits and a cup of water, causing him to experience a weight loss of 30 % from 85 to 58 kg. He subsequently developed painless gradually worsening horizontal binocular diplopia for 1 week and eventually decided to present to the hospital.

He denied smoking or drinking alcohol. There was no family history of malignancy, tuberculosis or neurological disorders. There were no symptoms indicating hyperthyroidism, and no symptoms suggestive of cancer such as headaches, vomiting, jaundice and alteration of bowel habits.

On examination, the patient was hypothermic, with a body temperature of 35°C–36°C and appeared severely deconditioned, lethargic and dehydrated. He was haemodynamically stable, and oriented, conversant and non-toxic. Multiple tophaceous deposits were noted over bilateral hands, wrists, knees and feet.

Thorough neurological examination revealed generalised muscle wasting over bilateral hands, thighs and calves (figure 1). A non-fatigable proximal myopathy pattern of muscle weakness was also noted, with the proximal myotomes in the upper and lower limbs (C5, C6, L2, L3) having a power of 2 out of 5 on the Medical Research Council scale, and the distal myotomes (C7, T1, L4, L5, S1) having a power of 4 out of 5. The patient was unable to ambulate. Bilateral lower limbs were areflexic, while upper limb reflexes were preserved. Fasciculations were noted over bilateral thighs and calves.

Figure 1 Clinical photographs of the patient's upper and lower limbs demonstrating generalised muscle wasting.

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Tan TXZ, Lim KC, Chan Chung C, et al. *BMJ Case Rep* 2019;**12**:e227412. doi:10.1136/bcr-2018-227412

Video 1 Clinical video of patient's eyes demonstrating limited abduction bilaterally with nystagmus.

Restriction of eye movement was also noted in the horizontal plane, with limited abduction bilaterally, nystagmus and compensatory movement of head (video 1). Convergence was also restricted, and passive movement and Doll's reflexes were preserved. Pupils were equal and reactive to light at 3 mm bilaterally and ptosis, anisocoria, relative afferent pupillary defect and fatigability were not observed. The rest of his neurological examination, including an Abbreviated Mental Test, was normal.

INVESTIGATIONS

Initial blood investigations revealed compensated lactic acidosis (venous blood gas pH 7.362 (reference range 7.350–7.450)), bicarbonate 16.8 mmol/L (19.0–29.0 mmol/L), anion gap 23.2 mEq/L, lactate 3.2 mmol/L (0.5–2.2 mmol/L), ketones 0.4 mmol/L (0.0–0.6 mmol/L), kidney disease: improving global outcomes stage 1 kidney injury¹ with raised urea 15 mmol/L (2.7–6.9 mmol/L) and creatinine 165 µmol/L (54–101 µmol/L), and iron and folate deficiency anaemia (haemoglobin 11.5 g/L (14–18 g/L), iron <2 µmol/L (11–32 µmol/L), total iron binding capacity 30 µmol/L (44–73 µmol/L), transferrin saturation below detection limit, low serum folate 4.5 nmol/L (>13.4 nmol/L) likely secondary to poor oral intake and dehydration. Neutrophilic leucocytosis (white cell count $21.2 \times 10^9/L$ ($4\text{--}10 \times 10^9/L$)) was also noted. Blood, stool and urine cultures, stool ova, cysts and parasites, stool virus panel, and hepatitis B, C and HIV screens returned negative. Thyroid and liver function tests, myeloma panel and glycated haemoglobin (HbA1c) returned normal.

Figure 2 MRI brain axial T2-weight FLAIR (T2 FLAIR) and postgadolinium T1-weighted MPRAGE (Gad T1) sequences demonstrating symmetrical high FLAIR signal and enhancement along the periaqueductal area, medial parts of bilateral thalami and hypothalamus suggestive of Wernicke's encephalopathy.

The nerve conduction study showed severe length-dependent predominantly axonal, sensorimotor polyneuropathy, right peroneal and right ulnar sensorimotor neuropathy and carpal tunnel syndrome bilaterally. Needle electromyography revealed neurogenic changes with chronic denervation. Further electro-myogram analysis was limited in view of joint pain from gout.

A chest X-ray and a CT of the thorax, abdomen and pelvis returned negative for malignancy. An MRI brain and orbits with contrast revealed symmetrical high T2-weighted FLAIR (fluid-attenuated inversion recovery) signal and enhancement along the periaqueductal grey and hypothalamus (figure 2) with symmetrical high T2-weighted FLAIR signal without enhancement at the bilateral medial thalami (figure 3). No restricted diffusion or mass effect was evident. The extra-ocular muscles were unremarkable. There was no space occupying lesion detected. In the clinical context of ophthalmoplegia, raised lactate and weight loss, the possibility of WE was raised based on imaging findings. This was likely due to thiamine deficiency secondary to poor nutrition, and explained the distal sensorimotor axonal polyneuropathy affecting lower more than upper limbs, a classical neurological manifestation present in our patient.² Concomitant folate deficiency also contributed synergistically to this patient's neurological symptoms.^{3,4}

The patient subsequently developed sinus tachycardia ranging from 100 to 140 beats per minute. This is probably due to underlying beriberi, secondary to thiamine deficiency. A Doppler ultrasonography of his lower limbs and a ventilation-perfusion scan reported no evidence of thromboembolism, and a trans-thoracic echocardiogram to assess for cardiac function returned normal.

DIFFERENTIAL DIAGNOSIS

In this patient presenting with severe weight loss and lethargy, it is crucial to rule out malignancy. Other differentials include chronic infectious diseases such as tuberculosis and AIDS, non-malignant gastrointestinal pathologies such as celiac disease and inflammatory bowel disease, and psychiatric conditions such as depression and eating disorders.

For symmetrical proximal myopathy, it is important to differentiate true muscle weakness from lassitude, weakness and pain. Differentials of true muscle weakness with diplopia would include inflammatory myopathies such as dermatomyositis and polymyositis and endocrinopathies such as thyroid disease. Rarer causes include malignancy-associated myositis and AIDS.

Myasthenia gravis can also present with proximal limb and ocular muscle weakness, with diplopia being one of the most common presenting complaints.^{4,5} However, ptosis and fatigability were absent. Diplopia is also more commonly asymmetric and fluctuates with diurnal variation,⁶ not present in this patient.

An intracranial space occupying lesion is another important differential to consider, for which an MRI brain was done. For MRI findings, the differential diagnoses of symmetric thalamic lesions include autoimmune encephalitis such as primary acute disseminated encephalomyelitis and Behçet disease, and infective encephalitis such as variant Creutzfeldt-Jakob disease, influenza A and cytomegalovirus encephalitis.⁷

Thiamine-responsive megaloblastic anaemia is a rare autoimmune recessive genetic condition characterised by hearing loss, diabetes and megaloblastic anaemia.^{8,9} However, this is unlikely to be the aetiology of this patient's anaemia, as it was microcytic (mean corpuscular volume 77 fL (78–98 fL)) and not congenital and therefore more likely to be secondary to iron and folate deficiency. Patients with nutritional deficiencies tend to have not one

Figure 3 MRI brain axial T2-weighted FLAIR (T2 FLAIR) sequences demonstrating symmetrical high T2-weighted periventricular signal in the medial thalami.

but multiple concomitant vitamin and mineral deficiencies.¹⁰ However, other than thiamine, iron and folate deficiency, there were no signs or symptoms of other vitamin or mineral deficits such as pellagra from niacin deficiency. There is also no direct relation between these deficiencies and gout.

TREATMENT

Beriberi and WE are medical emergencies. Proper diagnosis is key as up to 29% of cases experience complete remission with appropriate treatment,^{11 12} without which 10%–20% die and many more left with permanent debilitating neurological sequelae.¹³ However, as its symptoms are non-specific and can easily mimic other differentials, diagnosis is often difficult and delayed leading to prolonged stay and increased costs.¹⁴ Hence, there is a paucity of clinical trials with regard to treatment, particularly the ideal dose of thiamine. Recommended regimens in literature range from 100 to 500 mg of intravenous thiamine three to four times a day for two consecutive days, followed by 250–500 mg once a day for an additional 2–5 days or until oral thiamine is tolerated.^{12 13 15} Duration of treatment is also uncertain. Treatment route should initially be parenteral, and subsequently continued per-oral 60–100 mg a day.¹² We treated this man with folic acid and intravenous thiamine 500 mg three times a day for 2 days, reducing it to 250 mg once a day for 2 days before switching to oral Princi-B fort 1 tablet once a day (a vitamin B supplement containing 250 mg of thiamine, 250 mg of vitamin B₆ and 1000 mg of vitamin B₁₂ per tablet).

Multidisciplinary treatment was also adopted, and the patient was referred to rheumatology for gout control, the dietician for nutrition management, and the occupational and physical therapists for management of muscle strength and rehabilitation. Medical social services was also referred for financial and care difficulties.

Video 2 Clinical video of patient's eyes 2 weeks after treatment with thiamine supplementation demonstrating complete resolution of ophthalmoplegia.

OUTCOME AND FOLLOW-UP

The patient's signs and symptoms markedly improved after commencement of thiamine and folic acid supplementation. He reported feeling better, with restoration of appetite and weight gain. After 2 weeks, his eye signs, kidney injury and muscle weakness had resolved ([video 2](#)). There were no signs or biochemical abnormalities of refeeding syndrome. One month later, the patient was transferred to a step-down rehabilitation facility for further care. At the time of writing, the patient is undergoing further rehabilitation and recovering well.

DISCUSSION

Thiamine, also known as vitamin B₁, is primarily sourced from foods such as brown rice, cereals and pork. It plays a crucial role in nerve impulse propagation and in the catabolism of carbohydrates and amino acids, where it serves as a cofactor in the conversion of pyruvate to acetyl coenzyme A. In acute deficiency, pyruvate accumulation may occur and get converted to lactate

by lactate dehydrogenase, resulting in lactic acidosis.¹⁶ Thiamine deficiency can also cause WE, beriberi and Korsakoff syndrome.

Beriberi is often categorised in the literature as 'dry' or 'wet'.¹⁴ Dry beriberi is a peripheral sensorimotor polyneuropathy predominantly affecting the distal extremities, while wet beriberi is characterised by cardiovascular involvement and include signs of high output heart failure such as cardiomegaly, tachycardia and peripheral oedema.^{13 16} Our patient did present with peripheral neuropathy and sinus tachycardia. In addition, hypothermia due to involvement of the posterior hypothalamus,¹³ acute kidney injury,¹⁶ leucocytosis and increased lactate¹⁴ also points to the diagnosis of WE. Korsakoff syndrome is a late manifestation of WE characterised by selective amnesia and apathy which were not observed in this patient.^{13 17}

These conditions used to be common in seafarers on long voyages without appropriate nutrition, but in resource-rich countries, they are most commonly diagnosed in alcoholics.¹³ Other aetiologies can be divided into those caused by nutrition deficits and those by absorption defects. Nutrition deficits occur when intake of thiamine is inadequate, such as in hyperemesis gravidarum, anorexia and self-neglect as in our present case.^{13 18} As absorption of thiamine is mostly carrier-mediated at the jejunum and ileum, absorption defects can be caused by gastrointestinal tract surgery and inflammatory bowel disorders.^{19–21}

WE is 1.7 times more common in males than females, with an estimated mortality of 17%.¹⁹ WE is classically described as an acute encephalopathy characterised by an acute triad of confusion, ophthalmoplegia and ataxia. However, this triad is only present in two-thirds of patients,^{11 16} and 19% have none of these three symptoms at presentation. This triad is also seen more frequently in alcoholic than in non-alcoholic cases such as in our patient.¹³ In view of the low sensitivity of this triad, Caine *et al*²¹ proposed a new diagnostic criteria applicable to both alcoholic and non-alcoholic patients which requires two of the four following criteria to be present: oculomotor signs, cerebellar dysfunction, altered mental state or dietary deficiencies. This patient does meet the above criteria, having presented with oculomotor signs and dietary deficiencies. Other possible symptoms of WE range from hallucinations and visual disturbances, to epileptic seizures and dysarthria.¹³ Investigations can be done to support the clinical diagnosis of WE such as serum thiamine and red blood cell transketolase activity. However, these are limited by low specificity and technical difficulty.¹³ MRI of the brain has a high specificity (93%) and moderate sensitivity (53%) with a positive predictive value of 89% and is thus considered the most ideal investigation to confirm a diagnosis of WE.^{12 22}

Reversible cytotoxic oedema of the medial thalamus and periventricular regions of the third ventricles is a typical MRI finding in WE which appears as bilateral symmetric T2-weighted hyperintense signal with corresponding T1-weight hypointense signal. Commonly associated symmetric signal changes and enhancement can also be seen in the mammillary bodies, periaqueductal area and tectal plate.^{2 23} Atypical MRI findings, which are more commonly present among non-alcoholic patients, include signal alterations in the cerebellum, cranial nerve nuclei, splenium and cerebral cortex.^{11 13 18}

The symptoms of WE are typically non-specific, and many investigations are required to eventually reach a diagnosis. Many clinicians are also unaccustomed to making such a diagnosis in non-alcoholic patients.¹³ As such, 75%–80% of WE cases diagnosed postmortem had been misdiagnosed.^{14 19} In this case, astute radiological analysis provided the crucial diagnosis. Treatment is well tolerated and easily initiated and should therefore be initiated as early as possible to alleviate the associated morbidity and mortality.

Patient's perspective

(Given on 9 June 2018)

When I was unable to walk or eat much, I thought it was because of the gout attack and did not think much about it. I thought it would get better but it did not. I actually did not notice the eye problem until my girlfriend, who had returned from overseas told me and brought me to the hospital. At first the doctors did not know what was wrong and did many tests on me. After the brain scan, they told me it was because I did not eat properly and gave me some medications which helped with my eyes and now I feel much better. I started to eat more and was also able to move about more, although I am still unable to walk without a walking frame. Now I feel much better and am looking forward to walking around again. But I am concerned about the costs of my hospital stay because I have been in hospital for around 1 month already. I hope the doctors or social worker can help me with my finances. I am happy and thankful that I have recovered and am excited to go home soon.

Learning points

- Wernicke's encephalopathy and beriberi are life-threatening diseases with high morbidity and mortality. Although they can be easily treated, they are also difficult to diagnose as they can present with a variety of non-specific symptoms. A high index of suspicion and radiological experience is therefore essential, especially in non-alcoholic cases.
- Beriberi is an uncommon pathology caused by thiamine deficiency which can present with neurological symptoms such as polyneuropathy and areflexia, cardiovascular symptoms such as tachycardia and other signs such as hypothermia, acute kidney injury, leucocytosis and lactic acidosis.
- There is no consensus regarding the ideal dose and duration of thiamine supplementation. Further clinical trials, although potentially difficult to implement, are required to guide future management.
- Multidisciplinary management is crucial when caring for a decompensated and malnourished patient. Care must also be taken to address the patient's various issues, medical or otherwise, including but not limited to financial, emotional and social concerns.

Contributors The corresponding author TXZT followed up with the patient, had full access to all data and information used in this case report, and drafted the initial manuscript and subsequent edition. TXZT therefore takes responsibility for the integrity of the submission and its contents. KCL reported the initial imaging findings and selected the radiological images which were eventually selected for use in this article. KCL also reviewed the article and provided necessary feedback and comments. CCC and TA were involved in the direct care of the patient, and gave recommendations and edits which contributed to the final draft. TA provided overall direction and oversaw the entire study.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kellum JA, Lamire N, Aspelin P, *et al*. Kidney disease: Improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl* 2012;2:1–138.

- 2 Sechi G, Sechi E, Fois C, *et al.* Advances in clinical determinants and neurological manifestations of B vitamin deficiency in adults. *Nutr Rev* 2016;74:281–300.
- 3 Sechi G, Fois C, Addis A, *et al.* Clinicopathologic features of folate-deficiency neuropathy. *Neurology* 2015;85:1090.2–1.
- 4 Gwathmey KG, Burns TM. Myasthenia Gravis. *Semin Neurol* 2015;35:327–39.
- 5 Yousuf UA, Yashodhara BM, Thanigasalam T, *et al.* Why do I always see double? A misdiagnosed case of ocular myasthenia gravis for 10 years. *BMJ Case Rep* 2014;2014.
- 6 Vinge L, Jakobsen J, Pedersen AR, *et al.* Diurnal and day-to-day variation of isometric muscle strength in myasthenia gravis. *Muscle Nerve* 2016;53:67–72.
- 7 Zuccoli G, Pipitone N. Neuroimaging findings in acute Wernicke's encephalopathy: review of the literature. *AJR Am J Roentgenol* 2009;192:501–8.
- 8 Galvin R, Bråthen G, Ivashynka A, *et al.* EFNS guidelines for diagnosis, therapy and prevention of Wernicke encephalopathy. *Eur J Neurol* 2010;17:1408–18.
- 9 Green R. Mystery of thiamine-responsive megaloblastic anemia unlocked. *Blood* 2003;102:3464–a–5.
- 10 Koike H, Takahashi M, Ohya K, *et al.* Clinicopathologic features of folate-deficiency neuropathy. *Neurology* 2015;84:1026–33.
- 11 Victor M, Adams RA, Collins GH. *The Wernicke-Korsakoff syndrome and related disorders due to alcoholism and malnutrition*. Philadelphia: FA Davis, 1989.
- 12 Cook CC, Hallwood PM, Thomson AD. B Vitamin deficiency and neuropsychiatric syndromes in alcohol misuse. *Alcohol Alcohol* 1998;33:317–36.
- 13 Oudman E, Postma A, Van der Stigchel S, *et al.* The Montreal Cognitive Assessment (MoCA) is superior to the Mini Mental State Examination (MMSE) in detection of Korsakoff's syndrome. *Clin Neuropsychol* 2014;28:1123–32.
- 14 Ramsi M, Mowbray C, Hartman G, *et al.* Severe lactic acidosis and multiorgan failure due to thiamine deficiency during total parenteral nutrition. *BMJ Case Rep* 2014;2014:bcr2014205264.
- 15 Sechi G, Serra A. Wernicke's encephalopathy: new clinical settings and recent advances in diagnosis and management. *Lancet Neurol* 2007;6:442–55.
- 16 Palacios-Marqués A, Delgado-García S, Martín-Bayón T, *et al.* Wernicke's encephalopathy induced by hyperemesis gravidarum. *BMJ Case Rep* 2012;2012:bcr2012006216.
- 17 Duca J, Lum CJ, Lo AM. Elevated lactate secondary to gastrointestinal Beriberi. *J Gen Intern Med* 2016;31:133–6.
- 18 Moulin V, Grandoni F, Castioni J, *et al.* Pancytopenia in an adult patient with thiamine-responsive megaloblastic anaemia. *BMJ Case Rep* 2018;2018:bcr-2018-225035.
- 19 Ogershok PR, Rahman A, Nestor S, *et al.* Wernicke encephalopathy in nonalcoholic patients. *Am J Med Sci* 2002;323:107–11.
- 20 Weidauer S, Nichtweiss M, Lanfermann H, *et al.* Wernicke encephalopathy: MR findings and clinical presentation. *Eur Radiol* 2003;13:1001–9.
- 21 Caine D, Halliday GM, Kril JJ, *et al.* Operational criteria for the classification of chronic alcoholics: identification of Wernicke's encephalopathy. *J Neurol Neurosurg Psychiatry* 1997;62:51–60.
- 22 Kudru CU, Nagiri SK, Rao S. Wernicke's encephalopathy in a patient with gastric carcinoma: a diagnosis not to miss. *BMJ Case Rep* 2015.
- 23 Zuccoli G, Gallucci M, Capellades J, *et al.* Wernicke encephalopathy: MR findings at clinical presentation in twenty-six alcoholic and nonalcoholic patients. *AJNR Am J Neuroradiol* 2007;28:1328–31.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow