Published in final edited form as:

J Dev Orig Health Dis. 2019 April ; 10(2): 176–187. doi:10.1017/S2040174418000430.

Perinatal exposures to phthalates and phthalate mixtures result in sex-specific effects on body weight, organ weights, and IAP DNA methylation in weanling mice

Kari Neier^{1,δ}, Drew Cheatham^{1,δ}, Leah D. Bedrosian², and Dana C. Dolinoy^{1,3}

¹University of Michigan School of Public Health, Department of Environmental Health Sciences, Ann Arbor, MI

²University of Michigan School of Public Health, Department of Epidemiology, Ann Arbor, MI

³University of Michigan School of Public Health, Department of Nutritional Sciences, Ann Arbor, MI

Abstract

Developmental exposure to phthalates has been implicated as a risk for obesity; however, epidemiological studies have yielded conflicting results and mechanisms are poorly understood. An additional layer of complexity in epidemiological studies is that humans are exposed to mixtures of many different phthalates. Here, we utilize an established mouse model of perinatal exposure to investigate the effects of three phthalates, diethylhexyl phthalate (DEHP), diisononyl phthalate (DINP), and dibutyl phthalate (DBP), on body weight and organ weights in weanling mice. In addition to individual phthalate exposures, we employed two mixture exposures: DEHP+DINP and DEHP+DINP+DBP. Phthalates were administered through phytoestrogen-free chow at the following exposure levels: 25mg DEHP/kg chow, 25 mg DBP/kg chow, and 75 mg DINP/kg chow. The *A^{vy}* mouse strain, along with measurement of tail DNA methylation, was used as a biosensor to examine effects of phthalates and phthalate mixtures on the DNA methylome. We found that female and male mice perinatally exposed to DINP alone had increased body weights at PND21, and that exposure to mixtures did not exaggerate these effects. Females exposed to DINP and DEHP+DINP had increased relative liver weights at PND21, and females exposed to a mixture of DEHP+DINP+DBP had increased relative gonadal fat weight. Phthalate-exposed *A^{vy/a}* offspring exhibited altered coat color distributions and altered DNA methylation at intracisternal a-particles (IAPs), repetitive elements in the mouse genome. These findings provide evidence that developmental exposures to phthalates influence body weight and organ weight changes in early life, and are associated with altered DNA methylation at IAPs.

Graphical abstract

^δThese authors contributed equally to this work

CONFLICT OF INTEREST

KN, DC, LDB, and DCD declare no conflicts of interest.

ETHICAL STANDARDS

The authors assert that all procedures contributing to this work comply with the ethical standards of the Guide for the Care and Use of Laboratory Animals and has been approved by the University of Michigan Institutional Animal Care and Use Committee (IACUC).

Keywords

Phthalates; chemical mixtures; epigenetics; DNA methylation

INTRODUCTION

Endocrine disrupting chemicals (EDCs) are a class of compounds that are capable of interfering with hormone activity. Exposure to EDCs during development has been linked to increased risk of developing disease in adulthood, following the Developmental Origins of Health and Disease (DOHaD) paradigm.¹ Phthalates are one such example. Phthalates are chemicals added to plastics and personal care products and are present in a wide variety of consumer products, resulting in ubiquitous human exposure.^{2,3} Exposure to high levels of phthalates during the period of reproductive tract development leads to malformations in the reproductive tract, including hypospadias and cryptorchidism in males as demonstrated by both animal and human studies,^{4,5} and dysregulated ovarian function in females in rodent studies.⁶ More recently, human and animal studies have suggested that developmental exposure to lower levels of phthalates may increase the risk of obesity and other features of metabolic syndrome, such as insulin resistance.^{7–10} However, epidemiological studies have presented conflicting evidence with some studies demonstrating associations between early-life phthalate exposure and increased risk for obesity and insulin resistance, while other studies found no associations or negative associations.^{11,12} One factor that may contribute to inconsistencies in epidemiological literature is difficulty in interpreting effects of exposures to phthalate mixtures.^{13,14} Humans are exposed to mixtures of several phthalates, as evidenced by data from the National Health and Nutrition Examination Survey (NHANES) indicating that metabolites from 13 different phthalates were detected in urine samples from a representative study population from the US.¹⁵ Thus, animal experiments utilizing a mixtures approach to characterize metabolic outcomes resulting from developmental phthalate exposures will aid in the interpretation of human epidemiological studies.

Phthalate exposure during development may influence risk of reproductive and metabolic effects by altering the epigenome. The epigenome consists of heritable factors that influence chromatin structure and gene expression without a change in the underlying DNA sequence. DNA methylation at CpG sites is the most widely studied type of epigenetic factor, and

consists of a methyl group attached to the 5' position of a cytosine adjacent to a guanine (C-G dinucleotide). DNA methylation marks are inherited across mitotic cellular divisions and are relatively stable in somatic cells. However, during development, the epigenome is extensively reprogrammed, undergoing waves of DNA de-methylation and re-methylation during pre-implantation and during gametogenesis.^{16,17} Thus, exposure to environmental chemicals that interfere with this process may result in an altered epigenome that persists throughout the life course.^{18,19} Developmental phthalate exposures have been shown to impact DNA methylation levels at loci near genes that are relevant to hormone signaling, including reproductive and metabolic signaling, in germ cells and offspring in both animal and human studies.^{20–24} For example, a recently published birth cohort study identified 27 differentially methylated regions (DMRs) in cord blood from newborns that were associated with gestational exposure to phthalates; many of these DMRs were associated with exposure to multiple phthalates.²³ However, no studies to date have examined the mixture effects of developmental phthalate exposures on DNA methylation.

Repetitive elements are particularly susceptible to epigenetic re-programming via developmental exposures, including intracisternal A-particle (IAP) retrotransposons in mice.^{18,25} IAP retrotransposon repetitive elements are present throughout the mouse genome (several thousand IAPs have been detected in the mouse genome),^{26,27} and contain long terminal repeats (LTRs) carrying CpG sites. DNA methylation at these CpG sites influences expression and regulation of nearby genes, which could have implications for disease risk. One such IAP is found in the cryptic promoter region of the *A^{vy}* allele in the viable yellow agouti mouse model (*A^{vy}*).²⁸ The *A^{vy}* mouse model and its use in environmental epigenetics have been thoroughly described in previous studies and literature reviews.^{29–31} Briefly, DNA methylation levels at CpG sites within the IAP LTR in the promoter region of the *A^{vy}* locus are highly correlated with *A^{vy}* expression and therefore with the coat color of the mouse; yellow fur corresponds to low DNA methylation and high ectopic *agouti* expression, and brown fur corresponds to high DNA methylation and low *agouti* expression. DNA methylation at the *A^{vy}* locus is stochastic and set during early development. Therefore, the coat colors of mice in a colony are approximately normally distributed. However, coat color distributions can be shifted towards brown or yellow in mice exposed to toxicants or nutrients during early development, acting as a biosensor for altered DNA methylation.^{28,29}

Here, we utilized an established mouse model of developmental exposure to evaluate effects of three individual phthalates and two phthalate mixtures on weaning body weight, organ weight, coat color shift, and tail DNA methylation at a specific IAP (*A^{vy}*) and IAPs on a global scale. We chose to examine dietary exposure of two high molecular weight (HMW) phthalates, diethylhexyl phthalate (DEHP) and diisononyl phthalate (DINP), and one low molecular weight (LMW) phthalate, dibutyl phthalate (DBP), as well as a mixture of the two HMW phthalates (DEHP+DINP) and a mixture of all three phthalates (DEHP+DINP+DBP). We evaluated offspring at postnatal day 21 (PND21), which is roughly equivalent to early childhood, in order to provide comparisons to human birth cohort studies that examine metabolic outcomes and DNA methylation in early childhood. To our knowledge, this is the first study to characterize the effects of developmental exposure to phthalate mixtures on weaning body weight, organ weights, and DNA methylation in a mouse model.

MATERIALS AND METHODS

Animals and Exposures

This study utilized viable yellow agouti (A^{vy}) mice from a colony maintained for >220 generations with sibling mating and forced heterozygosity for the A^{vy} allele that is passed paternally, resulting in a genetically invariant background, 93% similar to C57BL/6.^{32,33} A^{vy}/a mice range in coat color from yellow to brown based on epigenetic modifications present at the A^{vy} locus.^{28,31,32,34} Mice were exposed to phthalates perinatally through maternal dietary consumption, with an exposure period that spanned pre-conception to weaning; Figure 1 contains a schematic describing the experimental design of the mouse perinatal exposure study. Two weeks prior to mating, virgin a/a dams at six to eight weeks of age were randomly assigned to one of six diets: 1) 7% corn oil control (phytoestrogen-free Teklad diet TD-95092, ENVIGO, Madison, WI), 2) 25mg DEHP/kg chow, 3) 25mg DBP/kg chow, 4) 75mg DINP/kg chow, 5) 25mg DEHP+75mg DINP/kg chow, or 6) 25mg DEHP +75mg DINP+25mg DBP/kg chow. Dams (a/a) were mated with A^{vy}/a males two weeks after being placed on the experimental diets. Dams remained on experimental diets throughout gestation and lactation until weaning. Birth was designated postnatal day 0 (PND0), and pups were weaned three weeks after birth at postnatal day 21 (PND21).

Exposure diets were based on ENVIGO Teklad diet TD-95092, with phthalate diesters (Sigma) mixed into the corn oil used for the chow. Concentrations in chow were intended to result in estimated maternal doses of 5 mg/kg-day for DEHP and DBP and 15 mg/kg-day for DINP, based on the assumption that pregnant and nursing female mice weigh approximately 25 grams and eat approximately five grams of chow per day.³⁵ These target doses were selected based on studies in the published literature that observed obesogenic effects, including increased body weight, in animals perinatally exposed to phthalates.^{36–38} At the beginning of the study, there were no such studies that found metabolic changes consistent with obesogenic effects for DINP; therefore, a three-fold higher target dose of DINP was selected based on its three-fold lower anti-androgenic potency relative to DBP and DEHP.³⁹ Although fetal exposure was not directly assessed in this study, a previously published study⁴⁰ administered 11 mg/kg-day of DEHP to pregnant rats and found mean levels of DEHP metabolites of 68 ng/mL in amniotic fluid. Amniotic fluid phthalate concentrations were likely lower than 68 ng/mL for the present study (estimated daily intake of 5 mg/kg-day), and would therefore be within the range of measured amniotic fluid levels in human populations,^{41–47} albeit on the higher end (Table 1).

A total of 98 litters were generated, with 17 control litters, 16 DEHP litters, 15 DBP litters, 15 DINP litters, 16 DEHP+DINP litters, and 19 DEHP+DINP+DBP litters. Approximately half of the total offspring were A^{vy}/a (50.5%, n=312) and half were a/a (49.5%, n=296). One DEHP litter (8 pups) and one DEHP+DINP+DBP litter (12 pups) were removed from further analyses based on high pup mortality between PND14 and PND21, which was determined to be due to poor maternal care after animal handlers examined pups. Maternal body weights were recorded at the start of exposure (two weeks prior to mating), at mating, and at time of weaning.

At weaning on PND21, body weights were recorded for all *a/a* and *A^{VY}/a* offspring, and tail tips were collected from all *A^{VY}/a* offspring. A subset of *a/a* mice (n=152) were sacrificed for organ and tissue collection, including tail tips, at PND21. Organs were not collected or analyzed from *A^{VY}/a* mice. For organ and tissue collection, mice were fasted during the light cycle for four hours beginning in the morning, and then underwent CO₂ euthanasia and cardiac puncture in the afternoon, followed by whole-body perfusion with saline (Sigma). Body weights were taken both pre- and post-fasting. Organs were weighed at the time of collection. On postnatal PND21, a single observer classified each *A^{VY}/a* offspring (n=311) into one of five categories based on the proportion of brown fur (yellow, <5% brown; slightly mottled, between 5 and 40% brown; mottled, between 40 and 70% brown; heavily mottled, between 70 and 95% brown; pseudoagouti, >95% brown). Coat color classifications were used as a visual proxy and preliminary assessment of DNA methylation at the *A^{VY}* allele.

All animals were given food and water ad libitum and kept on a 12-hour light/dark cycle. The daily health of the animals was monitored by the University Unit for Laboratory Animal Medicine (ULAM). Mice were treated humanely, and the guidelines for the use and care of laboratory animals were followed throughout the study. The University of Michigan Institutional Animal Care and Use Committee (IACUC) approved all animal procedures used for this project.

Hepatic Triglyceride Levels

Hepatic triglyceride levels were measured in wildtype non-agouti *a/a* mice that were sacrificed at PND21. Frozen liver tissue (30-35 mg) was homogenized in lysis buffer containing 10% NP-40 (ThermoFisher Scientific), 50 mM Tris-HCl (Sigma), and 100 mM NaCl, and then triglycerides were extracted using chloroform (Fisher). Triglyceride levels were measured using spectrometry; Infinity Triglycerides reagent (ThermoFisher Scientific, Waltham, MA) was added to each sample and then read at 550nm and 660nm and compared to glycerol standards to estimate triglycerides per milligram of liver tissue.

DNA Isolation and Methylation Analysis

Genomic DNA was isolated from PND21 tail tissue, flash frozen, and stored at -80°C from all *A^{VY}/a* and *a/a* offspring. Approximately 0.5cm of thawed distal tail tissue was placed in a tube containing 173.32 µL of Buffer ATL and 26.67 µL of Proteinase K and then incubated overnight at 50°C on a shaker. Samples were then cooled to room temperature before transferring to the Maxwell® 16 Mouse Tail DNA Purification Kit cartridges (Promega Corporation, Madison, WI) using manufacturers' standard protocol. One modification was made in which only 250 µL of elution buffer was added to the final elution tube to maximize final DNA concentration. DNA concentration was quantified using the NanoDrop 2000 Spectrophotometer (ThermoFisher Scientific, Waltham, MA); concentrations ranged from 50-100 ng/µL.

Genomic DNA was bisulfite converted using the EZ-96 DNA Methylation kit protocol (Zymo Research Corp, Irvine, CA), which converts all unmethylated cytosines to uracils and methylated cytosines are left unaffected.⁴⁸ For *A^{VY}/a* samples (n=302), 500 ng of DNA was

used as input for conversion and 1000 ng was inputted for all *a/a* samples (n=152); both 500 ng and 1000 ng are considered acceptable according to the EZ-96 DNA Methylation Kit, but 500 ng was used for *A^{VY}/a* samples due to lower tissue amounts and subsequent DNA concentrations extracted from some samples. After bisulfite conversion, amplification of *A^{VY}* and global IAP regions was performed by standard PCR in order to attain enough DNA to pyrosequence. Each PCR reaction well included 19.4 μ L of HotStarTaq DNA Polymerase (Qiagen Inc., Germantown, MD), 0.12 μ L forward primer (Invitrogen, Waltham, MA), 0.12 μ L biotinylated-reverse primer (Invitrogen, Waltham, MA), and 13.36 μ L H₂O. Two microliters of bisulfite converted DNA were added to each well to complete the 35 μ L reaction. Each PCR run included a no-template control and laboratory methylation controls (0%, 25%, 50%, 75%, 100%). Assay specifications are included in Table 2. PCR amplification lengths were confirmed by the Qiagen QIAxcel Advanced System (Qiagen) and QIAxcel ScreenGel software.

Pyrosequencing with the PyroMark Q96 ID (Qiagen) was used to assess DNA methylation percent at four CpG sites of interest for *A^{VY}* and IAP regions on a global scale. The global IAP assay covers all IAPs on the mouse genome with intact LTRs, including the *A^{VY}* IAP. The protocol followed is outlined in the PyroMark Q96 ID User Manual. PyroMark Software determines the percentage of methylation by calculating the fraction of methylated cytosines (read as cytosine after PCR) over the total sum of methylated and unmethylated cytosines (read as thymines after PCR). The amount of input DNA used for both assays was 16 μ L. All *A^{VY}* and global IAP assay specifications and pyrosequencing information are available in Table 2.

Statistical Analysis

All statistical analyses were conducted using R version 3.3.1 (www.R-project.org). Exposure-related differences in sex ratio, genotype ratio, and survival rate were determined by Fisher's exact test, comparing each exposure group to the control group. For sex and genotype ratios, exposure groups were also compared to expected values of 50%, since these are the values expected based on standard genetic probabilities. Coat color distribution across exposure groups was compared via the χ^2 goodness-of-fit test with the control coat color distribution as the reference. Coat color distribution analyses could not be stratified by sex because doing so yields some coat color categories within exposure groups having a count of zero, making it difficult to make interpretable statistical comparisons. Differences in litter size across the six experimental groups were compared using an ANOVA with a post-hoc Tukey test. Body weight, organ weights, and DNA methylation for each exposure group were compared to controls using linear mixed effects models with the lme4 and lmeTest packages in R. Mixed effects models were used to account for inherent correlations between littermates, and were also used to account for correlations between DNA methylation levels across nearby CpG sites. Number of pups in each litter was added into models evaluating exposure effects on body weight to control for litter size. Tail *A^{VY}* methylation levels were bimodally distributed, and we therefore split *A^{VY}* methylation data into high and low methylation groups, based on the median methylation of the control group (i.e., low methylation = methylation level below the control median, and high methylation = methylation level above the control median) in order to make statistical comparisons.³² All

analyses were stratified by genotype so that comparisons were made across isogenic mice. Analyses were also stratified by sex if initial analyses indicated that sex modified exposure effects.

RESULTS

Litter Parameters

Perinatal exposure to phthalates did not significantly alter genotype ratio, sex ratio, or mean pups per litter compared to controls, but did impact mortality rate in pups through weaning at PND21 (Table 3). Overall mortality rate in controls was 8.3%. Compared to controls, there was a reduced pup mortality rate in the DBP exposure group (mean=1.1%, $p=0.02$), and increased pup mortality rate in the DEHP exposure group at a level trending towards significance (mean=15.5%, $p=0.10$). One litter exposed to DEHP and one exposed to a mixture of all three phthalates were removed from subsequent analyses due to high pup mortality between PND14 and PND21 which was attributed to poor maternal care and hyperactive maternal behavior.

Body Weight

Mice that were perinatally exposed to phthalates and phthalate mixtures had increased body weights at PND21 (Figure 2). Different phthalate exposures exhibited significant impacts on body weight, depending on sex and genotype. Perinatal exposures to DINP and a mixture of all three phthalates were associated with increased body weights in females (Figure 2A). Female *a/a* and *A^{vy}/a* mice exposed to DINP weighed more than female *a/a* and *A^{vy}/a* control mice ($p = 0.03$, $p = 0.01$, respectively). Wild-type *a/a* and *A^{vy}/a* females exposed to DINP weighed an average of 8.44 and 9.29 grams, respectively, whereas *a/a* and *A^{vy}/a* control females weighed an average of 7.65 grams and 7.77 grams, respectively. Females of both genotypes exposed to a mixture of DEHP+DINP+DBP also had increases in body weight at PND21 that approached statistical significance ($p = 0.08$ for both genotypes); *a/a* females exposed to all three phthalates weighed 8.42 grams on average and *A^{vy}/a* females exposed to all three phthalates weighed 9.13 grams on average.

In males, phthalate-related body weight effects at PND21 were more pronounced in *A^{vy}/a* mice than in *a/a* mice, with more exposure groups exhibiting significant differences versus controls in *A^{vy}/a* mice than in *a/a* mice (Figure 2B). Male *A^{vy}/a* mice that were exposed to DEHP, DINP, and DEHP+DINP weighed more than male *A^{vy}/a* controls ($p=0.02$, $p=0.02$, and $p=0.03$, respectively). *A^{vy}/a* males exposed to DEHP weighed an average of 9.38 grams, and those exposed to DINP and DEHP+DINP weighed an average of 9.26 grams, while control *A^{vy}/a* males weighed an average of 8.22 grams. In addition, increased body weights observed in male *A^{vy}/a* mice exposed to DEHP+DINP+DBP (mean = 9.32 grams) approached statistical significance ($p=0.06$). On the other hand, the only difference in body weight across exposure groups in *a/a* males was an increase in body weight in DINP *a/a* males relative to control *a/a* males that trended towards significance (mean = 8.91 and 7.74 grams, respectively; $p=0.07$).

Maternal (F₀) body weights were not significantly different in exposure groups versus the control group at the time of exposure, time of mating, or time of weaning, with or without controlling for number of pups given birth to ($p>0.10$) (Supplemental Table S1). However, dams exposed to DINP gained more body weight between mating and weaning than controls at a level near statistical significance ($p=0.06$) when controlling for the number of pups per litter; dams exposed to DINP gained 9.77 grams whereas control dams gained 8.16 grams. Dams exposed to DINP also gained more weight across the duration of the study (from exposure onset to weaning) than controls ($p=0.01$), and dams exposed to DBP gained more weight than controls between exposure and weaning at levels nearing statistical significance ($p=0.08$).

Organ and Tissue Weights

Several organs and tissues were collected from fasted *a/a* mice at PND21; full details on organ and tissue weights can be found in Supplemental Table S2. The most prevalent effect of developmental phthalate exposures on organ weight was observed in female livers (Figure 3; Supplemental Table S2). Females exposed to DINP, and DEHP+DINP had increased relative liver weights, expressed as percent of body weight, compared to control females ($p=0.02$ and $p=0.006$, respectively). DEHP+DINP+DBP exposed female offspring also exhibited increased relative liver weights vs. controls at a level near statistical significance ($p=0.08$). Mean relative liver weights of DINP, DEHP+DINP, and DEHP+DINP+DBP exposed females were an average of 4.17%, 4.25%, and 3.99%, respectively, whereas control females had a mean relative liver weight of 3.61%. Hepatic triglyceride levels were measured to determine whether differences in relative liver weights were due to an increase in lipid accumulation in the liver; however, we did not observe any significant differences in hepatic triglycerides across exposure groups (data not shown).

Relative gonadal fat weight was also significantly higher in females exposed to all three phthalates compared to control females ($p=0.02$); gonadal fat from females exposed to DEHP+DINP+DBP weighed 0.15% of body weight whereas gonadal fat from control females weighed 0.10% of body weight. Relative brain weights were lower in females exposed to DEHP+DINP+DBP compared to controls ($p=0.04$), and decreases observed in relative brain weights in DINP and DEHP+DINP+DBP exposed males approached statistical significance ($p=0.07$ for both). Absolute brain weights, however, were not significantly different across exposure groups, with mean brain weights that ranged from 288.7 grams to 305.3 grams for all *a/a* mice. Males exposed to all three phthalates had lower absolute pancreas weights than controls ($p=0.05$), and females exposed to all three phthalates had lower absolute pancreas weights than controls at nearly statistically significant levels ($p=0.07$), but alterations observed in relative pancreas weights did not reach statistical significance. An increase observed in absolute spleen weights between females exposed to all three phthalates compared to controls approached statistical significance ($p=0.08$), but relative spleen weights were not significantly different across exposure groups. Lastly, kidney weights were not impacted by developmental phthalate exposure ($p>0.10$).

Coat Color Distribution

Developmental exposure to phthalates resulted in altered coat color distributions in A^{vy}/a offspring. Exposure to DEHP, DBP, DEHP+DINP, and DEHP+DINP+DBP significantly altered coat color distributions ($p=0.02$, $p<0.0001$, $p=0.02$, $p<0.0001$, respectively; Figure 4), while exposure to DINP resulted in only a modest change in coat color distribution compared to controls ($p=0.10$). Developmental exposure to DBP alone (Figure 4C) and in the mixture (DEHP+DINP+DBP, Figure 4F) exhibited a pronounced increase in the proportion of pseudoagouti offspring; 6.12% of the control offspring were classified as pseudoagouti, compared with 32.7% of the DBP offspring and 17.9% of the DEHP+DINP+DBP offspring. DEHP exposure (Figure 4B) showed the most pronounced increase in proportion of slightly mottled offspring, represented by 44.9% compared to 26.5% of the control offspring. There was also a relative increase in slightly mottled offspring among the DEHP+DINP exposed group (32.7%), and the DBP exposed group (36.7%).

A^{vy} DNA Methylation

Molecular analyses of DNA methylation at A^{vy} were used to corroborate the coat color assignments. Mean DNA methylation percent at four CpG sites within the 5' IAP LTR in the promoter of the A^{vy} allele corresponded appropriately to the coat color phenotypes (Supplemental Figure S1). Mice classified as having a yellow coat color had methylation levels at or approaching 0% (mean=0.73%), and increasing amounts of brown fur corresponded with increasing amounts of methylation. Pseudoagouti offspring had the highest amount of methylation (mean=74.4%). Mean A^{vy} methylation levels across exposure groups, with and without sex stratification, can be found in Table 4. Distributions of mean methylation for each exposure were not normally distributed, and in many exposure groups followed a bimodal distribution (Supplemental Figure S2). Thus, methylation for each exposure group was dichotomized at the median methylation value of the control group (29.9=low-methylation, >29.9=high-methylation) to more appropriately compare mean methylation values. The high-methylation group in DBP exposed females was significantly greater than control females, with a mean A^{vy} methylation of 67.37% in highly methylated DBP exposed females and a mean A^{vy} methylation of 50.03% in highly methylated control females ($p=0.006$). The high-methylation group in females perinatally exposed to DEHP+DINP+DBP also had modestly increased A^{vy} DNA methylation compared to control females (60.45% versus 50.03%, respectively; $p=0.07$). There were no significant differences ($p<0.10$) observed in males, or among the low-methylation categories.

Global IAP DNA Methylation

After observing phthalate-related differences in DNA methylation patterns at the A^{vy} IAP, we measured global IAP methylation levels in tail tips from A^{vy}/a ($n=302$) and a/a mice ($n=152$) to further explore whether developmental phthalate exposures are capable of altering DNA methylation on a global level. Developmental phthalate exposure altered IAP methylation on a global scale, in a sexually dimorphic manner. Developmental phthalate exposure tended to result in decreased IAP DNA methylation in males and increased IAP DNA methylation in females (Figure 5). Compared to controls, male A^{vy}/a mice exposed to DBP had a 1.4% decrease in mean global IAP methylation and DEHP+DINP had a 1.5%

decrease in mean global IAP methylation across four CpG sites assayed by pyrosequencing ($p=0.03$, $p=0.01$, respectively). In contrast, female A^{VY}/a mice exposed to DEHP had a 1.9% increase and DEHP+DINP had a 2.2% increase mean DNA methylation relative to controls ($p=0.03$, $p=0.01$, respectively). Female A^{VY}/a mice exposed to DINP had increased global IAP DNA methylation at levels approaching statistical significance when compared to controls ($p=0.09$). Trends in mean and site-specific DNA methylation levels were similar in a/a mice, but were mostly non-significant, possibly due to lower statistical power (Supplemental Figure S3).

DISCUSSION

Perinatal exposure to phthalates resulted in sex-specific and phthalate-specific effects on weaning body weight, organ weights, and tail DNA methylation at IAPs. Exposure to phthalates in mixture form did not appear to have an exaggerated effect for most outcomes tested, with the exception of certain organ and tissue weights. In general, developmental phthalate exposure was associated with an increase in PND21 body weight, especially in mice exposed to DINP. The DINP exposure group showed the most consistent effects on body weight, with both genotypes and both sexes having increased body weight relative to controls; in a/a males, the only group that was notably different compared to controls was the group exposed to only DINP. To our knowledge, this study was the first to report on body weight effects in mice following developmental exposures to DINP. Thus, the consistent increases in PND21 body weight we observed in mice developmentally exposed to DINP is a novel finding. One study in rats reported an association between *decreased* body weight gain in adult rats that were exposed to DINP from preconception to weaning; however, the exposure levels investigated in that study were much higher than those used in the present study (0.5% to 1.5% DINP in chow vs. 0.075% in our study).⁴⁹ Another study in rats similarly found that exposure to DINP from GD15 to PND10 resulted in decreased weight gain between PND2 and PND10, but the exposure levels examined were also relatively high (20,000 ppm or 2% in chow).⁵⁰ EDCs have been demonstrated to exhibit non-monotonic responses and low-dose effects.⁵¹ It is possible that increased body weight is only associated with lower-level developmental exposures of DINP, such as the levels used in this study.

Only A^{VY}/a males had increased body weight following developmental exposure to DEHP alone. Other studies on developmental DEHP exposure in mice have demonstrated mixed effects on body weight, with some studies showing an increase and some a decrease.^{36,52} Mice that were exposed to DBP alone did not have significantly different body weights from controls for either genotype or sex. This study is the first to examine body weight effects following developmental DBP exposure in mice, but studies in rats have also demonstrated mixed effects on offspring body weight.^{37,53,54}

This is the first animal study to examine the impact of developmental exposures to mixtures of phthalates on body weight. Exposure to phthalate mixtures did not appear to result in an exaggerated effect on PND21 body weight; females exposed to DEHP+DINP did not weigh significantly more than controls, and those exposed to DEHP+DINP+DBP weighed more than controls to a moderate degree of statistical significance. In addition, male A^{VY}/a mice

exposed to DEHP+DINP during development had increased PND21 body weights relative to controls, and those exposed to DEHP+DINP+DBP had a modest trend towards increased body weight compared to controls, but male *a/a* mice exposed to phthalate mixtures did not have increased body weight relative to controls. It is possible that mixture effects would be more apparent at lower exposures, especially if increased body weight is more prominent at lower exposures of phthalates. Since the mice exposed to mixtures of phthalates had higher total phthalate exposures, it is possible that any additive effects, or lack thereof, could be due to dose effects rather than mixture effects.

Because body composition was not assessed in this analysis, it is unknown whether the observed increases in body weight in phthalate-exposed mice were due to an increase in body fat, lean mass, or both. Additional studies are needed with inclusion of different dosing levels to confirm whether developmental exposures to phthalate mixtures have exaggerated effects on body weight, as well as on body composition. A follow-up study of a subset of wildtype non-agouti *a/a* mice that were aged to 10 months and includes body composition analysis is currently underway in order to assess long-lasting impacts and to determine whether perinatal phthalate exposures impact body fat or lean mass.

Multiple organ and tissue weights were altered by developmental exposures to phthalates in *a/a* mice, and females appeared to be more sensitive than males in this respect. Relative liver weights were increased in females exposed to DINP and DEHP+DINP compared to controls, and were increased in females exposed to DEHP+DINP+DBP at levels trending towards statistical significance. Phthalates have been demonstrated to activate PPAR- α , the dominant PPAR in the liver, in other animal and *in vitro* studies,^{55–57} and PPAR- α activation has been associated with increased relative liver weights in investigations of other chemicals.⁵⁸ The observed increases in relative liver weights of phthalate-exposed female offspring were likely not due to an increase in lipid accumulation in the liver since hepatic triglyceride levels did not differ across exposure groups. Thus, these findings are consistent with the idea that developmental exposure to DINP and DINP mixed with DEHP activate PPAR- α in the liver, and indicate that females may be more sensitive at this early developmental time point. The mixture of DEHP+DINP had a slightly larger impact on relative liver weights, but not at a statistically significant level, and the mixture of DEHP+DINP+DBP had only a modest effect on relative liver weights compared to controls. Thus, phthalate mixtures at the exposure levels tested in this study do not appear to have exaggerated effects on relative liver weights at PND21.

On the other hand, the DEHP+DINP+DBP exposure group was the only group that exhibited an alteration in relative gonadal fat weight at PND21, with females showing an increase relative to controls. This observation is consistent with the theory that phthalate exposure increases the risk of obesity, and indicates that a mixture of HMW and LMW phthalates have the largest impact on body fat at PND21, with females being more sensitive than males. A recent meta-analysis of developmental DEHP exposure in animal studies also indicated that early life phthalate exposures results in increased fat weight.⁵⁹ Absolute pancreas weights were increased in mice perinatally exposed to DEHP+DINP+DBP compared to controls, but this difference was mitigated when comparing pancreas weights relative to body weight, and therefore the increased pancreas weights might have been due to the

modest increased body weights observed in DEHP+DINP+DBP-exposed females. Decreased relative brain weights were observed in females exposed to a mixture of all three phthalates, and to a lesser degree of significance in males exposed to all three phthalates. However, these findings are skewed by the significant and non-significant increases in body weights observed in the mice in these exposure groups, since absolute brain weights did not differ by exposure.

This study was the first to utilize the A^{VY} mouse model as a biosensor to determine whether developmental exposures to phthalates and phthalate mixtures are capable of altering epigenetic marks at repetitive elements in the genome, such as the IAP located in the A^{VY} promoter region. Coat color distributions were significantly altered in all groups exposed to phthalates, except for the group exposed to only DINP. Mixtures did not appear to have an exaggerated effect on offspring coat color. Coat color distributions for DEHP and DEHP +DINP were extremely similar to one another, with marked increases in slightly mottled mice in both exposure groups. Coat color distributions for DBP and DEHP+DINP+DBP were also similar to one another, with prominent increases in pseudoagouti mice in both groups. The shift towards pseudoagouti coat color may indicate a protective effect against the yellow obese phenotype in mice perinatally exposed to DBP and DEHP+DINP+DBP; however, we did not follow the $A^{VY/a}$ mice into adulthood, when coat color-associated metabolic phenotypes emerge, to confirm this. In addition, developmental exposures to phthalates may influence metabolism through other mechanisms distinctive from altered DNA methylation at the A^{VY} locus. It should be noted, however, that while A^{VY} hypermethylation in the viable yellow agouti mouse model is likely protective against obesity, hypermethylation in humans is context dependent, and could result in protective, adaptive, or null effects depending on which regulatory or genic region is affected. These data suggested that DEHP and DBP both influenced shifts in coat color, and that DEHP, a HMW phthalate, had different effects from DBP, a LMW phthalate. Consistent with previous studies utilizing the A^{VY} mouse model to investigate environmental exposures, ^{29,60,61} we confirmed that coat colors were representative of methylation at the A^{VY} locus via tail DNA. Our analyses of DNA methylation at the A^{VY} locus indicated that there was an increase in methylation in the DBP exposure group, which was consistent with the observed increase in pseudoagouti mice.

Tail DNA methylation levels at IAPs were also altered by developmental phthalate exposures on a global level, and in a sexually dimorphic manner. We observed a general trend towards increased global IAP methylation in females exposed to phthalates, but in males exposed to phthalates we observed a general trend towards decreased global IAP methylation. In addition, different individual phthalates had more pronounced effects on global IAP methylation in females versus males. In $A^{VY/a}$ mice, developmental exposure to DEHP, and DINP to a lesser degree of significance, resulted in altered global IAP DNA methylation in females, whereas developmental exposure to DBP resulted in altered DNA methylation in males. Phthalate mixtures effects were complex. The DEHP+DINP mixture group had altered DNA methylation in both sexes, while the DEHP+DINP+DBP mixture group did not exhibit significant differences in either sex. Effect sizes for exposure-related changes in global IAP methylation were between 1.4 and 2.2%, and while small, are comparable to effect sizes for DNA methylation at repetitive elements frequently associated

with developmental exposures to other environmental chemicals.⁶² Similar trends in tail global IAP DNA methylation were observed in *a/a* mice, but we observed less statistically significant differences across exposure groups, likely due to decreased power resulting from lower sample size. It should also be noted that phthalate-mediated alterations in DNA methylation at repetitive elements are unlikely to be the sole drivers of phthalate-mediated metabolic impacts. Because phthalate metabolites have been demonstrated to activate PPARs,^{55,56} additional studies investigating DNA methylation at the promoter region of PPAR- α and PPAR- γ target genes in liver and adipose, respectively, would provide crucial mechanistic insights.

A^{VY} methylation levels were not predictive of global IAP methylation levels in this study ($R^2 = 0.001$, $p=0.22$ via Pearson correlation test). However, this is not unexpected. The *A^{VY}* locus is just one IAP out of thousands of IAPs that are present in the mouse genome, and previous studies that have demonstrated high variation in DNA methylation across different IAPs.²⁷ Another study examining perinatal lead (Pb) exposure in mice found different impacts of Pb on individual IAPs,⁶³ thus phthalates may also have IAP-specific effects that are not fully detected by examining *A^{VY}* and global IAP methylation levels alone. Based on the findings of this study, evaluation of methylation at additional specific IAPs may be warranted, especially those that are near promoter regions of genes that are known to play a role in metabolism.

Phthalates are ubiquitous chemicals that present a significant risk for human exposure, and recent research has suggested that exposure to physiologically relevant levels may influence risk of obesity and metabolic syndrome. Further, phthalates are known EDCs, and high exposures during development have been linked to permanent reproductive tract abnormalities. However, little is known regarding metabolic health outcomes resulting from developmental exposures to phthalates, and even less is known about risks from developmental exposures to phthalate mixtures. In this study, developmental exposures to phthalates in the perinatal window influenced body weight, liver weight, and gonadal fat weight in mice at weaning. Interestingly, females appeared to be more sensitive than males. Tail IAP DNA methylation at the *A^{VY}* locus and on a global level were altered by developmental phthalate exposures, presenting a possible mechanistic link between early-life exposures and potential later-life health outcomes; research focused on characterizing long-term metabolic impacts on a subset of *a/a* mice is currently underway in order to evaluate this potential. The results from this study are the first to describe metabolic and epigenetic effects of developmental exposure to phthalate mixtures. Mixture effects were complex in that they did not necessarily result in exaggerated effects, and in some cases, phthalate mixtures exhibited attenuated effects. However, this may be attributed to dose effects rather than mixture effects, since the groups receiving mixtures of phthalates received higher exposures to total phthalates. Future work investigating phthalate mixtures at multiple dose levels is needed to better understand these effects, which will aid in the interpretation of human epidemiological studies and further our understanding of molecular mechanisms.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to sincerely thank Christine Rygiel and Nicole Urdahl for their assistance with carrying out the global IAP pyrosequencing assays, as well as Elizabeth H. Marchlewicz for her guidance in measuring hepatic triglycerides.

FINANCIAL CONTRIBUTIONS

This work was supported by the University of Michigan (UM) NIEHS/EPA Children's Environmental Health and Disease Prevention Center P01 ES022844/RD83543601, the Michigan Lifestage Environmental Exposures and Disease (M-LEEAD) NIEHS Core Center (P30 ES017885), as well as the UM NIEHS Institutional Training Grant T32 ES007062 and NICHD Institutional Training Grant T32 079342 (KN).

References

1. Haugen AC, Schug TT, Collman G, Heindel JJ. Evolution of DOHaD: the impact of environmental health sciences. *J Dev Orig Health Dis*. 2015; 6(2):55–64. [PubMed: 25471238]
2. Schettler T. Human exposure to phthalates via consumer products. *Int J Androl*. 2006; 29(1):134–139. [PubMed: 16466533]
3. Heudorf U, Mersch-Sundermann V, Angerer J. Phthalates: toxicology and exposure. *Int J Hyg Environ Health*. 2007; 210(5):623–34. [PubMed: 17889607]
4. Foster P. Disruption of reproductive development in male rat offspring following in utero exposure to phthalate esters. *Int J Androl*. 2006; 29(1):140–147. [PubMed: 16102138]
5. Hsieh MH, Breyer BN, Eisenberg ML, Baskin LS. Associations among hypospadias, cryptorchidism, anogenital distance, and endocrine disruption. *Curr Urol Rep*. 2008; 9(2):137–142. [PubMed: 18419998]
6. Hannon PR, Flaws JA. The effects of phthalates on the ovary. *Front Endocrinol (Lausanne)*. 2015; 6:8. [PubMed: 25699018]
7. Hao C, Cheng X, Guo J, Xia H, Ma X. Perinatal exposure to diethyl-hexyl-phthalate induces obesity in mice. *Front Biosci (Elite Ed)*. 2013; 5:725–733. [PubMed: 23277027]
8. Yang TC, Peterson KE, Meeker JD, et al. Bisphenol A and phthalates in utero and in childhood: association with child BMI z-score and adiposity. *Environ Res*. 2017; 156:326–333. [PubMed: 28390300]
9. Rajesh P, Balasubramanian K. Phthalate exposure in utero causes epigenetic changes and impairs insulin signalling. *J Endocrinol*. 2014; 223(1):47–66. [PubMed: 25232145]
10. Strakovsky RS, Lezmi S, Shkoda I, et al. In utero growth restriction and catch-up adipogenesis after developmental di (2-ethylhexyl) phthalate exposure cause glucose intolerance in adult male rats following a high-fat dietary challenge. *J Nutr Biochem*. 2015; 26(11):1208–20. [PubMed: 26188368]
11. Buckley JP, Engel SM, Braun JM, et al. Prenatal Phthalate Exposures and Body Mass Index Among 4- to 7-Year-old Children: A Pooled Analysis. *Epidemiology*. 2016; 27(3):449–58. [PubMed: 26745610]
12. Shoaff J, Papandonatos GD, Calafat AM, et al. Early-life phthalate exposure and adiposity at 8 years of age. *Environ Health Perspect*. 2017; 125(9):97008.
13. Braun JM, Gennings C, Hauser R, Webster TF. What can epidemiological studies tell us about the impact of chemical mixtures on human health? *Environ Health Perspect*. 2016; 124(1):A6–A9. [PubMed: 26720830]
14. Braun JM, Just AC, Williams PL, et al. Personal care product use and urinary phthalate metabolite and paraben concentrations during pregnancy among women from a fertility clinic. *J Expo Sci Environ Epidemiol*. 2014; 24(5):459–466. [PubMed: 24149971]
15. Carlson KR, Garland SE. Estimated Phthalate Exposure and Risk to Pregnant Women and Women of Reproductive Age as Assessed Using Four NHANES Biomonitoring Data Sets (2005/2006, 2007/2008, 2009/2010, 2011/2012). 2015
16. Reik W, Dean W, Walter J. Epigenetic reprogramming in mammalian development. *Science*. 2001; 293(5532):1089–93. [PubMed: 11498579]

17. Jirtle RL, Skinner MK. Environmental epigenomics and disease susceptibility. *Nat Rev Genet.* 2007; 8(4):253–62. [PubMed: 17363974]
18. Dolinoy DC, Das R, Weidman JR, Jirtle RL. Metastable Epialleles, Imprinting, and the Fetal Origins of Adult Diseases. *Pediatr Res.* 2007; 61(5 Part 2):30R–37R.
19. Faulk C, Dolinoy DC. Timing is everything: the when and how of environmentally induced changes in the epigenome of animals. *Epigenetics.* 2011; 6(7):791–7. [PubMed: 21636976]
20. Zhang X-F, Zhang L-J, Li L, et al. Diethylhexyl phthalate exposure impairs follicular development and affects oocyte maturation in the mouse. *Environ Mol Mutagen.* 2013; 54(5):354–361. [PubMed: 23625783]
21. Martinez-Arguelles DB, Papadopoulos V. Identification of hot spots of DNA methylation in the adult male adrenal in response to in utero exposure to the ubiquitous endocrine disruptor plasticizer di-(2-ethylhexyl) phthalate. *Endocrinology.* 2015; 156(1):124–33. [PubMed: 25330100]
22. Huen K, Calafat AM, Bradman A, et al. Maternal phthalate exposure during pregnancy is associated with DNA methylation of LINE-1 and Alu repetitive elements in Mexican-American children. *Environ Res.* 2016; 148:55–62. [PubMed: 27019040]
23. Solomon O, Yousefi P, Huen K, et al. Prenatal phthalate exposure and altered patterns of DNA methylation in cord blood. *Environ Mol Mutagen.* 2017; 58(6):398–410. [PubMed: 28556291]
24. Wu H, Estill MS, Shershebnv A, et al. Preconception urinary phthalate concentrations and sperm DNA methylation profiles among men undergoing IVF treatment: a cross-sectional study. *Hum Reprod.* 2017; 32(11):2159–2169. [PubMed: 29024969]
25. Waterland RA, Jirtle RL. Early nutrition, epigenetic changes at transposons and imprinted genes, and enhanced susceptibility to adult chronic diseases. *Nutrition.* 2004; 20(1):63–8. [PubMed: 14698016]
26. Qin C, Wang Z, Shang J, et al. Intracisternal A particle genes: Distribution in the mouse genome, active subtypes, and potential roles as species-specific mediators of susceptibility to cancer. *Mol Carcinog.* 2010; 49(1):54–67. [PubMed: 20025072]
27. Faulk C, Barks A, Dolinoy DC. Phylogenetic and DNA methylation analysis reveal novel regions of variable methylation in the mouse IAP class of transposons. *BMC Genomics.* 2013; 14(1):48. [PubMed: 23343009]
28. Dolinoy DC. The agouti mouse model: An epigenetic biosensor for nutritional and environmental alterations on the fetal epigenome. *Nutr Rev.* 2008; 66(SUPPL.1)
29. Dolinoy DC, Huang D, Jirtle RL. Maternal nutrient supplementation counteracts bisphenol A-induced DNA hypomethylation in early development. *Proc Natl Acad Sci U S A.* 2007; 104(32):13056–61. [PubMed: 17670942]
30. Dolinoy DC, Jirtle RL. Environmental epigenomics in human health and disease. *Environ Mol Mutagen.* 2008; 49(1):4–8. [PubMed: 18172876]
31. Dolinoy DC, Weinhouse C, Jones TR, Rozek LS, Jirtle RL. Variable histone modifications at the A(vy) metastable epiallele. *Epigenetics.* 2010; 5(7):637–44. [PubMed: 20671424]
32. Waterland RA, Jirtle RL. Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol Cell Biol.* 2003; 23(15):5293–300. [PubMed: 12861015]
33. Weinhouse C, Anderson OS, Bergin IL, et al. Dose-dependent incidence of hepatic tumors in adult mice following perinatal exposure to bisphenol A. *Environ Health Perspect.* 2014; 122(5):485–91. [PubMed: 24487385]
34. Miltenberger RJ, Mynatt RL, Wilkinson JE, Woychik RP. The role of the agouti gene in the yellow obese syndrome. *J Nutr.* 1997; 127(9):1902S–1907S. [PubMed: 9278579]
35. Vandenberg LN, Welshons WV, Vom Saal FS, Toutain P-L, Myers JP. Should oral gavage be abandoned in toxicity testing of endocrine disruptors? *Environ Health.* 2014; 13(1):46. [PubMed: 24961440]
36. Schmidt J-S, Schaedlich K, Fiandanese N, Pocar P, Fischer B. Effects of di(2-ethylhexyl) phthalate (DEHP) on female fertility and adipogenesis in C3H/N mice. *Environ Health Perspect.* 2012; 120(8):1123–9. [PubMed: 22588786]
37. Mylchreest E, Wallace DG, Cattley RC, Foster PM. Dose-dependent alterations in androgen-regulated male reproductive development in rats exposed to Di(n-butyl) phthalate during late gestation. *Toxicol Sci.* 2000; 55(1):143–51. [PubMed: 10788569]

38. de Jesus MM, Negrin AC, Taboga SR, Pinto-Fochi ME, Góes RM. Histopathological alterations in the prostates of Mongolian gerbils exposed to a high-fat diet and di-n-butyl phthalate individually or in combination. *Reprod Toxicol*. 2015; 52:26–39. [PubMed: 25697569]
39. Hannas BR, Lambright CS, Furr J, et al. Dose-response assessment of fetal testosterone production and gene expression levels in rat testes following in utero exposure to diethylhexyl phthalate, diisobutyl phthalate, diisooheptyl phthalate, and diisononyl phthalate. *Toxicol Sci*. 2011; 123(1): 206–16. [PubMed: 21633115]
40. Calafat AM, Brock JW, Silva MJ, et al. Urinary and amniotic fluid levels of phthalate monoesters in rats after the oral administration of di(2-ethylhexyl) phthalate and di-n-butyl phthalate. *Toxicology*. 2006; 217(1):22–30. [PubMed: 16171919]
41. Lorber M, Calafat AM. Dose reconstruction of di(2-ethylhexyl) phthalate using a simple pharmacokinetic model. *Environ Health Perspect*. 2012; 120(12):1705–10. [PubMed: 23010619]
42. Wittassek M, Angerer J, Kolossa-Gehring M, et al. Fetal exposure to phthalates—a pilot study. *Int J Hyg Environ Health*. 2009; 212(5):492–8. [PubMed: 19423389]
43. Wittassek M, Angerer J. Phthalates: metabolism and exposure. *Int J Androl*. 2008; 31(2):131–138. [PubMed: 18070048]
44. Chang J-W, Lee C-C, Pan W-H, et al. Estimated Daily Intake and Cumulative Risk Assessment of Phthalates in the General Taiwanese after the 2011 DEHP Food Scandal. *Sci Rep*. 2017; 7:45009. [PubMed: 28327585]
45. Silva MJ, Reidy JA, Herbert AR, et al. Detection of phthalate metabolites in human amniotic fluid. *Bull Environ Contam Toxicol*. 2004; 72(6):1226–31. [PubMed: 15362453]
46. Huang P-C, Tsai C-H, Liang W-Y, et al. Early Phthalates Exposure in Pregnant Women Is Associated with Alteration of Thyroid Hormones. *PLoS One*. 2016; 11(7):e0159398. [PubMed: 27455052]
47. HUANG P, KUO P, CHOU Y, LIN S, LEE C. Association between prenatal exposure to phthalates and the health of newborns☆. *Environ Int*. 2009; 35(1):14–20. [PubMed: 18640725]
48. Grunau C, Clark SJ, Rosenthal A. Bisulfite genomic sequencing: systematic investigation of critical experimental parameters. *Nucleic Acids Res*. 2001; 29(13):E65–5. [PubMed: 11433041]
49. Waterman SJ, Keller LH, Trimmer GW, et al. Two-generation reproduction study in rats given diisononyl phthalate in the diet. *Reprod Toxicol*. 14(1):21–36.
50. Masutomi N, Shibutani M, Takagi H, et al. Impact of dietary exposure to methoxychlor, genistein, or diisononyl phthalate during the perinatal period on the development of the rat endocrine/reproductive systems in later life. *Toxicology*. 2003; 192(2–3):149–70. [PubMed: 14580783]
51. Vandenberg LN, Colborn T, Hayes TB, et al. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. 2012; 33(3):378–455.
52. Pocar P, Fiandanese N, Secchi C, et al. Exposure to di(2-ethyl-hexyl) phthalate (DEHP) in utero and during lactation causes long-term pituitary-gonadal axis disruption in male and female mouse offspring. *Endocrinology*. 2012; 153(2):937–48. [PubMed: 22147016]
53. Jiang J-T, Xu H-L, Zhu Y-P, et al. Reduced Fgf10/Fgfr2 and androgen receptor (AR) in anorectal malformations male rats induced by di-n-butyl phthalate (DBP): A study on the local and systemic toxicology of DBP. *Toxicology*. 2015; 338:77–85. [PubMed: 26514922]
54. Okayama Y, Wakui S, Wempe MF, et al. *In Utero* Exposure to Di(*n*-butyl)phthalate Induces Morphological and Biochemical Changes in Rats Postpuberty. *Toxicol Pathol*. 2017; 45(4):526–535. [PubMed: 28641505]
55. Oshida K, Vasani N, Thomas RS, et al. Identification of Modulators of the Nuclear Receptor Peroxisome Proliferator-Activated Receptor α (PPAR α) in a Mouse Liver Gene Expression Compendium. *PLoS One*. 2015; 10(2):e0112655. [PubMed: 25689681]
56. Sarath Josh MK, Pradeep S, Vijayalekshmi Amma KS, et al. Phthalates efficiently bind to human peroxisome proliferator activated receptor and retinoid X receptor α , β , γ subtypes: an in silico approach. *J Appl Toxicol*. 2014; 34(7):754–65. [PubMed: 23843199]
57. Hayashi Y, Ito Y, Yamagishi N, et al. Hepatic peroxisome proliferator-activated receptor α may have an important role in the toxic effects of di(2-ethylhexyl)phthalate on offspring of mice. *Toxicology*. 2011; 289(1):1–10. [PubMed: 21354252]

58. Laughter AR, Dunn CS, Swanson CL, et al. Role of the peroxisome proliferator-activated receptor α (PPAR α) in responses to trichloroethylene and metabolites, trichloroacetate and dichloroacetate in mouse liver. *Toxicology*. 2004; 203(1–3):83–98. [PubMed: 15363585]
59. Wassenaar PNH, Legler J. Systematic review and meta-analysis of early life exposure to di(2-ethylhexyl) phthalate and obesity related outcomes in rodents. *Chemosphere*. 2017; 188:174–181. [PubMed: 28886551]
60. Anderson OS, Nahar MS, Faulk C, et al. Epigenetic Responses Following Maternal Dietary Exposure to Physiologically Relevant Levels of Bisphenol A. *Environ Mol Mutagen*. 2012; 53(5): 334–342. [PubMed: 22467340]
61. Faulk C, Barks A, Liu K, Goodrich JM, Dolinoy DC. Early-life lead exposure results in dose- and sex-specific effects on weight and epigenetic gene regulation in weanling mice. *Epigenomics*. 2013; 5(5):487–500. [PubMed: 24059796]
62. Breton CV, Marsit CJ, Faustman E, et al. Small-Magnitude Effect Sizes in Epigenetic End Points are Important in Children's Environmental Health Studies: The Children's Environmental Health and Disease Prevention Research Center's Epigenetics Working Group. *Environ Health Perspect*. 2017; 125(4):511–526. [PubMed: 28362264]
63. Montrose L, Faulk C, Francis J, Dolinoy DC. Perinatal lead (Pb) exposure results in sex and tissue-dependent adult DNA methylation alterations in murine IAP transposons. *Environ Mol Mutagen*. 2017; 58(8):540–550. [PubMed: 28833526]
64. National Academies of Sciences Engineering and Medicine. Application of systematic review methods in an overall strategy for evaluating low-dose toxicity from endocrine active chemicals. 2017

Figure 1.

Experimental Design. Virgin a/a dams were given one of six types of chow containing either corn oil or phthalates dissolved in corn oil two weeks prior to mating with an A^{vy}/a male. They remained on this chow throughout gestation and lactation until weaning at PND21. Body weights at PND21 and tail global IAP DNA methylation were analyzed from both A^{vy}/a and a/a offspring. Coat color distribution and A^{vy} DNA methylation from tail tips were analyzed in only A^{vy}/a offspring, and organ and tissue weights were analyzed from a subset of a/a mice. A schematic of the Avy and global IAP assays can be found in Faulk et al. (2013)⁶¹ and Montrose et al. (2017)⁶³, respectively.

Figure 2.

PND21 body weights in females (A) and males (B) developmentally exposed to phthalates and phthalate mixtures. Body weights of mice in each exposure group were compared to those of control mice using linear mixed effects models controlling for number of pups per litter and within litter effects. Analyses were stratified by sex and genotype. Bars represent the mean for each group and error bars represent standard error (SE). ^p < 0.10, *p < 0.05.

Figure 3.

Relative liver weights in a subset of PND21 *a/a* mice (n=152) exposed to phthalates and phthalate mixtures. Relative liver weights are liver weight normalized to body weight, expressed as a percentage. Relative liver weights in exposure groups were compared to controls using linear mixed effects models controlling for within litter effects, and analyses were stratified by sex. Bar height represents mean relative liver weights for each group and error bars represent standard error (SE). ^p < 0.10, *p < 0.05, **p < 0.01.

Figure 4.

Coat color distributions of A^{vy}/a offspring across exposure groups. Coat color distributions for each exposure group were compared to the control using a χ^2 goodness of fit test. (A) Control coat color distribution. n=16 litters, 50 animals. (B) DEHP coat color distribution. p=0.016. n=14 litters, 49 animals. (C) DBP coat color distribution. p<0.0001. n=14 litters, 49 animals. (D) DINP coat color distribution. p=0.10. n=15 litters, 47 animals. (E) DEHP +DINP coat color distribution. p=0.016. n=16 litters, 55 animals. (F) DEHP+DBP+DINP coat color distribution. p<0.0001. n=17 litters, 61 animals.

Figure 5.

Mean tail global IAP DNA methylation across four CpG sites from PND21 *Avy/a* mice across exposure groups. Mean methylation in exposure groups was compared to controls via linear mixed effects models, which take methylation levels at nearby CpG sites into account, as well as within litter effects. Analyses were stratified by sex since initial analyses indicated a significant interaction between exposure and sex. Lines within boxes represent medians and whiskers represent 1.5*IQR. Gray dots are values outside of 1.5*IQR. ^p < 0.10, *p < 0.05.

Table 1

Comparison of DEHP daily intakes and DEHP metabolites in amniotic fluid in human and rodent studies

Population	Daily Intake (mg/kg-day)	Amniotic Fluid Levels (ng/mL)
Human Studies	0.0006 (min) 0.002 - 0.005 (range of medians) 0.042 (max)	<LOD (min) 1.6 - 22.1 (range of medians) 100.6 (max)
Calafat et al. (2006) (Rodents)	11 100	68 766
Present Study	5	<68

This table was adapted from the recent National Academies of Sciences, Engineering and Mathematics (NASEM) report on implementing systematic reviews to evaluate low-dose effects of EDCs.⁶⁴ Data from human studies includes a variety of populations.^{41–47} One rat study⁴⁰ examined concentrations of DEHP metabolites in amniotic fluid following oral doses of 11 and 100 mg/kg-day and was used to estimate potential amniotic fluid concentrations in the present mouse study.

Table 2

PCR and Pyrosequencing Assay Conditions

	<i>A^{py}</i> Assay ¹	Global IAP Assay ²
Primer/Sequence Location (5' to 3')	Chr2: 154761458 strand = reverse	Repetitive Element
Forward PCR Primer	ATTTTITAGGAAAAGAGTAAGAAGTAAG	GTGTTATTTTTTGATTGGTTGTAGTTT
Reverse PCR Primer (biotinylated)	CTACAAAACTCAAAAACTCA	ACCAAAAATATCTTATACTACTTATACT
Sequencing Primer	TAGAATATAGGATGTTAG	ATTTTTTGATTGGTTGTAGTTTA
Sequence to Analyze	YGTTATTTTGTGAYGGYGAATGTGGGGYGGTT	TYGGTYGAGTTGAYGTTAYGGGGAAAGTAGAGTATAAGTAGTTA
Amplicon Length (bp)	294	87
Annealing Temperature	53°C	56°C
Number of Cycles	54	45
Number of CpG Sites	4	4

¹ Assay adapted from Faulk et al. (2013);

² Assay adapted from Montrose et al. (2017)

Table 3

Litter outcomes: litter size, mortality rates, genotype ratio and sex ratio across exposure groups

Exposure Group	# Litters	Pup born	Pups died	Mean pups/litter (+/- SD)	Pups mortality rate (%)	A ^{W/a} offspring (%)	Female offspring (%)
Control	17	120	10	7.06 +/- 2.5	8.3	45.0	51.4
DEHP	16	110	17	6.88 +/- 1.8	15.45[^]	51.1	48.9
DBP	15	93	1	6.20 +/- 1.6	1.1[*]	53.3	42.4
DINP	15	101	4	6.73 +/- 2.1	4.0	48.0	48.0
DEHP+DINP	16	119	15	7.00 +/- 2.6	12.6	52.9	47.1
DEHP+DINP+DBP	19	123	15	6.47 +/- 2.2	12.2	53.4	51.5

* p 0.05 vs. control,

[^] p 0.10 vs. control. Genotype and sex ratios were determined for pups that survived until PND21 and do not include pups not surviving until PND21.

Table 4

Methylation Percent for A^{19}

A^{19} Mean Methylation Percent Across 4 CpG Sites									
Exposure	All Pups	Male A^{19}	Female A^{19}	Low-Methylation	High-Methylation	Male Low-Methylation	Female Low-Methylation	Male High-Methylation	Female High-Methylation
Control	31.8 (25.6)	26.8 (25.8)	37.4 (24.5)	10.2 (9.1)	52.4 (17.9)	11.0(9.7)	8.5 (7.9)	56.7(18.8)	50.0 (17.6)
DEHP	28.5 (24.8)	25.0 (25.2)	32.1 (24.5)	11.8(9.2)	54.9 (17.5)	11.9(9.6)	11.6 (9.1)	58.9 (20.4)	52.6 (16.0)
DBP	40.5 (30.1)	37.7 (29.6)	43.9 (31.1)	10.1 (7.2)	65.2 (14.7)[*]	10.1 (6.2)	10.1 (8.9)	63.3 (16.1)	67.4 (13.4)^{**}
DINP	34.7 (27.3)	35.6 (26.7)	33.4 (28.8)	11.5 (8.4)	60.1 (14.9)	13.7(8.5)	8.1 (7.5)	61.1 (14.5)	58.8 (16.2)
DEHP+DINP	36.1 (27.0)	35.3 (27.1)	37.4 (27.6)	11.5 (8.3)	58.8(15.8)	11.2 (8.2)	12.0 (8.9)	59.3 (14.6)	58.1 (18.1)
DEHP+DINP-DBP	35.2 (26.9)	33.0 (27.3)	37.3 (26.8)	11.2 (8.1)	59.3 (14.3)	9.9 (8.5)	12.5 (7.8)	58.0(15.5)	60.4 (13.6)[^]

Reported as percent mean methylation (standard deviation).

^{**} p<0.01 versus control,^{*} p<0.05 versus control,[^] p<0.10 versus control.