Published in final edited form as:

Curr Protoc Stem Cell Biol. 2019 February ; 48(1): e64. doi:10.1002/cpsc.64.

Highly efficient CRISPR/Cas9 mediated genome editing in human pluripotent stem cells

Jean Ann Maguire¹, Fabian L. Cardenas-Diaz², Paul Gadue^{1,2,3,4,*}, and Deborah L. French^{1,3,4,5,*}

¹Center for Cellular and Molecular Therapeutics, The Children's Hospital of Philadelphia

²Department of Biology, University of Pennsylvania

³Department of Pathology and Laboratory Medicine, The Children's Hospital of Philadelphia

⁴Department of Pathology and Laboratory Medicine, University of Pennsylvania

Abstract

Human PSCs offer tremendous potential for both basic biology and cell-based therapies for a wide variety of diseases. The ability to manipulate the genome of these cells using the CRISPR/Cas9 technology has expanded this potential by providing a valuable tool to engineer or correct disease-associated mutations. Because of the high efficiency by which CRISPR/Cas9 creates targeted double strand breaks, a major challenge has been the introduction of precise genetic modifications on one allele, without indel formation on the non-targeted allele. To overcome this obstacle, we describe the use of two oligonucleotides; one expressing the sequence change with the other maintaining the normal sequence. In addition, we have streamlined both the transfection and screening methodology to make this protocol efficient with small numbers of cells, and limit the amount of labor intensive clone passaging. This protocol provides a streamlined and technically simple approach for generating valuable tools to model human disease in stem cells.

Keywords

Human pluripotent stem cells (PSCs); genome editing; homology directed repair; CRISPR/Cas9

Introduction

Pluripotent stem cells (PSCs) are valuable tools for the study of human biology and *in vitro* disease modeling. The ability to introduce and/or correct disease-associated mutations using the CRISPR/Cas9 system promotes greater tractability within these specialized cells (Brookhouser et. al., 2017; Hockemeyer & Jaenisch, 2016). However, engineering of highly specific alterations has been challenging, prompting the development of new methodologies to enhance the efficiency and ease of genome engineering in stem cells. A limiting factor in genome editing includes generating precise mono-allelic sequence modifications, without

⁵Correspondence: Deborah L. French, PhD, The Children's Hospital of Philadelphia, 3501 Civic Center Blvd, CTRB 5014, Philadelphia, PA 19104, 267-426-9673, frenchd@email.chop.edu.

*PG and DLF contributed equally as senior authors

the introduction of nonspecific indels (Richardson et. al., 2016; Paquet et. al., 2016). To overcome this obstacle, we demonstrate the use of a double oligonucleotide (ODN) recombination system to introduce and/or correct mono-allelic genetic mutations. When using a single-stranded ODN (ssODN) template to generate a mutation, we found virtually all clones will harbor indel mutations in the non-targeted allele. To overcome this problem, we introduced two ssODNs, both of which contain the necessary blocking mutations to prevent guide RNA (gRNA) recognition and re-cutting, but only one of which includes the desired genetic alteration. We show that this methodology results in introduction of the desired genetic alteration in one allele, without occurrence of indel generation in the other allele. Data collected from thirteen unique gene editing experiments demonstrated an average editing efficiency of 9.5% (six shown in Figure 6).

This unit begins with a description of gRNA design, cloning and efficiency testing in PSCs. The method utilizes transient transfection of plasmid DNA, and evaluation of gRNA cleavage efficiency (Basic Protocol 1). This is followed by a protocol for the introduction of precise mutations by the recombination of two ssODNs (Basic Protocol 2). This method reduces the occurrence of indel formation in the second allele that often occurs when using a single ODN approach for homologous recombination.

NOTE: PSCs for CRISPR/Cas9 editing are cultured in human embryonic stem cell (hESC) media (see Reagents and Solutions) with irradiated mouse embryonic fibroblasts (MEFs) in a laminar-flow hood according to standard practices, and maintained in a humidified incubator at 37°C, 5% CO₂, 5% O₂, and 90% N₂ (Paluru, et. al., 2013).

BASIC PROTOCOL 1: Implementation of CRISPR based genome editing in human PSCs

This protocol outlines the design, cloning and efficiency testing of gRNAs required for CRISPR/Cas9 genome editing. Determination of guide cutting efficiency is performed in the target stem cell using a lipid based transfection method.

Materials

Plasmids

pCas9_GFP (Addgene #44719)
gRNA_Cloning Vector (Addgene #41824)

Tissue Culture

ThermoFisher® Lipofectamine Stem Reagent (STEM00001)
TrypLE (ThermoFisher 12605036)
Y-27632 Dihydrochloride/Rock inhibitor (ROCKi) (Tocris 1254)
Falcon polystyrene 6 well tissue culture dishes (Corning 353046)
Falcon 5 mL polystyrene round-bottom tube with cell strainer-cap (Corning 352235)

Falcon 10-cm² tissue culture dishes (Corning 353003)

Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix (354230)

Iscove's Modified Dulbecco's Medium (IMDM) (Corning 12440053)

Murine Embryonic Fibroblasts (MEFs)-irradiated

Kits

Phusion High Fidelity DNA Polymerase (New England Biolabs M0530)

dNTPs (Roche 11969064001)

Gibson Assembly Kit (New England Biolabs E2611)

Purelink™ Quick Plasmid Miniprep Kit (Invitrogen K210011)

Nucleospin® Gel and PCR Clean-up Kit (Macherey-Nagel 740609)

Miscellaneous

Agarose (VWR N605)

Gel extraction kit (Macherey-Nagel 740609)

Afl (New England Biolabs R0520)

Stellar™ competent cells (Takara 636763)

Kanamycin LB agar plates (50µg/ml Kanamycin)

Proteinase K Solution (Qiagen 19133)

Equipment

Orbital shaking incubator

Variable oxygen control CO₂ incubator

Fluorescence-Activated Cell Sorter

Guide RNA Design

1. Identify three to four 23 bp 5'-G(N₁₉)NGG-3' sequences within 20 bp of the target site on either strand of DNA (Figure 1A & B).

The protocol suggests designing 3–4 gRNAs, but situations do arise when fewer gRNA sequences are identified. If this occurs, perform efficiency testing on the gRNAs that are identified.

If you are unable to find a G (guanine) at the first position (20th nucleotide preceding the PAM), it can be engineered into the gRNA (Figure 1B). However, the optimal condition is for this G to be within the target sequence.

2. Evaluate the gRNA sequences for genomic redundancy and off-target probability.

There are multiple resources that can be accessed for gRNA evaluation. We use the CRISPOR website (<http://crispor.tefor.net/>).

3. Incorporate the 20 bp target sequence (excluding the PAM) into two 60-mer oligos as shown in Figure 1C (sequences are 5' to 3' and red and green are reverse compliments).
4. Order the two 60-mer oligos from IDT (www.idtdna.com) as 25 nmole DNA oligos (not ultramers).
5. Anneal the two oligos and generate a 100 bp dsDNA fragment using Phusion High Fidelity DNA polymerase.
  - a. Resuspend forward and reverse oligos to a final concentration of 100 μ M in water. Make a working stock of 10 μ M.
  - b. Combine the following from working stocks:

5 μ l 10 μ M forward oligo + 5 μ l 10 μ M reverse oligo
  - c. Incubate at 95°C for 5 min.
  - d. Cool at RT for 10 min.
  - e. Add the following to each primer set (20 μ l total volume):

0.4 μ l 10mM dNTPs

4 μ l 5x Phusion HF buffer

0.2 μ l Phusion polymerase

5.4 μ l ddH₂O
  - f. Perform PCR amplification using the following parameters:

98°C $\times$ 30 sec

30 cycles: 98°C $\times$ 10 sec

55°C $\times$ 20 sec

72°C $\times$ 30 sec

72°C $\times$ 5 min

4°C hold
6. Run the PCR product on a 1% agarose gel.
7. Excise a 100 bp band and purify with a gel extraction kit.

A nonspecific band at ~200 bp is sometimes observed.

Design of PCR Primers for Screening

1. Design forward and reverse PCR primers to amplify the region to be edited (Figure 1A).

The primers should asymmetrically flank the gRNA cleavage sites. The resulting amplicon should be ~300–400 bp.

2. Optimize the PCR on control genomic DNA, and confirm a clean amplicon by agarose gel electrophoresis.
3. A nested primer should also be designed for future sequencing of clones.

Preparation of gRNA_Cloning Vector Plasmid

1. Digest the gRNA_Cloning Vector with *Afl*II:
  - 10 µg gRNA_Cloning Vector
  - 4 µl 10X CutSmart Buffer
  - 24 µl water
  - 1.5 µl *Afl*II
2. Incubate at 37°C O/N.
3. Run digest on a 1% agarose gel and excise bands (expected band size is ~3519 bp).
4. Purify with a gel extraction kit.

Gibson Assembly of gRNA Vector

1. Using the NEB Gibson Assembly Kit, set-up each reaction at a 1:5 ratio of *Afl*III digested gRNA_Cloning Vector to 100 bp gel purified Phusion insert.
  - 50–100 ng *Afl*III digested vector
  - 250–500 ng (5x) 100bp insert
  - 10 µl 2x master mix
  - ddH₂O to 20 µl
2. Incubate at 50°C for 15 min.
3. Dilute Gibson Assembly reaction (20 µl) 1:3 in ddH₂O and proceed to plasmid transformation.

Transformation of gRNA_Cloning Vector

1. Add 3 µl of the diluted Gibson Assembly reaction mixture to 50 µl of competent cells.
2. Incubate cells on ice for 20 minutes, heat shock at 42°C for 45 seconds, and return to ice for 2 minutes.
3. Add 900 µl of SOC media to each sample and shake at 37°C for 1 hour in an orbital shaking incubator.
4. Plate 500 µl of each sample on Kan^R LB plates and grow O/N at 37°C.
5. Pick 3–5 colonies per gRNA. Inoculate each colony in 4 mL LB and grow O/N at 37°C on an orbital shaking incubator.

6. Purify plasmid DNA using a miniprep plasmid isolation kit.
7. Sequence each gRNA using the following primers:
Forward: GTACAAAAAAGCAGGCTTTAAAGG
Reverse: TGCCAACTTTGTACAAGAAAGCT
8. If the gRNA sequence is correct, continue with cell transfection. If the sequence is not found, cloning was unsuccessful, and additional bacterial colonies need to be screened.

Testing gRNA Cleavage Efficiency (Figure 2)

Multiple kits and assays are available to evaluate gRNA cleavage on pooled cells (i.e. Surveyor Assay), but we have had greater success screening individual clones as described below.

1. Plate target cell line in a 6-well dish on irradiated MEFs to reach 70–80% confluency after an overnight incubation. Each gRNA is tested in duplicate requiring 2 wells of cells.
Each cell line will require optimization to achieve the desired plating density.
2. Change the hESC medium (1.5 mL/well) at least 4 hours before transfection.
3. For each transfection, add the following reagents to a 1.5 mL eppendorf tube:
50 μ L base media (normal base media with no additions)
0.5 μ g pCas9_GFP vector
0.5 μ g guide RNA miniprep
3 μ L Lipofectamine Stem - added last
4. Mix by pipetting and incubate at RT for 15 minutes.
5. Add dropwise to the cells and incubate for 48 hours.
6. Harvest cells for sorting after 48 hours.
  - a. Remove MEFs with a 3 min RT TrypLE incubation.
  - b. Rinse cells 1 $\times$ with HESC medium and scrape into hESC medium + 10 μ M ROCKi, combining duplicate wells.
  - c. Pellet cells at 300 $\times$ g for 3 min and resuspend in 0.5 mL hESC medium + 10 μ M ROCKi.
  - d. Filter into a 5 mL tube through a cell-strainer cap.
7. Using FACS, sort GFP positive cells.
The percent of GFP+ cells is typically between 1–6%.
Use hESC + ROCKi for sorting and collection medium.

8. Transfer the sorted cells directly to a 10 cm² dish (previously coated with 1:3 matrigel and 1×10⁶ irradiated MEFs, see Reagents and Solutions) in growth medium containing ROCKi.

Do not pellet cells before plating on 10 cm² dish.

Plate a maximum of 15k cells per 10 cm² dish. A minimum of ~5k GFP + cells appears to be required to achieve adequate clone formation.
9. Change medium daily using hESC medium without ROCKi and manually pick clones after 10–15 days, when colonies are ~1 mm in diameter.
  - a. Using a P200 pipette and microscope, carefully scrape a single clone and draw cells into the pipette.
  - b. Disperse the cells by gently pipetting 3–4 times in a 96 well plate in the medium drawn up with the colony.
  - c. Dispense into PCR strip tubes for screening.

For gRNA screening, pick 20 clones per guide.

Clone Screening

1. Pellet cells in PCR strip tubes at 10,000×g for 5 min and remove supernatant.

Pellets can be stored at –20°C to screen at a later time.
2. Resuspend cell pellets in 20 µl of Proteinase K Buffer (see Reagents and Solutions).
3. Incubate at 55°C for 1 hour, 95°C for 10 min.
4. Vortex vigorously and centrifuge at 10,000×g for 5 min.
5. Analyze 5 µl of the Proteinase K digest (dirty miniprep) in a 20 µl PCR reaction using the primer set designed to flank the edited region. Include genomic DNA from the parental cell line to use as a control.
6. Analyze PCR products on a 2.5% agarose gel.

Size changes and doublets compared to the control are indicative of cleavage (Figure 3). Evaluate each gRNA and calculate the efficiency of cleaved/total clones picked. Proceed to Basic Protocol 2 with the most efficient gRNA. A cleavage efficiency of >25% is optimal, but not required.

BASIC PROTOCOL 2: Genetic modification of human PSCs using a double ODN CRISPR/Cas9 recombination system

This protocol is used to introduce precise mutations within a gene of interest and to allow efficient generation of a mutation in a single allele without generating an indel in the second allele. Using a prescreened gRNA (see Basic Protocol 1), two ssODNs are designed with

and without the desired base alteration(s). Additional silent mutations are introduced into the ssODNs to prevent gRNA cutting and to streamline screening of clones.

Materials

Plasmids

pCas9_GFP (Addgene #44719)

Tissue Culture

ThermoFisher® Lipofectamine Stem Reagent (STEM00001)

TrypLE (ThermoFisher 12605036)

Y-27632 Dihydrochloride/Rock inhibitor (ROCKi) (Tocris 1254)

Falcon polystyrene 6 well tissue culture dishes (Corning 353046)

Falcon 10-cm² tissue culture dishes (Corning 353003)

Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix (354230)

Iscove's Modified Dulbecco's Medium (IMDM) (Corning 12440053)

Murine Embryonic Fibroblasts (MEFs) - irradiated

Kits

Nucleospin® Gel and PCR Clean-up Kit (Macherey-Nagel 740609)

Miscellaneous

Agarose (VWR N605)

Proteinase K Solution (Qiagen 19133)

Equipment

Variable oxygen control CO₂ incubator

Fluorescence-Activated Cell Sorter

Oligonucleotide Design

1. Design 100 bp ssODNs centered around the most efficient gRNA sequence determined in protocol 1 (Figure 4A).
2. Introduce a silent mutation in the PAM sequence of the oligo, but if not possible, introduce 3–4 silent mutations within the gRNA sequence.

These mutations prevent gRNA “re-cleavage” of the recombined ODN.
3. As a screening strategy, introduce a unique restriction site within ~25 bp of the gRNA sequence using silent mutations. Targeted clones will be identified by restriction enzyme digestion.

4. Design one ODN with the desired base change(s) to create the mutation of interest and one ODN without the base change(s).

These changes should be no more than ~25 bp from the predicted CRISPR/Cas9 cut site, as recombination drops off considerably at greater distances (Paquet et. al., 2016).
5. Order the two ssODNs from IDT as 4 nmole Ultramer® DNA Oligos with standard desalting purification. The maximum length for this category is 200 bases.
6. Resuspend in water to make 1 µg/µl stocks. Store stocks at –20°C.

Transfection Setup

1. Plate the target cell line in a 6-well dish on irradiated MEFs to reach 70–80% confluency after an overnight incubation.
2. Change medium at least 4 hours before transfection.

Additional bFGF (75–100 ng/ml) can be added to the media (1.5 mL/per well) to increase proliferation.
3. For each transfection, add the following reagents to a 1.5 mL eppendorf tube:
  - 50 µl base media
  - 0.5 µg pCas9_GFP vector
  - 0.5 µg guide RNA
  - 1 µg ssODN (if using 2 ODNs, use 0.5 µg each)
  - 3 µl Lipofectamine Stem - added last
4. Mix by pipetting and incubate at RT for 15 minutes.
5. Add dropwise to the cells, and incubate O/N in a low oxygen incubator (37°C, 5% CO₂, 5% O₂, and 90% N₂.)

The MEF feeder layer will also exhibit a low efficiency of transfection. This is to be expected and will not have an effect on isolating edited clones as the MEFs are removed prior to sorting.
6. Harvest cells for sorting after 48 hours.
  1. Remove MEFs with a 3 min RT TrypLE incubation.
  2. Rinse cells 1× with media, and scrape into hESC medium + 10µM ROCKi, combining duplicate wells.
  3. Pellet cells at 300×g for 3 min and resuspend in 0.5 mL hESC medium + 10µM ROCKi.
  4. Filter into a 5 mL tube through a cell-strainer cap.
7. Using FACS, sort GFP positive cells.

Use hESC + ROCKi for sorting and collection media.

8. Transfer the sorted cells directly to a 10 cm² dish (coated with 1:3 matrigel, see Reagents and Solutions, and 1×10⁶ irradiated MEFs, containing hESC medium + ROCKi..

Plate a maximum of 15k cells per 10 cm² dish. A minimum of ~5k GFP + cells appears to be required to achieve adequate clone formation.

9. Change medium daily with hESC media and manually pick colonies after 10–15 days, when colonies are ~1 mm in diameter.
  - a. Using a P200 pipette and microscope, carefully scrape a single clone and draw cells into the pipette.
  - b. Disperse the cells by gently pipetting 3–4 times in a 96 well plate in the medium drawn up with the colony.
  - c. Distribute:
    - Half to a 24 or 48 well dish (pre-coated with 1:3 matrigel, see Reagents and Solutions, and 1×10⁶ irradiated MEFs), containing hESC medium + ROCKi.
    - Half to PCR strip tubes for screening.

Isolate 24–48 clones per condition.

Clone Screening

1. Pellet cells in PCR strip tubes at 10,000×g for 5 min and remove supernatant.

Pellets can be stored at –20°C to screen at a later time.
2. Resuspend cells in 20 µl of Proteinase K Buffer (see Reagents and Solutions).
3. Incubate at 55°C for 1 hour, 95°C for 10 min.
4. Vortex vigorously and centrifuge at 10,000×g for 5 min.
5. Add 5 µl of the Proteinase K digest to a 30 µl PCR reaction using primers flanking the gene edited region.
6. Run 5 µl of the PCR reaction on a 1.5% agarose gel to ensure product formation, and purify the remaining PCR product using a PCR clean-up kit.
7. Digest half of the PCR product using the restriction enzyme specific for the unique site designed in the ODN and run on a gel to select clones with recombination events (Figure 4B).

In this example, an uncleaved band represents no recombination (WT). Evidence of both uncleaved and cleaved bands indicates recombination with only one allele (One Allele). In our experience, these are not desirable clones, as the uncut allele most likely contains an indel. Complete cleavage of the PCR product indicates recombination in both

alleles (Two Allele). Sequencing is required to determine which ssODN was used as a template.

8. Sequence clones using the nested sequencing primer previously designed.
Focus on “Two Allele” clones due to the high probability that the “One Allele” clones may have an indel in the unrecombined allele.
9. Align sequences with the WT gene to ensure proper editing.
10. After confirmation, expand cells stocks and confirm normal karyotype of clones.

REAGENTS AND SOLUTIONS

Proteinase K Buffer

50 mM Tris-HCl
15 mM ammonium sulfate pH 9.3
2.5 mM MgCl₂
0.1% Tween 20
100 µg/mL Proteinase K

1:3 Matrigel

Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix diluted 1:3 in IMDM medium. Keep all components on ice.

hESC Media

DMEM/F12
15% knock-out serum replacement
100 µM non-essential amino acids
1 mM sodium pyruvate
2 mM glutamine, 50 U/ml penicillin
0.1 mM β-mercaptoethanol (Sigma, St Louis, MO)
10 ng/ml human bFGF (R&D Systems)

NOTE: All incubations are performed in a humidified incubator at 37°C, 5% CO₂, 5% O₂, and 90% N₂.

COMMENTARY

Background Information

Many genetic disorders have been studied by reprogramming induced PSCs (iPSC) from patients with a specific disease, but this approach is time consuming and severely limited by patient availability. Studies are also confounded by variable genetic backgrounds of patients.

These problems can be circumvented by introduction of the disease mutations in an iPSC line already generated by reprogramming of cells from a normal individual (Anderson & Francis, 2018). The process can be completed in a much shorter period of time, and provides an isogenic control for subsequent studies. In order to introduce these mutations, the CRISPR/Cas9 system is increasingly preferred, based on its high efficiency and ease of use compared with zinc-finger nucleases and transcription activator-like effector nucleases (TALENs) (Ding et. al., 2013; Hou et. al., 2013). When combined with the highly efficient ssODN template repair that occurs via homologous recombination, the introduction of specific mutations can be readily achieved (Richardson et. al., 2017).

Critical Parameters

This protocol has been optimized for PSCs cultured on a feeder layer of irradiated MEFs. Although feeder-free growth conditions may result in significantly higher transfection efficiency, cell survival post-transfection and editing efficiency may be reduced compared to cells grown on MEFs. Transfection efficiency of PSCs grown on MEFs is often <5%, but we have found that desired results can be adequately achieved. In addition, this protocol was optimized using iPSCs grown in hypoxic culture conditions. CRISPR/Cas gene editing has been successful using iPSCs grown in normoxic culture conditions (Giacalone et. al., 2018), so it would be anticipated that these conditions would apply to this protocol.

The design of optimal PCR primers for amplification of the edited sequence is important for both gRNA efficiency testing and clone screening. Without a clean PCR product, both gRNA cleavage and restriction enzyme screening will be difficult to assess. In addition, the screening procedure used in this protocol allows for expedited screening of clones so identification of possible targets prior to splitting of picked clones can occur. This method allows the investigator to save a significant amount of effort by only splitting and expanding clones that are positive in the initial screen. Some projects may not work with the dirty PCR prep and require expansion of all clones, and preparation of a high quality genomic DNA preparation.

Location of gRNAs must reside within coding regions, as the silent/blocking mutations introduced to the ssODN can have an unknown effect in non-coding regions. This protocol could be used to target non-coding, single nucleotide polymorphisms with two caveats. First, the nucleotide change must be within the PAM sequence. Second, only homozygous changes can be made efficiently without indel formation (data not shown).

In this protocol, transfection of plasmid DNA is accomplished with ThermoFisher® Lipofectamine Stem reagent. This protocol can work with other lipid transfection reagents, but will require further optimization. The use of lipid transfection allows the generation of sufficient numbers of transfected cells from two wells of a 6-well dish of PSCs. Lipid transfection is also less stressful on the cells compared to electroporation, and is technically easier to perform.

This protocol can also be amended to introduce deletions into a gene of interest. By designing two gRNAs flanking the region to be deleted, we have successfully deleted sequences up to ~10k bp. The cutting efficiency of each gRNA should be tested individually,

then added to the transfection mixture at equal concentrations. Screening is accomplished using PCR primers flanking the deletion. With larger deletions, these primers will only create a product if the sequence has been removed. To confirm deletion or identify single allele deletions, a primer within the deleted region is designed and combined with the opposing flanking primer.

Troubleshooting

For troubleshooting, please refer to Table 1.

Anticipated Results

Introduction of two ODNs (WT and mutant) will result in three possible recombination outcomes (Figure 5).

Outcome 1 shows homologous recombination with one ssODN. In our experience, when one allele recombines with an ODN, the second allele will harbor an indel. Either the mutant or WT ssODN may recombine with the one allele, which can be identified by TOPO PCR Cloning and sequence analysis.

Outcome 2 shows homologous recombination on both alleles with either the mutant or WT ssODNs.

Outcome 3 shows heterozygous recombination on both alleles with the mutant and WT ssODNs. One allele contains the desired base change and the other allele contains the wild type sequence. Both are identified by sequence analysis.

Using the two ssODN approach described in this protocol, we have found recombination efficiencies for Outcome 3 to be 0–21%. Figure 6 shows six representative editing experiments. Selected clones were karyotyped and all were found to be normal. Significant differences in editing success are evident for each gene. Of note, low numbers of clones or an inability to identify gRNAs that cut may be due to chromatin structure and/or gene lethality.

Time Considerations

The total amount of time for the described protocols largely depends on the success of gRNA cleavage. The following outlines an estimation of the time required for each task to be completed by a researcher familiar with PSC culture and molecular biology.

Design and cloning of gRNAs can take 4–5 days (not including time for shipment of ODNs). The gRNA efficiency testing of the protocol is time consuming, but an important required component of the assay. Single colonies can take up to 2 weeks to grow big enough for testing. With practice, 24–48 clones can be manually isolated within 1–2 hours. PCR screening of the clones will take 1–2 days. If an efficient gRNA is not identified in the first round of screening, additional gRNAs will need to be designed and cloned. This could delay the protocol by multiple weeks.

Once an optimal gRNA is identified, ssODN design can be accomplished in 1 day. Again, after a 48 hour transfection of gRNA and ODNs, clone expansion will take approximately 2 weeks. After manual isolation of clones, clone screening (PCR amplification and restriction enzyme digestion) will take 2–3 days. Potential edited clones can be sent for overnight sequencing, and upon confirmation of genome editing, cells can be expanded. It can be expected to take 2–3 weeks to expand and freeze stocks.

Acknowledgments

This research was supported by funding from U01HL099656 (PG and DLF) and U01HL134696 (PG and DLF) from the National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health.

Literature Cited

- Anderson, RH; Francis, KR. Modeling rare diseases with induced pluripotent stem cell technology. Mol And Cell Probes Article. 2018. in Press
- Brookhouser N, Raman S, Potts C, Brafman DA. 2017; May I cut in? Gene Editing approaches in human induced pluripotent stem cells. Cells. 6(1):5.
- Ding Q, Regan SN, Xia Y, Oostrom LA, Cowan CA, Mu-sunuru K. 2013; Enhanced efficiency of human pluripotent stem cell genome editing through replacing TALENs with CRISPRs. Cell Stem Cell. 12:393–394. [PubMed: 23561441]
- Giacalone JC, Sharma TP, Burnight ER, Fingert JF, Mullins RF, Stone EM, Tucker BA. 2018; CRISPR-Cas9-based genome editing of human induced pluripotent stem cells. Current Protocols in Stem Cell Biology. 44:5B.7.1–5B.7.22.
- Hockemeyer D, Jaenisch R. 2016; Induced Pluripotent Stem Cells Meet Genome Editing. Cell stem cell. 18:573–586. [PubMed: 27152442]
- Hou Z, Zhang Y, Propson NE, Howden SE, Chu LF, Sontheimer EJ, Thomson JA. 2013; Efficient genome engineering in human pluripotent stem cells using Cas9 from *Neisseria meningitidis*. Proc Natl Acad Sci. 110(39):15644–15649. [PubMed: 23940360]
- Paluru P, Hudock KM, Cheng X, Mills JA, Ying L, Galvao AM, Lu L, Tiyaaboonchai A, Sim X, Sullivan SK, et al. 2013; The negative impact of wnt signaling on megakaryocyte and primitive erythroid progenitors derived from human embryonic stem cells. Stem Cell Res. 12:441–451. [PubMed: 24412757]
- Paquet D, Kwart D, Chen A, Sproul A, Jacob S, Teo S, Olsen KM, Gregg A, Noggle S, Tessier-Lavigne M. 2015; Efficient introduction of specific homozygous and heterozygous mutations using CRISPR/Cas9. Nature. 533:125.
- Ran F, Hsu P, Wright J, Agarwala V. 2013; Genome engineering using the CRISPR-Cas9 system. Nat Protoc. 8:2281–2308. [PubMed: 24157548]
- Richardson CD, Ray GJ, DeWitt MA, Curie GL, Corn JE. 2016; Enhancing homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor DNA. Nat Biotech. 34:339–345.
- Richardson CD, Kazane KR, Feng SJ, Bray NL, Schäfer AJ, Floor S, Corn JE. 2017; CRISPR-Cas9 genome editing in human cells works via the Fanconi Anemia pathway. bioRxiv preprint.

Significance Statement

Human pluripotent stem cells (PSCs) offer great hope for the study and treatment of human disease. These cells can grow in culture almost indefinitely and can be used to manufacture many cell and tissue types found in the body. A major breakthrough in technology now enables these cells to be genetically manipulated to introduce or correct a mutation of interest. A highly efficient and technically simple protocol for genetically manipulating these cells using the CRISPR/Cas9 system is described in this report.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Guide RNA Design

A. Identify gRNAs within ± 20 bp of the target region (red box). gRNAs can be designed on either the sense or antisense strand. Design PCR primers to amplify ~300–400 bp flanking the region to be edited (green arrows). A nested sequencing primer should also be designed for future screening of clones (yellow arrow). **B.** 23 bp gRNA structure guideline with PAM at 3' end. **C.** Incorporation of gRNA sequence (minus the PAM site) into 60-mer ODNs.

Figure 2. CRISPR-Cas9 transfection and clone selection overview

PSCs are transfected using a lipid mediated reagent, and sorted for GFP 48 hours post-transfection. Clones are manually isolated 1–2 weeks after plating sorted cells, and screened by PCR for gRNA cleavage confirmation.

	# Clones picked	Cleavage	Efficiency (%)
gRNA 1	14	7	45
gRNA 2	5	2	40
gRNA 3	3	0	0
gRNA 4	8	3	38
gRNA 5	16	1	6

Figure 3. Guide RNA Efficiency Testing

To illustrate gRNA cleavage efficiency, H9 ESCs were co-transfected with Cas9-GFP and 5 individual gRNAs. Cells were sorted for GFP fluorescence 48 hours post-transfection, and plated at low density. After 1–2 weeks, clones were manually isolated and screened by PCR and agarose gel electrophoresis for gRNA cleavage. Blue dots indicate cleavage compared to control (C) samples. Efficiency of gRNA cleavage is determined by the ratio of cleaved samples to total clones evaluated. This figure depicts the agarose gel of the most efficient gRNA (#1). After identification of the optimal gRNA, genome editing with oligonucleotides can be performed.

Figure 4. Oligonucleotide Design and Recombination Screening

A. 100 bp ssODNs are designed flanking the optimal gRNA (blue/aqua shaded sequence). Silent/blocking mutation(s) are introduced in the PAM site or throughout the gRNA sequence to prevent subsequent cleavage after recombination (red shaded nucleotide). Introduction of a unique restriction site within the PCR amplicon allows for simplified screening of clones (green box). This example depicts an EcoRI site formed through introduction of a silent mutation in the PAM site. However, the restriction site does not need to be made within the gRNA sequence. Finally, wild-type (WT) and mutant ODNs are designed incorporating the endogenous sequence (WT) or a mutation of interest (mutant) (red box). **B.** Potential screening outcomes after PCR amplification and restriction digest include no recombination (WT), which shows an uncleaved band, one allele recombination, which shows both uncleaved and cleaved bands, and two allele recombination, which shows only cleaved product.

Outcome 1: One allele recombination **Outcome 2: Two allele recombination (homozygous)** **Outcome 3: Two allele recombination (heterozygous)** **Figure 5. Recombination Outcomes**

Potential outcomes include 1) single allele recombination (recombination in one allele and the presence or absence of indel formation in the second allele), 2) two allele recombination with either WT or MUT ODN (homozygous), or 3) two allele recombination with WT AND mutant ODN (heterozygous).

Gene	# clones picked	One allele recombination		Two allele recombination		
		With indel	Without indel	WT/WT	MUT/MUT	WT/MUT
Gene 1	24	5 (21%)	0	0	1 (4%)	5 (21%)
Gene 2	48	5 (10%)	0	0	0	6 (13%)
Gene 3	22	4 (18%)	0	1 (5%)	0	2 (9%)
Gene 4	12	5 (42%)	0	1 (8%)	3 (25%)	1 (8%)
Gene 5	24	7 (29%)	0	0	0	2 (8%)
Gene 6	24	3 (13%)	0	0	0	2 (8%)

Figure 6. Recombination Efficiencies

Table displays results from multiple genes edited with described protocol. When using one ssODN (one allele recombination), only one allele recombination with indel formation occurs. Introduction of two ssODN results in varying efficiencies of the three outcomes described in Figure 5.

Table 1

Troubleshooting

Technique	Problem	Possible Cause	Solution
Transformation of gRNA cloning vector	No gRNA sequence in vector	Inefficient cloning	Screen additional clones or repeat
Cleavage efficiency of Cas9 gRNA	Poor to undetectable cleavage	SNPs	Sequence target region in cells to be edited to confirm matching sequence
		Refractory chromosomal region	Design guides in another region
		Lethal mutation	None
ssODN Design	Cannot create unique restriction site with silent mutations	Limited sequence availability	A unique restriction site can be removed with silent mutations, but gRNA cleavage sometimes creates false positives when screening
Plasmid transfections	Poor cell survival	Toxic load of DNA	Decrease concentrations of plasmid/ssODN
	Low efficiency	Cell density	Determine the cell number for 70–80% confluency
		Plasmid ratios incorrect	Test multiple plasmid concentrations to achieve optimal GFP expression
		Lipofectamine Stem volume	Optimize volume needed for transfection
Clone screening	Heterogeneity of clones	Clone proximity too close upon manual isolation	Plate sorted cells at a higher dilution (no more than 15,000 sorted cells per 10 cm ² dish)
	No PCR product or dirty PCR product	Proteinase K minipreps may not be adequately clean for PCR reaction	Prepare purified genomic DNA