Published in final edited form as:

Drug Alcohol Rev. 2019 January ; 38(1): 42–49. doi:10.1111/dar.12882.

Prevalence of reagent test-kit use and perceptions of purity among ecstasy users in an electronic dance music scene in New York City

Joseph J. Palamar, PhD, MPH¹ [Associate Professor] and Monica J. Barratt, PhD^{2,3,4} [Research Fellow]

¹New York University Langone Medical Center, New York, USA

²Drug Policy Modelling Program, National Drug and Alcohol Research Centre, UNSW Sydney, Sydney, Australia

³National Drug Research Institute, Faculty of Health Sciences, Curtin University, Perth, Australia

⁴Behaviours and Health Risks Program, Burnet Institute, Melbourne, Australia

Abstract

Introduction and Aims: Ecstasy users in the electronic dance music scene are at high risk for using ecstasy adulterated with new psychoactive substances and/or methamphetamine. We examined self-reported testing of ecstasy among users in this scene.

Design and Methods: We surveyed individuals (aged 18–40 years) entering electronic dance music parties in New York City in 2017. Past-year ecstasy users (n=351) were asked if they had tested their ecstasy in the past year. We estimated prevalence and correlates of having tested one's ecstasy.

Results: 23.1% reported having tested their ecstasy in the past year. Those with some college (adjusted prevalence ratio [aPR]=0.49, $P=0.014$) or a college degree (aPR=0.41, $P=0.025$) were less likely to test their ecstasy than those with a high school diploma or less. Using ecstasy pills (aPR=1.89, $P=0.036$) or crystals (aPR=1.90, $P=0.006$) 3 times in the past year was associated with increased likelihood of testing one's ecstasy, and purchasing from an unknown or untrustworthy dealer was associated with decreased likelihood (aPR=0.63, $P=0.034$) of testing one's ecstasy. Half (51.1%) of ecstasy users reported finding out or suspecting their ecstasy had contained a drug other than MDMA. Of these, 49.2% reported finding out their ecstasy contained methamphetamine or speed/amphetamine. Most ecstasy users reported that they would be less likely to use again upon learning their ecstasy contained "bath salts" (54.8%) or methamphetamine (54.3%).

Corresponding author: Dr Joseph J. Palamar, New York University Langone Medical Center, 180 Madison Avenue, Room 1752, New York, NY 10016, USA, Phone: 1-646-501-3555, joseph.palamar@nyu.edu.

Declaration of interests

There are no relevant interests to declare.

Discussion and Conclusions: Drug testing appears to help ecstasy users detect adulterants and results can help inform harm reduction efforts. Less frequent users in particular may require education about adulteration and drug-testing.

Keywords

Drug Users; Designer Drugs; MDMA; Hallucinogens; Harm Reduction

Introduction

Prohibited drugs are supplied through unregulated channels and thus tend to lack quality control. One specific drug that has historically been associated with adulteration is 3,4-methylenedioxymethamphetamine (MDMA) [1], also known as ecstasy [2] or “Molly” when in powder or crystalline form [3]. Its recreational use first occurred in the mid-1970s [4]. According to Analysis Anonymous, a confidential drug-checking service in the 1970s and 1980s in the United States (US), through 1985, most submitted samples said to be MDMA tested positive for MDMA (and/or similar MDx substances such as MDA) [2]. However, after its scheduling in 1985 in the US, adulteration of the drug appeared to increase. The proportion of samples that contain only MDx has varied over time and place, with notably more adulteration detected globally in the late 1990s and the late 2000s [5–10]. In the last decade, the number of novel/new psychoactive substances identified in purported MDMA has increased exponentially: by 2014 in Spain, over 30 adulterants were identified compared with less than 5 in 2000 [8]. The most common adulterants detected in European ecstasy markets between 2008–2013 included: synthetic cathinones (e.g. mephedrone, methylone, ethylone), psychedelics (e.g. 2C-B), novel amphetamines (e.g. PMA, PMMA, 4-FA), dissociatives (e.g. DXM) and piperazines (e.g. mCPP) [11]. The unwitting consumption of synthetic cathinones, or ‘bath salts’ among Americans in the electronic dance music (EDM) party scene appears to be particularly prevalent. Specifically, recent US studies found that around half of ‘ecstasy’-using EDM party attendees tested positive for novel substances they did not report using, most commonly synthetic cathinones, including ethylone, methylone, butylone, pentylone and alpha-PVP [12–14].

Decades of research has associated the consumption of pure MDMA with a range of health harms dependent on dose and context (potentiated by dancing in hot, crowded environments), including thermal stress and overheating, impaired thermoregulatory control and serotonin syndrome [15]. But over and above the management of symptoms attributable to recreational consumption of *pure* MDMA, people who consume *adulterated* MDMA may experience psychoactive effects that are discrepant with their expectations. The unwitting use of a psychedelic adulterant (e.g. 2C or NBOMe series) may result in psychosis-like symptoms [16]; the unwitting use of a substance with a delayed onset time may lead to re-dosing too soon and subsequent fatal or nonfatal overdose (e.g. PMA, PMMA) [17], and the unwitting use of any highly potent substance (e.g. fentanyl or its analogs) can lead to fatal or non-fatal overdose [18]. Synthetic cathinones in particular have resulted in deaths attributed to hyperthermia, hypertension and cardiac arrest [19], and use was associated with over 20,000 emergency room visits in the US in 2011 alone [20]. Additionally, unintentionally using the synthetic cathinone mephedrone sold as ecstasy can more easily result in

dependence as this drug has a quicker onset and shorter dose duration than MDMA which can lead to compulsive re-dosing [21].

Many people who consume ecstasy/Molly are aware that what they consume may not actually be entirely MDMA or contain any MDMA at all, and studies have found that most users desire to know the content and dose of their drug prior to consumption [22–26]. In 2017, there were ‘drug checking’ or ‘pill testing’ services available in 20 countries, where drug consumers can provide samples of substances to be analysed using laboratory equipment to determine content and/or purity [27]. Field-ready analytical methods are also utilised on-site at some dance events by harm reduction organisations to provide this information directly to consumers to inform their drug consumption choices [27]. However, the operating of ‘drug checking’ services is legally and politically fraught in many countries. In the US, promoters of dance events are discouraged from supporting drug checking services on-site due to legal liability issues, because official support for such services implies prior knowledge by promoters that prohibited drugs will be consumed at their event [28].

Therefore, in the US, ecstasy consumers rarely have access to laboratory-grade analysis of purported MDMA samples. The main option available at consumer level is the use of colour reagent test kits or chemical spot tests. These are presumptive tests primarily used for forensic applications to determine the presence or absence of specific substances [29]. A drop of reagent fluid is placed onto a shaving of the drug sample (or a pin-head-sized amount of the drug is added to the reagent mixture), then a colour reaction (or lack thereof) is observed and compared with a chart. For example, with some reagents, if the reaction is black or dark purple this may indicate presence of MDMA or another MDx, but a yellow reaction may indicate presence of a synthetic cathinone such as methylenedioxymethamphetamine. Costs are low (e.g. often \$100) compared to laboratory instruments which can cost thousands of dollars; training is simple, and the tests can easily be used in field settings, while many other drug tests require machinery [30]. If test results are interpreted with their limitations in mind, they are considered accurate at detecting the presence or absence of specific psychoactive compounds [29,30]. Popular with toxicologists and law enforcement agents, these tests have also been adopted by a significant minority of people who consume drugs, especially ecstasy [23–26,31].

A limitation of the existing literature focusing on prevalence and correlates of use of drug testing is that it has been conducted with convenience samples, and in many cases, the literature is now dated. Studies of representative samples of those at risk are needed in order to more accurately estimate more generalisable results. The increase in detections of novel substances in the US, especially synthetic cathinones among clubbers [12,13], suggests we need up-to-date and locally-relevant information about ecstasy misrepresentation, attempts by consumers to determine adulteration, and how they respond to discovering that their drugs are adulterated.

Aims

We aimed to determine: (i) prevalence and correlates of testing ecstasy with reagent test-kits; (ii) perceptions of ecstasy purity; and (iii) intentions to use less ecstasy following

hypothetical positive test results for other drugs among a representative sample of EDM attendees in New York City (NYC).

Method

Participants and procedure

We surveyed 943 individuals about to enter EDM parties at nightclubs and festivals in NYC from June through September of 2017. Time-space sampling [32] was utilised in which parties were randomly selected each week, as this allowed us to maintain elements of randomness to the selection process while surveying this targeted population. Individuals about to enter these parties were approached to be surveyed and they were eligible if they were ages 18–40 and were about to attend the randomly selected party. Trained recruiters approached passersby (who were alone or in groups) and confirmed they were about to enter the randomly selected party. Those confirmed eligible were asked if they would like to take a survey about drug use. Surveys were conducted on tablets and participants provided informed consent on the first page of the survey. The response rate was 74%. This study was approved by the institutional review board at New York University Langone Medical Center.

Measures

Participants first provided information regarding demographics, including gender, age, race/ethnicity, educational attainment, weekly income, and sexual orientation. Age was split into young adults (ages 18–24) and older adults (ages 25–40) and income was median-split into <\$500 and \$500 per week.

Participants were then asked about their use of various drugs with a focus on ecstasy. We asked whether they used ecstasy/MDMA/Molly in the past year. “Molly” was added to the definition of ecstasy in this American sample because previous research has found that not all users of this powder or crystalline form of the drug are aware that it is ecstasy [33,34]. Those reporting past-year use were asked follow-up questions about use. Past-year users were asked whether they had used in the past 30 days, and they were also asked to check off which form(s) of ecstasy were used in the past year, which included pills, powder, and crystals. Those indicating use of a particular form of ecstasy were asked to report past-year frequency of use of that form of the drug via an ordinal item with the following response options: 1–2 times, 3–5 times, 6–9 times, 10–19 times, 20–39 times, and 40 times. Frequency of use for each form was coded into tertiles: 0 times, 1–2 times, and 3 times for statistical models.

Past-year ecstasy users were then asked whether they had ever found out their ecstasy/Molly contained a drug other than MDMA and then they were asked if they had ever suspected that their ecstasy/Molly contained a drug other than MDMA. Answer options for these two questions were “yes”, “no”, and “not sure”, and those who answered “yes” were asked to type in which drug(s) they found out or suspected their ecstasy had contained. Participants were able to type in names of multiple drugs. Type-in responses were double-coded to ensure that responses accurately reflected the drugs reported and iterations continued until

100% agreement was reached. We coded the names of drugs typed in into indicator variables for descriptive purposes.

Independent of questions asking about adulteration, past-year users were then asked if they or someone else had ever tested their ecstasy/Molly using a drug testing kit in the past year. Those answering affirmatively were asked how often they test their ecstasy (or get it tested) to make sure it is really MDMA and answer options were “rarely”, “sometimes”, “usually”, and “always”. All past-year ecstasy users were then asked how often they get their ecstasy/Molly from a dealer they do not know well or trust. Answer options were “never”, “sometimes”, “often”, and “always” and we dichotomised responses into “often/always” vs. “sometimes/never”. We asked users if on average, they think their ecstasy/Molly is as pure as others’ ecstasy/Molly and answer options were “more pure”, “just as pure”, and “less pure”, and finally, we asked users if there is anything that would make them less likely to use ecstasy/MDMA/Molly again. We asked specifically if they would be less likely to use again if they found out it contained “bath salts”, methamphetamine, or MDA.

Participants were also queried about past-year use of various other drugs including ketamine, amphetamine (nonmedical use), opioids (nonmedical use), and synthetic cathinones (“bath salts”). Specifically, participants were asked about 17 different opioids (e.g. Vicodin, fentanyl) and 25 “bath salts” (e.g. methylone, “bath salt” unknown) and if the participant reported use of any, he or she was coded as using a drug in the corresponding class.

Analyses

Analyses focused on the 351 participants (of 943 surveyed in the full sample) who reported past-year ecstasy use. We first examined descriptive statistics for all relevant variables. We then compared those who reported testing their ecstasy to those who reported not testing their ecstasy along each covariate using Rao-Scott chi-square, and then fit all covariates into a multivariable model. Specifically, covariates were fit into a generalised linear model using Poisson and log link with the binary drug-testing variable as the outcome. This model produced adjusted prevalence ratios (aPR) for each covariate. Finally, we examined descriptive statistics regarding adulterant drugs participants reported finding out and/or suspecting were in their ecstasy.

We created sample weights to account for our complex survey design. Weights were calculated based on reported frequency of party attendance and response rates for each night of recruitment. Participant selection probability was computed and we weighted our prevalence estimates by the inverse of that probability [32,35]. Weighted analyses accounted for clustering of participants in each party and differential selection probability using Taylor series estimation, which allowed us to obtain accurate standard errors [36]. Data were analysed using Stata 13 SE.

Results

Prevalence of reagent test-kit use

The majority (62.9%) of this past-year ecstasy-using sample identified as white, and over half (61.8%) had at least a college degree (Table 1). Half (51.3%) of past-year users reported

past-month use. Past-year pill use (73.4%) was most common, followed by past-year use of crystals (69.6%) and powder (62.9%). A quarter (24.5%) reported using all three forms in the past year; 29.2% used pills only, 18.5% used crystals only, 8.3% used powder only, 11.9% used powder and crystals, 5.7% used pills and powder, and 1.9% used pills and crystals.

Over one-fifth (23.1%) reported having tested their ecstasy in the past year. Most of these participants reported always testing (31.2%) or usually testing (32.3%) while a quarter (24.3%) reported sometimes and 12.2% reported rarely testing their ecstasy. As shown in Table 2, compared to white participants, Asian participants were less likely to test their ecstasy (aPR=0.44, $P=0.042$). Compared to those reporting a high school education or less, those reporting having attended some college (aPR=0.49, $P=0.019$) or having earned a college degree (aPR=0.42, $P=0.049$) were also less likely to have reported testing their ecstasy. Compared to those earning <\$500 per week, those earning \$500 were more likely to report having tested their ecstasy (aPR=1.83, $P=0.005$) and compared to those identifying as heterosexual, those identifying as bisexual (aPR=2.11, $P=0.005$) or identified as another sexuality (aPR=2.75, $P=0.043$) were more than twice as likely to report having tested their ecstasy. Regarding form(s) of ecstasy used, compared to those reporting no past-year use of pills and crystals, those reporting use of pills (aPR=1.93, $P=0.043$) or crystals (aPR=1.95, $P=0.005$) 3 times in the past year were more likely to report having tested their ecstasy. Finally, compared to those who report usually not purchasing their ecstasy from an unknown or untrustworthy dealer, those who did report usually purchasing their ecstasy from such dealers were less likely to report having tested their ecstasy (aPR=0.61, $P=0.018$).

Perceptions of purity

Half (51.1%) reported that they had found out or suspected their ecstasy had contained a drug other than MDMA. Specifically, 22.8% reported both finding out and suspecting adulteration, 9.9% reporting only finding out and 18.2% reported only suspecting adulteration. Table 3 presents drugs past-year ecstasy users (independent of having tested their ecstasy in the past year) found out or suspected their ecstasy had contained. Half (49.2%) reported having found out their ecstasy had ever contained methamphetamine or speed/amphetamine, and nearly half (46.5%) suspected these drugs had been present in their ecstasy. Almost a quarter (23.9%) reported finding out their ecstasy had contained cocaine and 11.2% suspected cocaine as an adulterant.

Close to a third (32.0%) reported they felt their ecstasy, on average, is purer than others' ecstasy; 42.6% felt theirs' is just as pure; while 25.4% felt their ecstasy, on average, is less pure than others' ecstasy. While we did not fit perceived purity into the model examining correlates of testing one's ecstasy because perception of purity may be dependent on testing, we did find that perceived purity differed ($\chi^2=20.90$, $P<0.001$) by whether the participant tested his or her ecstasy. Specifically, those who had tested their ecstasy were more likely to perceive their ecstasy as purer (67.5%) compared to those believing their ecstasy is just as pure (24.0%) or less pure (8.5%).

Behavioural intentions following test results

Regarding what would make participants less likely to use ecstasy again, 54.8% reported they would be less likely to use again if they learned their ecstasy contained “bath salts”, 54.3% said they would be less likely to use again if they learned it contained methamphetamine, and 36.2% said they would be less likely to use again if MDA was detected.

Discussion

Previous research has determined that ecstasy users in NYC are at high risk for using ecstasy adulterated with “bath salts”, other novel/new psychoactive substances and/or methamphetamine [13,14], and many of these drugs can be more dangerous than MDMA, especially when ingested unknowingly or in combination with MDMA. ‘Drug checking’ has become an important harm reduction measure to help users determine content of the ecstasy or other drugs they use, but more research was needed to determine prevalence estimates and correlates of testing in this population. Further, this paper sought to fill in the gap in research regarding what adulterants users believe or found out they had unintentionally used.

We estimate that over a fifth (23.1%) of past-year ecstasy users in the NYC EDM scene have tested their ecstasy in the past year, and of these, most (63.6%) usually or always test their ecstasy. We did not detect gender or age differences regarding who was more or less likely to test their ecstasy; however, prevalence of testing was still almost 12% higher among males compared to females. While further research is needed to continue to examine user characteristics, these results do suggest that female ecstasy users may require further education about adulteration and drug-testing. Findings were somewhat mixed regarding common indicators of socioeconomic status. Specifically, while those who earned more money per week were at higher odds for testing their ecstasy, many attendees who attended or had graduated from college were at lower odds of testing their ecstasy. While it is unknown which attendees purchased their own test kits, test kits can be expensive for an individual with lower income. For example, a full set of eight reagents can cost about \$100 USD and these reagents expire after 12 months. We also found that ecstasy users with high school education were more likely to report testing their ecstasy than those with some college or completed college degrees. Further research is needed to determine the extent to which use of reagents is a function of cost, as well as knowledge about adulterants and the limitations of test kits.

More frequent ecstasy pill and crystal users were at higher odds for having tested their ecstasy in the past year. This finding corroborates previous studies finding that more regular or frequent users are more likely to report using a test kit [23,24]. Higher frequency of ecstasy use may result in more experiences of adulteration and greater knowledge of mitigation strategies. Less experienced users, however, may have less knowledge about adulteration and may also use in a more opportunistic manner (e.g. purchasing from unknown individuals), which may further place them at risk for obtaining adulterated substances. Thus, it appears more awareness about drug-testing may be needed among less frequent or novice users.

Another finding was that individuals who usually or always purchased their ecstasy from an unknown or ‘untrustworthy’ dealer were at lower odds for testing their ecstasy. The circumstances of these transactions however are unknown. Not testing and purchasing from such dealers may indicate, for example, either a lack of education about adulterants (or about ecstasy in general [34]), or simple indifference. However, purchasing ecstasy from an unknown dealer at a party may not allow the opportunity to test the drug even if one desires to do so. Other research has identified an association between obtaining drugs before the drug use event and higher likelihood of testing those drugs [26]. More research is needed to determine how risky purchasing practices relate to use of harm reduction tactics such as reagent testing.

Half (51.1%) of ecstasy users reported that they had either found out or suspected their ecstasy had contained a drug other than MDMA, and half (49.2%) of these individuals found out or suspected adulteration with methamphetamine/”speed”. These results add to results from a previous study of this population in which among past-year ecstasy users whose hair tested positive for methamphetamine, over half (55.2%) denied known use [13]. Interestingly, only 2.9% reported finding out their ecstasy contained “bath salts” (and only 5.1% suspected adulteration with “bath salts”) as previous studies of this population have found that among ecstasy users whose hair tested positive for “bath salts”, 50.0–68.0% denied known use of such compounds [13,14]. Therefore, ecstasy users in NYC may be underestimating their exposure to such compounds.

Behavioral intentions regarding future ecstasy use also appear to be affected according to hypothetical situations of individuals learning their ecstasy contains or contained adulterants. For example, over half of users said they would be less likely to use ecstasy again if they learned their ecstasy contained “bath salts” or methamphetamine. Thus, learning about the extent of adulteration or learning one’s ecstasy is adulterated may be preventative against future use or encourage users to engage in harm reduction practices such as ecstasy-testing, as suggested by prior research [23,37,38].

Limitations

Results may not be fully generalisable to the EDM scene outside of NYC. Frequency of ecstasy use was queried separately for pills, powder, and crystals, and the ordinal nature of the items (and lack of data regarding whether separate forms were used on the same or on separate occasions), we were unable to combine these items into a single frequency of use variable. High-frequency use (e.g. use 10+ times) for any single form was rare (5%) when considering those who used a particular form 0 times, and after extensive sensitivity testing, we concluded that our current trichotomous frequency variables best represented the data and provided the best fit in models.

Conclusions

Individuals in the NYC EDM scene are at high risk of consuming ecstasy that contains other psychoactive substances, including ‘bath salts’ which are associated with greater dependence liability and higher overdose rates (fatal and non-fatal) than MDMA alone [19,21]. Evidence from other countries demonstrates that drug checking services can prevent

such unintentional consumption of novel/new psychoactive substances [38–40]. Drug checking services are preferable to self-administration of reagent test kits because more sophisticated equipment can be utilised and staff are trained in equipment use, interpretation, and provision of harm-reducing advice [41]. These data reiterate an urgent need for policy changes that facilitate development of drug checking services. The operation of such services in over 20 countries [27] provides multiple blueprints from which to design an appropriate service to cater for the needs of this population.

Acknowledgements

Research reported in this publication was supported by the National Institute on Drug Abuse of the National Institutes of Health under Award Number K01DA038800 (PI: Palamar). Barratt is supported by a fellowship from Australia's National Health and Medical Research Council (APP1070140). The National Drug and Alcohol Research Centre and the National Drug Research Institute receive funding from the Australian Government under the Substance Misuse Prevention and Service Improvement Grants Fund. We also acknowledge the contribution of the Victorian Operational Infrastructure Support Program received by the Burnet Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or any of the other funders listed.

References

- Hayner GN. MDMA misrepresentation: An unresolved problem for ecstasy users. *J Psychoactive Drugs* 2002;34:195–8. [PubMed: 12691209]
- Renfro CL. MDMA on the street: Analysis Anonymous. *J Psychoactive Drugs* 1986;18:363–9. [PubMed: 2880953]
- Palamar JJ. There's something about Molly: The underresearched yet popular powder form of ecstasy in the United States. *Subst Abus* 2017;38:15–7. [PubMed: 27925866]
- Shulgin A. The background and chemistry of MDMA. *J Psychoactive Drugs* 1986;18:291–304. [PubMed: 2880943]
- Vogels N, Brunt TM, Rigter S, van Dijk P, Vervaeke H, Niesink RJM. Content of ecstasy in the Netherlands: 1993–2008. *Addiction* 2009;104:2057–66. [PubMed: 19804461]
- Brunt TM, Niesink RJM, van den Brink W. Impact of a transient instability of the ecstasy market on health concerns and drug use patterns in The Netherlands. *Int J Drug Policy* 2012;23:134–40. [PubMed: 21741814]
- Parrott AC. Is ecstasy MDMA? A review of the proportion of ecstasy tablets containing MDMA, their dosage levels, and the changing perceptions of purity. *Psychopharmacology (Berl)*. 2004;173:234–41. [PubMed: 15007594]
- Vidal-Giné C, Ventura-Vilamala M, Fornís-Espinosa I, Gil-Lladanosa C, Calzada-Álvarez N, Fitó-Fruitós A, et al. Crystals and tablets in the Spanish ecstasy market 2000–2014: Are they the same or different in terms of purity and adulteration? *Forensic Sci Int* 2016;263:164–8. [PubMed: 27129144]
- Chen C, Kostakis C, Harpas P, Felgate PD, Irvine RJ, White JM. Marked decline in 3,4-methylenedioxymethamphetamine (MDMA) based on wastewater analysis. *J Studies Alcohol Drugs* 2011;72:737–40.
- Tanner-Smith EE. Pharmacological content of tablets sold as “ecstasy”: Results from an online testing service. *Drug Alcohol Depend* 2006;83:247–54. [PubMed: 16364567]
- Brunt TM, Nagy C, Bucheli A, Martins D, Ugarte M, Beduwe C, et al. Drug testing in Europe: Monitoring results of the Trans European Drug Information (TEDI) project. *Drug Test Anal* 2017;9:188–98. [PubMed: 26888408]
- Mohr ALA, Friscia M, Yeakel JK, Logan BK. Use of synthetic stimulants and hallucinogens in a cohort of electronic dance music festival attendees. *Forensic Sci Int* 2018;282:168–78. [PubMed: 29216523]

13. Palamar JJ, Salomone A, Gerace E, Di Corcia D, Vincenti M, Cleland CM. Hair testing to assess both known and unknown use of drugs amongst ecstasy users in the electronic dance music scene. *Int J Drug Policy* 2017;48:91–8. [PubMed: 28810159]
14. Palamar JJ, Salomone A, Vincenti M, Cleland CM. Detection of ‘bath salts’ and other novel psychoactive substances in hair samples of ecstasy/MDMA/‘Molly’ users. *Drug Alcohol Depend* 2016;161:200–5. [PubMed: 26883685]
15. Parrott AC. Human psychobiology of MDMA or ‘ecstasy’: An overview of 25 years of empirical research. *Hum Psychopharmacol* 2013;28:289–307. [PubMed: 23881877]
16. Hill SL, Doris T, Gurung S, Katebe S, Lomas A, Dunn M, et al. Severe clinical toxicity associated with analytically confirmed recreational use of 25I-NBOMe: Case series. *Clin Toxicol (Phila)* 2013;51:487–92. [PubMed: 23731373]
17. Caldicott DGE, Edwards NA, Kruys A, Kirkbride KP, Sims DN, Byard RW, et al. Dancing with “death”: P-methoxyamphetamine overdose and its acute management. *J Toxicol Clin Toxicol*. 2003;41:143–54. [PubMed: 12733852]
18. Suzuki J, El-Haddad S. A review: Fentanyl and non-pharmaceutical fentanyls. *Drug Alcohol Depend* 2017;171:107–16. [PubMed: 28068563]
19. Zaami S, Giorgetti R, Pichini S, Pantano F, Marinelli E, Busardo FP. Synthetic cathinones related fatalities: An update. *Eur Rev Med Pharmacol Sci* 2018;22:268–74. [PubMed: 29364498]
20. Substance Abuse and Mental Health Services Administration. “Bath salts” were involved in over 20,000 drug-related emergency department visits in 2011. The DAWN Report Data Spotlight, Drug Abuse Network; 2013.
21. Green AR, King MV, Shortall SE, Fone KC. The preclinical pharmacology of mephedrone; Not just MDMA by another name. *Br J Pharmacol* 2014;171:2251–68. [PubMed: 24654568]
22. Hansen D, Maycock B, Lower T. ‘Weddings, parties, anything...’, a qualitative analysis of ecstasy use in Perth, Western Australia. *Int J Drug Policy* 2001;12:181–99. [PubMed: 11399420]
23. Johnston J, Barratt MJ, Fry CL, Kinner S, Stoové M, Degenhardt L, et al. A survey of regular ecstasy users’ knowledge and practices around determining pill content and purity: Implications for policy and practice. *Int J Drug Policy* 2006;17:464–72.
24. Davis AK, Rosenberg H. Specific harm reduction strategies employed by 3,4-methylenedioxymethamphetamine/ ecstasy users in the United States and the United Kingdom. *Drug Science, Policy and Law* 2017;3:doi:2050324517711069.
25. Day N, Criss J, Griffiths B, Gujral SK, John-Leader F, Johnston J, et al. Music festival attendees’ illicit drug use, knowledge and practices regarding drug content and purity: A cross-sectional survey. *Harm Reduct J* 2018;5;15:1.
26. Barratt MJ, Bruno R, Ezard N, Ritter A. Pill testing or drug checking in Australia: Acceptability of service design features. *Drug Alc Review* 2018;37:226–36.
27. Barratt MJ, Kowalski M, Maier LJ, Ritter A. Global review of drug checking services operating in 2017. *Drug Policy Modelling Program Bulletin No. 24*. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney; 2018.
28. Levy BA. When cute acronyms happen to bad legislation: The Reducing Americans’ Vulnerability to Ecstasy “RAVE” act. *Northwestern University Law Review* 2004;98:1251–89.
29. O’Neal CL, Crouch DJ, Fatah AA. Validation of twelve chemical spot tests for the detection of drugs of abuse. *Forensic Sci Int* 2000;109:189–201. [PubMed: 10725655]
30. Harper L, Powell J, Pijl EM. An overview of forensic drug testing methods and their suitability for harm reduction point-of-care services. *Harm Reduction J* 2017;14:52.
31. Duterte M, Jacinto C, Sales P, Murphy S. What’s in a label? Ecstasy sellers’ perceptions of pill brands. *J Psychoactive Drugs* 2009;41:27–37. [PubMed: 19455907]
32. MacKellar DA, Gallagher KM, Finlayson T, Sanchez T, Lansky A, Sullivan PS. Surveillance of HIV risk and prevention behaviors of men who have sex with men—A national application of venue-based, time-space sampling. *Public Health Rep* 2007;122(s1):39–47. [PubMed: 17354526]
33. Palamar JJ, Keyes K, Cleland CM. Underreporting of ecstasy use among high school seniors in the US. *Drug Alcohol Depend* 2016;165:279–82. [PubMed: 27296977]
34. Palamar JJ. What’s in a name? Correlates of ecstasy users knowing or agreeing that Molly is ecstasy/MDMA. *J Psychoactive Drugs* 2017;22:1–6.

35. Jenness SM, Neaigus A, Murrill CS, Gelpi-Acosta C, Wendel T, Hagan H. Recruitment-adjusted estimates of HIV prevalence and risk among men who have sex with men: Effects of weighting venue-based sampling data. *Public Health Rep.* 2011;126:635–42. [PubMed: 21886323]
36. Heeringa SG, West BT, Berglund PA. *Applied survey data analysis*. London: Chapman and Hall; 2010.
37. Benschop A, Rabes M, Korf DJ. *Pill testing, ecstasy and prevention A scientific evaluation in three European cities*. Amsterdam: Rozenberg Publishers; 2002.
38. Martins D, Barratt MJ, Vale Pires C, Carvalho H, Ventura Vilamala M, Fornís Espinosa I, et al. The detection and prevention of unintentional consumption of DOx and 25x-NBOMe at Boom Festival. *Hum Psychopharmacol* 2017;32:e2608.
39. Royal Society for Public Health. *Drug safety testing at festivals and night clubs*. London: Royal Society for Public Health; 2017.
40. Munn MB, Lund A, Golby R, Turris SA. Observed benefits to on-site medical services during an annual 5-day electronic dance music event with harm reduction services. *Prehospital Disaster Med* 2016;31:228–34. [PubMed: 26831043]
41. Grabenhofer S, Kociper K, Nagy C, Luf A, Schmid R. Drug checking und aufklärung vor ort in der niedrigschwelligen präventionsarbeit [Drug checking and on-site reconnaissance in low-threshold prevention work]. In: von Heyden M, Jungaberle H, Maji T, editors *Handbuch Psychoaktive Substanzen*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2016 p. 327–338.

Table 1.Sample characteristics of past-year ecstasy users ($N=351$)

Characteristic	<i>n</i>	Weighted %
<i>Sex</i>		
Male	191	52.0
Female	160	48.0
<i>Age</i>		
18–24	147	41.0
25–40	204	59.0
<i>Race/ethnicity</i>		
White	222	62.8
Black	11	2.5
Hispanic	61	19.9
Asian	30	8.8
Other/mixed	27	6.1
<i>Education</i>		
High school or less	38	11.7
Some college	89	26.5
College degree	172	50.2
Graduate school	52	11.6
<i>Weekly income</i>		
<\$500	158	46.9
\$500	193	53.1
<i>Sexual orientation</i>		
Heterosexual	263	73.8
Gay/lesbian	36	10.6
Bisexual	43	12.1
Other sexuality	9	3.5
<i>Number of times used ecstasy pills in past year</i>		
0 Times	119	38.8
1–2 Times	110	34.8
3 times	122	26.4
<i>Number of times used ecstasy powder in past year</i>		
0 Times	129	49.4
1–2 Times	102	28.5
3 Times	120	22.1
<i>Number of times used ecstasy crystals in past year</i>		
0 Times	136	43.1
1–2 Times	95	30.8
3 Times	120	26.2
<i>Purchase ecstasy from unknown or untrustworthy dealer</i>		
Sometimes or Never	190	60.2

Characteristic	<i>n</i>	Weighted %
Usually or Always	160	39.8
<i>Past-12 month other drug use</i>		
Ketamine	103	16.4
Amphetamine (nonmedical)	143	30.5
Opioids (nonmedical)	58	18.3
"Bath salts"	14	2.8

Table 2.

Characteristics of past-year ecstasy users who report having tested their ecstasy

	Have not tested (76.9%, N=248), Weighted %	Have tested (23.1%, N=103), Weighted %	χ^2	aPR	(95% CI)
<i>Sex</i>			2.38		
Male	71.3	28.7		1.00	
Female	83.0	17.0		0.95	(0.57, 1.60)
<i>Age</i>			0.29		
18–24	79.0	21.0		1.00	
25–40	75.4	24.6		1.40	(0.77, 2.55)
<i>Race/ethnicity</i>			2.05		
White	76.5	23.5		1.00	
Black	56.6	43.4		0.84	(0.34, 2.07)
Hispanic	69.7	30.3		1.22	(0.68, 2.20)
Asian	91.2	8.8		0.45 *	(0.21, 0.97)
Other/mixed	92.4	7.6		0.55	(0.12, 2.63)
<i>Education</i>			3.47 *		
High school or less	58.9	41.1		1.00	
Some college	79.3	20.7		0.49 *	(0.28, 0.86)
College degree	82.9	17.1		0.41 *	(0.19, 0.89)
Graduate school	63.2	36.8		0.55	(0.21, 1.46)
<i>Weekly income</i>			3.30		
<\$500	83.3	16.7		1.00	
\$500	71.2	28.8		1.83 **	(1.19, 2.82)
<i>Sexual orientation</i>			2.98 *		
Heterosexual	83.1	16.9		1.00	
Gay/Lesbian	58.9	41.1		2.01	(0.80, 5.07)
Bisexual	57.4	42.6		2.11 **	(1.28, 3.49)
Other Sexuality	68.9	31.1		2.42	(0.94, 6.30)
<i>Past-year ecstasy pill use</i>			4.62 *		
0 times	77.6	22.4		1.00	
1–2 times	88.6	11.4		0.63	(0.32, 1.24)
3+ times	60.5	39.5		1.89 *	(1.04, 3.41)
<i>Past-year ecstasy powder use</i>			0.47		
0 times	78.8	21.3		1.00	
1–2 times	78.2	21.8		1.43	(0.93, 2.18)
3+ times	71.2	28.8		0.94	(0.52, 1.67)
<i>Past-year ecstasy crystal use</i>			3.16 *		
0 times	80.5	19.5		1.00	
1–2 times	84.7	15.3		0.85	(0.33, 2.17)
3+ times	61.7	38.3		1.90 **	(1.22, 2.96)

	Have not tested (76.9%, N=248), Weighted %	Have tested (23.1%, N=103), Weighted %	χ^2	aPR	(95% CI)
<i>Purchase from unknown dealer</i>			3.29		
No	71.9	28.1		1.00	
Yes	84.2	15.8		0.63 *	(0.41, 0.96)
<i>Past-12 month drug use</i>					
Ketamine	74.5	25.5	0.12	1.18	(0.64, 2.18)
Amphetamine (nonmedical)	79.4	20.6	0.26	0.70	(0.37, 1.35)
Opioids (nonmedical)	73.9	26.1	0.22	0.68	(0.40, 1.16)
"Bath salts"	35.6	64.4	6.74 *	1.16	(0.55, 2.47)

aPR, adjusted prevalence ratio; CI, confidence interval. The adjusted model contained all covariates presented in the table.

* $P < 0.05$,

** $P < 0.01$.

Table 3.

Drugs participants reported finding out or suspecting were adulterants in their ecstasy

	Found out Weighted %	Suspected Weighted %
<i>Methamphetamine/"speed"/amphetamine</i>	49.2	46.5
Methamphetamine	30.1	30.9
Speed	12.2	10.9
Amphetamine	8.1	5.6
Cocaine	23.9	11.2
MDMA ⁺	3.8	1.5
MDA	3.1	2.2
<i>Bath salts</i>	2.9	5.1
Methylone	2.5	2.3
Methadone ⁺⁺	1.8	0.4
Caffeine	1.4	0.8
2C	1.4	0.2
LSD	1.2	1.0
Heroin	1.0	9.3
Ketamine	0.0	1.8
Non-specified	7.9	13.6
Don't know	7.2	12.1
Miscellaneous product/drug	2.5	1.6

Note: While "speed" most commonly refers to methamphetamine, sometimes it refers to amphetamines in general so we assigned it its own subcategory.

⁺ Some participants may have misunderstood the questions or were unaware that MDMA is (or is supposed to be) ecstasy/Molly,

⁺⁺ Some participants likely confused "methadone" with either methylone or methamphetamine.