CASE REPORT

Solid papillary thyroid carcinoma with Hashimoto's thyroiditis: description of a further case with challenging cytological features

Franco Fulciniti,¹ Jessica Barizzi,¹ Pierpaolo Trimboli,² Luca Giovannella²

¹Pathology, Institute of Pathology, Locarno, Switzerland
²Department of Nuclear Medicine and Thyroid Centre, Oncology Institute of Southern Switzerland, Bellinzona, Switzerland

Correspondence to

Dr Franco Fulciniti,
 franco.fulciniti@gmail.com

Accepted 17 December 2018

SUMMARY

Solid papillary thyroid carcinoma (SV-PTC) is a rare variant which is mainly observed in young patients with a history of exposure to ionising radiations. Neoplasms belonging to such category generally carry RET-PTC (REarranged during Transfection- Papillary Thyroid Carcinoma) fusions and seem to have a slightly worse prognosis with respect to classical and follicular variants of papillary thyroid carcinoma (PTC), though consistent prognostic and survival data are scarce. SV-PTC should be differentiated from trabecular-insular poorly differentiated thyroid carcinomas, which occur in a different age group and carry a dismal prognosis. These latter tumours do not show the typical nuclear features of PTC and show tumour necrosis with an high mitotic activity. In this report a further case of SV-PTC is described which was associated to Hashimoto's thyroiditis, a finding never described in the cytological literature up to now for SV-PTC; this association created further differential diagnostic problems. The neoplasm displayed RET-PTC1 fusion.

BACKGROUND

Solid variants account for ~3% of all papillary thyroid carcinoma (PTC) cases.¹ These are formed by differentiated thyrocytes which have the nuclear features of classic PTC but generally show minor (<50%) or absent papillary architecture, both in fine-needle aspirates and in tissue samples. Two major case series have been published in the histopathological literature.^{2,3} Four extensive case series of solid papillary thyroid carcinoma (SV-PTC)⁴⁻⁷ have been published in the cytopathological literature, plus some case reports.⁸⁻¹¹ Several of these cases occur in the paediatric age and may have a history of ionising radiation exposure.^{1-3,11} Epidemiological data are not always coherent: some authors underline a slightly worse prognosis of SV-PTC as compared with classical PTC and its follicular variants.

In this report we describe a further case of SV-PTC carrying a RET-PTC fusion and in which an association with Hashimoto's thyroiditis was documented cytohistologically and clinically.

CASE PRESENTATION

A 21-year-old man with chronic thyroiditis presented for an endocrinological check-up. During the clinical interview, he declared to suffer from slight tachycardia and shortness of breath while training in a body-building fitness programme. He displayed diffuse muscular hypertrophy and admitted the chronic

assumption of anabolising hormones plus off-label testosterone tablets. An ultrasound examination of the thyroid performed at the time of the initial consultation disclosed an unhomogeneous gland structure, in keeping with chronic thyroiditis. His TSH was 2.6 UI/mL under L-thyroxin therapy. At a difference with an ultrasound examination performed 1 year before, a hypoechogenic solid nodule with innumerable microcalcifications, measuring 15×15 mm, was now found in the apical paraisthmic region of the left lobe (figure 1). An ultrasound-guided fine needle aspiration biopsy (FNAB) was performed on this nodule (three passes with a 25 gauge needle) and several smears were obtained, which were stained with May-Grünwald-Giemsa (MGG) and Papanicolaou (PAP). Needle washings were also performed and kept in CytoLyt solution. A liquid-based cytological preparation was also obtained, which was stained with PAP.

The cytological picture was very cellular. Numerous epithelial cell sheets and clusters could be noticed in a background containing abundant lymphoid cells in various degrees of maturation and mature plasma cells. Both normal Hürthle-type cells and atypical thyrocytes could be seen: these latter also displayed oxyphilic morphology and were present as large syncytial sheets or three-dimensional clusters with ragged borders and architectural disorder, without an evident follicular or papillary aggregation (figure 2). They had irregularly shaped, oval nuclei, larger than those of usual PTC, with marked anisonucleosis, granular chromatin and frequent nuclear grooves or infoldings. Their cytoplasm were granular and cyanophylic on PAP staining (figures 3A and 4). Occasional mitotic figures could be observed. The neoplastic cells showed diffuse immunocytochemical expression of CK 19 (figure 3B). The background also showed numerous dystrophic microcalcifications.

INVESTIGATIONS

Two MGG-stained smears and the needle washings were sent for molecular pathological examination. Next generation sequencing was performed by using a Ion Ampliseq Colon & Lung Cancer Panel (Life Technology, Ion Torrent PGM). PCR was also performed for RNA study; a RET-PTC1 fusion was found. A diagnosis of positive for malignancy (Tir 5 according to the 2014 modification of the Italian consensus for classification and reporting of thyroid

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Fulciniti F, Barizzi J, Trimboli P, et al. *BMJ Case Rep* 2019;**12**:e226153. doi:10.1136/bcr-2018-226153

Figure 1 Transversal ultrasound scan of the thyroid gland showing a hypoechoic solid nodule with innumerable fine calcifications.

cytology)¹² was formulated. Further comments to the cytological diagnosis indicated the possibility of a non-classical variant of PTC.

TREATMENT

Total thyroidectomy was performed. One left neck lymph node was also excised. The thyroid weighed 40 g and showed a left apical nodular area of firm consistency, 15 mm in diameter, with a gitting cut surface.

OUTCOME AND FOLLOW-UP

Histopathological findings

The nodular area of the left lobe showed a neoplastic proliferation of thyrocytes with solid-trabecular and syncytial growth. The solid cell nests were separated by thin fibrocapillary septa and/or by thicker fibrous bands. (figure 5). The neoplasm was immersed in a florid lymphoid proliferation with formation of

Figure 2 Fine-needle aspiration biopsy sample. Direct smear showing a sheet of atypical epithelial cells with evident phenomena of anisocytosis and anisonucleosis. The cells show irregular chromatin structure, with folded nuclear membrane and dense, granular bluish cytoplasm. Numerous lymphoid cells can be seen in the background as well as some normal looking thyrocytes, (lower centre) May-Grünwald-Giemsa, ×40, original magnification.

Figure 3 (A) Fine needle aspiration biopsy sample: liquid-based cytology sample, showing an epithelial cell cluster with ragged borders. The cells show pleomorphic nuclei, with folded nuclear membranes and evident chromocentres, with dense cytoplasm, mostly of polygonal shape. More than occasional lymphoid cells can be seen in the background. Papanicolaou, ×630, original magnification. (B) (inset) immunocytochemical staining, showing diffuse cytoplasmic expression of CK19 by the neoplastic cells (alkaline phosphatase, ×630, original magnification).

lymphoid follicles. The neoplastic cells nests were formed by medium sized thyrocytes having the classic nuclear features of PTC and oxyphylic differentiation (figures 6–7).

Well-formed papillary or follicular structures were almost completely absent (both being <5% of the neoplasm). The neoplastic cells expressed diffusely TTF1, PAX8 (figure 8A,B), CK19, CK7, Galectin3 and HBME1. The Ki67 proliferation rate was 2%. There was no necrosis. A histopathological diagnosis of SV-PTC (pT1a, pN0) was made.

Figure 4 Fine needle aspiration biopsy sample. Direct smear showing an irregular 3-D three dimensional cluster of neoplastic cells showing pleomorphic nuclei with irregular chromatin structure, folded nuclei and some intranuclear inclusions (right half of picture). A mitotic figure may be seen in the upper left portion of the picture. Notice the absence of branching phenomena, or of papillary or follicular structures. The cytoplasm is well defined, granular, cyanophylic and tend to the polygonal shape. Papanicolaou, ×630, original magnification.

Figure 5 Surgical sample. Low-power microscopic magnification, showing a neoplastic cell growth with solid and syncytial pattern. The neoplastic cell solid nests are separated by thin fibrovascular septa. There is no definite papillary or follicular pattern and a lymphoid infiltrate is evident in the background. H&E, ×100, original magnification.

DISCUSSION

SV-PTC is a rare variant of thyroid carcinoma with solid-trabecular or syncytial growth, having the classic nuclear features of PTC but with scant or absent papillary architecture. The extent of the solid growth should be >50% of its total volume.¹⁻⁴ SV-PTC should be considered a further variant of PTC, differing from all of the other ones because of its peculiar pattern of growth. Nikiforov *et al*³ analysed a series of 20 cases of SV-PCT, with a survival rate of 90% after a mean follow-up of 18.7 years, slightly worse than that observed for classic PTC (93.5% with a mean follow-up of 16.6 years). RET/PCT rearrangement was demonstrated in 3/7 SV-PCT (43%), of which two cases were RET/PTC1 and 1 was RET/PTC3.

Numerous SV-PTCs have been reported in the cytopathological literature.^{4-11 13-16} In FNAB samples, they were found to have syncytial, microfollicular or dyshesive growth in more or less one-third of the cases each.⁵

Figure 6 Surgical sample. Medium-power microscopic magnification, showing a neoplastic growth with trabecular and solid pattern to which a lymphoplasmacellular infiltrate is intimately admixed. Notice the irregularly-cleft nuclei and the granular, eosinophilic cytoplasm of the neoplastic cells. H&E, ×400, original magnification.

Figure 7 Surgical sample. High-power microscopic magnification, showing the nucleo-cytoplasmic features of the epithelial component. Notice the presence of distinct nuclear atypia, somewhat superior to that of ordinary variants of papillary thyroid carcinoma and the well-defined, granular, eosinophilic cytoplasm. An accompanying lymphoid infiltrate is admixed to the neoplastic cell population. H&E, ×630, original magnification.

Despite the numerous reports, it is surprising that the association with Hashimoto's thyroiditis is only present in an histopathological series examining exclusively paediatric cases from Japan and Ukraine, in which it was respectively demonstrated in 25.9% and 29.4% of tumours with dominant solid-trabecular growth.¹⁷ To our knowledge, the current case seems to be the first one observed in the cytological literature in which this association has been described preoperatively.

We were perplexed by this case because, despite the evident lymphocytic infiltration, which indicated concurrent Hashimoto's disease, this neoplasms, unlike the so-called 'Warthin-like' PTC,¹⁸⁻²⁵ did not display lymphocytic infiltration of the papillary stalks of the carcinomatous component, as these latter were completely absent.

Figure 8 (A) Surgical sample. Immunohistochemical stain for TTF1 showing diffuse expression by the neoplastic cells. The stain also enhances the solid and alveolar pattern of growth of the neoplasm. Immunoperoxidase, ×200, original magnification. (B) Surgical sample. Immunohistochemical stain, showing expression of PAX8 in most neoplastic nuclei. Immunoperoxidase, ×400, original magnification.

In this aspect, the deep mingling of the lymphoid component with the carcinomatous component was, instead, somewhat more reminiscent of some lymphoepithelial carcinomas of other organs. Moreover, due to the characteristic nuclear features of the neoplasm and to the absence of mitotic figures or necrosis, the possibility of a poorly differentiated thyroid carcinoma had already been excluded. So, the possibility of a solid variant of PTC or that one of a tall cell variant was left open, the latter one due to the oxyphylic appearance of the cytoplasm and the higher than average degree of atypia of the neoplastic cells.^{15 16 26 27} A final comment to the cytopathological diagnosis was issued, stating that a non-classical variant of PTC could not be excluded.

Based on the criteria exposed in the literature, we believe that the diagnosis of the SV-PTC could be entertained in cases showing solid-trabecular/syncytial pattern of growth, as the one here described. SV-PTC could also be considered in cases with dyshesive pattern, in which the absence of papillary stalks and of 'branching' features in the cell clusters could lead to a correct diagnosis. SV-PTC cases with microfollicular pattern may be more difficult to differentiate from the Follicular Variant of Papillary Carcinoma (FVPC), with frequent nuclear overlapping and clear-cut nuclear features of PTC, as opposed to weaker nuclear criteria addressing for FVPC, being the major differential criteria.

Molecular pathology studies of the SV-PTC have found RET/PTC1 or RET/PTC3 rearrangements, as the current case, or B-RAF (B-Raf protein) mutations, with at least two novel mutations: 1) the B-RAFK600-1E¹³ and 2) B-RAFK600 delinsAL.¹⁴

Moreover, 43% of the SV-PCT studied by Prasad *et al*²⁸ showed ETV6/NTRK3 fusions (as in secretory breast and salivary gland carcinoma). NTRK fusions in SV-PCT become, in this way, the second most common molecular pathological abnormality in this neoplasm.

Interestingly, in a recent Japanese study,²⁹ the expression of two cancer-stem cell markers: ABCG2 (ATP binding cassette 2) and multidrug resistance associated protein 1 was statistically more frequent in SV-PTC as compared with usual PTC and was significantly associated to adverse clinical outcomes.

We believe that the study of more extensive case series may still add significant prognostic information on this singular subtype of PTC.

Learning points

- Solid papillary thyroid carcinoma (SV-PTC) is a rare variant of papillary thyroid carcinoma that may raise diagnostic problems on fine needle aspiration biopsy samples if associated to lymphocytic thyroiditis.
- Some SV-PTC may host, besides to RET-PTC fusions, supplementary genetic alterations, like novel or uncommon BRAF mutations and 43% of them also show ETV6-NTRK3 fusions:
- the prognosis of SV-PTC may be slightly worse than classical variant/usual papillary thyroid carcinoma.
- SV-PTC associated to lymphocytic thyroiditis seems to be more frequent in Asian paediatric patients with or without a previous history of radiation to the neck.

Contributors FF and PT saw the patient and made the diagnosis. FF and JB edited the manuscript. PT and LG were responsible for the patient's treatment and coauthored this paper.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Rosai J, Alboreo Saavedra J, Ascoli S. Papillary thyroid carcinoma. In: Lloyd RV, Osamura RY, Klöppel G, Rosai J. eds. *WHO classification of tumors of endocrine organs*. 4th edn. Lyon: International Agency for Research on Cancer, 2017:81–91.
- 2 Carcangiu ML, Zampi G, Pupi A, *et al*. Papillary carcinoma of the thyroid. A clinicopathologic study of 241 cases treated at the University of Florence, Italy. *Cancer* 1985;55:805–28.
- 3 Nikiforov YE, Erickson LA, Nikiforova MN, *et al*. Solid variant of papillary thyroid carcinoma: incidence, clinical-pathologic characteristics, molecular analysis, and biologic behavior. *Am J Surg Pathol* 2001;25:1478–84.
- 4 Chang H, Kim SM, Chun KW, *et al*. Clinicopathologic features of solid variant papillary thyroid cancer. *ANZ J Surg* 2014;84:380–2.
- 5 Giorgadze TA, Scognamiglio T, Yang GC. Fine-needle aspiration cytology of the solid variant of papillary thyroid carcinoma: a study of 13 cases with clinical, histologic, and ultrasound correlations. *Cancer Cytopathol* 2015;123:71–81.
- 6 Higuchi M, Hirokawa M, Suzuki A, *et al*. Cytological features of solid variants of papillary thyroid carcinoma: a fine needle aspiration cytology study of 18 cases. *Cytopathology* 2017;28:268–72.
- 7 Ohashi R, Murase Y, Matsubara M, *et al*. Fine needle aspiration cytology of the papillary thyroid carcinoma with a solid component: a cytological and clinical correlation. *Diagn Cytopathol* 2017;45:391–8.
- 8 Nguyen GK, Lee MW. Solid/trabecular variant papillary carcinoma of the thyroid: Report of three cases with fine-needle aspiration. *Diagn Cytopathol* 2006;34:712–4.
- 9 Troncone G, Russo M, Malapelle U, *et al*. Cytological and molecular diagnosis of solid variant of papillary thyroid carcinoma: a case report. *Cytojournal* 2008;5:2.
- 10 Damle N, Ramya S, Bal C, *et al*. Solid variant of papillary thyroid carcinoma with no history of radiation exposure. *Indian J Nucl Med* 2011;26:196–8.
- 11 Abdul Raman WF, Mn MH, Win TT, *et al*. Solid variant of papillary thyroid carcinoma in a 14-year-old girl. *BMJ Case Rep* 2013.
- 12 Nardi F, Basolo F, Crescenzi A, *et al*. Italian consensus for the classification and reporting of thyroid cytology. *J Endocrinol Invest* 2014;37:593–9.
- 13 Trovisco V, Vieira de Castro I, Soares P, *et al*. BRAF mutations are associated with some histological types of papillary thyroid carcinoma. *J Pathol* 2004;202:247–51.
- 14 Chiosea S, Nikiforova M, Zuo H, *et al*. A novel complex BRAF mutation detected in a solid variant of papillary thyroid carcinoma. *Endocr Pathol* 2009;20:122–6.
- 15 Yang GCH. Papillary thyroid carcinoma, uncommon variants and mimicker. Yang CGH, ed. *Thyroid Fine Needle Aspiration*. London, UK: Cambridge University Press, 2013:18–23.
- 16 Hawk WA, Hazard JB. The many appearances of papillary carcinoma of the thyroid. *Cleve Clin J* 1976;43:207–16.
- 17 Bogdanova TI, Saenko VA, Hirokawa M, *et al*. Comparative histopathological analysis of sporadic pediatric papillary thyroid carcinoma from Japan and Ukraine. *Endocr J* 2017;64:977–93.
- 18 Apel RL, Asa SL, LiVolsi VA. Papillary Hürthle cell carcinoma with lymphocytic stroma. "Warthin-like tumor" of the thyroid. *Am J Surg Pathol* 1995;19:810–4.
- 19 Baloch ZW, LiVolsi VA. Warthin-like papillary carcinoma of the thyroid. *Arch Pathol Lab Med* 2000;124:1192–5.
- 20 D'Antonio A, De Chiara A, Santoro M, *et al*. Warthin-like tumour of the thyroid gland: RET/PTC expression indicates it is a variant of papillary carcinoma. *Histopathology* 2000;36:493–8.
- 21 Yeo M, Bae JS. The Warthin-like variant of papillary carcinoma: a comparison with classic type in the patients with co-existing Hashimoto's thyroiditis. *International J of Endocrinol* 2015;456027.
- 22 Chong Y, Suh S, Kim TJ, *et al*. Fine needle aspiration cytology of warthin-like papillary thyroid carcinoma: a brief case report. *Korean J Pathol* 2014;48:170–3.
- 23 Paliogiannis P, Attene F, Trogu F, *et al*. Warthin-like papillary carcinoma of the thyroid gland: case report and review of the literature. *Case Rep Oncol Med* 2012;2012:1–4.
- 24 Erşen A, Durak MG, Canda T, *et al*. Warthin-like papillary carcinoma of the thyroid: a case series and review of the literature. *Türk Patoloji Derg* 2013;29:150–5.
- 25 González-Colunga KJ, Loya-Solis A, Ceceñas-Falcón LÁ, *et al*. Warthin-Like Papillary thyroid carcinoma associated with lymphadenopathy and Hashimoto's thyroiditis. *Case Rep Endocrinol* 2015;2015:251898.
- 26 Guan H, Vandenbussche CJ, Erozan YS, *et al*. Can the tall cell variant of papillary thyroid carcinoma be distinguished from the conventional type in fine needle aspirates? A cytomorphologic study with assessment of diagnostic accuracy. *Acta Cytol* 2013;57:534–42.
- 27 Lastra RR, LiVolsi VA, Baloch ZW. Aggressive variants of follicular cell-derived thyroid carcinomas: a cytopathologist's perspective. *Cancer Cytopathol* 2014;122:484–503.
- 28 Prasad ML, Vyas M, Horne MJ, *et al*. NTRK fusion oncogenes in pediatric papillary thyroid carcinoma in northeast United States. *Cancer* 2016;122:1097–107.
- 29 Ohashi R, Kawahara K, Namimatsu S, *et al*. Expression of MRP1 and ABCG2 is associated with adverse clinical outcomes of papillary thyroid carcinoma with a solid component. *Hum Pathol* 2017;67:11–17.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow