Published in final edited form as:

Oncogene. 2017 February 16; 36(7): 899–911. doi:10.1038/onc.2016.267.

PIPKI γ and talin couple phosphoinositide and adhesion signaling to control the epithelial to mesenchymal transition

N Thapa, X Tan, S Choi, T Wise, and RA Anderson

Molecular and Cellular Pharmacology Program, University of Wisconsin-Madison, School of Medicine and Public Health, Madison, WI, USA.

Abstract

Epithelial cells acquire migratory/invasive and stemness traits upon conversion to the mesenchymal phenotype. The expression of E-cadherin is a key to this transition; yet precise understanding of the pathways involved in integrating E-cadherin loss to the gain of mesenchymal traits remains poorly understood. Here, we show that phosphoinositide-generating enzyme, PIPKI γ , expression is upregulated upon epithelial–mesenchymal transition (EMT) and together with the cytoskeletal protein talin assemble into a signaling complex upon E-cadherin loss. PIPKI γ and talin together control the adhesion and phosphoinositide signaling that regulates conversion to the mesenchymal phenotypes. PIPKI γ and talin regulate the stability of E-cadherin transcriptional repressors, snail and slug, induced by transforming growth factor- β 1 or extracellular matrix protein. Loss of PIPKI γ or talin or their interaction impaired EMT and the acquisition of cell motility and stemness. This demonstrates a mechanism where a phosphoinositide-generating enzyme PIPKI γ couples with a cytoskeletal protein talin to control the acquisition of mesenchymal phenotypes.

INTRODUCTION

A prevailing theme in cancer specifies that the loss of E-cadherin-mediated adherens junctions and acquisition of migratory/invasive traits in conjunction with self-replicating stemness property determine the success of tumor metastasis.^{1–4} Epithelial–mesenchymal transition (EMT) is a normal embryonic development program often hijacked by metastasizing tumor cells, whereby tumor cells acquire different traits required for metastasis.^{3,4} However, the precise understanding of signaling molecules that couple E-cadherin loss to gain of migratory/invasive and stemness traits remains poorly understood.¹ Uncovering the role of molecules and signaling pathways that are involved is key to the development of effective therapeutic approaches in cancer treatment as the majority of carcinomas originate from epithelial cells.^{3,5} Arguably, the signaling pathways commonly

Correspondence: Professor RA Anderson, Molecular and Cellular Pharmacology Program, School of Medicine and Public Health, University of Wisconsin-Madison, 3750 Medical Science Center, 1300 University Ave., Madison, WI 53706, USA. raanders@wisc.edu.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

Supplementary Information accompanies this paper on the *Oncogene* website (<http://www.nature.com/onc>)

deregulated in cancer are responsible for orchestrating these processes, thus provoking us to interrogate the role of molecules in phosphoinositide signaling.

Phosphatidylinositol-4,5-bisphosphate (PIP₂) is a lipid messenger and a substrate for the generation of other messengers (PIP₃, DAG and IP₃), all of which regulate cell polarity and motility.^{6,7} PIP₂ is synthesized by type I phosphatidylinositol 4-phosphate kinase (PIPKI) enzymes encoded by three genes in mammalian cells, PIPKI α , PIPKI β and PIPKI γ .^{8,9} In epithelial cells, different splice variants of PIPKI γ colocalize and associate with E-cadherin at adherens junctions, and they also regulate E-cadherin trafficking and epithelial morphogenesis.^{10–12} PIPKI γ is also found over-expressed in triple-negative breast cancer,¹³ as it regulates cell migration/anchorage-independent growth of tumor cells^{14–17} and functions as a proximal regulator of PI3K/Akt signaling.¹⁸ PIPKI γ i2, a focal adhesion targeting variant of PIPKI γ , interacts with talin and regulates adhesion signaling by generating PIP₂ that modulates the assembly of adhesion complexes.^{19,20} Talin, an FERM-domain containing cytoskeletal protein, is the structural and functional unit of integrin-mediated adhesion complexes that mediate ‘inside-out’ and ‘outside-in’ signaling at cell–matrix interaction sites.²¹ Although EMT is accompanied by a profound increase in adhesive and migratory activity of the transitioning cells, roles for talin and PIPKI γ in EMT are not defined.

Here, we show that upon E-cadherin loss, PIPKI γ couples with talin to form a signaling complex that regulates the adhesion-stimulated PI3K/Akt signaling required for epithelial cells undergoing EMT. PIPKI γ /PIPKI γ i2 expression and PI3K/Akt signaling were increased in mesenchymal cells induced by transforming growth factor- β 1 (TGF β 1) treatment. The integrity of PIPKI γ and talin complex was required for the stability of E-cadherin transcriptional repressors and the gain of mesenchymal traits, highlighting the integrative role of adhesion and PI3K/Akt signaling in EMT. The assembly of PIPKI γ /PIPKI γ i2 with talin and their collaborative functions provide the signaling platform for the regulation of PI3K/Akt signaling downstream of extracellular matrix (ECM) proteins and growth factors. These are required for the stability of EMT-regulating transcription factors and the maintenance of mesenchymal phenotypes, including cell motility and stemness properties. This demonstrates that E-cadherin loss in EMT is coupled with the assembly of PIPKI γ and talin for regulation of adhesion and PI3K/Akt lipid signaling required for gain of mesenchymal phenotypes.

RESULTS

Mesenchymal cells displays increased PI3K/Akt signaling

Epithelial cells acquire properties essential for cancer progression upon transition into the mesenchymal state.³ We used the EMT model of murine mammary epithelial cells, NMuMG, that can be progressively transformed into mesenchymal state by TGF β 1 treatment or by culturing on ECM protein or E-cadherin knockdown as illustrated in this study. EMT was assessed by loss of epithelial markers and increased expression of mesenchymal marker proteins (Figure 1a) and change in cell morphology (e.g. loss of organized compact cell islands and gain of front–back polarity) (Figure 1b). The progressive changes in the morphology of NMuMG cells undergoing EMT upon TGF β 1 treatment is

demonstrated in Supplementary Figure S1. Consistent with previous studies^{3,5} epithelial cells converted into mesenchymal state showed dramatically increased adhesive and migratory activity (Figures 1c and d).

Next, the activation level of signaling molecules in epithelial vs mesenchymal cells were interrogated by an array kit, which detects the phosphorylation levels of 26 different kinases. As shown in Figure 1e, the phosphorylation level of molecules in the phosphoinositide signaling pathway (e.g. Akt1, total Akt and GSK3 β) was significantly higher in mesenchymal cells over epithelial cells. We also demonstrated the upregulated activation level of Akt during TGF β 1-induced EMT (Figure 1a). This was further confirmed by demonstrating the higher activation level of Akt in mesenchymal cells in response to fetal bovine serum (FBS) or growth factor (e.g. epidermal growth factor (EGF), hepatocyte growth factor (HGF)) stimulation of the cells (Figures 1f–h), and upon adhesion to ECM protein (Figure 1i). This indicates that EMT is accompanied by increased PI3K/Akt lipid signaling. Consistently, the inhibition of PI3K/Akt signaling (e.g. by PI3K inhibitor) impaired the transition of epithelial cells into mesenchymal cells and acquisition of cell motility (Figures 1j and k), indicating that PI3K/Akt signaling is an integral part of the EMT process.^{22,23}

PIPKI γ expression is increased in mesenchymal cells

After demonstrating the increased PI3K/Akt lipid signaling in mesenchymal cells and its role in EMT, the expression levels of type I PIPKI and PI3K enzymes during EMT were examined. As shown in Figure 2a, the expression level of PIPKI γ and its focal adhesion targeting variant, PIPKI γ 2, showed a progressive increase, whereas PIPKI α progressively decreased during TGF β 1-induced EMT. However, no obvious changes were observed in the expression level of PIPKI β . Although the expression levels of PIPKI α and PIPKI γ appeared inversely related, their precise role in the regulation of EMT remains undefined. The expression level of PI3K or PTEN also remained largely unchanged. PIPKI γ knockdown affected the phosphorylation level of the different molecules in the phosphoinositide signaling pathway (e.g. Akt1, total Akt and GSK3 β) as indicated by the array kit (Figure 2b). This was further substantiated by demonstrating impaired activation level of Akt in PIPKI γ knocked down cells (Figure 2c), and is consistent with a previous study.¹⁸ Furthermore, the expression level of ectopically expressed PIPKI γ 2 and PIPKI γ 4 splice variants was upregulated in NMuMG cells upon induction of EMT by TGF β 1 treatment (Supplementary Figure S2A), indicating post-transcriptional control of PIPKI γ expression during EMT.

In addition to TGF β 1, ECM proteins and growth factors (e.g. EGF and HGF) are common regulators of phosphoinositide signaling and EMT.^{5,24,25} NMuMG cells cultured into type I collagen (Col. I)-coated culture plates in the presence or absence of EGF or HGF demonstrated a conversion into the mesenchymal state, which was also accompanied by strongly induced PIPKI γ expression (Figures 2d and e). Upregulated expression of PIPKI γ was also observed in MDA-MB-231 and SUM159 breast cancer cells upon culturing into ECM protein-coated culture plates or treatment with TGF β 1 (Supplementary Figure S2B). As PIPKI γ and the PIP₂ lipid messenger are integral parts of multiple cellular functions,

PIPKI γ appeared important for EMT as loss of PIPKI γ diminished EMT (Figures 2f and g), consistent with a loss of Akt activation (Figure 2c).

PIPKI γ displays increased association with talin and other pro-migratory molecules during EMT

During EMT, epithelial cells lose E-cadherin-mediated cell–cell adherens junction and simultaneously gain increased interactions with ECM proteins.^{3,5} Different variants of PIPKI γ , including PIPKI γ 2, associate with E-cadherin¹⁰ and are localized at cell–cell contact sites of adherens junctions in epithelial cells (Supplementary Figure S3). PIPKI γ 2 also associate with talin, and is localized at integrin-mediated cell–matrix interaction sites, focal adhesions.¹⁹ Upon transition of these cells into mesenchymal phenotype by TGF β 1 treatment, PIPKI γ 2, but not PIPKI γ 1, becomes extensively localized at the interface of cell–matrix interaction sites along with talin (Figure 3a). Furthermore, with a decreased expression level of E-cadherin and loss of PIPKI γ 2 interaction with E-cadherin during EMT, PIPKI γ 2 showed a progressive increase in its association with various pro-migratory and pro-invasive molecules, including talin, N-cadherin, Src and IQGAP1 (Figure 3b).

Among the PIPKI γ variants, PIPKI γ 2 showed the strong association with talin, N-cadherin, moesin and Src upon EMT (Figure 3c). This was further demonstrated in NMuMG cells transitioning into mesenchymal cells by culturing into the ECM proteins (Figure 3d). E-cadherin, N-cadherin, talin, IQGAP1 and Src are previously identified interacting partners of PIPKI γ /PIPKI γ 2.^{10,14,17} Moesin, an FERM-domain containing protein is involved in cell polarity.²⁶ Moesin was identified by mass spectrometry in the immuno-complex of PIPKI γ 2 isolated from NMuMG cells induced to mesenchymal state by TGF β 1 treatment and associates with PIPKI γ 2 via talin (Supplementary Figures S4A and B). However, unlike pan-PIPKI γ , the knockdown of PIPKI γ 2 alone was not sufficient to block EMT induced by TGF β 1 treatment, indicating the collective role of PIPKI γ variants in the EMT process (Figure 3e). Alternatively, a less efficient knockdown of PIPKI γ 2 might have also contributed to the lack of effect on TGF β 1-induced EMT.

Remarkably, the knockdown of talin, a key cytoskeletal protein at integrin-mediated adhesion complex, diminished EMT induced by TGF β 1 (Figure 3e) and more significantly by ECM proteins (see below). In NMuMG cells fully converted into mesenchymal state by TGF β 1 treatment, the knockdown of PIPKI γ or PIPKI γ 2 or its associating partners (such as talin and Src) significantly impaired cell migration, one of the key phenotypes of mesenchymal cells (Figures 3f and g). Consistently, all of these molecules are also known to play essential role in cell migration of different tumor cells.^{14–16} These results indicate that upon the loss of PIPKI γ interaction with E-cadherin, PIPKI γ shows increased association with talin and other pro-migratory molecules to regulate EMT and gain of cell motility.

E-cadherin loss promotes PIPKI γ association with talin and PI3K/Akt signaling

E-cadherin and talin, cornerstones of cell–cell adherens junctions and integrin-mediated adhesion complexes, respectively, are the direct interacting partners of PIPKI γ /PIPKI γ 2, which appear to shuttle between these two molecules in epithelial and mesenchymal cells as indicated in Figure 3a. Consistently, talin knockdown promoted PIPKI γ 2 association with

E-cadherin (Figure 4a). Conversely, E-cadherin knockdown promoted PIPKI γ 2 association with talin (Figure 4b), and PIPKI γ 2 was extensively reorganized to the leading edge adhesion complexes along with talin in E-cadherin knocked down cells (Figure 4c). These results indicate that E-cadherin loss promotes the assembly and reorganization of PIPKI γ /PIPKI γ 2 with talin in transitioning epithelial cells.

The loss of E-cadherin is a hallmark of EMT^{5,27} and the knockdown of E-cadherin was sufficient to convert NMuMG cells from epithelial into mesenchymal state, which was further substantiated by culturing into Col. I-coated culture plates (Figure 4d). However, knockdown of talin and PIPKI γ /PIPKI γ 2 abrogated the cell motility gained by E-cadherin knockdown cells (Figure 4e). Furthermore, E-cadherin knockdown cells acquired increased PI3K/Akt signaling, which was severely impaired upon PIPKI γ and talin knockdown (Supplementary Figures S5A–D), suggesting that the PIPKI γ and talin assembly provides a platform for the regulation of PI3K/Akt signaling in mesenchymal cells. As expected, the pharmacological inhibition of PI3K blocked cell motility and the expression of mesenchymal marker proteins in the E-cadherin knockdown cells (Figures 4f and g). Together, these results indicate that mesenchymal traits acquired by epithelial cells upon E-cadherin loss depend upon the integrity of the talin and PIPKI γ /PIPKI γ 2 complex assembled in the vicinity of the cell–matrix interface, and their signaling.

PIPKI γ and talin interaction is required for EMT

During EMT, epithelial cells lose E-cadherin-mediated cell–cell adherens junctions and simultaneously gain the increased adhesive interaction with ECM proteins through the integrin-mediated adhesion complex.^{2,5} Talin, a cytoskeletal protein in the cytosolic compartment of the integrin-mediated adhesion complex, provides the platform for bidirectional signaling (‘outside-in’ and ‘inside-out’).^{21,28} The interaction of cells with ECM proteins and derived cellular signals appears pivotal for acquisition of mesenchymal traits as knockdown of talin abrogated the transition of epithelial cells into mesenchymal state induced by ECM proteins (Figures 5a and b) or E-cadherin knockdown (Figures 5c and d). Talin knockdown cells appeared similar to epithelial NMuMG control cells and the expression level of E-cadherin and N-cadherin was similar to that of epithelial control cells, although the expression level of FN was higher. Similarly, EMT induced by E-cadherin knockdown was also effectively blocked by the knockdown of talin as indicated by lack of induced N-cadherin expression and impaired cell motility (Figure 4e), although, the expression level of FN was subtly affected. In contrast, moesin knockdown promoted EMT, which was accompanied by increased talin association with integrin (Figure 5e), suggesting that moesin controls EMT by regulating talin assembly with integrins at adhesion complexes, which is in agreement with reported study.²⁹ Furthermore, the knockdown of talin, but not moesin, promoted reversion of NMuMG cells from mesenchymal (after TGF β 1 treatment) into the epithelial state (Figures 5f and g). These results indicate that talin, a cytoskeletal protein at integrin-mediated adhesion complex, plays crucial role in the initiation of EMT as well as in the maintenance of mesenchymal phenotypes.

Next, we tried to define the functional role of talin interaction with PIPKI γ in EMT and the acquisition of mesenchymal traits as talin association with PIPKI γ 2 is significantly

increased during EMT. Furthermore, the lipid messenger PIP₂ regulates the recruitment and activation of talin at the adhesion complex required for bidirectional signaling.³⁰ The PIP₂ is putatively provided by PIPKIγi2 associated with talin^{16,19,21} (Figure 5h). To explore this, GFP-fusion proteins of the F2 or F3 subdomains of talin FERM domain were constructed and expressed into NMuMG cells by lentiviral infection. The F3 subdomain harbors the binding sites for PIP₂, PIPKIγi2 and integrins.^{20,21} As expected, the expression of talin F3 subdomain abrogated the co-immunoprecipitation between talin and PIPKIγi2 (Supplementary Figures S6A and B). Furthermore, when NMuMG cells expressing F2 or F3 subdomains of talin FERM domain were induced to undergo EMT by E-cadherin knockdown, the expression of talin F3 subdomain completely blocked talin association with PIPKIγi2, this also impaired adhesion (as indicated by phosphorylation level of FAK) and PI3K/Akt lipid signaling (Figure 5i). These cells showed defects in acquiring mesenchymal traits as indicated by impaired N-cadherin expression and cell motility (Figure 5j). The expression of talin F3-subdomain also impaired adhesion and PI3K/Akt lipid signaling as well as cell motility of highly aggressive MDA-MB-231 tumor cells (Supplementary Figures S6C and D). These results indicate that talin interaction with PIPKIγ is required for gain of mesenchymal properties in transitioning epithelial cells.

PIPKIγ and talin regulate the stability of snail and slug

Different agonists inducing EMT, such as TGFβ1, ECM proteins, growth factors, WNT and NOTCH, converge to inducing E-cadherin transcriptional repressors (e.g. snail, slug and twist) to orchestrate EMT.^{5,31,32} EMT induced by ECM or TGFβ1 in NMuMG cells is associated with increased expression levels of snail and slug (Figure 6a). Consistently, knockdown of snail and/or slug severely impaired the induction of EMT by TGFβ1 treatment (not shown). These transcriptional repressors are short-lived proteins requiring cellular signaling emanating from plasma membrane for their stability³¹ and PI3K/Akt signaling regulates the stability of these transcriptional repressors by inhibiting the GSK3β, which promotes ubiquitination and proteasomal degradation of these transcriptional repressors.^{31,33} This suggests that PIPKIγ and talin might regulate the stability of these E-cadherin transcriptional repressors during EMT. In alignment with this notion, the expression level and half-life of snail and slug, but not β-catenin, were dramatically reduced upon PIPKIγ or talin knockdown in NMuMG cells transformed into mesenchymal state by TGFβ1 treatment (Figure 6b). However, the treatment of cells with proteasomal inhibitor but not lysosomal inhibitors (not shown) completely replenished the expression level of snail and slug in talin or PIPKIγ knockdown cells (Figures 6c–e), indicating that PIPKIγ and talin are required for the stability of snail and slug by preventing proteasomal degradation of these transcription factors. This fully reconciled with increased ubiquitinated snail upon knockdown of PIPKIγ or talin in NMuMG cells overexpressing snail and also in cancer cells (Figure 6f). Consistently, the expression level of mesenchymal markers and mesenchymal attributes (e.g. migratory) acquired by ectopic expression of snail in NMuMG cells was significantly abrogated upon knockdown of talin or PIPKIγ (Figures 6g–i). A decreased expression level of ectopically expressed snail in talin or PIPKIγ knockdown cells concurs with increased E-cadherin expression and decreased mesenchymal phenotypes. All of these results indicate that PIPKIγ and talin provide a critical signaling nexus for the stability of snail and slug during EMT.

PIPKI γ regulates the stemness property of mesenchymal and tumor cells

Stemness trait is one of the key properties acquired by epithelial cells upon conversion into mesenchymal cells.^{34,35} The ability to form mammospheres/tumorspheres in suspension culture is the commonly used assay to measure stemness property.³⁴ Induction of EMT by TGF β 1 treatment or E-cadherin knockdown or snail overexpression (not shown) substantially increased the ability of NMuMG cells to form mammospheres/tumorspheres, which was significantly compromised upon talin or PIPKI γ knockdown (Figures 7a and b). The overexpression of PIPKI γ further substantiated the ability of E-cadherin knockdown cells to form mammospheres/tumorspheres (Figures 7c and d). However, kinase dead PIPKI γ 2-expressing cells or knockdown of talin in PIPKI γ 2-overexpressing cells all formed significantly fewer mammospheres/tumorspheres upon E-cadherin knockdown (Supplementary Figure S7A). Together, these results indicate the key role of PIPKI γ and talin complex in acquisition of self-replicating stemness property upon EMT.

To further explore PIPKI γ functions, mammospheres/tumor-spheres formed by different tumor cells were isolated. The expression level of PIPKI γ /PIPKI γ 2 and level of activated Akt were higher in mammospheres/tumorspheres formed by different tumor cells (Figure 7e). The expression of pluripotent stem cell marker genes was highly upregulated in isolated mammospheres/tumorspheres as indicated by the pluripotent stem cell array kit (Supplementary Figure S7B). In a complementary assay, anoikis-resistant tumor cells isolated at different time points after culturing in non-adherent condition also showed the increased expression of PIPKI γ that maintained a higher activation level of Akt and mammospheres/tumorspheres forming capacity (Supplementary Figures S7C and D). These results indicate that PIPKI γ and PI3K/Akt signaling is increased in the subpopulation of tumor cells enriched in stemness property. Corroborating this, the loss of PIPKI γ or PIPKI γ 2 or talin (not shown) expression in different tumor cells (e.g. Cal51, A431 and T47D) significantly compromised their ability to form mammospheres/tumorspheres (Figure 7f). With these results, we conclude that PIPKI γ /PIPKI γ 2 and the cytoskeletal protein talin provide an important signaling nexus for EMT and acquisition of mesenchymal phenotypes as summarized in the schematic diagram (Figure 8).

DISCUSSION

The present study indicates that coupling of adhesion and PI3K/Akt signaling by talin and PIPKI γ provides the integral signaling platform for acquisition of mesenchymal traits by transitioning epithelial cells. The conversion of epithelial into mesenchymal phenotype is accompanied by profound changes in adhesion and PI3K/Akt signaling that are initiated and sustained by ECM proteins and growth factors known to constitute key components of tumor microenvironment.^{2,4,5} Furthermore, concomitant with E-cadherin loss, talin and PIPKI γ gained increased expression/interaction and reorganization from cell–cell contact sites to adhesion complexes. This provides a critical signaling nexus for adhesion and PI3K/Akt signaling in transitioning epithelial cells.

The loss of talin or PIPKI γ expression or abrogation of their molecular interaction by overexpression of talin F3 fragment profoundly affected PI3K/Akt and adhesion signaling in transitioning epithelial cells, and these cells retained epithelial-like morphology even after

E-cadherin knockdown or culturing on the ECM protein. These cells also failed to acquire full-fledged mesenchymal attributes, including cell motility and stemness properties. This shows that E-cadherin loss in EMT is coupled with a gain of increased PI3K/Akt and adhesion signaling regulated by PIPKI γ and, the cytoskeletal protein, talin that are required for mesenchymal attributes. This is fully consistent with the prevailing notion that spatial generation of PIP₂ lipid messenger by PIPKI γ regulates the development of integrin-mediated adhesion complex at cell–matrix interface, where talin assumes key roles in executing inside-out and outside-in bidirectional signaling. However, the precise mechanism and dynamics of molecular interaction among talin, PIPKI γ 2 and integrins at the adhesion complex and their regulation by PIP₂ lipid messenger still remain poorly understood.²¹ In alignment with increased adhesive/invasive property of transitioning epithelial cells, the predominant changes observed in EMT are the profound increase in the synthesis of ECM proteins and integrin expressions, which probably occurs irrespective of agonists used for inducing EMT (e.g. TGF β 1, oncogenes or overexpression of EMT-inducing transcription factors).^{27,36–38} Again, these events also result in initiation and sustenance of more profound PI3K/Akt lipid signaling downstream of integrins/growth factors in transitioning epithelial cells indicating that all of these are interconnected events of the EMT program, and the integrity of PIPKI γ and talin is essential for acquisition as well as sustenance of mesenchymal attributes in transitioning epithelial cells.

In mesenchymal cells, talin, PIPKI γ and their interaction appear to have pleiotropic roles that range from the initiation and sustenance of adhesion and PI3K/Akt signaling to regulation of cell motility and the stability of EMT-regulating transcription factors. In different physio-pathological conditions, upregulated expression of the transcription factors snail and slug block E-cadherin expression promoting EMT. Different agonists that induce EMT, such as TGF β 1, ECM protein, growth factors, WNT and NOTCH, all converge into inducing these E-cadherin transcriptional repressors.^{5,31} In the absence of talin or PIPKI γ , the expression of EMT-regulating transcription factors snail and slug was compromised. The integrity of talin and PIPKI γ , and their cellular signaling, appears to play a pivotal role in the stability of snail and slug by preventing their proteosomal degradation, possibly by inhibiting the GSK3 β phosphorylation and ubiquitination, as number of studies indicate crucial roles of cellular signals, such as PI3K/Akt and Erk1/2, in the stability of these EMT-regulating transcription factors by inhibiting GSK3 β .^{33,39}

Along with increased motility, another key facet of EMT and many cancer cells is increased stemness property, which dictates the ability of cancer cells to metastasize and colonize to distant places.^{34,35,40} PIPKI γ expression and PI3K/Akt signaling are selectively enriched in anoikis-resistant cell populations with stemness traits. The loss of PIPKI γ or disruption of the PIPKI γ and talin interaction and their signaling nexus affected the gain of stemness traits by transitioning epithelial cells, and also impacted the stemness property of many tumor cells. In summary, we illustrate the comprehensive mechanism how coupling of phosphoinositide kinase PIPKI γ and cytoskeletal protein talin provides a platform for the acquisition of mesenchymal traits by transitioning epithelial cells.

MATERIALS AND METHODS

Cell culture

Mouse mammary epithelial cells, NMuMG were cultured in Dulbecco's modified Eagle's medium-containing 10% FBS supplemented with 10 µg/ml insulin (Sigma, St Louis, MO, USA). MDA-MB-231, Cal51, A431, MCF-7, HCC1954 and HEK293FT cells were cultured into Dulbecco's modified Eagle's medium-containing 10% FBS. T47D cells were cultured in RMPI-1640 containing 10% FBS. SUM159 cells were culture in Ham's F12 supplemented with 5% FCS. All cells were grown at 37 °C in a 5% CO₂ incubator.

For the induction of EMT, NMuMG cells were treated with TGFβ1 (2 ng/ml), and harvested at different time points. The mesenchymal cells indicated in this study refer to NMuMG cells treated with TGFβ1 for 3–5 days. Similarly, for studying the effect of PI3K inhibitor, NMuMG cells were treated with LY294002 (2 µg/ml) during induction of EMT by TGFβ1 treatment. For induction of EMT by ECM protein, cells were seeded into Col. I-coated culture plate for 2–3 days in the presence or absence of EGF or HGF.

For examination of activated phosphoinositide signaling (PI3K/Akt) by FBS or different growth factors, cells were serum- and insulin-starved overnight before stimulating with FBS or indicated growth factor and then harvested at different time points. Similarly, cells, after resuspension in serum-free medium, were seeded into Col. I-coated culture plate and incubated for 15–20 min. Non-attached cells were removed by phosphate-buffered saline washing before cell harvesting.

For examining the stability of EMT-regulating transcription factors, NMuMG cells fully converted into mesenchymal state by TGFβ1 treatment were treated with cyclohexamide (50 µg/ml) before harvesting them at different time periods. Similarly, for inhibition of proteosomal degradation of EMT-regulating transcription factors, cells were treated with MG132 (50 µg/ml) before harvesting them at different time points.

Transfection or lentiviral infection

For transfection of siRNA, LipofectamineRNAiMAX (Invitrogen, Carlsbad, CA, USA) was used following the protocol provided by the manufacturer, and cells were assayed 48–72 h post-transfection. For transient transfection into HEK293 cells, Lipofectamine-2000 (Invitrogen) was used. Cells were harvested 24 h post-transfection. For the expression of PIPKIγ variants, snail or talin fragments into NMuMG or other cells, the lentiviral system was used as described previously.¹⁶ Cells were harvested 48–72 h post-infection and 70–80% of infection efficiency was achieved for the experiments.

Immunoprecipitation and immunoblotting

Cells were lysed using lysis buffer (50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.5% Triton-X100, 1 mM EDTA, 10 mM NaF, 5 mM Na₃VO₄ and protease inhibitors). Clear supernatants were incubated with indicated antibodies for 3–4 h to overnight at 4 °C followed by isolation of immunocomplexes using protein G Sepharose 4B beads (Amersham, Uppsala, Sweden). Beads were washed three times with lysis buffer before

eluting the immuno-complexes with $2 \times$ sample buffer, run through sodium dodecyl sulfate–polyacrylamide gel electrophoresis gel and subjected to immunoblotting using specific antibodies.

Mammospheres/tumorspheres formation

For mammospheres/tumorspheres formation, different cell types at the density of 5×10^3 cells/ml were suspended into the complete mammoCults medium (StemCell Tech, Vancouver, BC, Canada) containing methylcellulose were seeded into 24-well ultra-low attachment plates (Corning, Kennebun, ME, USA). The mammospheres/tumorspheres formed were isolated or counted after 3–5 days of incubation. For examination of protein expression by the pluripotent stem cell array kit, the clear cell lysates (200 μ g) prepared from isolated mammospheres/tumorspheres were used. For isolation of anoikis-resistant tumor cell subpopulation, single cells suspended at the density of 1×10^5 cells/ml in Dulbecco's modified Eagle's medium medium-containing 0.5% FBS were seeded into 10 cm dish culture plates precoated with 0.5% agar for preventing the cell attachment. Cells surviving under this condition were isolated at different time points and harvested.

Cell adhesion and migration

Cell adhesion and migration assays were carried out as described previously.^{7,16} In brief, cells were serum- and insulin-starved overnight before resuspension in Dulbecco's modified Eagle's medium medium containing 0.2% bovine serum albumin. For cell adhesion, cells were seeded into Col. I-coated culture plate and incubated for 15–30 min before removing the non-adherent cells by phosphate-buffered saline washing. Adherent cells were fixed with paraformaldehyde and cells counted from different areas of the plate. For cell migration, cells were allowed to migrate towards 0.25% FBS as chemoattractant. Time allowed for cell migration was 5–6 h. The cell migrated to the underside of membrane were fixed, stained and counted.

Immunofluorescence microscopy

For immunofluorescence study, colonies developed in the soft agar were fixed with 3.7% paraformaldehyde followed by cell permeabilization with 0.1% Triton-X and blocked with 3% bovine serum albumin in phosphate-buffered saline. Cells were incubated with a primary antibody overnight at 4 °C followed by incubation with Alexa555- and/or Alexa488-conjugated secondary antibodies (Molecular Probes, Eugene, OR, USA) for 1 h at room temperature. Slides were mounted using Vectashield and visualized with a Nikon TE2000-U microscope, Japan. Images were acquired using MetaMorph and processed using Adobe Photoshop.

Statistical analysis

The data are presented mean $\pm$ s.d. from at least three-independent experiments. Unpaired t-test was conducted to determine the *P*-value and the statistical significance between two groups (*P*-value less than 0.05 were considered significant).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Health (NIH) grants (CA104708 and GM057549) to RAA; American Heart Association (AHA) grants to NT (10POST4290052) and SC (13PRE14690057); and Howard Hughes Medical Institute (HHMI) International Student Research Fellowship to XT.

REFERENCES

1. Davis FM, Stewart TA, Thompson EW, Monteith GR. Targeting EMT in cancer: opportunities for pharmacological intervention. *Trends Pharmacol Sci* 2014; 35: 479–488. [PubMed: 25042456]
2. Jung HY, Fattet L, Yang J. Molecular pathways: linking tumor microenvironment to epithelial-mesenchymal transition in metastasis. *Clin Cancer Res* 2015; 21: 962–968. [PubMed: 25107915]
3. Polyak K, Weinberg RA. Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat Rev Cancer* 2009; 9: 265–273. [PubMed: 19262571]
4. Valastyan S, Weinberg RA. Tumor metastasis: molecular insights and evolving paradigms. *Cell* 2011; 147: 275–292. [PubMed: 22000009]
5. Thiery JP, Sleeman JP. Complex networks orchestrate epithelial-mesenchymal transitions. *Nat Rev Mol Cell Biol* 2006; 7: 131–142. [PubMed: 16493418]
6. Heck JN, Mellman DL, Ling K, Sun Y, Wagoner MP, Schill NJ et al. A conspicuous connection: structure defines function for the phosphatidylinositol-phosphate kinase family. *Crit Rev Biochem Mol Biol* 2007; 42: 15–39. [PubMed: 17364683]
7. Thapa N, Anderson RA. PIP2 signaling, an integrator of cell polarity and vesicle trafficking in directionally migrating cells. *Cell Adh Migr* 2012; 6: 409–412. [PubMed: 23076053]
8. Choi S, Thapa N, Tan X, Hedman AC, Anderson RA. PIP kinases define PI4,5P signaling specificity by association with effectors. *Biochim Biophys Acta* 2015; 1851: 711–723. [PubMed: 25617736]
9. Sun Y, Thapa N, Hedman AC, Anderson RA. Phosphatidylinositol 4,5-bisphosphate: targeted production and signaling. *Bioessays* 2013; 35: 513–522. [PubMed: 23575577]
10. Ling K, Bairstow SF, Carbonara C, Turbin DA, Huntsman DG, Anderson RA. Type I gamma phosphatidylinositol phosphate kinase modulates adherens junction and E-cadherin trafficking via a direct interaction with mu 1B adaptin. *J Cell Biol* 2007; 176: 343–353. [PubMed: 17261850]
11. Schill NJ, Hedman AC, Choi S, Anderson RA. Isoform 5 of PIPKIgamma regulates the endosomal trafficking and degradation of E-cadherin. *J Cell Sci* 2014; 127: 2189–2203. [PubMed: 24610942]
12. Xiong X, Xu Q, Huang Y, Singh RD, Anderson R, Leof E et al. An association between type Igamma PI4P 5-kinase and Exo70 directs E-cadherin clustering and epithelial polarization. *Mol Biol Cell* 2012; 23: 87–98. [PubMed: 22049025]
13. Sun Y, Turbin DA, Ling K, Thapa N, Leung S, Huntsman DG et al. Type I gamma phosphatidylinositol phosphate kinase modulates invasion and proliferation and its expression correlates with poor prognosis in breast cancer. *Breast Cancer Res* 2010; 12: R6. [PubMed: 20074374]
14. Choi S, Thapa N, Hedman AC, Li Z, Sacks DB, Anderson RA. IQGAP1 is a novel phosphatidylinositol 4,5 bisphosphate effector in regulation of directional cell migration. *EMBO J* 2013; 32: 2617–2630. [PubMed: 23982733]
15. Sun Y, Ling K, Wagoner MP, Anderson RA. Type I gamma phosphatidylinositol phosphate kinase is required for EGF-stimulated directional cell migration. *J Cell Biol* 2007; 178: 297–308. [PubMed: 17635937]
16. Thapa N, Sun Y, Schramm M, Choi S, Ling K, Anderson RA. Phosphoinositide signaling regulates the exocyst complex and polarized integrin trafficking in directionally migrating cells. *Dev Cell* 2012; 22: 116–130. [PubMed: 22264730]

17. Thapa N, Choi S, Hedman A, Tan X, Anderson RA. Phosphatidylinositol phosphate 5-kinase Igamma2 in association with Src controls anchorage-independent growth of tumor cells. *J Biol Chem* 2013; 288: 34707–34718. [PubMed: 24151076]
18. Thapa N, Choi S, Tan X, Wise T, Anderson RA. Phosphatidylinositol phosphate 5-kinase Igamma and PI3K/Akt signaling couple to promote oncogenic growth. *J Biol Chem* 2015; 290: 18843–18854. [PubMed: 26070568]
19. Ling K, Doughman RL, Firestone AJ, Bunce MW, Anderson RA. Type I gamma phosphatidylinositol phosphate kinase targets and regulates focal adhesions. *Nature* 2002; 420: 89–93. [PubMed: 12422220]
20. Ling K, Doughman RL, Iyer VV, Firestone AJ, Bairstow SF, Mosher DF et al. Tyrosine phosphorylation of type Igamma phosphatidylinositol phosphate kinase by Src regulates an integrin-talin switch. *J Cell Biol* 2003; 163: 1339–1349. [PubMed: 14691141]
21. Calderwood DA, Campbell ID, Critchley DR. Talins and kindlins: partners in integrin-mediated adhesion. *Nat Rev Mol Cell Biol* 2013; 14: 503–517. [PubMed: 23860236]
22. Bakin AV, Tomlinson AK, Bhowmick NA, Moses HL, Arteaga CL. Phosphatidylinositol 3-kinase function is required for transforming growth factor beta-mediated epithelial to mesenchymal transition and cell migration. *J Biol Chem* 2000; 275: 36803–36810. [PubMed: 10969078]
23. Derynck R, Muthusamy BP, Saeteurn KY. Signaling pathway cooperation in TGF-beta-induced epithelial-mesenchymal transition. *Curr Opin Cell Biol* 2014; 31: 56–66. [PubMed: 25240174]
24. Park J, Schwarzbauer JE. Mammary epithelial cell interactions with fibronectin stimulate epithelial-mesenchymal transition. *Oncogene* 2014; 33: 1649–1657. [PubMed: 23624917]
25. Shintani Y, Fukumoto Y, Chaika N, Svoboda R, Wheelock MJ, Johnson KR. Collagen I-mediated up-regulation of N-cadherin requires cooperative signals from integrins and discoidin domain receptor 1. *J Cell Biol* 2008; 180: 1277–1289. [PubMed: 18362184]
26. Ben-Aissa K, Patino-Lopez G, Belkina NV, Maniti O, Rosales T, Hao JJ et al. Activation of moesin, a protein that links actin cytoskeleton to the plasma membrane, occurs by phosphatidylinositol 4,5-bisphosphate (PIP2) binding sequentially to two sites and releasing an autoinhibitory linker. *J Biol Chem* 2012; 287: 16311–16323. [PubMed: 22433855]
27. Onder TT, Gupta PB, Mani SA, Yang J, Lander ES, Weinberg RA. Loss of E-cadherin promotes metastasis via multiple downstream transcriptional pathways. *Cancer Res* 2008; 68: 3645–3654. [PubMed: 18483246]
28. Nieves B, Jones CW, Ward R, Ohta Y, Reverte CG, LaFlamme SE. The NPIY motif in the integrin beta1 tail dictates the requirement for talin-1 in outside-in signaling. *J Cell Sci* 2010; 123: 1216–1226. [PubMed: 20332112]
29. Vitorino P, Yeung S, Crow A, Bakke J, Smyczek T, West K et al. MAP4K4 regulates integrin-FERM binding to control endothelial cell motility. *Nature* 2015; 519: 425–430. [PubMed: 25799996]
30. Saltel F, Mortier E, Hytonen VP, Jacquier MC, Zimmermann P, Vogel V et al. New PI (4,5)P2- and membrane proximal integrin-binding motifs in the talin head control beta3-integrin clustering. *J Cell Biol* 2009; 187: 715–731. [PubMed: 19948488]
31. Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol* 2014; 15: 178–196. [PubMed: 24556840]
32. Medici D, Hay ED, Olsen BR. Snail and Slug promote epithelial-mesenchymal transition through beta-catenin-T-cell factor-4-dependent expression of transforming growth factor-beta3. *Mol Biol Cell* 2008; 19: 4875–4887. [PubMed: 18799618]
33. Kao SH, Wang WL, Chen CY, Chang YL, Wu YY, Wang YT et al. GSK3beta controls epithelial-mesenchymal transition and tumor metastasis by CHIP-mediated degradation of Slug. *Oncogene* 2014; 33: 3172–3182. [PubMed: 23851495]
34. Brabletz T To differentiate or not—routes towards metastasis. *Nat Rev Cancer* 2012; 12: 425–436. [PubMed: 22576165]
35. Puisieux A, Brabletz T, Caramel J. Oncogenic roles of EMT-inducing transcription factors. *Nat Cell Biol* 2014; 16: 488–494. [PubMed: 24875735]
36. Chen YS, Mathias RA, Mathivanan S, Kapp EA, Moritz RL, Zhu HJ et al. Proteomics profiling of Madin-Darby canine kidney plasma membranes reveals Wnt-5a involvement during oncogenic H-

Ras/TGF-beta-mediated epithelial-mesenchymal transition. *Mol Cell Proteomics* 2011; 10: M110001131.

37. Granato DC, Zanetti MR, Kawahara R, Yokoo S, Domingues RR, Aragao AZ et al. Integrated proteomics identified up-regulated focal adhesion-mediated proteins in human squamous cell carcinoma in an orthotopic murine model. *PLoS One* 2014; 9: e98208. [PubMed: 24858105]
38. Seguin L, Desgrosellier JS, Weis SM, Cheresh DA. Integrins and cancer: regulators of cancer stemness, metastasis, and drug resistance. *Trends Cell Biol* 2015; 25: 234–240. [PubMed: 25572304]
39. Zhou BP, Deng J, Xia W, Xu J, Li YM, Gunduz M et al. Dual regulation of Snail by GSK-3beta-mediated phosphorylation in control of epithelial-mesenchymal transition. *Nat Cell Biol* 2004; 6: 931–940. [PubMed: 15448698]
40. Malanchi I, Santamaria-Martinez A, Susanto E, Peng H, Lehr HA, Delaloye JF et al. Interactions between cancer stem cells and their niche govern metastatic colonization. *Nature* 2011; 481: 85–89. [PubMed: 22158103]

Figure 1.

EMT is associated with increased PI3K/Akt signaling. **(a, b)** NMuMG cells cultured into complete growth medium were treated with TGFβ1 (2 ng/ml) before harvesting the cells at different time points. For culturing the cells for more than 3–4 days, cells were subcultured into new culture plates and the TGFβ1 concentration reduced to half (1 ng/ml). Transition to mesenchymal state was examined by downregulation of E-cadherin expression and gain of mesenchymal marker proteins by immunoblotting **(a)** and immunofluorescence study/change in cell morphology **(b)** (images taken using $\times 10$ and $\times 40$ objective lenses). **(c, d)** NMuMG cells in epithelial state, NMuMG **(e)** or after conversion into mesenchymal state by TGFβ1 treatment for 3–4 days, NMuMG **(M)** were cultured in serum- and insulin-free medium overnight before adhesion and migration assays as described in ‘Materials and methods’. For cell adhesion, attached cells to Col. I-coated culture plates after 15-min of incubation were counted. For cell migration, migrated cells towards underside of membrane after 5–6 h of incubation were stained and counted (results are represented as mean $\pm$ s.d. from three independent experiments and error bars represents s.d.). **(e)** The activation of

cellular signaling in NMuMG (E) or NMuMG (M) were analyzed by the MAPK array kit following the protocol provided by the manufacturer (1, Akt1; 2, Akt2; 3, total Akt; 4, GSK3 β). (f–i) Examination of Akt activation in response to FBS or growth factors (e.g. EGF, HGF) stimulation of the cells, and upon adhesion to ECM protein. NMuMG (E) or NMuMG (M) cells were cultured in serum- and insulin-free medium overnight before cell stimulation with indicated factors. Cells were harvested at different time points for examining the activation level of Akt by immunoblotting. (j, k) NMuMG cells were treated with PI3K inhibitor (LY294002) along with TGF β 1 (2 ng/ml) and cultured for 2-days. The effect of PI3K/Akt inhibition on EMT was examined by gain of mesenchymal marker proteins and cell motility. Results are represented as mean $\pm$ s.d. from three independent experiments and error bars represents s.d.

Figure 2.

PIPKIγ expression is increased during EMT. **(a)** The expression levels of type I PIPKI (PIPKIα, PIPKIβ and PIPKIγ) and PI3K enzymes in NMuMG cells during TGFβ1-induced EMT were examined. Cells were harvested at different time points following TGFβ1 treatment and immunoblotted with specific antibodies. **(b)** siRNA was used to knockdown PIPKIγ from the NMuMG cells transformed into mesenchymal state by TGFβ1 treatment. Then, the effect of PIPKIγ knockdown on cellular signaling was examined by the MAPK array kit as described above (1, Akt1; 2, Akt2; 3, total Akt; 4, GSK3β). **(c)** NMuMG cells converted into mesenchymal cells by TGFβ1 treatment were transfected with siRNA for PIPKIγ knockdown. At 48–72 h after transfection, cells were cultured in serum- and insulin-free medium overnight before stimulating with 10% FBS. The cells were harvested at different time points and the activation level of Akt examined by immunoblotting using phosphor-Akt antibody. **(d, e)** The examination of PIPKIγ expression in NMuMG cells

converted into mesenchymal cells by ECM protein. Cells were cultured into type I collagen (Col. I)-coated culture plates in the presence or absence of EGF or HGF for 2–3 days before harvesting for the expression of mesenchymal marker proteins by immunoblotting or examination by immunofluorescence study (image taken at $\times 40$; bright field at $\times 10$). **(f, g)**, the effect of PIPKI γ knockdown on TGF β 1-induced EMT and gain of cell motility. At 24–36 h post-transfection with siRNA for the knockdown of PIPKI γ , NMuMG cells were treated with TGF β 1. Induction of EMT was assessed 36–48 h later either by immunoblotting for the expression of mesenchymal marker proteins or by immunofluorescence study (image taken at $\times 40$).

Figure 3.

PIP KIγ displays increased association with talin and other pro-migratory molecules during EMT. (a) Localization of HA-tagged and ectopically expressed PIP KIγ1 and PIP KIγ2 in epithelial and mesenchymal cells were examined by immunofluorescence study. Upon induction of EMT, PIP KIγ2 showed extensive localization at cell–matrix interaction sites along with talin. (b) Endogenous PIP KIγ2 was immunoprecipitated from NMuMG cells at different time points following TGFβ1 treatment and associated molecules (e.g. talin, N-cadherin, moesin, IQGAP1 and Src) examined by immunoblotting. (c) NMuMG cells expressing different variants of HA-tagged PIP KIγ were transformed into mesenchymal state by TGFβ1 treatment. PIP KIγ was immunoprecipitated using anti-HA antibody and associated molecules were examined by immunoblotting. (d) NMuMG cells expressing different variants of HA-tagged PIP KIγ were transformed into mesenchymal cells by

culturing into Col. I-coated culture plates. PIPKI γ was immunoprecipitated using anti-HA antibody and associated molecules were examined by immunoblotting. The increased association of PIPKI γ 2 with talin, N-cadherin, moesin and Src was observed. (e), NMuMG cells were treated with TGF β 1 for 36–48 h after siRNA transfection for knockdown of PIPKI γ 2 or talin. Cells were harvested 48 h after TGF β 1 treatment. The conversion of NMuMG cells into mesenchymal state was assessed by examining the expression level of mesenchymal marker proteins. The knockdown of PIPKI γ 2 unlike talin was not sufficient to impair EMT induced by TGF β 1 treatment. (f, g), PIPKI γ or PIPKI γ 2 or their associating partners (e.g. talin and Src) were knocked down using siRNA in NMuMG cells fully converted into mesenchymal state by TGF β 1 treatment. Cells were cultured in serum- and insulin-free medium overnight before cell migration assay. Results are represented as mean $\pm$ s.d. from three-independent experiments and error bars represent s.d.

Figure 4.

E-cadherin loss promotes PIPKI γ association with talin and PI3K/Akt signaling. **(a)** PIPKI γ 2 was immunoprecipitated from control or talin knockdown cells, and co-immunoprecipitation of E-cadherin with PIPKI γ 2 was examined by immunoblotting. **(b, c)** PIPKI γ 2 was immunoprecipitated from NMuMG cells after E-cadherin knockdown and co-immunoprecipitation of talin with PIPKI γ 2 was examined. The localization of PIPKI γ 2 and talin in E-cadherin knockdown cells was examined by immunofluorescence study. E-cadherin knockdown induced extensive reorganization of PIPKI γ 2 to leading edge adhesion complexes along with talin. **(d)** siRNA was used to knockdown E-cadherin expression in NMuMG cells. Cells were harvested 48–72 h post-transfection to examine the expression level of mesenchymal marker proteins (fibronectin and N-cadherin). The cell motility was examined as described above. The knockdown of E-cadherin was sufficient to promote EMT of NMuMG cells and gain of cell motility. The culture into Col.I-coated culture plates

further substantiated the EMT process. (e) NMuMG cells were co-transfected with siRNA for knockdown of E-cadherin along with talin or PIPKI γ or PIPKI γ i2. At 48–72 h after transfection, cell motility was assessed as described above. However, cell motility gained by E-cadherin knockdown was abrogated upon talin or PIPKI γ /PIPKI γ i2 knockdown. (f, g) NMuMG cells were transfected with siRNA for the knockdown of E-cadherin. Cells were treated with DMSO or LY294002 compound 36–48 h post-transfection, and allowed to grow for another 24–36 h. The activation level of Akt and expression level of mesenchymal marker proteins were examined by immunoblotting. Cell motility was examined as described above. Mesenchymal phenotype and cell motility gained by E-cadherin knockdown cells were significantly compromised upon inhibition of PI3K/Akt signaling by LY294002 compound. Results are represented as mean $\pm$ s.d. from three-independent experiments and error bars represent s.d.

Figure 5.

The PIPKI γ -talin interaction is required for EMT. (**a**, **b**) NMuMG cells seeded into a Col. I-coated culture plate and induced to undergo EMT were transfected with siRNA for the knockdown of talin or moesin. Cell morphology was examined 48 h post-transfection (image taken using $\times 10$ objective lens), and the expression level of E-cadherin and mesenchymal marker proteins were examined by immunoblotting after harvesting the cells. (**c**, **d**) NMuMG cells were co-transfected with siRNA for the knockdown of E-cadherin along with talin or moesin. Cell morphology was examined 48 h post transfection (image taken using $\times 10$ objective lens). Gain of mesenchymal marker protein expression was examined by immunoblotting after harvesting the cells. (**e**) Endogenous talin was immunoprecipitated

from NMuMG cells cultured into Col. I-coated culture plates and/or after the knockdown of moesin. Co-immunoprecipitation of $\beta 1$ integrin with talin was examined by immunoblotting. Moesin knockdown promoted EMT and the association of talin with integrin. **(f, g)** NMuMG cells fully converted into mesenchymal state by TGF $\beta 1$ treatment was transfected with siRNA for knockdown of talin or moesin. At 48–72 h post-transfection, the expression level of E-cadherin and N-cadherin was examined by immunofluorescence study and immunoblotting. **(h)** Schematic diagram depicting talin and PIPKI γ as a platform for bidirectional signaling at cell–matrix interaction sites for epithelial cells transitioning into mesenchymal state. Increased association/expression of talin and PIPKI γ is essential for transitioning epithelial cells which is accompanied by increased adhesion and PI3K/Akt signaling. **(i, j)** NMuMG cells infected with lentiviral vector for expression of GFP-fusion protein of F2 or F3 subdomains of talin FERM domain were induced to undergo EMT by E-cadherin knockdown/culture into Col. I-coated culture plates. Cells were harvested 48–72 h post-transfection. Talin was immunoprecipitated and co-immunoprecipitation of PIPKI $\gamma 2$ was examined to determine the inhibition of talin interaction with PIPKI $\gamma 2$ by F3 subdomain. The effect on focal adhesion and PI3K/Akt signaling by F3 subdomain were determined by examining the phosphorylated FAK and Akt. The effect of F3 subdomain on acquisition of mesenchymal phenotypes were assessed by N-cadherin expression and cell motility. Results are represented as mean $\pm$ s.d. from three-independent experiments and error bars represent s.d.

Figure 6.

PIPKI γ and talin are required for the stability of EMT-regulating transcription factors. (a) Induction of E-cadherin transcriptional repressors (e.g. snail, slug and twist) during transition of NMuMG cells into mesenchymal state by TGF β 1 treatment or culture into ECM protein was examined by immunoblotting. (b) NMuMG cells transformed into mesenchymal cells by TGF β 1 treatment were transfected with siRNA for knockdown of talin or PIPKI γ expression. At 48 h post-transfection, cells were treated with cyclohexamide (CHX, 50 μ g/ml) to block protein synthesis. Cells were harvested at different time points following cyclohexamide treatment to examine the expression level of EMT-regulating transcription factors by immunoblotting. The protein level of snail, slug and β -catenin at different time points is shown as % of control (before cyclohexamide treatment). The knockdown of PIPKI γ and talin is shown by immunoblotting. (c-e) PIPKI γ or talin knockdown cells were treated with proteasomal inhibitor, MG132 (50 μ g/ml), before

harvesting the cells at different time points. The expression level of snail, slug and β -catenin was examined by immunoblotting. (e) The expression level of snail and slug in PIPKI γ or talin knockdown cells before MG132 treatment was quantified by measuring the intensity of immunoblot using ImageJ. The data are presented as mean $\pm$ s.d. from three-independent experiments. (f) siRNA was used to knockdown PIPKI γ /PIPKI γ i2 or talin expression in NMuMG cells overexpressing snail or from Cal51 cells. At 48–72 h post-transfection, cells were treated with MG132 for 3 h before immunoprecipitating the snail. Immunoprecipitates were resolved by sodium dodecyl sulfateS–polyacrylamide gel electrophoresis followed by immunoblotting with HRP-conjugated ubiquitin antibody. (g–i) siRNA was used to knockdown talin or PIPKI γ in NMuMG cells overexpressing snail. At 72-h post-transfection, cell morphology and E-cadherin expression were examined. The effect of talin or PIPKI γ knockdown on cell motility of snail-overexpressing cells was examined as described above. Results are represented as mean $\pm$ s.d. from three independent experiments and error bars represent s.d.

Figure 7.

PIPKI γ and talin control mammosphere/tumorsphere formation of mesenchymal and tumor cells. **(a)** NMuMG cells were transfected with siRNA to knock down talin or PIPKI γ . At 36–48 h after siRNA transfection, cells were treated with TGF β 1 (2 ng/ml) to induce EMT. Cells were allowed to grow in non-adherent suspension condition 48 h after the TGF β 1 treatment. Mammospheres/tumorspheres formed were counted 3–5 days after incubation. **(b)** NMuMG cells were co-transfected with siRNA for the knockdown of E-cadherin and talin or PIPKI γ . 48–72 h after siRNA transfection, cells were allowed to grow in non-adherent condition. Mammospheres/tumorspheres formed were counted 3–5 days after incubation. **(c, d)** NMuMG cells ectopically overexpressing PIPKI γ 1 or PIPKI γ 2 were transfected with siRNA for knockdown of E-cadherin. At 48–72 h post-transfection, cells were resuspended and cultured in non-adherent condition to examine their ability to form mammospheres/tumorspheres. The representative images of mammospheres/tumorspheres formed are shown (images taken using $\times 10$ objective lens). **(e)** Different tumor cells were cultured in non-adherent condition and mammospheres/tumorspheres formed by these tumor cells were isolated and harvested to examine the expression level of PIPKI γ /PIPKI γ 2 and phosphorylated Akt by immunoblotting. **(f)** siRNA was used to knockdown PIPKI γ or

PIPKI γ 2 in different tumor cells (e.g. Cal51, A431 and T47D). 48–72 h post-transfection, cells were harvested and cultured in non-adherent condition to examine their ability to form mammospheres/tumorspheres. Results are represented as mean $\pm$ s.d. from three independent experiments and error bars represent s.d.

Figure 8.

Schematic diagram depicting PIPKI γ and talin control of EMT and mesenchymal phenotypes. During EMT, loss of E-cadherin expression promotes the assembly of PIPKI γ with talin at adhesion complex, which is required for the increased adhesion and PI3K/Akt signaling in the transitioning epithelial cells. PIPKI γ and talin control the stability of E-cadherin transcriptional repressors, snail and slug by preventing their ubiquitination and proteosomal degradation. Acquisition of mesenchymal phenotypes (e.g. cell motility and stemness property) further depends upon the integrity of PIPKI γ and talin (ECM, extracellular matrix; GF, growth factor; E-cad., E-cadherin; MET, mesenchymal epithelial transition).