Published in final edited form as:

Antiviral Res. 2019 January ; 161: 163–168. doi:10.1016/j.antiviral.2018.11.017.

A potent prolyl tRNA synthetase inhibitor antagonizes Chikungunya and Dengue viruses

Jesse Hwang^{a,#}, Alfred Jiang^a, and Erol Fikrig^{a,b}

^aSection of Infectious Diseases, Department of Internal Medicine, Yale University School of Medicine, New Haven, Connecticut, USA

^bHoward Hughes Medical Institute, Chevy Chase, Maryland, USA

Abstract

Arboviruses represent a group of pathogens that can spread efficiently throughout human populations by hematophagous arthropod vectors. The mosquito-borne (re)emerging Chikungunya and Dengue viruses belong to the alphavirus and flavivirus genus, respectively, with no approved therapeutics or safe vaccines for humans. Transmitted by the same vector *Aedes spp.*, these viruses cause significant morbidity and mortality in endemic areas. Due to the increasing likelihood of co-circulation and co-infection with viruses, we aimed to identify a pharmacologically targetable host factor that can inhibit multiple viruses and show that a potent antagonist of prolyl tRNA synthetase (halofuginone) suppresses both Chikungunya and Dengue viruses. Host tRNA synthetase inhibition may signify an additional approach to combat present and future epidemic pathogens.

Keywords

Chikungunya; Dengue; arbovirus; antiviral; halofuginone

Arthropod-borne pathogens are spread from blood-feeding invertebrates to host vertebrate species, and mosquitos constitute a major vector group responsible for transmission of arboviruses. Chikungunya virus (CHIKV) and Dengue virus (DENV) are positive-sense RNA viruses in the alphavirus and flavivirus genera, respectively, that are spread by *Aedes spp.* mosquitos (Higgs and Vanlandingham, 2015; Shaw and Catteruccia, 2018). Both cause fever, rash, headaches, and general malaise, but CHIKV can cause chronic joint

[#]Correspondence: Jesse Hwang, Section of Infectious Diseases, Department of Internal Medicine, Yale University School of Medicine, The Anlyan Center for Medical Research and Education, 300 Cedar Street, New Haven, CT 06520. Phone: (203) 737-5635, jesse.hwang@yale.edu.

Contributions

J.H., E.F.: conceived and designed the project. J.H., A.J.: performed the experiments. J.H. analyzed the data. J.H., E.F.: drafted and revised the manuscript.

Conflict of interests

The authors declare that they have no conflict of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

inflammation and arthritis-like symptoms beyond the acute phase (Schwartz and Albert, 2010), while DENV can lead to hemorrhage and death (Wiwanitkit, 2010).

CHIKV had been confined to Africa and Asia, but in the past decade, the virus has spread to the western hemisphere to cause widespread epidemics (Burt et al., 2017). The typical vector responsible for transmission of CHIKV was *Aedes aegypti*, but a mutation in the viral envelope led to enhanced replication in the *Aedes albopictus* mosquitos present in temperate regions, potentially expanding the range of transmission (Tsetsarkin et al., 2007). CHIKV-induced arthropathy is due to viral replication and the ensuing host inflammatory response (Suhriebier et al., 2012), including inflammasome activation (Chen et al., 2017) and T-cell mediated pathogenesis (Amdekar et al., 2017).

DENV has four serotypes, and antibody-dependent enhancement caused by cross-reactive but sub-neutralizing antibodies has hampered the development of safe vaccines that are equipotent against all four serotypes (Acosta and Bartenschlager, 2016a; Rey et al., 2018). In 2010, there were an estimated 96 million symptomatic cases of dengue infection worldwide, and a significant portion of the human population, especially in the tropics, remains at risk (Bhatt et al., 2013). Unlike CHIKV, normal clinical sequelae of DENV includes mortality.

For both CHIKV and DENV, there are no approved therapeutics aside from palliative care and pain reduction (Fischer et al., 2014; Lee et al., 2017), although many direct virus targeting compounds have been investigated (Boldescu et al., 2017; Powers, 2018). Due to a potent innate immune response and sterilizing immunity, the acute phase of CHIKV and DENV infection is relatively short compared to the duration of chronic infection by other viruses, such as HIV or hepatitis C virus, so the risk of developing escape mutants to a given antiviral compound is lessened. However, direct virus-targeting pharmacophores may be less effective against multiple strains of viruses, or even ineffective across genera. In an epidemic setting where multiple arboviruses might co-circulate, it would be advantageous to find a means to inhibit viruses from different families. Furthermore, homotypic reinfections with genetically similar DENV have been documented (Forshey et al., 2016), suggesting natural infection or vaccination may not offer complete protection in all cases.

Host-targeting antivirals are another class of therapeutics. For instance, metabolism, subcellular trafficking, and cytoskeletal organization are avenues that can be discretely manipulated to hinder various steps of the viral life cycle (Acosta and Bartenschlager, 2016b; Ching et al., 2017; Dohner and Sodeik, 2005; Pombo and Sanyal, 2018). Aside from the decreased chances of emergence of resistance compared to a pathogen-specific approach, hosttargeting compounds have the potential to simultaneously inhibit multiple viruses, provided that the host pathway is an essential component of the replication strategies (Lou et al., 2014). In pragmatic terms, host-targeting compounds designated to be safe may be advantageous in the event of an outbreak caused by a previously uncharacterized virus where vaccines would not be readily available.

Viruses must hijack the host translational machinery to synthesize viral proteins (Hoang et al., 2018). For this reason, a common host response to viral infection is shutting down translation, such as through protein kinase R activation and inhibition of translation initiation

(Garcia et al., 2007). Another endogenous pathway that controls translation initiation is the amino acid response. When the intracellular concentration of one or more amino acids is low, the cognate tRNA remains uncharged and binds to GCN2 kinase (also known as EIF2AK4) (Dong et al., 2000). The phosphorylated, active GCN2 then blocks translation initiation in a manner similar to protein kinase R (Pakos-Zebrucka et al., 2016).

Halofuginone is a potent, orally-available synthetic derivative of the plant compound febrifugine (Pines et al., 2000). By binding to and inhibiting the prolyl tRNA synthetase domain of the bifunctional enzyme EPRS (glutamyl-prolyl-tRNA synthetase), halofuginone causes buildup of uncharged prolyl tRNAs, forcing the cell to trigger the amino acid response and to suppress bulk translation even when proline levels are sufficient (Keller et al., 2012; Yao and Fox, 2013).

To determine the effect of EPRS inhibition on CHIKV replication, we chose to test human foreskin fibroblasts (HFFs) because fibroblasts are physiologically relevant *in vivo* targets (Couderc et al., 2008). HFFs were infected with CHIKV and treated with different concentrations of halofuginone starting at 1 hpi. Supernatants were harvested and cells were fixed with 2% paraformaldehyde at 24 hpi. Halofuginone exhibited dose-dependent inhibition of viral progeny production (Figure 1A), leading to an approximately 3 log₁₀ fold reduction at 100 nM. Likewise, indirect immunofluorescence assay to visualize viral proteins E2 and nsP2 showed dose-dependent reduction in replication-permissive cells (Figure 1B). Pretreatment with halofuginone for 16 hours led to even greater inhibition of viral titer (results not shown), but we performed all subsequent studies with posttreatment alone. There was no cytotoxicity as assessed by LDH release assay (Supplemental Figure 1).

Western blot analysis also showed that halofuginone dose-dependently inhibited viral protein synthesis. HFFs were infected as described above, and whole-cell lysates were harvested at 6, 12, and 24 hpi. As expected, halofuginone treatment led to an increase in phosphorylated GCN2 with maximal induction at 100 nM (Figure 1C), consistent with accumulation of uncharged prolyl tRNA. Both the early non-structural nsP2 protein and late structural capsid protein were repressed by halofuginone, and as a consequence, Rpb1, which is normally degraded by nsP2 in order to cause host transcriptional shutoff (Akhrymuk et al., 2012), remained steady. SQSTM1/p62, the receptor for proteins targeted for autophagosomal degradation, such as CHIKV capsid (Judith et al., 2013), also remained stable in the presence of halofuginone, whereas DMSO treated cells showed gradual downregulation of SQSTM1/p62. Thus, halofuginone efficiently inhibited CHIKV replication at a relatively early step.

Halofuginone binds to EPRS, and proline, the endogenous substrate of EPRS, should compete with halofuginone for interaction. To confirm that the antiviral effect of halofuginone is due to competitive disruption of aminoacylation, infected HFFs were treated with halofuginone or DMSO in the presence or absence of excess proline. Supernatant titration at 24 hpi showed that excess proline completely rescued the defect in virus replication (Figure 1D), and the rescue was further verified by western blot analysis of CHIKV nsP1 and capsid levels (Figure 1E).

DENV is another globally prevalent arbovirus. To determine whether halofuginone is effective against this flavivirus, we infected the permissive Huh7 cell line (Lin et al., 2000) and observed that the compound also led to dose-dependent inhibition of viral progeny release into the culture media (Figure 2A). Indirect immunofluorescence assay showed that treated cells could not form fluorescent foci, indicative of failure to spread cell-to-cell proficiently (Figure 2B). Viral protein levels were drastically reduced after exposure to halofuginone (Figure 2C), including the NS1 protein that is responsible for vascular leakage during severe disease (Beatty et al., 2015). Similar to what was observed with CHIKV, proline co-treatment permitted DENV protein synthesis (Figure 2D). These results show that halofuginone inhibits both arboviruses through EPRS inhibition. We tested additional arboviruses and confirmed that halofuginone also inhibited replication of O'nyong'nyong virus (ONNV), an alphavirus closely related to CHIKV, and Zika virus (ZIKV) in co-infected HFFs (Figure 3), which were infected at a higher MOI due to lower susceptibility.

Although halofuginone is active against both virus families, targeting multiple host pathways may lead to even greater antiviral effects. Interferon signaling is an endogenous mechanism to hinder viral replication, and although injected PEGylated interferon has been utilized in different pathogenic settings (Palumbo, 2009), such an approach may not be practical in areas with less developed healthcare infrastructure. In contrast, RO8191 is a synthetic, orally-available IFNAR2 agonist that engenders an antiviral state like *bona fide* interferon, and importantly, this compound induces interferon stimulated genes (ISGs) without significant elevation of pro-inflammatory genes (Konishi et al., 2012).

To evaluate whether halofuginone can be used in combination with RO8191, we first confirmed the efficacy of the compound in our system and showed that RO8191 dose-dependently increased STAT1 and ISG15 levels (Figure 4A, C). Next, we pretreated HFFs or Huh7 cells with the lowest dose of RO8191 (0.33 μ M) or DMSO for 24 hours and then infected HFFs with CHIKV or Huh7 cells with DENV. After 1 hour post infection, cells were washed and incubated with or without halofuginone in the absence of any RO8191. Viral titration at 24 hpi showed that while each compound inhibited replication individually, successive exposure to both led to an even greater effect (Figure 4B, D). Co-treatment of compounds post-infection also displayed augmented effects, although total efficacy was less (Supplemental Figure 2). These results show that even if a single compound fails to reach maximal effective dose, targeting multiple pathways consecutively may still yield protective antiviral effect.

CHIKV and DENV continue to pose a significant challenge for multiple reasons. There is a lack of safe vaccine against the four serotypes of DENV due to complications of antibody-dependent enhancement (Acosta and Bartenschlager, 2016a), and even for CHIKV which has a single serotype (Sahadeo et al., 2015), there is reduced impetus for human vaccine clinical trials, with one factor being that CHIKV is usually not lethal and infection leads to protective immunity.

Another complicating factor that delays response to an arbovirus epidemic is unpredictability of emergence. CHIKV acquired mutations that allowed replication in a mosquito species that previously exhibited low vector competence (Tsetsarkin et al., 2016),

broadening the potential range of transmission. Although this particular virus lineage was not responsible for the large outbreaks in the Americas (2013-2015), the threat continues to persist in the Pacific, Southeast Asia, and Latin America (Pyke et al., 2018). Most recently, the formerly obscure Zika virus caused a large global epidemic. Two key mutations in the viral genome have been identified, which increased virus acquisition by *Aedes* mosquitoes (NS1 mutation) (Liu et al., 2017) and enhanced infection of neural progenitor cells to contribute to fetal microcephaly (prM mutation) (Yuan et al., 2017).

Thus, every arbovirus has unique biological and practical elements that need to be taken into account to limit spread, and this is compounded by increased likelihood of co-circulation and co-infection of pathogens. Because viruses require the host cell resources for replication, therapeutically targeting a common essential factor could be beneficial to inhibit multiple viruses even if their replication strategies are otherwise divergent.

Host translational machinery is an absolute requirement for virus production, and one way to inhibit translation is to activate an endogenous mechanism that suppresses global protein synthesis. GCN2 is a sensor of uncharged tRNAs which accumulate during times of amino acid deprivation, and this kinase shuts down bulk translation to prevent wasteful energy expenditure (Dong et al., 2000). In fact, limitation of a single amino acid can activate GCN2 (Peng et al., 2012).

We tested halofuginone, an inhibitor of EPRS (Keller et al., 2012), against CHIKV and DENV because it is potent and orally-available, which may be practical criteria in clinical settings. Halofuginone indeed inhibited both alphaviruses and flaviviruses in a proline-dependent manner (Figures 1-3). It was previously shown that GCN2 knockout cells and mice were more susceptible to Sindbis virus replication (Berlanga et al., 2006), but thus far, pharmacological activation has not been tested against mosquito-borne viruses. We demonstrate that a single compound that prevents aminoacylation inhibits RNA viruses of epidemic potential, and it is likely that halofuginone would inhibit replication of other emerging arboviruses, such as those spread by ticks (e.g., tick-borne encephalitis virus or Powassan virus).

Halofuginone can also exert anti-inflammatory effects. Recently, it was shown that treatment of murine macrophages with halofuginone leads to sequestration of IL-1 β mRNA into translationally inactive riboclusters, thereby suppressing the inflammasome (Battu et al., 2018). Since inflammasome activation plays a significant role in CHIKV pathogenesis (Chen et al., 2017), halofuginone may be efficacious against both viral replication and the subsequent arthralgia-inducing host inflammation. Furthermore, halofuginone downregulates NF- κ B signaling pathway (Luo et al., 2018), which contributes to DENV-induced hemorrhage (Lin et al., 2014). Interestingly, both protein-deficient diet and halofuginone protected against ischemia reperfusion injury, leading to improved organ function concomitant with lower inflammatory markers (Peng et al., 2012). It remains possible that modifying dietary protein intake or composition can ameliorate virus-induced inflammation and shock.

Although inhibiting host translation may lead to toxic effects, halofuginone and similar derivatives have been shown in murine models to inhibit malaria (Herman et al., 2015), reduce cardiac stress (Qin et al., 2017), suppress viral myocarditis (Sun et al., 2016), and improve hepatitis B virus-induced liver inflammation (Zhan et al., 2017). Furthermore, commercial preparations of halofuginone (Stenorol and Halocur) are used in livestock to inhibit parasite growth (Pines and Spector, 2015), and a clinical trial for a slow-release form of halofuginone has been approved by the FDA to treat Duchenne muscular dystrophy (Pines and Spector, 2015). Thus, with proper dosing or synergy with other antiviral compounds, such as an orally-available interferon agonist (Figure 4), targeting host aminoacylation may be an effective strategy against arboviruses that gain a foothold in naive human populations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

E.F. is an investigator with the Howard Hughes Medical Institute. J.H. is supported by a training grant from the NIH 4T32HL007974-15.

References

- Acosta EG, Bartenschlager R, 2016a Paradoxical role of antibodies in dengue virus infections: considerations for prophylactic vaccine development. *Expert Rev Vaccines* 15, 467–482. [PubMed: 26577689]
- Acosta EG, Bartenschlager R, 2016b The quest for host targets to combat dengue virus infections. *Current opinion in virology* 20, 47–54. [PubMed: 27648486]
- Akhrymuk I, Kulemzin SV, Frolova EI, 2012 Evasion of the innate immune response: the Old World alphavirus nsP2 protein induces rapid degradation of Rpb1, a catalytic subunit of RNA polymerase II. *J Virol* 86, 7180–7191. [PubMed: 22514352]
- Amdekar S, Parashar D, Alagarasu K, 2017 Chikungunya Virus-Induced Arthritis: Role of Host and Viral Factors in the Pathogenesis. *Viral Immunol* 30, 691–702. [PubMed: 28910194]
- Battu S, Afroz S, Giddaluru J, Naz S, Huang W, Khumukcham SS, Khan RA, Bhat SY, Qureshi IA, Manavathi B, Khan AA, August A, Hasnain SE, Khan N, 2018 Amino acid starvation sensing dampens IL-1 β production by activating riboclustering and autophagy. *PLoS Biol* 16, e2005317. [PubMed: 29621237]
- Beatty PR, Puerta-Guardo H, Killingbeck SS, Glasner DR, Hopkins K, Harris E, 2015 Dengue virus NS1 triggers endothelial permeability and vascular leak that is prevented by NS1 vaccination. *Sci Transl Med* 7, 304ra141.
- Berlanga JJ, Ventoso I, Harding HP, Deng J, Ron D, Sonenberg N, Carrasco L, de Haro C, 2006 Antiviral effect of the mammalian translation initiation factor 2 α kinase GCN2 against RNA viruses. *EMBO J* 25, 1730–1740. [PubMed: 16601681]
- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Hoen AG, Sankoh O, Myers MF, George DB, Jaenisch T, Wint GR, Simmons CP, Scott TW, Farrar JJ, Hay SI, 2013 The global distribution and burden of dengue. *Nature* 496, 504–507. [PubMed: 23563266]
- Boldescu V, Behnam MAM, Vasilakis N, Klein CD, 2017 Broad-spectrum agents for flaviviral infections: dengue, Zika and beyond. *Nat Rev Drug Discov* 16, 565–586. [PubMed: 28473729]

- Burt FJ, Chen W, Miner JJ, Lenschow DJ, Merits A, Schnettler E, Kohl A, Rudd PA, Taylor A, Herrero LJ, Zaid A, Ng LFP, Mahalingam S, 2017 Chikungunya virus: an update on the biology and pathogenesis of this emerging pathogen. *Lancet Infect Dis* 17, e107–e117. [PubMed: 28159534]
- Chen W, Foo SS, Zaid A, Teng TS, Herrero LJ, Wolf S, Tharmarajah K, Vu LD, van Vreden C, Taylor A, Freitas JR, Li RW, Woodruff TM, Gordon R, Ojcius DM, Nakaya HI, Kanneganti TD, O'Neill LAJ, Robertson AAB, King NJ, Suhrbier A, Cooper MA, Ng LFP, Mahalingam S, 2017 Specific inhibition of NLRP3 in chikungunya disease reveals a role for inflammasomes in alphavirus-induced inflammation. *Nat Microbiol*.
- Ching KC, L FPN, Chai CLL, 2017 A compendium of small molecule direct-acting and host-targeting inhibitors as therapies against alphaviruses. *J Antimicrob Chemother* 72, 2973–2989. [PubMed: 28981632]
- Couderc T, Chretien F, Schilte C, Disson O, Brigitte M, Guivel-Benhassine F, Touret Y, Barau G, Cayet N, Schuffenecker I, Despres P, Arenzana-Seisdedos F, Michault A, Albert ML, Lecuit M, 2008 A mouse model for Chikungunya: young age and inefficient type-I interferon signaling are risk factors for severe disease. *PLoS Pathog* 4, e29. [PubMed: 18282093]
- Dohner K, Sodeik B, 2005 The role of the cytoskeleton during viral infection. *Curr Top Microbiol Immunol* 285, 67–108. [PubMed: 15609501]
- Dong J, Qiu H, Garcia-Barrio M, Anderson J, Hinnebusch AG, 2000 Uncharged tRNA activates GCN2 by displacing the protein kinase moiety from a bipartite tRNA-binding domain. *Mol Cell* 6, 269–279. [PubMed: 10983975]
- Fischer M, Staples JE, Arboviral Diseases Branch, N.C.f.E., Zoonotic Infectious Diseases, C.D.C., 2014 Notes from the field: chikungunya virus spreads in the Americas - Caribbean and South America, 2013-2014. *MMWR Morb Mortal Wkly Rep* 63, 500–501. [PubMed: 24898168]
- Forshey BM, Stoddard ST, Morrison AC, 2016 Dengue Viruses and Lifelong Immunity: Reevaluating the Conventional Wisdom. *J Infect Dis* 214, 979–981. [PubMed: 26984147]
- Garcia MA, Meurs EF, Esteban M, 2007 The dsRNA protein kinase PKR: virus and cell control. *Biochimie* 89, 799–811. [PubMed: 17451862]
- Herman JD, Pepper LR, Cortese JF, Estiu G, Galinsky K, Zuzarte-Luis V, Derbyshire ER, Ribacke U, Lukens AK, Santos SA, Patel V, Clish CB, Sullivan WJ, Jr., Zhou H, Bopp SE, Schimmel P, Lindquist S, Clardy J, Mota MM, Keller TL, Whitman M, Wiest O, Wirth DF, Mazitschek R, 2015 The cytoplasmic prolyl-tRNA synthetase of the malaria parasite is a dual-stage target of febrifugine and its analogs. *Sci Transl Med* 7, 288ra277.
- Higgs S, Vanlandingham D, 2015 Chikungunya Virus and Its Mosquito Vectors. *Vector Borne Zoonotic Dis* 15, 231–240. [PubMed: 25674945]
- Hoang HD, Graber TE, Alain T, 2018 Battling for Ribosomes: Translational Control at the Forefront of the Antiviral Response. *J Mol Biol* 430, 1965–1992. [PubMed: 29746850]
- Judith D, Mostowy S, Bourai M, Gangneux N, Lelek M, Lucas-Hourani M, Cayet N, Jacob Y, Prevost MC, Pierre P, Tangy F, Zimmer C, Vidalain PO, Couderc T, Lecuit M, 2013 Species-specific impact of the autophagy machinery on Chikungunya virus infection. *EMBO Rep* 14, 534–544. [PubMed: 23619093]
- Keller TL, Zocco D, Sundrud MS, Hendrick M, Edenius M, Yum J, Kim YJ, Lee HK, Cortese JF, Wirth DF, Dignam JD, Rao A, Yeo CY, Mazitschek R, Whitman M, 2012 Halofuginone and other febrifugine derivatives inhibit prolyl-tRNA synthetase. *Nat Chem Biol* 8, 311–317. [PubMed: 22327401]
- Konishi H, Okamoto K, Ohmori Y, Yoshino H, Ohmori H, Ashihara M, Hirata Y, Ohta A, Sakamoto H, Hada N, Katsume A, Kohara M, Morikawa K, Tsukuda T, Shimma N, Foster GR, Alazawi W, Aoki Y, Arisawa M, Sudoh M, 2012 An orally available, smallmolecule interferon inhibits viral replication. *Scientific reports* 2, 259. [PubMed: 22355771]
- Lee TH, Lee LK, Lye DC, Leo YS, 2017 Current management of severe dengue infection. *Expert Rev Anti Infect Ther* 15, 67–78. [PubMed: 27786589]
- Lin JC, Lin SC, Chen WY, Yen YT, Lai CW, Tao MH, Lin YL, Miaw SC, Wu-Hsieh BA, 2014 Dengue viral protease interaction with NF-kappaB inhibitor alpha/beta results in endothelial cell apoptosis and hemorrhage development. *J Immunol* 193, 1258–1267. [PubMed: 24973451]

- Lin YL, Liu CC, Lei HY, Yeh TM, Lin YS, Chen RM, Liu HS, 2000 Infection of five human liver cell lines by dengue-2 virus. *J Med Virol* 60, 425–431. [PubMed: 10686026]
- Liu Y, Liu J, Du S, Shan C, Nie K, Zhang R, Li XF, Zhang R, Wang T, Qin CF, Wang P, Shi PY, Cheng G, 2017 Evolutionary enhancement of Zika virus infectivity in *Aedes aegypti* mosquitoes. *Nature* 545, 482–486. [PubMed: 28514450]
- Lou Z, Sun Y, Rao Z, 2014 Current progress in antiviral strategies. *Trends Pharmacol Sci* 35, 86–102. [PubMed: 24439476]
- Luo L, Gao Y, Yang C, Shao Z, Wu X, Li S, Xiong L, Chen C, 2018 Halofuginone attenuates intervertebral discs degeneration by suppressing collagen I production and inactivating TGFbeta and NF-small ka, CyrillicB pathway. *Biomed Pharmacother* 101, 745–753. [PubMed: 29524883]
- Mainou BA, Zamora PF, Ashbrook AW, Dorset DC, Kim KS, Dermody TS, 2013 Reovirus cell entry requires functional microtubules. *MBio* 4.
- Pakos-Zebrucka K, Koryga I, Mnich K, Ljubic M, Samali A, Gorman AM, 2016 The integrated stress response. *EMBO Rep* 17, 1374–1395. [PubMed: 27629041]
- Palumbo E, 2009 PEG-interferon in acute and chronic hepatitis C: a review. *American journal of therapeutics* 16, 573–578. [PubMed: 19433975]
- Peng W, Robertson L, Gallinetti J, Mejia P, Vose S, Charlip A, Chu T, Mitchell JR, 2012 Surgical stress resistance induced by single amino acid deprivation requires Gcn2 in mice. *Sci Transl Med* 4, 118ra111.
- Pines M, Spector I, 2015 Halofuginone - the multifaceted molecule. *Molecules* 20, 573–594. [PubMed: 25569515]
- Pines M, Vlodavsky I, Nagler A, 2000 Halofuginone: From veterinary use to human therapy. *Drug Develop Res* 50, 371–378.
- Pombo JP, Sanyal S, 2018 Perturbation of Intracellular Cholesterol and Fatty Acid Homeostasis During Flavivirus Infections. *Frontiers in immunology* 9, 1276. [PubMed: 29915602]
- Powers AM, 2018 Vaccine and Therapeutic Options To Control Chikungunya Virus. *Clinical microbiology reviews* 31.
- Pyke AT, Moore PR, McMahon J, 2018 New insights into chikungunya virus emergence and spread from Southeast Asia. *Emerging microbes & infections* 7, 26. [PubMed: 29535302]
- Qin P, Arabacilar P, Bernard RE, Bao W, Olzinski AR, Guo Y, Lal H, Eisennagel SH, Platchek MC, Xie W, Del Rosario J, Nayal M, Lu Q, Roethke T, Schnackenberg CG, Wright F, Quail MP, Halsey WS, Hughes AM, Sathe GM, Livi GP, Kirkpatrick RB, Qu XA, Rajpal DK, Faeltz Savitski M, Bantscheff M, Joberty G, Bergamini G, Force TL, Gatto GJ, Jr., Hu E, Willette RN, 2017 Activation of the Amino Acid Response Pathway Blunts the Effects of Cardiac Stress. *J Am Heart Assoc* 6.
- Rey FA, Stiasny K, Vaney MC, Dellarole M, Heinz FX, 2018 The bright and the dark side of human antibody responses to flaviviruses: lessons for vaccine design. *EMBO Rep* 19, 206–224. [PubMed: 29282215]
- Sahadeo N, Mohammed H, Allicock OM, Auguste AJ, Widen SG, Badal K, Pulchan K, Foster JE, Weaver SC, Carrington CV, 2015 Molecular Characterisation of Chikungunya Virus Infections in Trinidad and Comparison of Clinical and Laboratory Features with Dengue and Other Acute Febrile Cases. *PLoS neglected tropical diseases* 9, e0004199. [PubMed: 26580074]
- Schwartz O, Albert ML, 2010 Biology and pathogenesis of chikungunya virus. *Nature reviews. Microbiology* 8, 491–500. [PubMed: 20551973]
- Shaw WR, Catteruccia F, 2018 Vector biology meets disease control: using basic research to fight vector-borne diseases. *Nat Microbiol*.
- Suhrbier A, Jaffar-Bandjee MC, Gasque P, 2012 Arthritogenic alphaviruses--an overview. *Nature reviews. Rheumatology* 8, 420–429. [PubMed: 22565316]
- Sun XH, Fu J, Sun DQ, 2016 Halofuginone alleviates acute viral myocarditis in suckling BALB/c mice by inhibiting TGF-beta1. *Biochem Biophys Res Commun* 473, 558–564. [PubMed: 27021682]
- Tsetsarkin KA, Chen R, Weaver SC, 2016 Interspecies transmission and chikungunya virus emergence. *Current opinion in virology* 16, 143–150. [PubMed: 26986235]
- Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S, 2007 A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS Pathog* 3, e201. [PubMed: 18069894]

- Wiwanitkit V, 2010 Dengue fever: diagnosis and treatment. *Expert Rev Anti Infect Ther* 8, 841–845. [PubMed: 20586568]
- Yao P, Fox PL, 2013 Aminoacyl-tRNA synthetases in medicine and disease. *EMBO molecular medicine* 5, 332–343. [PubMed: 23427196]
- Yuan L, Huang XY, Liu ZY, Zhang F, Zhu XL, Yu JY, Ji X, Xu YP, Li G, Li C, Wang HJ, Deng YQ, Wu M, Cheng ML, Ye Q, Xie DY, Li XF, Wang X, Shi W, Hu B, Shi PY, Xu Z, Qin CF, 2017 A single mutation in the prM protein of Zika virus contributes to fetal microcephaly. *Science* 358, 933–936. [PubMed: 28971967]
- Zhan W, Kang Y, Chen N, Mao C, Kang Y, Shang J, 2017 Halofuginone ameliorates inflammation in severe acute hepatitis B virus (HBV)-infected SD rats through AMPK activation. *Drug Des Devel Ther* 11, 2947–2955.

Highlights

- Specific aminoacylation inhibitor suppresses replication of globally relevant arboviruses
- Phenotype is completely reversed by excess endogenous substrate
- Antiviral effect is synergistic with synthetic IFNAR2 agonist

Figure 1.

Halofuginone inhibits CHIKV replication. (A) HFFs were infected with CHIKV (MOI 1) and treated with indicated concentrations of halofuginone at 1 hpi. Supernatants were titrated at 24 hpi. (B) Similar to (A) except cells were processed for indirect immunofluorescence to detect CHIKV E2 (red) and CHIKV nsP2 (green). Nuclei are stained with Hoechst. (C) HFFs were infected with CHIKV (MOI 1) and treated with halofuginone at 1 hpi. Whole-cell lysates were harvested at 6, 12, and 24 hpi for western blot analysis. (D) HFFs were infected with CHIKV (MOI 1) and treated with halofuginone (100nM) or proline (4 mM). Supernatants were titrated at 24 hpi. (E) Similar to (D) except whole-cell lysates were

subjected to western blot analysis. Values are the means $\pm$ standard deviation (SD). Significant values are defined by $*P < 0.05$.

Figure 2.

Halofuginone inhibits DENV replication. (A) Huh7 cells were infected with DENV (MOI 0.5) and treated with halofuginone at 1 hpi. Supernatants were titrated at 24 and 48 hpi. (B) Similar to (A) except infected cells were detected by indirect immunofluorescence assay using pan-flavivirus monoclonal antibody (clone 4G2). (C) Huh7 cells were infected with DENV (MOI 0.5) and treated with halofuginone. Protein samples were analyzed at 24 and 48 hpi to detect viral protein accumulation. (D) Similar to (C) except cells were treated with halofuginone (50 nM) or proline (4 mM). Protein samples were harvested at 48 hpi for detection of DENV NS5 and NS1. Values are the means $\pm$ standard deviation (SD). Significant values are defined by $*P < 0.05$.

Figure 3.

Effect of halofuginone on co-infection with related viruses. HFFs were co-infected with ONNV (MOI 10) and ZIKV (MOI 10) and treated with halofuginone at 2 hpi. Supernatants were titrated at 24 hpi. Fluorescent focus units (FFU) of ONNV and ZIKV were measured by using cross-reactive monoclonal antibody against CHIKV E2 and pan-flavivirus monoclonal antibody, respectively. Values are the means $\pm$ standard deviation (SD). Significant values are defined by * $P < 0.05$.

Figure 4.

Synergistic activity of halofuginone with IFNAR2 agonist. (A) HFFs were treated with RO8191 for 24 hours, and protein samples were immunoblotted against interferon stimulated genes. (B) HFFs were pretreated 24 hours with RO8191 (0.33 μM) or DMSO and inoculated with CHIKV (MOI 1). After 1 hour viral adsorption period, cells were washed and replaced with media containing halofuginone (50 nM) or DMSO. Protein samples and supernatants were harvested at 24 hpi. (C) Similar to (A) except Huh7 cells were treated with RO8191 for 24 hours. (D) Similar to (B) except Huh7 cells were infected with DENV (MOI 0.5). Values are the means ± standard deviation (SD). Significant values are defined by * $P < 0.05$.