Published in final edited form as:

Trends Endocrinol Metab. 2019 February ; 30(2): 93–105. doi:10.1016/j.tem.2018.11.004.

The Role of Immature and Mature Adipocytes in Hair Cycling

Ilja L. Kruglikov¹, Zhuzhen Zhang², and Philipp E. Scherer²

¹Scientific Department, Wellcomet GmbH, Karlsruhe, Germany

²Touchstone Diabetes Center, Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390-8549, USA

Abstract

Hair follicles (HFs) strongly interact with adipocytes within the dermal white adipose tissue (dWAT), suggesting a strong physiological dependence on the content of immature and mature adipocytes in this layer. This content is regulated by processes of proliferation and differentiation of adipose precursors and preadipocytes, as well as by dedifferentiation of mature existing adipocytes. Spatially, long-range interactions between HFs and dWAT involve the exchange of extracellular vesicles which are differently released by precursors, preadipocytes and mature adipocytes. Different exogenous factors, including light irradiation, are likely to modify the release of adipocyte-derived exosomes in dWAT, which can lead to aberrations of the HF cycle. Consequently, dWAT should be considered as a potential target for the modulation of hair growth.

Keywords

hair follicle cycle; dermal white adipose tissue; immature adipocytes; mature adipocytes; exosomal exchange

Introduction

There is a rapidly growing body of data that suggests that hair follicle (HF) cycling strongly correlates with spatiotemporal behavior of the dermal white adipose tissue (dWAT) located in rodents above the *panniculus carnosus* and in humans in the superficial layer of the subcutaneous adipose tissue adjacent to the dermis [1-5].

Different correlations between HFs and dWAT were observed in various murine knockout models. Myelin protein zero-like 3 (*Mpzl3*^{-/-}) knockout mice [6], which develop alopecia

Corresponding author: Scherer, P.E., philipp.scherer@utsouthwestern.edu.

Author Contributions

Conceptualization: Ilja L. Kruglikov, Zhuzhen Zhang and Philipp E. Scherer.

Formal analysis: Ilja L. Kruglikov.

Funding acquisition: Philipp E. Scherer.

Methodology: Ilja L. Kruglikov.

Writing - original draft: Ilja L. Kruglikov.

Writing - review & editing: Zhuzhen Zhang and Philipp E. Scherer.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

soon after birth, demonstrate no significant changes in the HF cycle, but a striking reduction in dWAT thickness as well as in the average size of adipocytes. The loss of *Mpl3* function may cause defects in the HF structures formed during anagen. Knockout mice for early B-cell factor-1 (*Ebfl*^{-/-}) are characterized by reduced dWAT and lack of adipocyte progenitors in the postnatal skin. HFs in these animals fail to re-enter anagen [1,7] which leads to the abortion of the HF cycle. It should be, however, mentioned that *Ebfl* is also highly expressed in dermal papilla fibroblasts and its defect can directly perturb HFs cycling.

Mice lacking platelet derived growth factor (PDGFA) demonstrate phenotypic similarities with the *Ebfl*^{-/-} model and produce a block in anagen [1]. At the same time, intradermal preadipocytes have a high expression level of PDGFA, and local injections of PDGFA were able to induce anagen in murine HFs [1,8]. Transgenic mice overexpressing human apolipoprotein CI in the skin, fatty acid transport protein (FATP)-4-deficient mice as well as *Dgat1*^{-/-} or *Dgat2*^{-/-} mice also have reduced dWAT layers and display hair loss 2.

Cyclosporine A (CsA) is a well-known HF stimulator which causes prolongation of anagen both *in vitro* and *in vivo* [9]. CsA downregulates the expression of secreted frizzled related protein 1 (SFRP1) in human HFs *ex vivo* [10]. On the one hand, SFRP1 is an antagonist of Wnt/ β -catenin signaling; while on the other hand, it is a known adipokine [11]. Suppression of SFRP1 with subsequent activation of Wnt/ β -catenin signaling in preadipocytes inhibits their differentiation, whereas both intra- and extracellular activation of SFRP1 and inactivation of Wnt/ β -catenin pathway induce adipogenesis [12]. The latter observation indicates that SFRP1 modulating HF development must not be of intracellular origin since SFRP1 is transported from differentiating preadipocytes located in dWAT to HFs. A more recent report indicates that CsA inhibits adipogenic differentiation in adipocyte precursors [13]. Generally, interactions between dWAT and HFs should be reciprocal, and the influence of epidermal Wnt/ β -catenin signaling on adipocyte differentiation was investigated in [14].

Whereas most of studies referenced above do not describe direct cellular pathways and interactions, periodical cycling of dWAT synchronized with the HF cycle [3,5] together with above results suggest that the modification of dWAT content can influence HF cycling. Here, we discuss possible mechanisms for these interactions and how their modification can lead to aberration of HF cycles.

Immature and mature adipocytes in HF cycling

Adipocytes are derived from mesenchymal stem cells which can differentiate in various cell lineages. These stem cells (also known as adipose-derived stem cells, ADSCs) mainly reside in the vascular stroma of adipose tissue and can undergo a multistep commitment process producing preadipocytes (immature adipocytes) which bear some typical adipocyte markers, among them PPAR γ . Transformation of precursors to the adipocyte lineage is initiated by the activation of several factors, including bone morphogenetic proteins (BMPs), as well as components of the Wnt and hedgehog signaling pathways [15]. These preadipocytes can further differentiate into mature, lipid-laden adipocytes (Fig. 1).

ADSCs, preadipocytes and mature adipocytes (Table 1) are differentially involved in HF cycling: while appearance of mature adipocytes inhibits HF progression, there are different indications that stimulation of immature adipocytes in dWAT can induce the telogen-to-anagen transition [2,5,16]. Non-committed ADSCs have stimulatory effects on dermal papilla cells and are able to promote HF cycling [17,18]. At the same time, defects in the generation of immature adipocytes block follicle stem cell activation and immature adipocytes thought to be not only necessary, but also sufficient for the proper HF cycling [1].

There are distinct roles of ADSCs versus preadipocytes in HF cycling. While almost normal HF cycling is observed in A-ZIP/F1 mice, which contain both ADSCs and preadipocytes, but lack mature white adipocytes system-wide, depletion of immature adipocytes (e.g., with PPAR γ antagonists or genetically in *Ebf1*^{-/-} mice) caused complete HF cycle aberration. It should be noted that the application of PPAR γ antagonists such as BADGE or GW9662 is postulated to deplete preadipocytes, but not ADSCs, which suggests a unique role for preadipocytes in HF cycling [1,5]. On the other hand, transplantation of ADSCs into the skin was able to induce anagen in both wild-type and *Ebf1*^{-/-} mice [1]. This clearly demonstrates that immature adipocytes in dWAT can serve as a target for modulation of the HF cycle.

Mature adipocytes can also influence the HF cycle through expression of BMPs. Adipocyte-derived BMP2 is reaching HF stem cells while they are in the quiescent state [5,19] and thus suppresses hair growth. BMP expression levels increase during anagen, along with the increased concentration of mature adipocytes around HFs. On the other hand, BMPs can significantly influence the appearance and behavior of immature adipocytes: BMP4 not only promotes adipogenic commitment and subsequent differentiation of adipose precursor cells, but it is also induced in precursor cells during their differentiation, as well as promoting the differentiation of preadipocytes and the activation of PPAR γ in these cells [20]. Deletion of BMPRI1 in myofibroblasts results in the lack of new adipocytes, which demonstrated that BMP signaling in myofibroblasts is necessary for adipocyte regeneration in wound healing [21]. Additionally, mature adipocytes can regulate some differentiation processes in HFs during late anagen [19,22].

The local volume of dWAT adjacent to HFs increases in anagen and significantly decreases in catagen and telogen [3,5]. Such remodeling of dWAT includes not only the processes of hyperplasia and hypertrophy. It is highly likely that some dedifferentiation and maybe transdifferentiation events are part of the process as well.

Transitions between immature and mature adipocytes as a hallmark of proper HF cycling

Modification of the dWAT volume and structure during HF cycling demonstrates two unique features. First, 20-40% of mature adipocytes in fully expanded dWAT (mid-anagen) are supposed to be newly formed cells derived from differentiated progenitors, and the remaining cells are thought to be pre-existing adipocytes [23,24]. This means that a proper HF cycling can take place only if a sufficient number of immature adipocytes in dWAT have the ability to proliferate and differentiate. Consequently, depletion of immature adipocytes in

dWAT should lead to aberration of HF cycle. However, the field is awaiting more detailed lineage tracing studies to see whether this is actually true.

Second, as HFs enter telogen, dWAT abruptly decreases to approximately half of its maximum local volume. Such quick transformation of the dWAT layer is not caused by apoptotic cell death, and it was assumed that this effect is facilitated by the stimulation of lipolysis [5]. However, as it was recently stated in [25], the relative contributions of hypertrophy/lipolysis and hyperplasia in modulation of dWAT volume during HF cycling is different in spontaneous and induced HF cycling. Whereas all these processes are evidently important for dWAT modulation during the HF cycle, another phenomenon may also significantly contribute to this process - namely, the dedifferentiation of adipocytes.

Mature adipocytes can indeed undergo trans- and dedifferentiation [26]. In the latter case, they produce a population of adipocyte-derived preadipocytes, ADPs, which have an intermediate fibroblast-adipocyte phenotype and bear the markers for both of these cell types, even though most ADPs exhibit a fibroblast-like morphology and are very similar to adipose tissue derived stem/stromal cells. They do express some specific markers *in vitro*, and can differentiate into adipogenic/osteogenic/chondrogenic lineages by culture. Among others, expressions of transforming growth factor beta (TGF- β 1) and different collagen genes are significantly higher in ADP cells compared with normal adipocytes; moreover, it was shown that stimulation of TGF- β 1 expression enhances adipogenic dedifferentiation [27]. Such dedifferentiation was reliably reproduced in different experiments *in vitro*, and dedifferentiated cells can either differentiate back into mature adipocytes or into different other lineages in tissue culture [28]. This awaits validation *in vivo*.

Another transformation of adipocytes was observed *in vivo* after subcutaneous injections of bleomycin known to induce systemic sclerosis in murine models. Bleomycin strongly suppresses the adipogenic commitment of adipocyte precursors and simultaneously increases the expression of TGF- β 1 in injected areas [29]. Such TGF- β 1 induction is thought to lead to an adipocyte-myofibroblast transition, which involves formerly adiponectin positive intradermal progenitors from dWAT [30]. Already twenty-four hours after stimulation with TGF- β 1, dermal adipocytes demonstrate an intermediate phenotype bearing both perilipin (typical for adipocytes) and alpha smooth muscle actin (typical for myofibroblasts) markers. Such an intermediate state is characteristic for adipocytes from the interfacial adipose layer adjacent to the skin. Similar effects were observed after chronic UV irradiation of murine skin (see discussion in [31]) and was described in cancer progression, where these cells were named cancer-associated adipocytes [32]. Recently, it was also argued that adipocytes with intermediate phenotypes may be involvement in the pathophysiology of androgenetic alopecia [33] and melanoma [34].

Dedifferentiation of adipocytes and induction of a quasi-fibroblast phenotype in dedifferentiated cells in murine model was postulated to be triggered by expression of resistin-like molecule α (RELMA/FIZZ1) [35,36]. Moreover, FIZZ1 knockout mice treated with bleomycin exhibit significantly impaired pulmonary fibrosis [37], which may be linked to the suppression of adipocyte dedifferentiation. The main sources of FIZZ1 in rodent skin are the adipocytes, macrophages and HF epithelial cells. FIZZ1 is much more highly

expressed in mature adipocytes compared to preadipocytes and is able to inhibit the differentiation of preadipocytes [35]. It is likely that the abrupt reduction of dWAT volume in murine model as HF's enter telogen is connected with a local activation of RELM α /FIZZ1 expression that may lead to dedifferentiation of existing mature adipocytes. Whereas the key trigger leading to adipocyte dedifferentiation in humans is unknown, it is possible that RELM β /FIZZ2 is involved as well.

Two other phenotypic changes - epithelial-mesenchymal and endothelial-mesenchymal transitions - were earlier reported to be reversible both *in vitro* and *in vivo* [38,39]. It was even postulated that the cells involved should have a very high flexibility in their phenotype, allowing them to undergo multiple rounds of conversions [38]. In analogy, one could envision that under proper conditions, not only hyperplasia, hypertrophy and lipolysis, but also multiple cycles of transitions between immature and mature adipocytes in dWAT could take place during HF cycling *in vivo* (Fig. 1). This important postulate still has to be formally demonstrated.

The ratio of immature-to-mature adipocytes in dWAT correlates with the HF cycle and reaches its maximum during the mid-anagen phase, connected with downward growth of HF's [1,5]. This ratio in a given adipose depot is, however, not constant and can be affected by different exogenous and endogenous factors.

Mechanisms for long-range spatial interactions between HF's and dWAT

Selective modulation of adipogenic differentiation is likely to solicit a reciprocal reaction in adjacent HF's inducing their cycling or bring this process to a halt. In order for this to occur, the induction of HF modulating factors should be spatially linked between dWAT and HF's.

Normally, proliferating stem cells divide asymmetrically, producing one stem cell and one adipogenic progenitor committed to differentiation. However, in some tissues, stem cells produce fast-cycling "transit amplifying cells" (TACs), which undergo differentiation only after several rounds of mitosis [40]. TACs have a very high capacity to amplify the number of differentiated cells and are typical in HF's. These cells are known as an early intermediate in tissue regeneration [41], and their appearance generally means that the regeneration processes initiated by stem cells has reached the "point of no return". These cells first appear in the HF matrix during anagen II, and their content sufficiently increases during anagen progression [41]. HF-TACs are strongly involved in the downgrowth of HF's and actively express *Sonic hedgehog* (Shh) in a HF-cycle dependent manner. Moreover, these cells may be responsible for a morphological and physiological coupling between HF's and dWAT through the regulation of proliferation of adipose precursors [23].

Hedgehog signaling is known to progressively decrease during adipocyte differentiation in humans [42]. Activation of this pathway (similar to Wnt signaling) impairs adipogenesis and lipid accumulation, whereas suppression of hedgehog signaling was reported to be a necessary condition for adipocyte differentiation [42]. However, the activation of the hedgehog pathway does not inhibit the early steps of adipogenesis connected with the

commitment and proliferation of precursors [23,42]. Nevertheless, it impairs the differentiation of preadipocytes [15].

Interactions between dWAT and HF s can be provided by different mechanisms, among them via an extracellular exchange of microRNA (miR) and some signaling proteins. One of the most important miRs involved in HF cycling is miR-214, demonstrating an inhibitory effect on β -catenin and thus modulating Wnt/ β -catenin signaling [43]. This miR also suppresses the osteogenic differentiation of bone marrow-derived mesenchymal stem cells, thereby effectively promoting adipogenic differentiation in these cells [44]. Additionally, miR-214 plays a dominant role in exosome-mediated signaling between endothelial cells [45].

Transportation of miRs in extracellular vesicles between the source and recipient cells provides a spatially long-ranged intercellular communication and can effectively alter the transcriptome of the recipient cells [46]. Extracellular vesicles can transport different factors, effectively modulating the physiological state of HF s and dWAT. Amongst these factors transported in extracellular vesicles is SFRP1 [47], which is a well-known antagonist of Wnt/ β -catenin signaling [10], and caveolin [48]. Primary vertebrate cells secrete Shh in distinct vesicular forms [49] which can significantly increase the spatial correlation of Shh signaling compared to simple diffusion. Adipose tissue-derived exosomes carry miRs to regulate TGF- β and Wnt/ β -catenin signaling [50,51] and thus can provide reciprocal regulation of HF s and dWAT.

Extracellular vesicles released by immature and mature adipocytes have different proteomic profiles *in vitro* [52] (Fig. 2). Moreover, the concentration of extracellular vesicles released by preadipocytes may be higher than for mature adipocytes [53], even though this may depend on the nutritional state of the system [54]. Adipocyte precursors also produce high concentrations of extracellular vesicles which, however, demonstrate very high interindividual variations [55] and which are regulated by PDGF [56].

It must be mentioned, that very recently effective vesicular exchange of caveolin regulated by systemic metabolic state was demonstrated between endothelial and adipose cells [54]. Such exchange was strong enough to replace the protein levels of caveolin-1 in genetically caveolin-1 deficient adipocytes.

Adipose tissue content specific profiling of adipocyte-derived vesicles may allow distinct interactions between HF s and dWAT during different stages of the HF cycle, however this awaits further elucidation in *in vivo* models.

Selective modification of immature and mature adipocytes by light irradiation

The absorption coefficients of light waves in human subcutaneous adipose tissue (sWAT) demonstrate a strong dispersion and were reported to be 2.3 cm^{-1} at a wavelength of 400 nm and of $1.0\text{--}1.2 \text{ cm}^{-1}$ at a wavelength of 600–1,200 nm [57]; much higher values of 16.0 cm^{-1} at 400 nm were measured in [58]. Absorption coefficients determined with the same method were higher in adipose tissue than in the adjacent skin [59]. Additionally, extinction

coefficients of porcine sWAT are significantly higher in the superficial than in the deep adipose tissue layers [60]. All of this suggests that the dWAT depot should have much higher light absorption than the classical deep sWAT.

Immature and mature adipocytes have substantially different light absorption properties [61]. Absorption spectra obtained with FTIR spectrometry in 3T3-L1 preadipocytes and corresponding mature adipocytes revealed that light absorption in some wavelength intervals was much higher in preadipocytes than in the corresponding mature adipocytes. We can infer that this effect may lead to a preferential reaction to the light of immature cells compared to mature adipocytes, at least in some intervals of the light spectrum.

Generally, the response of adipocyte precursors to light irradiation is biphasic: whereas low light intensity demonstrates stimulatory effects on the proliferation of these cells [62], high intensity normally produces inhibitory effects. Such inhibition was observed, for example, at fluences of 10-15 J/cm² in [63] and over 20 J/cm² in [64]. Visible light can also effectively interact with adipocytes: irradiation of mature 3T3-L1 adipocytes demonstrated that the light with wavelengths of 500-600 nm was able to significantly reduce lipid droplets and increase the efflux of glycerol from these cells, thus effectively reducing their volumes [65].

Taking together, immature and mature adipocytes differentially react to the same light irradiation, and selective absorption of the light energy in these cells should be able to modify their content in dWAT and thus regulate HF cycling.

The relationship between HF pigmentation and content of immature and mature adipocytes in dWAT

Hair pigmentation is strongly dependent on the number and activity of melanocytes present in HFs. Black hair (containing eumelanin) demonstrates the highest light sensitivity, followed by blond, red (containing more pheomelanin) and gray hairs (containing less melanin). In accordance with the theory of selective photothermolysis, this effect was connected with a reduced concentration of light sensitive eumelanin in red and gray HFs. At the same time, when the light sensitivity of HFs depends on the content and physiological state of adjacent dWAT, distinctly pigmented HFs should have different dWAT structures, specifically, different ratios of immature-to-mature adipocytes.

The biosynthetic pathways for eumelanin and pheomelanin are mainly determined by the signaling activity of melanocortin receptor 1 (MC1R) and Wnt/ β -catenin signaling [66]. Two different MC1R ligands demonstrate opposing effects on this receptor: α -melanocyte stimulating hormone (α -MSH) is an agonist, while agouti signaling protein (ASIP in human) is an antagonist. Activation of MC1R stimulates production of eumelanin, whereas inactivation of this receptor leads to a reduction of eumelanin synthesis and to a corresponding increase of pheomelanin content in HFs [67]. On the other hand, the MC1R/ α -MSH signaling pathway was found to be strongly activated in cutaneous wound repair as well as in hypertrophic scarring [68], demonstrating that this pathway has other functions in the skin in addition to pigment production.

As discussed earlier, both cutaneous wound repair and formation of hypertrophic scars are connected with activity of dWAT [4]. From this point of view, it is not surprising that a genetic link involving α -MSH/ASIP signaling was recently found between adipose tissue and pigmentation [34]. Human adipose tissue indeed demonstrates high expression levels of ASIP [69] and MC1R [70]. Expression of MC1R is much higher in mature adipocytes than in preadipocytes [70], and stimulation of MC1R in adipose precursors inhibits their proliferation [71]. Additionally, precursors can directly affect the behavior of melanocytes, inhibiting their proliferation, differentiation and even melanogenesis. This effect was demonstrated in an engineered skin-equivalent model [72] and it was connected with high expression of TGF- β 1 which maintained melanocytes in an immature state and thus prohibited melanin synthesis.

On the other hand, TGF- β signaling is tightly connected with plasma membrane structures known as caveolae, which are characteristic Ω -shaped plasma membrane invaginations. Caveolae contain different types of caveolins, one of which, Cav-1, is highly expressed in mechanically stressed cells, including in mature adipocytes. Cav-1 physically interacts with TGF- β membrane receptors [73] and participates in their internalization [74]. Internalized TGF- β receptors undergo rapid degradation, thereby leading to an effective reduction of TGF- β signaling. At the same time, Cav-1 expression was reported to be very low in adipose precursors, gradually increasing in preadipocytes and reaching high values in mature adipocytes [75,76]. This explains how the relative content of adipocyte precursors, preadipocytes and mature adipocytes can influence TGF- β expression, and thus modulate the local activity of melanocytes. In this context, it should be once more noted that both Wnt/ β -catenin signaling and TGF- β can be spatially regulated through exosomal exchange [50].

MC1R expression in HFs and dWAT should demonstrate a long-range spatiotemporal correlation which is highly likely regulated by extracellular vesicles. We can infer that a long-lasting reduction of MC1R activity in HFs leading to modulation of their pigmentation must correspond to a modified activity of this receptor and/or of its ligands in adjacent dWAT. Taking together, HF pigmentation correlates with the structure of adjacent dWAT, especially with its content of immature and mature adipocytes. Variation of cellular content in dWAT adjacent to diversely pigmented HFs should cause a distinct response of these dWAT structures to the same light irradiation.

Possible applications for the reduction of unwanted hair growth

Removal of unwanted hair remains one of the most important fields in aesthetic medicine and dermatology. Photoepilation, currently the leading method in hair removal, is based on the widely accepted theory of selective photothermolysis originally formulated for the light absorption in micro-vessels and melanosomes [77,78]. Extension of this theory to the light absorption in HFs restricted the effect of the light to its absorption in melanin. Melanin content is variable in different stages of HF cycling, being generally higher in anagen and correspondingly lower in catagen and telogen phases. According to the theory of selective photothermolysis, light sensitivity of a given HF should be higher in anagen than in other phases of HF cycle.

These observations were widely integrated into clinical applications of photoepilation, but led to some contradictions between the theory and experimental/clinical results which were discussed in detail in [79,80]. The most serious of them is the well-known reduced photosensitivity of HFs in hirsutism. Hirsutism is generally connected with an elongation of the late anagen. Thus, according to the theory of selective photothermolysis, the hirsute HFs should theoretically demonstrate higher photosensitivity than normal HFs.

As discussed in [79], this contradiction can be resolved if the HFs are much less photosensitive in the sub-phase anagen VI than in early- and mid-anagen sub-phases. During anagen VI, only the hair shaft growth, but not the downgrowth of HFs takes place. This is typically found not only in rodents [81], but also in humans. A direct comparison between the stages of hair growth is shown in Table 2. Additionally, the processes of mesenchymal remodeling at the distal end of HF are known to be strongly reduced or even completely suppressed in anagen VI. This leads to the conclusion that not photothermolysis in HFs *per se*, but the modulation of the mesenchymal remodeling at the distal ends of down-growing HFs is mainly responsible for the aberration or even the abortion of the HF cycle after sufficiently intensive light exposure [79,80]. In line with that, focusing microwave energy to the interface of the dermis/subcutis can produce an axillary hair reduction of up to 75% one year after irradiation. This effect is independent of hair color [82].

Interactions between HFs and dWAT depend on the content of immature and mature adipocytes. The selective character of light absorption in these cells allows for an effective modulation of the ratio of immature-to-mature adipocytes in dWAT. Therefore, we can assume that interference or even the complete abortion of the HF cycle, which can often be observed after light irradiation, is at least partly connected with absorption of the light energy not in HFs, but in dWAT (Fig. 3). At the same time, interference in the HF cycle after light irradiation is known to be strongly dependent on hair pigmentation. Since dWAT is involved in this process, the relative content of immature and mature adipocytes in dWAT adjacent to differently pigmented hairs should be different.

Conclusions

HFs and dWAT strongly interact with each other, demonstrating a highly coordinated behavior. This interaction is substantially dependent on the content of immature and mature adipocytes in dWAT adjacent to HFs. Such content should be regulated not only by processes of proliferation and differentiation of adipose precursors and preadipocytes, but also include the processes of dedifferentiation of mature adipocytes. Thus, this involves mechanisms and factors that promote such dedifferentiation, among others, TGF- β signaling.

Interactions between HFs and dWAT can be provided through the exchange of extracellular vesicles. Since vesicles released by precursors, preadipocytes and mature adipocytes probably have different proteomic profiles and are released in different concentrations, such mechanism can provide a fine regulation of the HF-dWAT interaction. However, this is mostly based on *in vitro* observations and awaits *in vivo* validation. dWAT exposure to different exogenic factors could lead to a quick direct modification of exosomal release in

this tissue or modify this release on a long-lasting scale through modulation of the ratio of immature-to-mature adipocytes. Both types of modification can lead to consequent changes in the HF cycle.

One such factor demonstrating selective effects on immature and mature adipocytes is the light irradiation in some specific wavelength intervals. Adipose precursors, preadipocytes and mature adipocytes distinctly react to the same light irradiation, demonstrating significantly higher light absorption in immature adipocytes. Since the tissue content of precursors, preadipocytes and mature adipocytes significantly varies over the course of the HF cycle (and even during different sub-phases of anagen), this effect leads to variable photosensitivity of HFs in different phases of the HF cycle. Altogether, this means that more attention should be paid to the dWAT as a potential target in modulation of hair growth.

Acknowledgments

ILK is the managing partner of Wellcomet GmbH. No methods or devices of Wellcomet were used in this publication. ILK received no financial support from Wellcomet GmbH. Wellcomet GmbH provided support in the form of salaries for ILK, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of ILK are articulated in the 'author contributions' section. P.E.S. was supported by US National Institutes of Health (NIH) grants R01-DK55758, R01-DK099110, P01-DK088761 and P01-AG051459 (PES; Project 3) as well as by an unrestricted grant from the Novo Nordisk Research Foundation.

References

1. Festa E et al. (2011) Adipocyte lineage cells contribute to the skin stem cell niche to drive hair cycling. *Cell* 146, 761–771. [PubMed: 21884937]
2. Schmidt B and Horsley V (2012) Unravelling hair follicle–adipocyte communication. *Exp. Dermatol* 21, 827–830. [PubMed: 23163647]
3. Kruglikov IL and Scherer PE (2016) Dermal adipocytes and hair cycling: is spatial heterogeneity a characteristic feature of the dermal adipose tissue depot? *Exp. Dermatol* 25, 258–262. [PubMed: 26781768]
4. Kruglikov IL & Scherer PE (2016) Dermal adipocytes: from irrelevance to metabolic targets? *Trends Endocrinol. Metab* 27, 1–10.
5. Guerrero-Juarez CF and Plikus MV (2018) Emerging nonmetabolic functions of skin fat. *Nat. Rev. Endocrinol* 14, 163–173. [PubMed: 29327704]
6. Leiva AG et al. (2014) Loss of Mpl3 function causes various skin abnormalities and greatly reduced adipose depots. *J. Invest. Dermatol* 134, 1817–1827. [PubMed: 24531688]
7. Hesslein DG et al. (2009) Ebf1-dependent control of the osteoblast and adipocyte lineages. *Bone* 44, 537–546. [PubMed: 19130908]
8. Tomita Y et al. (2006) PDGF isoforms induce and maintain anagen phase of murine hair follicles. *J. Dermatol. Sci* 43, 105–115. [PubMed: 16725313]
9. Hawkshaw NJ et al. (2015) Re-evaluating cyclosporine A as a hair growth-promoting agent in human scalp hair follicles. *J. Invest. Dermatol* 135, 2129–2132. [PubMed: 25826423]
10. Hawkshaw NJ et al. (2018) Identifying novel strategies for treating human hair loss disorders: Cyclosporine A suppresses the Wnt inhibitor, SFRP1, in the dermal papilla of human scalp hair follicles. *PLOS Biol.* 16, e2003705. [PubMed: 29738529]
11. Ehlund A et al. (2013) Characterization of the Wnt inhibitors secreted frizzled-related proteins (SFRPs) in human adipose tissue. *J. Clin. Endocrinol. Metab* 98, E503–E508. [PubMed: 23393180]
12. Lagathu C et al. (2010) Secreted frizzled-related protein 1 regulates adipose tissue expansion and is dysregulated in severe obesity. *Int. J. Obes* 34, 1695–1705.

13. Qu Y et al. (2018) Cyclosporin A inhibits adipogenic differentiation and regulates immunomodulatory functions of murine mesenchymal stem cells. *Biochem. Biophys. Res. Commun* 498, 516–522. [PubMed: 29510137]
14. Donati G et al. (2014) Epidermal Wnt/ β -catenin signaling regulates adipocyte differentiation via secretion of adipogenic factors. *Proc. Nat. Acad. Sci* 11, E1501–E1509.
15. Tang QQ and Lane MD (2012) Adipogenesis: from stem cell to adipocyte. *Annu. Rev. Biochem* 81, 715–736. [PubMed: 22463691]
16. Shook B et al. (2016) The role of adipocytes in tissue regeneration and stem cell niches. *Annu. Rev. Cell. Develop. Biol* 32, 609–631.
17. Won CH et al. (2010) Hair growth promoting effects of adipose tissue-derived stem cells. *J. Dermatol. Sci* 57, 134–137. [PubMed: 19963355]
18. Jin SE and Sung JH (2016) Hair regeneration using adipose-derived stem cells. *Histol. Histopathol* 31, 249–256. [PubMed: 26536569]
19. Plikus MV et al. (2008) Cyclic dermal BMP signalling regulates stem cell activation during hair regeneration. *Nature* 451, 340–344. [PubMed: 18202659]
20. Gustafson B et al. (2013) Restricted adipogenesis in hypertrophic obesity: the role of WISP2, WNT, and BMP4. *Diabetes* 62, 2997–3004. [PubMed: 23970518]
21. Plikus MV et al. (2017) Regeneration of fat cells from myofibroblasts during wound healing. *Science* 355, 748–752. [PubMed: 28059714]
22. Zhang P et al. (2014) A review of adipocyte lineage cells and dermal papilla cells in hair follicle regeneration. *J. Tissue Eng* 5, 1–10.
23. Zhang B et al. (2016) Hair follicles' transit-amplifying cells govern concurrent dermal adipocyte production through Sonic Hedgehog. *Genes Dev.* 30, 2325–2338. [PubMed: 27807033]
24. Rivera-Gonzalez GC et al. (2016). Skin adipocyte stem cell self-renewal is regulated by a PDGFA/AKT-signaling axis. *Cell Stem Cell* 19, 738–751. [PubMed: 27746098]
25. Foster A et al. (2018) Dermal white adipose tissue undergoes major morphological changes during the spontaneous and induced murine hair follicle cycling: A reappraisal. *Arch. Dermatol. Res* 310, 453–462. [PubMed: 29704126]
26. Matsumoto T et al. (2008) Mature adipocyte- derived dedifferentiated fat cells exhibit multilineage potential. *J. Cell Physiol* 215, 210–222. [PubMed: 18064604]
27. Côté JA et al. (2017) Role of the TGF- β pathway in dedifferentiation of human mature adipocytes. *FEBS Open Bio.* 7, 1092–1101.
28. Wie S et al. (2013) Cellular and molecular implications of mature adipocyte dedifferentiation. *J. Genomics* 1, 5–12. [PubMed: 25031650]
29. Ohgo S et al. (2013) Bleomycin inhibits adipogenesis and accelerates fibrosis in the subcutaneous adipose layer through TGF- β 1. *Exp. Dermatol* 22, 769–771. [PubMed: 24118261]
30. Marangoni RG et al. (2015) Myofibroblasts in murine cutaneous fibrosis originate from adiponectin- positive intradermal progenitors. *Arth. Rheumatol* 67, 1062–1073. [PubMed: 25504959]
31. Kruglikov IL and Scherer PE (2016) Skin aging: are adipocytes the next target? *Aging (Albany NY)* 8, 1457. [PubMed: 27434510]
32. Deng T et al. (2016) Obesity, inflammation, and cancer. *Annu. Rev. Pathol. Mech. Disease* 11, 421–449.
33. Kruglikov IL and Scherer PE (2017) Adipocyte- myofibroblast transition as a possible pathophysiological step in androgenetic alopecia. *Exp. Dermatol* 26, 522–523. [PubMed: 28482134]
34. Clement E et al. (2017) Obesity and melanoma: could fat be fueling malignancy? *Pigment Cell. Melan. Res* 30, 294–306.
35. Martins V et al. (2015) FIZZ1-induced myofibroblast transdifferentiation from adipocytes and its potential role in dermal fibrosis and lipoatrophy. *Am. J. Pathol* 185, 2768–2776. [PubMed: 26261086]
36. Liu T et al. (2015) A role for FIZZ1 in a model of scleroderma. *FASEB J* 29 (Suppl 1), 927–10.

37. Liu T et al. (2014) The in vivo fibrotic role of FIZZ1 in pulmonary fibrosis. *PloS One* 9, e88362. [PubMed: 24516640]
38. Lamouille S et al. (2014) Molecular mechanisms of epithelial–mesenchymal transition. *Nat. Rev. Mol. Cell Biol* 15, 178–196. [PubMed: 24556840]
39. Ubil E et al. (2014) Mesenchymal–endothelial transition contributes to cardiac neovascularization. *Nature* 514, 585–590. [PubMed: 25317562]
40. Rangel-Huerta E and Maldonado E (2017) Transit-amplifying cells in the fast lane from stem cells towards differentiation. *Stem Cell Int.* 2017, 7602951.
41. Hsu YC et al. (2014) Transit-amplifying cells orchestrate stem cell activity and tissue regeneration. *Cell* 157, 935–949. [PubMed: 24813615]
42. Fontaine C et al. (2008) Hedgehog signaling alters adipocyte maturation of human mesenchymal stem cells. *Stem Cells* 26, 1037–1046. [PubMed: 18258719]
43. Ahmed MI et al. (2014) MicroRNA-214 controls skin and hair follicle development by modulating the activity of the Wnt pathway. *J. Cell Biol* 207, 549–567. [PubMed: 25422376]
44. Guo Y et al. (2017) miR-214 suppresses the osteogenic differentiation of bone marrow-derived mesenchymal stem cells and these effects are mediated through the inhibition of the JNK and p38 pathways. *Int. J. Mol. Med* 39, 71–80. [PubMed: 27959394]
45. Van Balkom BW et al. (2013) Endothelial cells require miR-214 to secrete exosomes that suppress senescence and induce angiogenesis in human and mouse endothelial cells. *Blood* 121, 3997–4006. [PubMed: 23532734]
46. Ibrahim A and Marbán E (2016) Exosomes: fundamental biology and roles in cardiovascular physiology. *Annu. Rev. Physiol* 78, 67–83. [PubMed: 26667071]
47. Rappa G et al. (2013) Biochemical and biological characterization of exosomes containing prominin-1/CD133. *Mol. Cancer* 12, 62. [PubMed: 23767874]
48. Das M and Das DK (2012) Caveolae, caveolin, and cavins: potential targets for the treatment of cardiac disease. *Annal. Med* 44, 530–541.
49. Vyas N et al. (2014) Vertebrate Hedgehog is secreted on two types of extracellular vesicles with different signaling properties. *Sci. Rep* 4, 7357. [PubMed: 25483805]
50. Ferrante SC et al. (2015) Adipocyte-derived exosomal miRNAs: a novel mechanism for obesity-related disease. *Pediatr. Res.* 77, 447–454. [PubMed: 25518011]
51. Zhang Y et al. (2017) miR-450a-5p within rat adipose tissue exosome-like vesicles promotes adipogenic differentiation by targeting WISP2. *J. Cell Sci* 130, 1158–1168. [PubMed: 28167681]
52. Gao X et al. (2017) Extracellular vesicles from adipose tissue - A potential role in obesity and type 2 diabetes? *Front. Endocrinol* 8, 202.
53. Connolly KD et al. (2015) Characterisation of adipocyte-derived extracellular vesicles released pre-and post-adipogenesis. *J. Extracell. Vesicles* 4, 29159. [PubMed: 26609807]
54. Crewe C et al. (2018) An endothelial to adipocyte extracellular vesicle axis governed by metabolic state. *Cell* 175, 695–708.e13. [PubMed: 30293865]
55. Kalinina N et al. (2015) Characterization of secretomes provides evidence for adipose-derived mesenchymal stromal cells subtypes. *Stem Cell Res. Ther* 6, 221. [PubMed: 26560317]
56. Lopatina T et al. (2014) Platelet-derived growth factor regulates the secretion of extracellular vesicles by adipose mesenchymal stem cells and enhances their angiogenic potential. *Cell Commun. Signal* 12, 26. [PubMed: 24725987]
57. Bashkatov AN et al. (2005) Optical properties of human skin, subcutaneous and mucous tissues in the wavelength range from 400 to 2000 nm. *J. Phys. D Appl. Phys* 38, 2543–2555.
58. Salomatina EV et al. (2006) Optical properties of normal and cancerous human skin in the visible and near-infrared spectral range. *J. Biomed. Opt* 11, 064026. [PubMed: 17212549]
59. Bashkatov AN et al. (2011) Optical properties of skin, subcutaneous, and muscle tissues: a review. *J. Innov. Opt. Health Sci* 4, 9–38.
60. Filatova SA et al. (2017) Optical properties of animal tissues in the wavelength range from 350 to 2600 nm. *J. Biomed. Optics* 22, 035009.

61. Masaki N and Okazaki S (2018) Selective delivery of laser energy to ester bonds of triacylglycerol in lipid droplets of adipocyte using a quantum cascade laser. *Biomed. Optic. Express* 9, 2095–2103.
62. Mvula B et al. (2008) The effect of low level laser irradiation on adult human adipose derived stem cells. *Laser Med. Sci* 23, 277–282.
63. Abrahamse H et al. (2010) Fluence and wavelength of low intensity laser irradiation affect activity and proliferation of human adipose derived stem cells: peer reviewed original article. *Med. Technol. SA* 24, 15–20.
64. Wang Y et al. (2017) Photobiomodulation of human adipose-derived stem cells using 810 nm and 980 nm lasers operates via different mechanisms of action. *Biochim. Biophys. Acta* 1861, 441–449.
65. Park PJ et al. (2017) Specific visible radiation facilitates lipolysis in mature 3T3-L1 adipocytes via rhodopsin-dependent β 3-adrenergic signaling. *Eur. J. Cell Biol* 96, 301–311. [PubMed: 28483278]
66. Lin JY and Fisher DE (2007) Melanocyte biology and skin pigmentation. *Nature* 445, 843–850. [PubMed: 17314970]
67. Nasti TH and Timares L (2015) MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem. Photobiol* 91, 188–200. [PubMed: 25155575]
68. Muffley LA et al. (2011) Spatial and temporal localization of the melanocortin 1 receptor and its ligand α -melanocyte-stimulating hormone during cutaneous wound repair. *J. Histochem. Cytochem* 59, 278–288. [PubMed: 21378282]
69. Voisey J and van Daal A (2002) Agouti: from mouse to man, from skin to fat. *Pigment Cell Melan. Res* 15, 10–18.
70. Hoch M et al. (2007) Expression and localization of melanocortin-1 receptor in human adipose tissues of severely obese patients. *Obesity* 15, 40–49. [PubMed: 17228030]
71. Hoch M et al. (2008) Weak functional coupling of the melanocortin-1 receptor expressed in human adipocytes. *J. Recept. Signal Trans* 28, 485–504.
72. Klar AS et al. (2017) Human adipose mesenchymal cells inhibit melanocyte differentiation and the pigmentation of human skin via increased expression of TGF- β 1. *J. Invest. Dermatol* 137, 2560–2569. [PubMed: 28774590]
73. Ito T et al. (2004) Hyaluronan regulates transforming growth factor- β 1 receptor compartmentalization. *J Biol Chem* 279, 25326–25332. [PubMed: 15084590]
74. Galdo FD et al. (2008) Decreased expression of caveolin 1 in patients with systemic sclerosis: crucial role in the pathogenesis of tissue fibrosis. *Arth. Rheumatol* 58, 2854–2865.
75. Scherer PE et al. (1994) Induction of caveolin during adipogenesis and association of GLUT4 with caveolin-rich vesicles. *J. Cell Biol* 127, 1233–1243. [PubMed: 7962086]
76. Fernández-Real JM et al. (2010) Study of caveolin-1 gene expression in whole adipose tissue and its subfractions and during differentiation of human adipocytes. *Nutr. Metab* 7, 20.
77. Anderson RR and Parrish JA (1983) Selective photothermolysis: precise microsurgery by selective absorption of pulsed radiation. *Science* 220, 524–527. [PubMed: 6836297]
78. Altshuler GB et al. (2001) Extended theory of selective photothermolysis. *Laser Surg. Med* 29, 416–432.
79. Kruglikov IL (2012) Melanin light absorption as the necessary but not sufficient condition for photoepilation: Intra-anagen variability of hair follicle light sensitivity. *Am. J. Cos. Surg* 29, 266–272.
80. Kruglikov IL (2013) Melanin light absorption as the necessary but not sufficient condition for photoepilation: miniaturization and eclipse phenomena. *Am. J. Cos. Surg* 30, 21–27.
81. Müller-Röver S et al. (2001) A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J. Invest. Dermatol* 117, 3–15. [PubMed: 11442744]
82. Brauer JA et al. (2017) A prospective study of axillary hair reduction in patients treated with microwave technology. *Dermatol. Surg* 43, 558–565. [PubMed: 28005625]

Box 1 : Hair Follicle Development

Hair follicles (HF) are skin appendages demonstrating remarkable level of cyclical development. HF cycling is traditionally subdivided in three phases: **anagen** (growth), **catagen** (regression), and **telogen** (rest). Morphologic and physiologic features of HFs significantly vary not only between the phases but also within a specific phase. As a result, anagen is further subdivided into 6 sub-phases, the first two of which (“early anagen”) are connected with modifications of the hair follicle structure, the sub-phases III – V (“mid-anagen”) – with downgrowth and full development of the hair follicle, and the sub-phase VI (“late anagen”) – with the development of a hair shaft.

Morphologic and physiologic variations of hair follicles demonstrate a strong spatiotemporal correlation with behavior of adjacent superficial adipose tissue known as dermal white adipose tissue. This suggests a long-range interaction between hair follicles and dermal adipocytes. Such correlations are observed not only as a function of varying thickness of the dermal adipose layer, but also at the level of modulation of the relative ratios of immature and mature dermal adipocytes during HF cycling.

Outstanding Questions

Are there phenotypical differences between adipocyte-derived preadipocytes and “virgin” preadipocytes that had not reached the adipocyte stage previously?

What is the difference between adipocyte-derived preadipocytes in dermal adipose tissue vs. other fat pads?

Do the adipocyte-derived preadipocytes play a role in obesity, weight loss and diabetes?

Is the number of “back and forth” transitions between immature and mature adipocytes in dWAT limited?

How does the pigmentation of hair follicles correlate with the structure of adjacent dermal adipose tissue?

Is there any difference in light sensitivity between cycling adipocytes and “virgin” adipocytes?

What’s the difference between the normal skin fibroblasts and dermal adipocyte precursor cells?

What is the natural turnover rate of dermal adipocytes in adult mice?

Are mature dermal adipocytes necessary for hair cycling?

Highlights

- Hair follicles strongly interact with adipocytes within the dermal white adipose tissue, suggesting a strong physiological dependence on the content of immature and mature adipocytes in this layer.
- This content is regulated by processes of proliferation and differentiation of adipose precursors and preadipocytes, as well as by dedifferentiation of mature existing adipocytes leading to production of adipocyte-derived preadipocytes.
- Long-range spatial interactions between hair follicles and dermal adipocytes involves the exchange of extracellular vesicles which are differentially released by precursors, preadipocytes and mature adipocytes and are likely to carry various proteomic fingerprints during different phases of the hair follicle cycle, effectively modulating the transcriptome profile of the recipient hair follicles.
- Different exogenous factors, including light irradiation and some chemical agents, can affect the immature and immature adipocytes and are likely enabled to modify the release of adipocyte-derived exosomes in dWAT, which can lead to aberrations of the HF cycle.
- Dermal adipocytes should be considered as a potential target both for the modulation of hair growth and for the removal of unwanted hair.

Fig. 1.

Precursors, preadipocytes and mature adipocytes can undergo different conversions. Mature adipocytes can dedifferentiate, producing adipocyte-derived preadipocytes (ADP). This mechanism can contribute to the modification of dWAT volume during the HF cycle.

Fig. 2.

Local content of dWAT results in a specific “fingerprint” of exosomes with specific content in this area. Precursors and preadipocytes may produce higher levels of extracellular vesicles than mature adipocytes. Concentration and loading of exosomes strongly vary according to the phase of HF cycle

Fig. 3. Different content of precursors, preadipocytes and mature adipocytes in dWAT in distinct phases of the HF cycle as well as the differential sensitivity of these cells to the same light irradiation is at least partly responsible for different light sensitivity of HFs in the corresponding phases of the HF cycle.

Table 1

Differences amongst skin adipose-derived stem cells, skin preadipocytes, and mature dermal adipocytes.

	Skin Adipocyte-Derived Stem Cells (ADSCs)	Skin Pre-Adipocytes	Mature dermal adipocytes
Location	Lower dermis	Lower dermis	<i>Rodents:</i> under reticular dermis and above the panniculus carnosus. <i>Human:</i> under the reticular dermis, superficial to subcutaneous adipose tissue
Morphology	Fibroblast-like	Fibroblast-like	Unilocular with large lipid droplets
Markers	Lin ⁻ , Sca1 ⁺ , CD34 ⁺ , CD29 ⁺ , CD24 ⁺ , Pdgfra ⁺	Lin ⁻ , Sca1 ⁺ , CD34 ⁺ , CD29 ⁺ , Pdgfra ⁺ , CD24 ⁻ , CEBPa ⁺	Common adipocyte markers

Table 2

Comparison of hair growth stages between rodents and humans.

	Mouse	Human
Main Cycle Stage	Anagen Catagen Telogen	Anagen Catagen Telogen
Length of Anagen	1-3 weeks	Can last for several years
Length of Catagen	About 2-4 days	Around 2-3 weeks
Length of Telogen	First cycle, 1-2 days, from second cycle: more than 2 weeks	A few months
Synchronization	Highly synchronize for the first two cycles. Only small patches of synchronous hair growth after the second cycle	Synchronized fetal hair follicle cycling becomes asynchronous soon after birth. Mosaic cycling pattern.