Published in final edited form as:

Clin Psychol Sci. 2018 January ; 6(1): 123–133. doi:10.1177/2167702617730889.

Callous-Unemotional Traits Moderate Genetic and Environmental Influences on Rule-Breaking and Aggression: Evidence for Gene $\times$ Trait Interaction

Frank D. Mann¹, Jennifer L. Tackett², Elliot M. Tucker-Drob^{1,3}, and K. Paige Harden^{1,3}

¹Department of Psychology, University of Texas at Austin

²Department of Psychology, Northwestern University

³Population Research Center, University of Texas at Austin

Abstract

Previous behavioral genetic research in children has found that conduct problems in the presence of high CU traits are more heritable than conduct problems in the presence of low CU traits -- a gene $\times$ trait interaction. The current study replicates and extends this finding using a sample of adolescent twins from the Texas Twin Project, who were assessed for rule-breaking and aggression. We find evidence that genetic influences on CU traits contribute to genetic liability for both rule-breaking and aggressive behavior. CU traits moderate genetic influences on aggressive behavior, such that the heritability of aggression is higher among youth with high levels of CU traits. However, we do not find evidence that CU traits moderate genetic influences on rule-breaking behavior. The continuum of callous-unemotionality and the aggression versus rule-breaking distinction continues to be meaningful and intersecting methods for characterizing heterogeneity in the etiology of antisocial behavior.

Keywords

callous-unemotional; rule-breaking; aggression; antisocial behavior; gene $\times$ trait interaction; behavior genetics

Callous-unemotional (CU) traits are a cluster of affective and interpersonal risk factors for antisocial behavior that include shallow affect, lack of remorse and a cruel disregard for others. An established corpus of studies indicates that antisocial behavior is unique for youth with high levels of CU traits, including distinct neurological and physiological profiles (reviewed by Frick, Ray, Thorton & Kahn, 2014; Viding, Fontaine & McCrory, 2012). To this end, “callousness” was included in the DSM-5 as a personality disorder trait facet, as well as a specifier for Conduct Disorder (American Psychological Association, 2013; Krueger & Markon, 2014).

A number of quantitative genetic studies have examined the etiology of CU traits and antisocial behavior. Twin and adoption studies have found evidence that CU traits are

moderately to highly heritable and index genetic liability for antisocial behavior (Hyde et al., 2016; Viding, Frick & Plomin, 2007; Waldman et al., 2011). That is, the same genes that influence CU traits also contribute to genetic risk for antisocial behavior. For example, Hyde et al. (2016) found that CU traits in adopted children (27-months-old) were significantly correlated with their biological mothers' antisocial behavior, which provides evidence of heritable risk for CU traits in early childhood. In a sample of 7- to 9-year-old twins, Viding, Frick & Plomin (2007) found that the genetic correlation between CU traits and conduct problems was statistically significant and moderate in magnitude for both boys and girls (.57 & .65, respectively). In a sample of child and adolescent twins ($N \sim 4,000$), Waldman et al. (2011) found that additive genetic influences on low levels of prosociality (similar to high levels of callousness) accounted for 20% of the variance in symptoms of conduct disorder. Moreover, additional twin studies of youth and young adults have found evidence of substantial genetic overlap between CU traits and antisocial behavior (Taylor et al, 2003; Larsson et al., 2006). Taken together, these results indicate that the phenotypic association between CU traits and antisocial behavior is driven partly by genetic factors shared by both phenotypes.

In addition to indexing genetic liability for antisocial behavior, there is also evidence that the presence of high CU traits differentiates a highly heritable form of antisocial behavior in childhood. Specifically, Viding, Blair, Moffitt & Plomin (2005) found that antisocial behavior was more highly heritable among 7-year-old children with high levels of CU traits, compared to children with average or low levels of CU traits (i.e. a gene $\times$ trait interaction; DeYoung, 2012). This gene $\times$ CU trait interaction was replicated by the same research group in a sample of 9-year-old children (Viding, Jones, Frick, Moffitt & Plomin, 2008), although the magnitude of the effect was less pronounced. Nonetheless, results suggest that antisocial behavior is more highly heritable among children who have high levels of CU traits, compared to children with low levels of CU traits.

It is currently unclear whether the moderating effect of CU traits on genetic risk for antisocial behavior extends to different subtypes of antisocial behavior or to antisocial behavior measured in developmental periods beyond childhood. One useful method for parsing heterogeneity in antisocial behavior centers upon the distinction between aggressive behavior (e.g., acts that harm or victimize others) and non-aggressive rule-breaking (e.g., theft and truancy) (Burt, 2009; Harden et al., 2015). Previous studies of whether CU traits moderate the heritability of antisocial behavior have used measurements that collapse across the aggression vs. rule-breaking distinction. For example, Viding et al. (2008) used a 5-item measure of antisocial behavior that included both aggression items ("I fight a lot") and rule-breaking items ("I am often accused of lying and cheating"). Similarly, the 5-item measure of antisocial behavior in Viding et al. (2005) included both "often fights with other children or bullies them" and "often lies and cheats." Therefore, it is unclear from previous studies whether the observed moderating effect of CU traits is specific to aggression or rule-breaking, or general across these different types of antisocial behavior.

Indeed, there are at least three reasons to believe that a gene $\times$ CU trait interaction may differ across development, as well as between aggressive versus non-aggressive subtypes of antisocial behavior. First, the rank-order stability and mean prevalence rates of aggression

and rule-breaking behaviors vary across development. As is the case for CU traits, rank-order stability of aggression is high beginning in early childhood, (Burt, 2012; Frick & White, 2008). In contrast, rank order stability of rule-breaking is low in childhood and increases in adolescence. Moreover, mean levels of aggression peak in early childhood, whereas mean levels of rule-breaking increase in adolescence. Second, aggression tends to be more heritable (~ 65%) than rule-breaking (~ 48%), and is subject to little or no shared environmental influences, particularly after childhood (Burt, 2009). Conversely, rule-breaking is more susceptible to shared environmental influences (~ 18%; Burt, 2009). This has been interpreted as reflecting a potentially more normative trajectory of rule-breaking behaviors compared to physical aggression, an interpretation also supported by differentiated personality correlates between these subtypes (Tackett, Daoud, De Bolle & Burt 2013). Third, the etiology of aggression and rule-breaking differs across development. The heritability of aggression remains stable from childhood to adolescence, whereas the heritability of rule-breaking increases with adolescent age and pubertal development (Burt & Klump, 2009; Burt & Neiderhiser, 2009; Harden et al., 2015). Given that aggression and rule-breaking have distinct etiologies and show differential patterns of phenotypic expression across development, it is important to establish whether the gene $\times$ CU trait interaction previously observed in children will generalize across subtypes and developmental periods.

Goals of the Current Study

The current study uses a sample of adolescent twins to test the hypothesis that antisocial behavior is highly heritable in the presence of high levels of CU traits, as has been previously found in childhood. One limitation of previous research is suboptimal measurement of the CU construct, defined by the use of questionnaires with few items, binary response options and poor internal consistencies (Waller, Gardner & Hyde, 2013). In the current study we measure CU traits in a population-based sample of adolescent twins using a well-validated, 24-item questionnaire. Furthermore, to help further delineate heterogeneity in the association between CU traits and antisocial behavior, we measure rule-breaking and aggression separately using a 25-item questionnaire.

Method

Sample

Texas Twin Sample.—The current sample consists of ($N = 824$) twins and multiples from ($N = 407$) families registered in the Texas Twin Project (Harden, Tucker-Drob, & Tackett, 2013), a school-aged twin registry that is representative of the greater Austin and Houston metropolitan areas. Twins and multiples in grades 9–12 were recruited using public school rosters and invited to participate in an on-going in-lab study, which involves a large battery of cognitive assessments, surveys and behavioral tasks. Trained research assistants tested all siblings separately. Surveys were computerized and administered to participants privately in a quiet room. All participants were either currently enrolled in grade school or had graduated within the past 3 months, but were still living at home with their co-twin.¹

The Texas Twin sample represents a socioeconomically diverse population. Approximately one third of families reported having received a needs-based form of public assistance since the twins were born. Sixty percent of the sample identified as non-Hispanic Caucasian, 25% identified as Hispanic/Latino, 10% identified as African American, and 5% identified as another race or ethnicity. Parental education varied widely with 6% of parents not completing high school, 28% completing high school or a vocational or technical program, 6% completing an associates degree, 36% a bachelors degree and 24% a graduate or professional degree.

Zygosity.—Twins and multiples were designated as monozygotic (MZ) or dizygotic (DZ) based on the results of a latent class analysis (LCA; Muthén, 2004), which assigns members to different subgroups in a population (e.g. MZ or DZ twins) on the basis of self- and informant-reports of physical similarity. Specifically, each twin, parents of twins and two research assistants per twin-pair reported on questionnaire items that assessed the twins physical similarity (e.g. “Do your twins have the same color eyes?” “How similar do the twins look in their facial appearance/hair color?”) and likelihood for being mistaken for each other (e.g. “Do family members ever mistake one twin for the other?”, “Do you have difficulty telling one twin from the other?”). Results of the LCA indicated that approximately 35% and 65% of the sample were MZ and DZ twins, respectively, and had an entropy statistics equal to 0.999, which indicates that there was a high level of certainty in assigning twin-pairs to zygosity groups. Note, previous research that compared zygosity classification obtained by genotyping to LCA of questionnaire data found that the latter had a misclassification rate of less than 1% (Heath et al., 2003).

Measures

Callous-Unemotional (CU) Traits.—CU traits were measured using adolescents’ responses to the Inventory of Callous-Unemotional Traits. This 24-item questionnaire was developed to measure individual differences in callous, careless, uncaring and unemotional dispositions (Kimonis et al., 2008). Items include “I do not care whom I hurt to get what I want”, “I do not care about doing things well”, “I am concerned about the feelings of others (R)” and “I do not show my emotions to others”, which are rated on a four point scale (1 = Disagree Strongly; 4 = Agree Strongly). Consistent with the results of a recent item response theory analysis (Ray, Frick, Thornton, Steinberg & Cauffman, 2015), we do not decompose the ICU into subscale scores and, instead, compute a mean composite score including responses from all items.

Aggression and Rule-Breaking.—Aggression and rule-breaking were measured using twins’ responses to externalizing items from the Youth Self-Report (YSR) version of Child Behavior Checklist (Lizotte, Chard-Wierschem, Loeber, Stern, 1992). Based on previously published factor analytic work (Harden et al., 2015), 13 items were used to measure aggression, and 12 items were used to measure rule-breaking. All items are rated on a 3-point scale (0 = Not True; 2 = Very True or Often True), and twins’ responses to items were

¹ Less than 5% of the sample was over 18 years of age at the time of participation.

summed and then divided by the total number of items to produce mean aggression and rule-breaking scores (see Table 1).

Data Analytic Procedures

Internal consistencies (Cronbach's alpha) and descriptive statistics were calculated using *R* version 3.1.2 (R Core Team, 2013). All inferential analyses were conducted within a structural equation modeling framework using *Mplus* version 7.1 (Muthén & Muthén, 2010). Missing data was handled by estimating models using full-information maximum likelihood with robust standard errors. The CU scale approximated a Gaussian distribution and rule-breaking and aggression scales were log-transformed to correct for positive skew. To control for potential covariates, log-transformed scales and the CU scale were residualized for the main effects of Caucasian, Hispanic, and African American race/ethnicity; age; sex; and age $\times$ sex interaction (McGue & Bouchard, 1984). Finally, residual scale scores were standardized before conducting quantitative genetic analyses. All phenotypic models implemented a sandwich estimator to account for the nonindependence of data that results from siblings being clustered within the same family. The absolute fit of models was evaluated using root mean squared error of approximation (RMSEA) and Tucker-Lewis Index (TLI), where RMSEA values $< .05$ and TLI values $> .90$ indicate good fit to the data. Comparative model fit was evaluated using Akaike Information Criteria (AIC), Bayesian Information criteria (BIC), sample-size adjusted Bayesian Information criteria (SSBIC) and Satorra-Bentler scaled loglikelihood chi-squared difference tests (χ^2).

Inferential analyses were conducted in three steps. First, zero-order and partial phenotypic correlations between focal constructs were estimated. Second, quantitative genetic models were used to estimate latent genetic and environmental correlations between CU traits, rule-breaking, and aggression. Finally, quantitative genetic models were extended to test whether CU traits moderated the latent genetic and environmental components of rule-breaking and aggression.

Results

Phenotypic Analyses.—Descriptive and reliability statistics are reported in Table 1. Zero-order correlations between CU traits and measures of antisocial behavior were similar in magnitude, .30 (with aggression) to .43 (with rule-breaking), and significantly different from zero at $p_{(\text{two-tailed})} < .001$. These correlations are reported below the diagonal in Table 1. Partial correlations controlling for the main effects of race, age, sex and age $\times$ sex interaction are reported above the diagonal. These correlations are slightly attenuated relative to the zero-order correlations, although the pattern of correlations remains unchanged.

Bivariate Quantitative Genetic Analyses.—Based on the results of previous behavior genetic studies of CU traits (Moore et al., 2016), rule-breaking and aggressive behavior (Burt, 2009), variance in CU traits was modeled as a function of latent additive genetic (*A*) and non-shared environmental factors (*E*), and variance in antisocial behavior was modeled as a function of latent additive genetic (*A*), shared environmental (*C*) and non-shared

environmental (E) factors. For each twin, we performed a Cholesky decomposition such that the second phenotype (i.e. antisocial behavior) was regressed on the latent genetic and environmental factors of the first phenotype (i.e. CU traits). This model is depicted in the top panel of Figure 1. (For details regarding the specification of twin models, see Neale and Maes, 2004.)

Parameter estimates from the Cholesky decompositions were used to compute additive genetic and non-shared environmental correlations between focal study constructs. The genetic correlation between CU traits and aggression was moderate in magnitude ($r_A = .38$, $CI_{.95\%} = .14, .62$, $p_{(two-tailed)} = .002$). On the other hand, the genetic correlation between CU traits and rule-breaking was large ($r_A = .80$, $CI_{.95\%} = .27, 1.33$, $p_{(two-tailed)} = .003$), similar in magnitude to the genetic correlation between aggression and rule-breaking ($r_A = .77$, $CI_{.95\%} = .60, .95$, $p_{(two-tailed)} < .001$). Non-shared environmental correlations between CU traits, aggression and rule-breaking were comparatively small in magnitude. Nonetheless, non-shared environmental correlations between CU traits and aggression ($r_E = .23$, $CI_{.95\%} = .08, .38$, $p_{(two-tailed)} = .002$), CU traits and rule-breaking ($r_E = .30$, $CI_{.95\%} = .15, .45$, $p_{(two-tailed)} < .001$) and, aggression and rule-breaking ($r_E = .30$, $CI_{.95\%} = .16, .43$, $p_{(two-tailed)} < .001$) were all significantly different than zero.

To test the hypothesis that antisocial behavior is more highly heritable in the presence of high levels of CU traits, compared to low levels of CU traits (i.e. a gene $\times$ trait interaction), bivariate twin models were specified to include a series of parameter constraints (see bottom panel of Figure 1). Specifically, latent genetic and non-shared environmental paths from CU traits to antisocial behavior, as well as the genetic, shared and non-shared environmental paths unique to antisocial behavior were specified to interact with individual differences in CU traits.

Models that included interaction terms were preferred over models with main effects only (see Table S1). Parameter estimates from gene $\times$ trait interaction models are reported in Table 2. Several results are noteworthy. First, CU traits significantly moderated ($p = .001$) additive genetic variance in aggression, such that aggression was more highly heritable among adolescents high on CU traits. Specifically, variance in aggression was 64% heritable for adolescents high (+2 SD) on CU traits. In contrast, for adolescents low (−2 SD) on CU traits, variance in aggression was only 9% heritable; for adolescents with average levels of CU traits, aggression was 34% heritable. Second, we found evidence that CU traits moderate shared environmental variance in rule-breaking ($p < .001$), such that shared environmental influences are magnified among adolescents with high levels of CU traits. For adolescents high (+2 SD) on CU traits, environments shared by siblings living in the same home (i.e. shared environments) accounted for 31% of the variance in rule-breaking. In contrast, for adolescents low (−2 SD) on CU traits, shared environmental variance in rule-breaking approximated zero. For adolescents with average levels of CU traits, shared environmental influences accounted for 18% of the variance in rule-breaking.

Third, CU traits moderated non-shared environmental variance in rule-breaking ($p < .05$), such that non-shared environmental influences are also magnified among adolescents with high levels of CU traits. Specifically, for adolescents high (+2 SD) on CU traits,

environments uniquely experienced by siblings accounted for 46% of the variance in rule-breaking. On the other hand, for adolescents low ($-2\ SD$) on CU traits, non-shared environmental influences accounted for 35% of the variance in rule-breaking. The magnitude of non-shared environmental influences for adolescents with high levels of CU traits was similar to adolescents with average levels of CU traits, accounting for 48% of the variance in rule-breaking. To help visualize moderation effects, the total additive genetic (A), shared environmental (C) and non-shared environmental (E) variance in aggression and rule-breaking were plotted across the continuous range ($\pm 2\ SD$) of CU traits (see Figure 2).

Multivariate Quantitative Genetic Analyses.—Consistent with results of the bivariate quantitative genetic models reported above, a previous meta-analysis of twin and sibling studies indicated that genetic influences on aggression overlap with genetic influences on rule-breaking (Burt, 2013). Also consistent with previous studies, rule-breaking and aggression evinced a moderate phenotypic correlation in the current study ($r = .50$). Therefore, similar to Harden et al. (2015), we fit a multivariate quantitative genetic, common and specific factor model, to decompose variance shared between, and unique to, aggression and rule-breaking into latent biometric components (i.e. A , C , E factors). This model clarifies whether the gene $\times$ CU trait interaction documented in the current study operates on variance unique to aggression, or variance common to aggression and rule-breaking. Similarly, this model also clarifies whether the environment $\times$ CU trait interaction on rule-breaking is on variance common to, or independent of, aggression.

Parameter estimates from this model are reported in Table S2. Several results are noteworthy (see Figure S1 in the online supplement). Only additive genetic variance *unique* to aggression was significantly moderated by CU traits ($b = .196$, $SE = .063$, $p = .002$); additive genetic variances unique to rule-breaking, and common to rule-breaking and aggression, were not significantly moderated by CU traits ($ps > .05$). Shared-environmental variances unique to rule-breaking and aggression were not significantly moderated by CU traits ($ps > .05$). However, shared-environmental variance *common* to rule-breaking and aggression was significantly moderated by CU traits ($b = .130$, $SE = .056$, $p = .022$). Finally, non-shared environmental variance unique to rule-breaking was also significantly moderated by CU traits ($b = .141$, $SE = .035$, $p < .001$); non-shared environmental variances unique to aggression, as well as common to rule-breaking and aggression, were not significantly moderated by CU traits ($ps > .05$).

Discussion

The current study replicates and extends previous research on the association between CU traits and antisocial behavior. First, we find evidence that CU traits index genetic liability for aggressive and non-aggressive expressions of antisocial behavior in adolescence. We also find evidence that higher levels of CU traits amplify additive genetic influences on aggressive behavior, as has been previously found in samples of children using composite indices of antisocial behavior, which collapse across the aggression vs. rule-breaking distinction. Moreover, results provide evidence that aggressive and (non-aggressive) rule-breaking have distinct etiologies, as CU traits did not moderate genetic influences on rule-

breaking behavior. Rather, environmental influences on rule-breaking were heightened among adolescents with high levels of CU traits, such that the heritability of rule-breaking was actually lower at higher levels of CU. Results of the multivariate model, which accounted for the shared etiology between rule-breaking and aggression, were largely consistent with the results of the bivariate models in which rule-breaking and aggression were analyzed independently of each other. Both modeling procedures indicated that genetic influences on aggression (but not rule-breaking), and non-shared environmental influences on rule-breaking (but not aggression) were positively moderated by CU traits. The multivariate model did, however, reveal that shared environmental influences common to both rule-breaking and aggression were positively moderated by CU traits.

These findings are broadly consistent with the results of a meta-analysis (Burt, 2013), which indicates that genetic and environmental influences on rule-breaking are partly distinct from those operating on aggression. The current results from gene $\times$ trait interaction models suggest that CU traits further contribute to the rule-breaking vs. aggression distinction. The finding that environmental influences on rule-breaking were significantly heightened among adolescents with high levels of CU traits was not predicted. Consequently, this finding should be considered tentative prior to replication. Nevertheless, to speculate on this unexpected finding, there is considerable evidence that adolescents who engage in risky behavior, including delinquency, do so in the presence of peers (Boivin, Vitaro, & Poulin, 2005; Bukowski, Brendgen, & Vitaro, 2006; Monahan, Steinberg, & Cauffman, 2009; Silva, Chein & Steinberg, 2016). Moreover, there is evidence that adolescents who engage in minor acts of delinquency are rated as more popular than their rule-abiding peers (Allen, Porter, McFarland, Marsh, & McElhaney, 2005) and unique variance in rule-breaking, after accounting for shared variance with aggression, is actually associated with higher levels of emotional stability (Tackett et al., 2013). Conversely, hostile aggressive behaviors, which may lead to more serious antisocial offenses, are less likely to be rewarded by peers (Allen et al., 2005) and are associated with higher levels of negative affectivity and disagreeableness (Tackett et al., 2013). In sum, there appears to be a strong social component that may be unique to rule-breaking expressions of antisocial behavior. The current study suggests that adolescents with a limited capacity to express prosocial emotions are especially susceptible to environmental sources of variation in rule-breaking behavior, specifically environments that are uniquely experienced by siblings living in the same home. Individual differences in peer groups may provide a non-heritable pathway through which CU traits contribute to risk for non-aggressive expressions of antisocial behavior. Although the current study did not examine individual differences in peer groups, future research efforts may benefit from exploring these and related constructs as they relate to CU traits and the etiology of antisocial behavior.

It also remains an open question as to whether intervention programs can leverage these distinct patterns of social influence on aggressive and nonaggressive behavior to develop more effective strategies for deterring different expressions of antisocial behavior. The present study is particularly notable in identifying underlying CU mechanisms interacting with etiological factors to account for individual differences in rule-breaking behaviors, which have often been characterized as more normative than aggression. Although CU traits are typically thought to be a heritable risk factor for severe expressions of antisocial

behavior, high levels of CU traits may contribute to less severe expressions of antisocial behavior by magnifying the salience of environmental pathways. Future research and intervention efforts stand to benefit from identifying the specific environmental mechanisms that account for the heightened latent environmental influences operating on rule-breaking documented in the current study.

CU traits magnification of genetic risk for aggressive behavior is consistent with the general consensus that, even among “adolescents who show serious antisocial behavior, CU traits designate a particularly aggressive subgroup” (Frick et al., 2014, p.534). However, the results of the current study should *not* be interpreted to suggest that CU traits only confer risk for aggressive, as opposed to non-aggressive antisocial behavior. Rather, results highlight distinct etiological pathways, both heritable and environmental, through which CU traits confer risk for different expressions of antisocial behavior. Indeed, CU traits were as highly correlated with rule-breaking as they were with aggression – a finding that has been observed in other samples, too (Burt, 2013). This pattern of phenotypic associations appears to be invariant to important sample characteristics, including severity of antisocial offending; documented in the current study using a population-representative sample of adolescents, rule-breaking was more highly correlated with CU than aggression ($|r| = .13$, $SE = .04$, $p = .006$), similar to associations observed in a sample of adjudicated adolescents (Kimonis et al., 2008). Although CU traits are most commonly thought of in relation to cruelty or callousness toward a victim, violating societal norms and rules – even violations that do not obviously harm another person – is also facilitated by a disregard for other people. For youth with low levels of CU traits, affectively-charged interpersonal concerns – such as fears about how one’s relationships with close or valued others (parents, teachers, friends) will be affected by one’s behavior – might discourage rule-breaking.

The gene $\times$ trait interaction documented in the current study may also shed light on the problem of missing heritability (Eichler et al., 2010; DeYoung, 2012; Manolio et al., 2009), which refers to the lack of correspondence between molecular genetic and quantitative genetic estimates of heritability. Molecular genetics techniques, specifically genome-wide complex trait analysis (GCTA; Yang, Lee, Goddard, & Visscher, 2011), can estimate the amount of variance in a phenotype collectively explained by thousands of measured variants across the genome. Traditional quantitative genetic techniques, alternatively, estimate heritability based on the known genetic relatedness of relatives. Interestingly, GCTA has failed to reproduce the heritability estimates of childhood behavior problems and associated traits found using quantitative genetic methodology. However, similar to gene $\times$ environment interactions, if the genetic underpinning of a phenotype varies across different levels of an associated phenotype, then efforts to identify genetic main effects will remain difficult, unless such moderating phenotypes are taken into account. Results of the current study suggest that CU traits are an example of such a phenotype.

There are a number of limitations to the current study. First, CU and antisocial constructs were measured using self-reports only. Furthermore, the sample used in the current study ($N = 824$) was moderate in size for quantitative genetic analyses of gene-by-trait interaction. This limitation is evident in the range of confidence intervals surrounding parameter estimates in quantitative genetic interaction models. The precision of the current results

would, therefore, stand to benefit from replication using a larger sample. While previous behavior genetic research has found evidence for a gene $\times$ CU trait interaction on antisocial behavior within tightly circumscribed ages (i.e. 7 and 9-year old children), the current study documented a gene $\times$ CU trait interaction in a sample of adolescents ranging 13 to 20 years old. It remains an open question whether the gene $\times$ CU trait interaction documented in the current study are further nuanced by developmental differences that manifest across adolescence.

There are also a number of strengths to the current study. First, to help overcome limitations of previous research, a well-validated questionnaire with high internal consistency was used to measure CU traits. Second, the majority of quantitative genetic studies on CU traits and antisocial behavior have analyzed racially homogenous samples. The current study used a racially and socio-economically diverse sample, taking care to residualize for main effects of race prior to model fitting. We also used well-validated instruments to measure different subtypes of antisocial behavior and data analytic procedures that modeled a continuous relationship between CU traits and antisocial behavior. As interdisciplinary research continues to elucidate the multi-level cascade of biological processes that bridge the gap between polygenic risk and psychopathology, CU traits continue to be helpful in further organizing and delineating the constellation of physiological, neurological and cognitive mechanisms that account for heterogeneity in the etiology of antisocial behavior.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Reference

- Allen JP, Porter MR, McFarland FC, Marsh P, & McElhaney KB (2005). The two faces of adolescents' success with peers: Adolescent popularity, social adaptation, and deviant behavior. *Child Development*, 76, 747–760. [PubMed: 15892790]
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed). Arlington, VA: American Psychiatric Publishing.
- Boivin M, Vitaro F, & Poulin F (2005). Peer relationships and the development of aggressive behavior in early childhood In Tremblay RE, Hartup WW, & Archer J (Eds.), *Developmental origins of aggression* (pp. 376–397). New York, NY: Guilford Press.
- Bukowski WM, Brendgen M, & Vitaro F (2006). Peer relations In Grusec JE, & Hastings PD (Eds.), *Handbook of Socialization* (pp. 355–381). New York, NY: Guilford Press.
- Burt SA (2009). Are there meaningful etiological differences within antisocial behavior? Results of a meta-analysis. *Clinical Psychology Review*, 29, 163–178. [PubMed: 19193479]
- Burt SA (2012). How do we optimally conceptualize the heterogeneity within antisocial behavior? An argument for aggressive versus non-aggressive behavioral dimensions. *Clinical Psychology Review*, 32, 263–279. [PubMed: 22459789]
- Burt SA (2013). Do etiological influences on aggression overlap with those on rule breaking? A meta-analysis. *Psychological Medicine*, 43(09), 1801–1812. [PubMed: 22909442]
- Burt SA, & Klump KL (2009). The etiological moderation of aggressive and nonaggressive antisocial behavior by age. *Twin Research and Human Genetics*, 12(04), 343–350. [PubMed: 19653834]
- Burt SA, & Neiderhiser JM (2009). Aggressive versus nonaggressive antisocial behavior: Distinctive etiological moderation by age. *Developmental Psychology*, 45, 1164–1176 [PubMed: 19586186]
- DeYoung CG, & Clark R (2012). The gene in its natural habitat: The importance of gene–trait interactions. *Development and psychopathology*, 24(04), 1307–1318. [PubMed: 23062299]

- Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, & Nadeau JH (2010). Missing heritability and strategies for finding the underlying causes of complex disease. *Nature Reviews Genetics*, 11(6), 446–450.
- Frick PJ, & Moffit TE (2010). A proposal to the DSM-V childhood disorders and the ADHD and disruptive behavior disorders work groups to include a specifier to the diagnosis of conduct disorder based on the presence of callous–unemotional traits. USA: American Psychiatric Association, 1–36.
- Frick PJ, Ray JV, Thornton LC, & Kahn RE (2014). Can callous-unemotional traits enhance the understanding, diagnosis, and treatment of serious conduct problems in children and adolescents? A comprehensive review. *Psychological Bulletin*, 140(1), 1–57. [PubMed: 23796269]
- Frick P, & White S (2008). Research review: The importance of callous–unemotional traits for developmental models of aggressive and antisocial behavior. *Journal of Child Psychology and Psychiatry*, 49, 359–375. [PubMed: 18221345]
- Harden KP, Patterson MW, Briley DA, Engelhardt LE, Kretsch N, Mann FD,... & Tucker Drob EM (2015). Developmental changes in genetic and environmental influences on rule breaking and aggression: age and pubertal development. *Journal of child psychology and psychiatry*, 56(12), 1370–1379. [PubMed: 25902931]
- Harden KP, Tucker-Drob EM, & Tackett JL (2013). The Texas twin project. *Twin Research and Human Genetics: the Official Journal of the International Society for Twin Studies*, 16(1), 385–390. [PubMed: 23111007]
- Heath AC, Nyholt DR, Neuman R, Madden PA, Bucholz KK, Todd RD,... & Martin NG (2003). Zygosity diagnosis in the absence of genotypic data: an approach using latent class analysis. *Twin Research*, 6(01), 22–26. [PubMed: 12626225]
- Hyde LW, Waller R, Trentacosta CJ, Shaw DS, Neiderhiser JM, Ganiban JM, Reiss D, Leve LD (2016). Heritable and Nonheritable Pathways to Early Callous-Unemotional Behaviors. *American Journal of Psychiatry*, 173, 903–910. [PubMed: 27056607]
- Kimonis ER, Frick PJ, Skeem JL, Marsee MA, Cruise K, Munoz LC, Aucoin KJ, & Morris AS (2008). Assessing callous-unemotional traits in adolescent offenders: validation of the inventory of callous-unemotional traits. *International Journal of Law and Psychiatry*, 31(3), 241–252. [PubMed: 18514315]
- Krueger RF, & Markon KE (2014) The Role of the DSM-5 Personality Trait Model in Moving Toward a Quantitative and Empirically Based Approach to Classifying Personality and Psychopathology. *Annual Review of Clinical Psychology*, 10, 477–501.
- Larsson H, Andershed H, & Lichtenstein P (2006). A genetic factor explains most of the variation in the psychopathic personality. *Journal of Abnormal Psychology*, 115, 221–230. [PubMed: 16737387]
- Lizotte AJ, Chard-Wierschem DJ, Loeber R, & Stern SB (1992). A shortened Child Behavior Checklist for delinquency studies. *Journal of Quantitative Criminology*, 8(2), 233–245.
- Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ,... & Cho JH (2009). Finding the missing heritability of complex diseases. *Nature*, 461(7265), 747–753. [PubMed: 19812666]
- McGue M, & Bouchard TJ, Jr (1984). Adjustment of twin data for the effects of age and sex. *Behavior Genetics*, 14(4), 325–343. [PubMed: 6542356]
- Monahan K, Steinberg L, & Cauffman E (2009). Affiliation with antisocial peers, susceptibility to peer influence, and desistance from antisocial behavior during the transition to adulthood. *Developmental Psychology*, 45, 1520–1530. [PubMed: 19899911]
- Moore AA, Carney D, Moroney E, Machlin L, Towbin KE, Brotman MA,... & Hettema JM (2016). The Inventory of Callous-Unemotional Traits (ICU) in Children: Reliability and Heritability. *Behavior Genetics*, 1–11. [PubMed: 26732815]
- Muthén B (2004). Latent variable analysis The Sage handbook of quantitative methodology for the social sciences. Thousand Oaks, CA: Sage Publications, 345–68.
- Muthén LK, & Muthén BO (2010). Mplus User's Guide: Statistical Analysis with Latent Variables: User's Guide. Muthén & Muthén.

- Neale MC, & Maes HH (2004). Methodology for genetic studies of twins and families. Dordrecht: Kluwer Academic Publishers B.V.
- Ray JV, Frick PJ, Thornton LC, Steinberg L, & Cauffman E (2015). Positive and Negative Item Wording and Its Influence on the Assessment of Callous-Unemotional Traits. *Psychological Assessment*, Advance Online Publication. 10.1037/pas0000183
- Silva K, Chein J, & Steinberg L (2016). Adolescents in peer groups make more prudent decisions when a slightly older adult is present. *Psychological Science*, 27, 322–330. [PubMed: 26791822]
- Tackett JL, Daoud SLS, De Bolle M, & Burt SA (2013) Is Relational Aggression Part of the Externalizing Spectrum? A Bifactor Model of Youth Antisocial Behavior. *Aggressive Behavior*, 39, 149–159. [PubMed: 23386551]
- Taylor J, Loney BR, Bobadilla L, Iacono WG, & McGue M (2003). Genetic and environmental influences on psychopathy trait dimensions in a community sample of male twins. *Journal of Abnormal Child Psychology*, 31, 633–645. [PubMed: 14658743]
- Viding E, Blair RJR, Moffitt TE, & Plomin R (2005). Evidence for substantial genetic risk for psychopathy in 7 year olds. *Journal of Child Psychology and Psychiatry*, 46(6), 592–597. [PubMed: 15877765]
- Viding E, Frick PJ, & Plomin R (2007). Aetiology of the relationship between callous–unemotional traits and conduct problems in childhood. *The British Journal of Psychiatry*, 190(49), s33–s38.
- Viding E, Jones AP, Frick JP, Moffitt TE, & Plomin R (2008). Heritability of antisocial behaviour at 9: do callous unemotional traits matter?. *Developmental Science*, 11(1), 17–22. [PubMed: 18171362]
- Viding E Fontaine NMG, McCrory EJ (2012) Antisocial behaviour in children with and without callous-unemotional traits. *Journal of the Royal Society of Medicine*, 105, 195–200. [PubMed: 22637770]
- Waldman ID, Tackett JL, Van Hulle CA, Applegate B, Pardini D, Frick PJ, & Lahey BB (2011). Child and adolescent conduct disorder substantially shares genetic influences with three socioemotional dispositions. *Journal of Abnormal Psychology*, 120, 57–70. [PubMed: 21142331]
- Waller R, Gardner F, & Hyde LW (2013). What are the associations between parenting, callous–unemotional traits, and antisocial behavior in youth? A systematic review of evidence. *Clinical Psychology Review*, 33(4), 593–608. [PubMed: 23583974]
- Yang J, Lee SH, Goddard ME, & Visscher PM (2011). GCTA: a tool for genome-wide complex trait analysis. *The American Journal of Human Genetics*, 88(1), 76–82. [PubMed: 21167468]

Figure 1. Path Diagram for Bivariate Quantitative Genetic Models

Note. Main effects models (top panel) were used to estimate latent genetic and environmental correlations. Moderation models (bottom panel) were used to test Gene $\times$ Trait interactions.

Figure 2. Gene × Callous-Unemotional Trait Interaction Effects on Antisocial Behavior

Note. Additive genetic (blue), shared environmental (gray) and non-shared environmental (black) variance in aggression and rule-breaking are plotted across the continuous range (+/- 2 SD) of callous-unemotional traits. Top panel: dashed and solid lines represent unstandardized estimates of variance in antisocial behavior that are common to and unique of callous-unemotional traits, respectively. Asterisks mark components of variance significantly moderated by CU traits. Bottom panel: shaded regions represent portions of total variance in aggression and rule-breaking. Portions of variance calculated from parameter estimates reported in Table 2 using the following equations:

$$A = \frac{(a_{12} + a'_{12}x)^2 + (a_2 + a'_2x)^2}{(a_{12} + a'_{12}x)^2 + (e_{12} + e'_{12}x)^2 + (a_2 + a'_2x)^2 + (c_2 + c'_2x)^2 + (e_2 + e'_2x)^2}$$

$$C = \frac{(c_2 + c'_{2x})^2}{(a_{12} + a'_{12x})^2 + (e_{12} + e'_{12x})^2 + (a_2 + a'_{2x})^2 + (c_2 + c'_{2x})^2 + (e_2 + e'_{2x})^2}$$

$$E = \frac{(e_{12} + e'_{12x})^2 + (e_2 + e'_{2x})^2}{(a_{12} + a'_{12x})^2 + (e_{12} + e'_{12x})^2 + (a_2 + a'_{2x})^2 + (c_2 + c'_{2x})^2 + (e_2 + e'_{2x})^2}$$

Table 1.
Descriptive Statistics, Correlations and Univariate Twin Models of Callous-Unemotional (CU) Traits, Aggression (AGG) and Rule-Breaking (RB)

Variables	Descriptive Statistics					Phenotypic Correlations			Twin-Pair Correlations		Univariate ACE Models		
	<i>M</i>	<i>SD</i>	<i>min</i>	<i>max</i>	α	CU	AGG	RB	<i>rMZ</i>	<i>rDZ</i>	h^2	c^2	e^2
CU	1.88	.31	1.12	2.79	.79	1	.29	.42	.45	.11	.39	.00	.61
AGG	.39	.24	0.00	1.62	.71	.30	1	.50	.36	.19	.36	.01	.63
RB	.28	.25	0.00	1.50	.75	.43	.50	1	.48	.32	.32	.16	.52

Note. Means (*M*), standard deviations (*SD*), minimum observed values (*min*) and maximum observed values (*max*) reported for non-transformed variables. α = Cronbach's alpha. Correlations calculated using log-transformed variables. Zero-order phenotypic correlations (reported below diagonal). Partial phenotypic correlations (reported above diagonal) and cross-twin within-trait correlations for monozygotic twins (*rMZ*) and dizygotic twins (*rDZ*) control for age, sex, age $\times$ sex and race. All correlations are significant at $p(\text{two-tailed}) < .001$, after accounting for the non-independence of data due to siblings being nested with in the same home. h^2 = additive genetic variance. c^2 = shared environmental variance. e^2 = non-shared environmental variance.

Table 2.
Parameter Estimates from Bivariate Gene $\times$ Trait Interaction Models

(N Pairs = 436)	Instrument:		YSR				
	Scales:		Aggression		Rule-Breaking		
	Parameters:	<i>b</i>	(CI.95%)	<i>p</i>	<i>b</i>	(CI.95%)	<i>p</i>
Variance in CU Traits							
Main genetic path (a_1)		.63	(.52, .75)	< .001	.64	(.47, .74)	< .001
Main nonshared-environmental path (e_1)		.78	(.69, .86)	< .001	.77	(.70, .89)	< .001
Common Variance							
Main genetic path (a_{12})		.26	(.11, .41)	.001	.37	(.22, .52)	< .001
Gene $\times$ Trait interaction (a_{12}')		-.01	(-.16, .13)	.839	.11	(-.05, .27)	.190
Main nonshared environmental path (e_{12})		.16	(.04, .28)	.007	.21	(.09, .33)	.001
Nonshared environment $\times$ Trait interaction (e_{12}')		.05	(-.05, .16)	.323	-.04	(-.15, .08)	.500
Unique Variance in Rule-Breaking/Aggression							
Main genetic effect (a_2)		.49	(.30, .67)	< .001	.41	(.11, .70)	.007
Gene $\times$ CU Trait interaction (a_2')		.21	(.09, .33)	.001	-.11	(-.29, .07)	.233
Main environmental path (c_2)		.22	(-.08, .52)	.146	.41	(.20, .62)	< .001
Shared Environment $\times$ Trait interaction (c_2')		-.12	(-.30, .06)	.192	.16	(.06, .26)	.001
Main nonshared environmental effect (e_2)		.72	(.66, .79)	< .001	.63	(.54, .72)	< .001
Nonshared environmental $\times$ Trait interaction (e_2')		-.03	(-.10, .04)	.399	.12	(.03, .21)	.019

Note. *b* = unstandardized parameter estimates. CI.95% = 95% confidence intervals. *p* = two-tailed probability of type-I error.