Cochrane
Library

Cochrane Database of Systematic Reviews

Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest (Review)

Arrich J, Holzer M, Havel C, Warenits AM, Herkner H

Arrich J, Holzer M, Havel C, Warenits AM, Herkner H.

Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest.

Cochrane Database of Systematic Reviews 2016, Issue 3. Art. No.: CD010570.

DOI: [10.1002/14651858.CD010570.pub2](https://doi.org/10.1002/14651858.CD010570.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	6
OBJECTIVES	7
METHODS	7
RESULTS	9
Figure 1.	10
Figure 2.	12
Figure 3.	13
Figure 4.	14
Figure 5.	15
DISCUSSION	16
AUTHORS' CONCLUSIONS	17
ACKNOWLEDGEMENTS	17
REFERENCES	18
CHARACTERISTICS OF STUDIES	21
DATA AND ANALYSES	34
Analysis 1.1. Comparison 1 Survival: pre-hospital cooling versus in-hospital cooling, Outcome 1 Survival.	34
Analysis 2.1. Comparison 2 Neurological outcome: pre-hospital cooling versus in-hospital cooling, Outcome 1 Good neurological outcome.	34
ADDITIONAL TABLES	35
APPENDICES	36
WHAT'S NEW	41
CONTRIBUTIONS OF AUTHORS	41
DECLARATIONS OF INTEREST	42
SOURCES OF SUPPORT	42
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	42
NOTES	42
INDEX TERMS	43

[Intervention Review]

Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest

Jasmin Arrich¹, Michael Holzer¹, Christof Havel¹, Alexandra-Maria Warenits¹, Harald Herkner¹

¹Department of Emergency Medicine, Medical University of Vienna, Vienna, Austria

Contact: Jasmin Arrich, Department of Emergency Medicine, Medical University of Vienna, Währinger Gürtel 18-20 / 6D, Vienna, 1090, Austria. jasmin.arrich@meduniwien.ac.at.

Editorial group: Cochrane Emergency and Critical Care Group.

Publication status and date: Edited (no change to conclusions), published in Issue 1, 2019.

Citation: Arrich J, Holzer M, Havel C, Warenits AM, Herkner H. Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest. *Cochrane Database of Systematic Reviews* 2016, Issue 3. Art. No.: CD010570. DOI: [10.1002/14651858.CD010570.pub2](https://doi.org/10.1002/14651858.CD010570.pub2).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Targeted temperature management (also known under 'therapeutic hypothermia', 'induced hypothermia', or 'cooling') has been shown to be beneficial for neurological outcome in patients who have had successful resuscitation from sudden cardiac arrest, but it remains unclear when this intervention should be initiated.

Objectives

To assess the effects of pre-hospital initiation of cooling on survival and neurological outcome in comparison to in-hospital initiation of cooling for adults with pre-hospital cardiac arrest.

Search methods

We searched CENTRAL, MEDLINE, EMBASE, CINAHL, BIOSIS, and three trials registers from inception to 5 March 2015, and carried out reference checking, citation searching, and contact with study authors to identify additional studies.

Selection criteria

We searched for randomized controlled trials (RCTs) in adults with out-of-hospital cardiac arrest comparing cooling in the pre-hospital setting to in-hospital cooling. Our primary outcomes were survival and neurological outcome; our secondary outcomes were adverse events, quality of life, and length of stay in the intensive care unit (ICU) and in the hospital.

Data collection and analysis

We used Cochrane's standard methodological procedures.

Main results

We included seven RCTs (2369 participants randomized) on the induction of pre-hospital cooling in comparison to in-hospital cooling. There was considerable methodological heterogeneity and risk of bias mainly due to deficits in the administration of cooling, therefore we refrained from pooling the results for survival and neurological outcome and we presented the results for each study separately. Adverse events were rare: based on four studies with 1713 adults pre-hospital induction of cooling may increase the risk of cardiac re-arrests. Risk of bias within the seven individual studies was generally moderate. Overall the quality of the evidence was very low. This was mainly driven by inconsistency and low precision.

Authors' conclusions

Currently, there is no convincing evidence to clearly delineate beneficial or harmful effects of pre-hospital induction of cooling in comparison to in-hospital induction of cooling. This conclusion is based on very low quality evidence.

PLAIN LANGUAGE SUMMARY

Should patients experiencing sudden cardiac death be cooled to lower their body temperature prior to or after admission to hospital?

Review question

We reviewed the current available evidence in order to answer the question of whether early cooling in people who receive basic life support for sudden cardiac death influences survival and brain damage compared to cooling that is started after their admission to hospital. Early cooling means the cooling of the person quickly by the ambulance staff, paramedics or doctors, in the field. We included seven studies meeting the Cochrane requirements in this review.

Background

Population

This review deals with people who receive basic life support for sudden cardiac death. Sudden cardiac death means that the heart and subsequently the circulation stops. If these people do not receive early cardiopulmonary resuscitation then their brain cells begin to be irreversibly damaged and subsequently they die. If basic life support is successful, one form of therapy that may help to prevent further cell damage is to cool the body for several hours to 32°C to 36°C. This therapy has been shown to be beneficial in reducing brain damage and is recommended in international guidelines for the treatment of people that have been brought back to life after sudden cardiac death.

Intervention

The optimal timing for the initiation of cooling is unclear. This review compares people who had their cooling therapy started before hospital admission to those who had their cooling therapy started after admission to a hospital.

Outcomes

The effects of the intervention were measured by survival and brain damage, together with side effects, quality of life, and length of hospital stay.

Search date

We completed the review searches in March 2015.

Study characteristics

The seven studies included had a total of 2369 participants and compared the effects of cooling before and after being admitted to the hospital. The mean age of the participants in the studies was between 59 and 68 years with the majority being male. People that were not included in the trials were generally those with trauma, those with a terminal disease, those at the natural end of their life, pregnant women, and those that already had a low body temperature.

Study funding sources

Two out of seven studies were funded by the medical industry, four received funding from the government or non-profit organizations, and one study did not receive any funding.

Key results

None of the studies found any evidence for a benefit of pre-hospital cooling versus in-hospital cooling. However, we discovered that in almost all studies a relevant amount of participants did not receive pre-hospital cooling or in-hospital cooling or cooling according to the guidelines at all. The reasons for this were not clearly stated. The question of whether the decision to cool participants may have been influenced by other factors cannot be reliably answered. Proper design and conduct of the included studies was of concern, therefore to avoid making misleading interpretations we did not pool the results of the single studies. We found that in adults that received pre-hospital cooling the heart was slightly more likely to stop again before they were admitted to the hospital.

Quality of the evidence

Many of the included studies were of limited use because they focused on the practicability and safety of pre-hospital cooling without specifically emphasising cooling therapy. Other factors that contributed to a downgrading of the quality of the evidence were that the

information came from different study populations and from different time points of applying pre-hospital cooling. In addition, there was risk of bias within the studies. The quality of the individual studies was moderate. In summary, the quality of the evidence to answer our review question was very low.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Pre-hospital cooling compared to in-hospital cooling for survival, neuroprotection, and adverse events after out-of-hospital cardiac arrest

Survival, neurological outcome, and adverse events: pre-hospital cooling compared to in-hospital cooling after out-of-hospital cardiac arrest

Patient or population: out-of-hospital cardiac arrest
Settings: emergency medicine and intensive care, worldwide
Intervention: pre-hospital cooling
Comparison: in-hospital cooling

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	In-hospital cooling	Pre-hospital cooling				
Survival and good neurological outcome	Study population		Not pooled	2369 randomized participants (7 RCTs) (Bernard 2010; Bernard 2012; Castren 2010; Debaty 2014; Kämäräinen 2009; Kim 2007; Kim 2014)	⊕⊕⊕⊕ VERY LOW 1,2,3,4,5,6,7	—
	Not pooled	Not pooled				
	Not pooled	Not pooled				
Adverse events - re-arrest after randomization	Study population		RR 1.23 (1.02 to 1.48)	1713 participants with available information (4 studies) (Castren 2010; Kämäräinen 2009; Kim 2007; Kim 2014)	⊕⊕⊕⊕ VERY LOW 1,2,3,4, 7	—
	183 per 1000	225 per 1000 (187 to 271)				
	Moderate					
	186 per 1000	229 per 1000 (190 to 276)				

We downgraded the quality of the evidence to 'very low' for the following reasons:

¹Inappropriate application of intervention or control, or both.

²Indirectness in the intervention: two studies evaluated intra-arrest cooling while all others evaluated post-arrest cooling (Castren 2010; Debaty 2014).

³Indirectness in the intervention: the rate of application of pre-hospital cooling varied over all studies; up to 50% of all participants did not receive the full intervention (Bernard 2010; Bernard 2012; Kim 2007; Kim 2014); up to 16% of participants did not receive the intervention at all (Bernard 2010; Bernard 2012; Kim 2007; Kim 2014).

⁴Indirectness in the comparator: the rate of application of in-hospital cooling varied over all studies; some studies did not provide information (Castren 2010; Kämäräinen 2009; Kim 2007; Kim 2014); in some only some of the participants received in-hospital cooling (Kämäräinen 2009; Kim 2007; Kim 2014); temperature curves in some studies indicated that a relevant proportion of participants were not cooled according to the then current guidelines (Bernard 2010; Bernard 2012); in some studies the target temperature was at the upper limit of the then current guidelines (Castren 2010).

⁵Indirectness in the population: one study included only adults with non-ventricular fibrillation cardiac arrest (Bernard 2010); another only ventricular fibrillation cardiac arrest (Bernard 2012); the others did not make restrictions.

⁶Imprecision: three studies were feasibility or pilot studies with sample sizes too small to evaluate clinical outcomes (Castren 2010; Kämäräinen 2009; Kim 2007). Due to the above described reasons we refrained from pooling the estimates, therefore we are left with the lower precision of the individual studies.

⁷Risk of bias within studies: three studies lacked blinding of outcome assessment, which may substantially bias the assessment of neurological outcome (Castren 2010; Kämäräinen 2009; Kim 2007); insufficient administration and continuation of the intervention and comparator/no information on administration and continuation of the intervention and comparator (see above).

BACKGROUND

Cooling the body compared with no cooling has been shown to be beneficial for neurological outcome and survival in patients who have had successful resuscitation from sudden cardiac arrest (Arrich 2016), but it remains unclear when this intervention should be initiated.

Description of the condition

The incidence of out-of-hospital sudden cardiac arrest is not easily determined. A review of emergency medicine services (EMS)-treated out-of-hospital cardiac arrest in Europe reports an incidence of 41 per 100,000 person-years (Berdowski 2010), while in North America around 52 per 100,000 person-years are reported (Nichol 2008). The survival rates of 9% to 11% for all-rhythm cardiac arrests and 19% to 21% for ventricular fibrillation cardiac arrests seem to have improved in recent decades, but are still disastrously low. When the circulation stops the brain cells are directly damaged through the lack of oxygen and adenosine triphosphate (ATP), but even after the circulation is re-established continuous damage of the brain cells happens through neuronal necrosis and apoptosis caused by a myriad of pathophysiological mechanisms, microcirculatory failure, and other factors such as pyrexia, hyperglycaemia, and seizures (Holzer 2010; Nolan 2008). After resuscitation and admission to a hospital, post-resuscitation care aims to reduce the secondary reperfusion injuries caused by the cardiac arrest. This treatment includes early treatment of the cause of the arrest (the majority have a coronary heart disease, thus they will undergo coronary angiography with subsequent percutaneous coronary intervention (PCI) if indicated), cooling, optimization of oxygenation, ventilation, circulation and metabolism, and early seizure detection and treatment following a standardized treatment protocol (Callaway 2015; Soar 2015).

Description of the intervention

Cooling the body as the principle of targeted temperature management (formerly 'therapeutic hypothermia' or 'induced hypothermia') may improve survival and reduce the amount of neurological damage after cardiac arrest (Arrich 2016). According to recent guidelines, comatose survivors of out-of-hospital ventricular fibrillation cardiac arrest should be cooled with internal or external cooling techniques to a target temperature of 32°C to 36°C (patients with in-hospital cardiac arrest or other primary rhythms may also benefit) (Callaway 2015; Soar 2015). This targeted temperature should be maintained for at least 24 hours, and after this cooling period the patients should be rewarmed at a rate of 0.25°C to 0.5°C/hour to normothermia. Pre-hospital application of cooling refers to the reduction of body core temperature for at least 12 hours by the healthcare professionals in the field, either during resuscitation or shortly after resuscitation. Several methods are available to decrease body temperature, including surface cooling methods like ice packs, cold-air mattresses, water-circulating cooling pads, and pre-cooled cooling pads. The core temperature may also be decreased by intravascular cooling catheters (Holzer 2010). Other methods include cooling caps or the application of coolant through the nose, which cools the nasal cavity and subsequently the whole body through evaporation (Castren 2010). The rapid infusion of large volumes of cold intravenous fluid immediately after return of spontaneous circulation was discouraged in the most recent guidelines (Callaway 2015; Soar 2015). However, not all methods

that are available after admission to a hospital will be available or practical in the field.

Formally the intervention in this review is the time point of application of cooling. Practically, application of cooling includes significant variation and we expected study protocols and clinical routines to differ substantially. Therefore we pragmatically compared early application in the pre-hospital phase to later application after hospital admission. However, cooling should be continued after hospital admission. Maintenance cooling and rewarming should be according to the current guidelines on resuscitation at that time and comparable to the control group.

How the intervention might work

Pathophysiologically, brain damage through hypoxia is due to two main mechanisms. The first is direct excitotoxic and ischaemic cell death leading to necrosis and apoptosis, and the second is reperfusion injury, which is also known as 'post-resuscitation disease or syndrome' (Holzer 2010; Nolan 2008). Cooling the body inhibits numerous pathways of these two mechanisms. In vivo studies comparing different starting points of cooling are scarce, but there are a number of animal studies that have indicated that an earlier start of cooling, even during the resuscitation phase, might lead to an improved neurological outcome (Abella 2008; Janata 2010; Kuboyama 1993; Sterz 1991).

Cooling compared to no cooling has been shown to be beneficial for neurological outcome and survival in adults who have had resuscitation from sudden cardiac arrest (Arrich 2016). This evidence is mainly based on three randomized controlled trials (Bernard 2002; HACA 2002; Hachimi-Idrissi 2001). In these trials cooling had been initiated as late as four hours after cardiac arrest. However, the question of whether an earlier cooling start may be more beneficial has been raised in many experimental and human studies. It is important to note that 'earlier' refers to different time points of cooling, before resuscitation is started (Janata 2010; Zhao 2008), during resuscitation (Abella 2004; Yannopoulos 2009), immediately after resuscitation (Kuboyama 1993) (as opposed to a 15-minute delay), and after one hour (as opposed to a four-hour delay) (Colbourne 1995). All of these studies showed that an earlier initiation of cooling was more beneficial. However, a recent animal study demonstrated that there seemed to be no difference as long as cooling was started less than four hours after cardiac arrest (Che 2011). Also, retrospective human data show conflicting results. Some studies showed a better neurological outcome with an earlier start of cooling or faster cooling (Sendelbach 2012; Wolff 2009), and others did not find that earlier cooling had a beneficial effect (Nielsen 2009). One cohort study even suggested that adults who reached the target temperature earlier also had a poorer prognosis (Haugk 2011). This finding can probably be explained by the fact that patients who are more severely harmed by the cardiac arrest might have compromised thermoregulation and therefore less resistance against induction of cooling (Suffoletto 2009). The dilemma of differentiating a diagnostic sign (an early outcome) from an effective intervention can only be challenged by randomized trials.

Why it is important to do this review

We have conducted this systematic review to explore the uncertainty arising from conflicting results in a number of animal and clinical studies.

Following the animal studies ([Abella 2008](#); [Janata 2010](#); [Kuboyama 1993](#); [Sterz 1991](#)), a few case series were published on the potential beneficial effect of an earlier versus a delayed start for cooling, with conflicting results ([Nielsen 2009](#); [Wolff 2009](#)).

A few randomized controlled trials have been published that evaluate the feasibility and effectiveness of cooling during resuscitation and shortly after resuscitation versus in-hospital cooling, which is the standard care in most centres. The results were conflicting, partly due to the small study size of the pilot studies and maybe partly due to methodological shortcomings. Some showed a significant effect in the subgroups ([Castren 2010](#); [Kim 2007](#)).

Cooling may improve neurological outcome after resuscitation from sudden cardiac arrest even when its application is delayed. It is important to find out if timing of cooling initiation has an effect on clinical outcomes. If a pre-hospital start of cooling provides a further improvement of neurological outcome, this effect should not be missed because of methodological shortcomings or small sample sizes of the preceding randomized controlled trials. If there is no further gain with a pre-hospital cooling start, the extra effort to start cooling in the field could be spared. This would help decision makers and guideline committees in advising healthcare professionals. Currently the guidelines comment that "Early cooling strategies, other than rapid infusion of large volumes of cold IV fluid, and cooling during CPR in the prehospital setting have not been studied adequately" ([Callaway 2015](#); [Soar 2015](#)). For the scientific community, a structured evaluation of the available evidence will provide a basis on which to decide whether further studies should be undertaken in this field and will serve as the basis for sample size calculations.

OBJECTIVES

To assess the effects of pre-hospital initiation of cooling on survival and neurological outcome in comparison to in-hospital initiation of cooling for adults with pre-hospital cardiac arrest.

METHODS

Criteria for considering studies for this review

Types of studies

We planned to include randomized and 'quasi-randomized' controlled trials. 'Quasi-randomized' refers to allocation procedures such as alternating days, odd and even days and the like, which we planned to include because of the relative novelty of the intervention and the expected low number of trials available to date. We planned to exclude cross-over studies given the condition and the nature of the outcomes assessed.

Types of participants

We included studies in adults with out-of-hospital cardiac arrest who received targeted temperature management.

Types of interventions

We compared pre-hospital induction of cooling to in-hospital induction of cooling.

Pre-hospital induction of cooling (intervention): cooling during resuscitation or shortly after resuscitation in the out-of-hospital setting.

Later induction of cooling (standard therapy; control): in-hospital cooling.

Types of outcome measures

Primary outcomes

1. Survival: we investigated short-term survival (closest to 30 days) and long-term survival (closest to six months).
2. Neurological outcome: ideally we expected the outcome to be reported as best neurological outcome during hospital stay and in cerebral performance categories (CPC) ([Cummins 1991](#); [Jennett 1975](#)). Good neurological outcome is usually considered if the CPC is 1 or 2. If authors grouped this outcome along with other categories or cut-offs, or used other instruments for neurological outcome assessment, like the modified Rankin Scale, we accepted this for our meta-analysis and planned to perform a sensitivity analysis based on the outcome definition.

We considered both survival and neurological outcome as dichotomous data at a given point in time.

Secondary outcomes

1. Adverse events, as reported by study authors (dichotomous data).
2. Adverse events related to cooling methods, as reported by study authors (dichotomous data).
3. Quality of life, as reported by study authors (data as reported).
4. Length of stay in the intensive care unit (ICU) and in the hospital, as proxies for economic outcomes (continuous data).

Availability of outcomes was not part of the study eligibility criteria. We planned to include studies that met the participant, intervention, and comparison criteria in the review even if they reported no relevant outcomes.

Search methods for identification of studies

Electronic searches

We searched the following databases (from inception to March 2015): the Cochrane Central Register of Controlled Trials (CENTRAL) ([Appendix 1](#)), MEDLINE ([Appendix 2](#)), EMBASE ([Appendix 3](#)), CINAHL ([Appendix 4](#)), and BIOSIS ([Appendix 5](#)), and three trials registers (EudraCT, [ClinicalTrials.gov](#), and the International Clinical Trials Registry Platform).

We did not apply any language restrictions.

We used a search strategy for identifying randomized controlled trials (RCTs) in MEDLINE and EMBASE ([Higgins 2011](#)).

Searching other resources

In an attempt to identify further studies we searched the primary clinical trials registers (March 2015) accepted by the International Committee of Medical Journal Editors (see <http://www.icmje.org/recommendations/browse/publishing-and-editorial-issues/clinical-trial-registration.html>). We also asked experts in the field whether they were aware of any ongoing, unpublished, or published trials on this subject. We also searched the reference lists of included studies and other reviews on the topic.

Data collection and analysis

Selection of studies

We imported all retrieved results into EndNote (version X7, Thomson Corporation) and eliminated any duplicates. Two authors each (CH + MH and A-M W + JA) independently scanned titles and abstracts for relevance. For all references that were not excluded by both authors at this stage, we retrieved the full texts and examined them for compliance with the inclusion criteria. We linked multiple reports of the same study. We resolved discrepancies by discussion or by involving a third author as an arbiter (HH).

Data extraction and management

Two authors each (CH + MH and A-M W + JA) independently extracted all relevant data into a predefined form ([Appendix 6](#)). We resolved discrepancies by discussion or by involving a third author as arbiter (HH). We then entered data into the Cochrane software program Review Manager ([RevMan 5.3](#)).

Assessment of risk of bias in included studies

To assess the internal validity of the identified trials, we used a domain-based evaluation, assessing random sequence generation, allocation concealment, blinding of outcome assessment (for neurological outcome and quality of life), incomplete outcome data (primary outcome), selective reporting (neurological outcome), and exclusion of randomized participants from the analysis (primary outcome).

Two authors (as named above) independently extracted and tabulated all relevant information. We resolved discrepancies by discussion or by involving a third author as arbiter.

We assumed that blinding of participants and personnel was not relevant, as participants are unconscious and cooling interventions are almost impossible to blind.

We assessed the domains as low risk of bias, unclear, or high risk of bias.

Measures of treatment effect

The primary measure of treatment effect for the primary outcomes was the risk ratio (relative risk) for surviving and achieving good neurological recovery in participants allocated to pre-hospital cooling when compared to in-hospital cooling. The same applied for adverse events. For quality of life and length of hospital stay data, we used mean differences as measures of treatment effect. In the case that several instruments were used for quality of life assessment across studies, we used standardized mean differences instead.

Unit of analysis issues

Cluster-randomized trials

For cluster-randomized trials we planned to use estimates that allow for the clustered structure of the data. In the absence of adequate estimates we would have used appropriate approximations.

Studies with multiple treatment groups

In the case of multiple treatments (for example intra-arrest cooling versus pre-hospital arrest cooling versus in-hospital cooling) we

planned to combine the groups to create a single pair-wise comparison, but to avoid overall estimates.

Dealing with missing data

For a negligible amount of missing data (or if they were convincingly missing at random), we analysed only the available data. We did not employ data imputation or data replacement methods.

Assessment of heterogeneity

We evaluated clinical and statistical heterogeneity. We assessed clinical heterogeneity by tabulating and informally inspecting relevant data. We only performed quantitative synthesis if clinical heterogeneity was negligible. We assessed statistical heterogeneity using the I^2 statistic ([Higgins 2003](#)). We considered statistical heterogeneity to be relevant if the I^2 statistic was $> 50\%$. To investigate heterogeneity we performed subgroup analyses and the test for subgroup differences. If possible, we employed meta-regression to further investigate heterogeneity.

Assessment of reporting biases

We planned to assess the presence of possible reporting bias using funnel plots (plotting the effect against precision) ([Egger 1997](#)), and to inspect them visually ([Higgins 2011](#)).

Data synthesis

We calculated risk ratios (RR) and their 95% confidence intervals (CI) for dichotomous variables. For continuous variables we planned to calculate mean differences or standardized mean differences depending on the comparability of measurement methods. Generally we used random-effects models. We used RevMan 5.3 for standard calculations and Stata 11 for meta-regression.

Subgroup analysis and investigation of heterogeneity

We planned to investigate the following subgroups:

1. Efficacy of cooling methods in the intervention group and in the control group.
2. Intra-arrest versus early post-arrest cooling in the intervention group.
3. Duration of cardiac arrest.
4. Primary cardiac rhythm.

Sensitivity analysis

1. We planned to perform sensitivity analyses on the effect estimate by omitting studies with an overall 'high risk of bias' to assess the robustness of our estimates against within-study bias.
2. If several methods of neurological outcome assessment were used we planned to investigate the robustness of the effect estimates.

Summary of findings

We used the principles of the GRADE system ([Guyatt 2008](#)), in order to assess the quality of the body of evidence associated with the specific outcomes of survival and neurological outcome in our review, and we constructed a 'Summary of findings' table using the GRADE software. The GRADE approach appraises the quality of a body of evidence based on the extent to which one can be confident that an estimate of effect or association reflects the item being

assessed. The quality of a body of evidence considers within-study risk of bias (methodological quality), the directness of the evidence, heterogeneity of the data, precision of effect estimates, and risk of publication bias.

RESULTS

Description of studies

Results of the search

The systematic search of databases (from inception to March 2015) resulted in 1575 hits (duplicates excluded, see [Figure 1](#)). From

these, we excluded 1565 papers according to our eligibility criteria, leaving 10 papers to which we added one additional paper found in another review ([Callaway 2002](#)). From these remaining 11 papers we excluded three papers after further evaluation and discussion ([Belohlavek 2013](#); [Busch 2010](#); [Taccone 2010](#)); see [Characteristics of excluded studies](#). Seven completed studies ([Bernard 2010](#); [Bernard 2012](#); [Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)), and one ongoing study ([Nordberg 2013](#)), remained for inclusion in our review.

Figure 1. Study flow diagram.

Included studies

We included seven studies in our review ([Bernard 2010](#); [Bernard 2012](#); [Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)).

The basic points and an assessment of the quality criteria of the included studies can be found in [Characteristics of included studies](#).

Design

All studies were randomized, controlled, parallel-group trials.

Sample sizes

The sample sizes of the included studies were highly variable, with one small study ([Kämäräinen 2009](#) with 43 participants), five middle-sized studies ([Kim 2007](#) with 125 participants; [Bernard 2012](#) with 163 participants; [Castren 2010](#) with 200 participants; [Bernard](#)

2010 with 234 participants; and [Debaty 2014](#) with 245 participants), and one bigger study ([Kim 2014](#) with 1359 participants). Four studies were aimed primarily at establishing safety and feasibility or differences in inflammation markers and calculated their sample sizes accordingly ([Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#)).

Setting

All studies included participants with out-of-hospital cardiac arrest, hence including ambulance services ([Bernard 2010](#); [Bernard 2012](#); [Castren 2010](#); [Debaty 2014](#); [Kim 2007](#); [Kim 2014](#)), or helicopter services and staff ([Kämäräinen 2009](#)), or emergency departments of intensive or acute care units in urban areas. All were multicentre studies.

Participants

Inclusion criteria for all studies were heterogenous. Most studies included participants with out-of-hospital cardiac arrest due to all causes, regardless of the primary cardiac rhythm ([Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)). One study only included participants with a witnessed collapse ([Castren 2010](#)). The two studies by Bernard and colleagues differentiated between participants with ventricular fibrillation ([Bernard 2010](#)) and asystole and pulseless electrical activity (PEA) ([Bernard 2012](#)) as primary cardiac rhythms but were similar otherwise. In six studies temperature at study inclusion was reported and ranged between 35.2°C to 35.9°C ([Bernard 2010](#); [Bernard 2012](#); [Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#)) (see also [Characteristics of included studies](#)).

Interventions

Pre-hospital cooling was comparable in five studies ([Bernard 2010](#); [Bernard 2012](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)), using up to 2 L of cold fluids as administered after resuscitation. In contrast, [Castren 2010](#) and [Debaty 2014](#) began with the intervention already during resuscitation, with [Castren 2010](#) using intranasal cooling and [Debaty 2014](#) cold fluids and external cooling. However, the actual administration of cold fluid varied considerably in all studies. The pre-hospital patients' temperature was measured by tympanic probes ([Bernard 2010](#); [Bernard 2012](#); [Castren 2010](#)), and oesophageal/nasopharyngeal temperature probes ([Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)). In many studies a relevant proportion of participants received no cooling either in the pre-hospital group ([Bernard 2010](#); [Bernard 2012](#); [Kim 2007](#); [Kim 2014](#)), or during the hospital phase ([Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)). Most of the studies focused on feasibility and safety in the pre-hospital phase and there were no reported specific protocols for 'in-hospital cooling' ([Castren 2010](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)), which was left to the discretion of the treating physicians. These studies also did not provide further information on the initiation of cooling, the methods, cooling rates, cooling durations, and rewarming rates in the hospital phase. Only three out of seven studies provided temperature curves showing that a relevant proportion of the participants did not receive cooling according to guidelines at the time of conduct ([Bernard 2010](#); [Bernard 2012](#);

[Debaty 2014](#)), which limits the generalizability of the results. In summary we considered methodological heterogeneity to be a relevant limitation in this review.

Outcome

For most of the studies, the primary outcome parameters were feasibility and safety, the differences in patients' temperatures, or differences in serum concentration of inflammation markers on admission to the hospital ([Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#); [Kim 2007](#)). All studies additionally provided hospital mortality data and some simple neurological outcome parameters evaluated at discharge like "discharge either to home or to a rehabilitation facility" ([Bernard 2010](#); [Bernard 2012](#)), "severe neurologic deficits" ([Kim 2007](#)), "full recovery, mildly to moderately impaired, severely impaired, comatose, or dead" ([Kim 2014](#)), or the cerebral performance categories (CPC) score ([Castren 2010](#); [Debaty 2014](#); [Kämäräinen 2009](#)).

Excluded studies

We excluded three studies ([Belohlavek 2013](#); [Busch 2010](#); [Taccone 2010](#)) (see [Characteristics of excluded studies](#)). The study by [Belohlavek 2013](#) and colleagues was not considered for inclusion in this review because combinations of treatments were investigated, which makes the estimation of the isolated effect of a single component impossible. The abstracts by [Busch 2010](#) and [Taccone 2010](#) were separate presentations of their centre-specific patients as part of [Castren 2010](#), with no additional information. We linked these references to the respective study.

Ongoing studies

We found one ongoing study that would meet our eligible criteria ([Nordberg 2013](#)). It was started in 2010 and is currently recruiting participants. The estimated completion is December 2016. For further details see [Characteristics of ongoing studies](#).

Studies awaiting classification

There are no studies awaiting classification.

Risk of bias in included studies

Risk of bias across the individual studies was generally moderate ([Figure 2](#); [Figure 3](#); [Characteristics of included studies](#)). This was mainly driven by deficits in blinding the neurological outcome assessment and 'other biases', addressing the inconsistent administration of the intervention and the control. None of the studies was sufficiently designed to show equivalence or non-inferiority, but failed to prove superiority only. For the feasibility studies this might have been appropriate, however the bigger studies used effectiveness measures like neurological outcome and survival as primary outcomes with a similar design to the feasibility studies. Here the danger of misinterpretation is considerable. Pooling of the results of these studies would have been inappropriate and would have led to an invalid interpretation of the result.

Figure 2. 'Risk of bias' graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. 'Risk of bias' summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias): Good neurological outcome	Blinding of outcome assessment (detection bias): Survival	Blinding of outcome assessment (detection bias): Adverse events	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bernard 2010	+	+	?	+	+	?	+	+	-
Bernard 2012	+	+	?	+	+	?	+	+	-
Castren 2010	+	+	?	-	+	?	+	?	-
Debaty 2014	+	+	?	+	+	?	+	+	?
Kämäräinen 2009	+	+	?	-	+	?	+	+	-
Kim 2007	+	+	?	-	+	?	+	?	-
Kim 2014	+	?	?	+	+	?	+	+	-

Allocation

Risk of bias from inadequate allocation concealment was generally low. Only in one study was allocation concealment not described (Kim 2014).

Blinding

By nature in these trials blinding of staff to treatment allocation (in ambulances, helicopters, and hospitals) would have been difficult and was not done in any of the included studies. Blinding of outcome assessment was reported in four studies (Bernard 2010; Bernard 2012; Debaty 2014; Kim 2014). For all others the outcome assessors were not blinded or not reported to be. Survival may not be sensitive against possible information bias from unblinded outcome assessment. For the assessment of the neurological outcome, however, we classified unblinded assessment and missing information for this item as 'high risk'.

Incomplete outcome data

Three studies had complete outcome data for all randomized participants (Bernard 2010; Bernard 2012; Kim 2007). The remaining four reported some loss to follow-up but numbers were either too small to have a relevant impact or numbers and reasons for exclusion were balanced between groups. We therefore rated all studies as 'low risk' of bias for incomplete outcome data.

Three studies reported on loss to follow-up after randomization (Castren 2010; Kämäräinen 2009; Kim 2014) (see [Characteristics of included studies](#)). All studies gave reasons for the exclusions, which were either balanced between the study groups or the proportion of missing outcomes compared to the number of observed events was too low to have a clinically relevant impact on the intervention effect estimate.

Selective reporting

All expected outcomes were reported. We rated all studies 'low risk' for this item.

Other potential sources of bias

An inappropriate administration of the intervention and control might have resulted in an underestimation of a potential effect. In four studies only between 20% and 50% of all participants received the full intervention of 2000 mL of cold fluids (Bernard 2010; Bernard 2012; Kim 2007; Kim 2014). In the same studies up to 16% of all participants assigned to pre-hospital cooling did not receive any intervention at all. In all studies the continuation of cooling or the initiation of cooling in the hospital was left to the discretion of the treating physician. In three studies only between 61% and 77% of all participants received cooling in the hospital; no detailed information was provided for these participants (Kämäräinen 2009; Kim 2007; Kim 2014). Further, only three studies provided data on actual patient temperature during treatment (Bernard 2010; Bernard 2012; Debaty 2014). Judging from the graphs in two studies the temperatures of the intervention groups rewarmed shortly after hospital admission with an average temperature of above 34°C in about half of participants in both the intervention and control groups (Bernard 2010; Bernard 2012). This indicates that around 50% of the participants did not receive cooling to the guidelines of that time. One study required a target temperature of 34°C, on the upper limit of the recommended 32°C to 34°C (Castren 2010). For most other studies no further information on the application of in-hospital cooling or any temperature curves for their participants was provided, so it was not possible to determine if was applied appropriately in these studies (Castren 2010; Kämäräinen 2009; Kim 2007; Kim 2014).

Effects of interventions

See: [Summary of findings for the main comparison](#) Pre-hospital cooling compared to in-hospital cooling for survival, neuroprotection, and adverse events after out-of-hospital cardiac arrest

We present the results of seven randomized controlled trials for the primary endpoints of this review separately in [Figure 4](#) and [Figure 5](#).

Figure 4. Forest plot of comparison: 2 Survival: pre-hospital cooling versus in-hospital cooling, outcome: 2.1 Survival.

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias): Survival
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Figure 5. Forest plot of comparison: 1 Neurological outcome: pre-hospital cooling versus in-hospital cooling, outcome: 1.1 Good neurological outcome.

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias): Good neurological outcome
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Primary outcomes

1. Survival

We investigated short-term survival (closest to 30 days) and long-term survival (closest to six months) (see [Analysis 1.1](#)). There was considerable methodological heterogeneity and risk of bias, mainly due to deficits in the administration of cooling (see [Characteristics of included studies](#)). We have therefore refrained from pooling the results for survival. Insufficient information on six-month survival was available. Accordingly we did not perform sensitivity or subgroup analyses. Pre-hospital cooling did not appear to have an effect on survival compared with in-hospital cooling.

2. Neurological outcome

There was considerable methodological heterogeneity and risk of bias, mainly due to deficits in the administration of cooling (see [Characteristics of included studies](#)). We have therefore refrained from pooling the results for neurological outcome (see [Analysis 2.1](#)). Accordingly we did not perform sensitivity or subgroup analyses. Pre-hospital cooling did not appear to have an effect on neurological outcome compared with in-hospital cooling in any of the included studies.

Secondary outcomes

1. Adverse events, as reported by study authors

We pooled the available data on adverse events. Four studies reported on re-arrests after resuscitation ([Castren 2010](#); [Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)). These showed a higher incidence in the pre-hospital cooling group (risk ratio (RR) 1.23; 95% confidence interval (CI) 1.02 to 1.48; P value = 0.03). There was no effect on the incidence of pulmonary oedema (RR 1.02; 95% CI 0.67 to 1.57; P value = 0.77) ([Kämäräinen 2009](#); [Kim 2007](#); [Kim 2014](#)). See [Table 1](#).

2. Adverse events related to cooling methods, as reported by study authors

For device-related adverse events there was a higher incidence of nasal whitening with the intranasal cooling device used for intra-arrest cooling (RR 29.30; 95% CI 1.77 to 486.02; P value = 0.02, see [Table 2](#)) ([Castren 2010](#)). For all other reported adverse events there was no significant difference between pre-hospital and in-hospital cooling.

3. Quality of life, as reported by study authors

There were no data on quality of life.

4. Length of stay in the intensive care unit (ICU) and in the hospital, as proxies for economic outcomes

For ICU length of stay, [Castren 2010](#) reported no significant difference between the pre-hospital and in-hospital cooling group (eight days for the pre-hospital cooling participants versus 11 days for control participants; no standard deviation (SD) given).

For in-hospital length of stay, [Castren 2010](#) reported no significant difference between the pre-hospital and in-hospital cooling group (24.1 days for the pre-hospital cooling participants versus 26 days for control participants; no SD given). [Kim 2007](#) reported that the median length of stay was similar for the intervention and control group (12.2 days for the pre-hospital cooling participants versus 9.9 days for control participants; P value = 0.71). [Kim 2014](#) reported that the median length of stay was similar for the intervention and control groups among those with ventricular fibrillation (9.1 days for the pre-hospital cooling participants versus 9.4 days for control participants; P value = 0.75) and among those without ventricular fibrillation (11.8 days for the pre-hospital cooling participants versus 10.5 days for control participants; P value = 0.45).

Overall quality of the evidence

Using the GRADE approach we downgraded the overall quality of the evidence to 'very low', mainly due to a considerable amount of inconsistency, but also due to risk of bias within the studies and low precision (for details see the footnotes of [Summary of findings for the main comparison](#)).

DISCUSSION

Summary of main results

Evidence from seven studies in 2369 people contributing data to the primary outcomes of this review was insufficient to show that pre-hospital induction of cooling in comparison to in-hospital cooling improved survival or neurological outcome after out-of-hospital cardiac arrest. Studies indicated a slightly increased number of re-arrests with the application of pre-hospital hypothermia.

However, relevant heterogeneity in terms of methods, interventions, and cohorts, as well as risk of bias in these seven included randomized controlled trials was too large to present a summary estimate of the effect of pre-hospital cooling on survival and neurological outcome. Overall, pre-hospital cooling did not appear to have an effect on neurological outcome and survival compared to in-hospital cooling.

Significant shortcomings in these studies make any inferences about any efficacy outcome difficult. For some studies the focus and primary endpoint was strictly on the pre-hospital phase so not much attention was given to the continuation of the cooling therapy. As a consequence some participants rewarmed after admission, and a significant number of participants did not receive any continuous cooling therapy.

Nonetheless, the intervention had an effect on procedural outcomes. In all studies the application of cooling in the pre-hospital phase by the ambulance and helicopter staff resulted in a significantly lower body temperature on admission when compared to the control group.

Overall completeness and applicability of evidence

Unfortunately, the included studies had major heterogeneity in the intervention and control group treatments. The focus of the majority of the studies was put on the pre-hospital phase; accordingly no common cooling management was in place across all studies. For the pre-hospital phase cooling in some studies was not applied in the prespecified manner or was omitted. In-hospital cooling was not required in all feasibility studies and in a relevant number of participants it was not initiated. This does not reflect common practice and adherence to cooling guidelines. Some studies did not document any data on the conduct of in-hospital cooling preventing an assessment of whether the therapy was applied in an effective way and according to any cooling guidelines at that time. Another main source of heterogeneity stems from the incorporation of intra-arrest and post-arrest pre-hospital cooling, which is evidenced by the large differences in outcome risks in the control groups. Additionally, common practice varies considerably across countries, likewise reflecting the uncertainty of the currently available evidence.

In 2013 one study was published that compared the effectiveness of temperature management at 33°C to temperature management at 36°C (Nielsen 2013). The authors found no difference in neurological outcome and survival. In the context of this review this result challenges the question of whether cooling to 32°C to 34°C is effective in the first place. Currently, there is no clear answer and this has been extensively discussed in the scientific community and by guidelines panels. The most recent guidelines have picked up on the Nielsen 2013 study and stated that temperature control between 32°C and 36°C is recommended, "whether certain

subpopulations of cardiac arrest patients may benefit from lower (32–34°C) or higher (36°C) temperatures remains unknown, and further research may help elucidate this" (Callaway 2015; Soar 2015). This may reflect on the fact that in some ways the Nielsen 2013 study differs from previous efficacy studies on hypothermia. Firstly, Nielsen 2013 was a pragmatic study (multicentre, different methods of cooling etc.), which may not be suitable as long as the cooling 'dose', which is mainly target temperature and cooling duration, is not unequivocally characterized in proof of concept studies. Additionally, Nielsen 2013 had a very short duration from collapse to resuscitation (on average one minute) compared to the other trials (around 10 minutes). This is of relevance since there are reports showing that the effect of cooling may depend this no-flow time (Testori 2012), and the very short times are unrealistic in many countries.

However, if heterogeneity between previous studies and the Nielsen 2013 study were ignored and they were put together, targeted temperature management may still improve neurological outcome by more than 50% (Arrich 2016).

Quality of the evidence

Risk of bias within the seven individual studies was generally moderate (see Figure 2; Figure 3; Characteristics of included studies). An exception was 'blinding of outcome assessment', which was absent in almost half of all studies and may have substantially biased outcomes like neurological state. 'Other sources of bias' (see Characteristics of included studies) formed the other exception and was the reason why we have not pooled the results. Using the GRADE approach we downgraded the quality of the overall evidence to 'very low' (see Summary of findings for the main comparison). This was mainly driven by a relevant amount of inconsistency due to the inconsistent application of the intervention and control, which significantly reduced confidence in the results and opens up the possibility that future studies with a more rigorous study conduct may have a different result. Adding to the inconsistency were the different study populations and the two different time points of intervention (intra-arrest and post-arrest). Imprecision also led to a further downgrading of the evidence as most studies were small, with three out of seven studies being feasibility or pilot studies (Castren 2010; Kämäräinen 2009; Kim 2007). The pre-hospital participants' temperature was measured by tympanic probes (Bernard 2010; Bernard 2012; Castren 2010), and oesophageal/nasopharyngeal temperature probes (Kämäräinen 2009; Kim 2007; Kim 2014), the former having been shown to be the least reliable method for temperature measurement during the cooling phase (Krizanac 2013).

In the absence of summary estimates for the primary outcome we are left with the low precision of the single study estimates.

Potential biases in the review process

We have strived to find all comparable studies in this field and presented their results separately, together with assessment of their methodology, strengths, and limitations, to give the most comprehensive information on the question of the effectiveness of pre-hospital cooling. We cannot exclude potential reporting bias due to the limited number of studies available. However, as is known from empirical evidence, reporting bias is usually driven by statistical significance and positive results (Hopewell 2009). It

is noteworthy that in our review we only identified non-significant studies.

Agreements and disagreements with other studies or reviews

We found four systematic reviews, which are at least partly comparable to our review. Two included meta-analyses, whereas another two abstained from presenting summary estimates.

Cullen and colleagues published a review and meta-analysis comparing pre-hospital cooling to a later induction of cooling (Cullen 2011). Up to 2011 they found the same four studies as in our review (Bernard 2010; Castren 2010; Kämäräinen 2009; Kim 2007). They pooled all data and found no difference between the two groups but they were not specific in the description of the outcome and did not present any evaluation of clinical heterogeneity or possible sources of bias in the included studies. They concluded, however, that cooling in the pre-hospital setting is feasible, but that it is unclear if it is beneficial in the long term, including for improving neurological outcomes.

Diao and colleagues presented a systematic review and meta-analysis comparing pre-hospital cooling to in-hospital cooling or no cooling (Diao 2013). They pooled five studies, which are all included in our review (Bernard 2010; Bernard 2012; Castren 2010; Kämäräinen 2009; Kim 2007). They found that pre-hospital cooling after cardiac arrest resulted in significantly lower body temperature on hospital admission. No differences were observed in survival to hospital discharge, favourable neurological outcome at hospital discharge, or re-arrests. The overall quality of the included studies was graded as very low and this coincides with our methodological and clinical heterogeneity assessment.

Scolletta and colleagues summarized studies on intra-arrest cooling (Scolletta 2012). Among the available randomized controlled trials they only included the study by Castren 2010 and did not otherwise pool data.

Cabanas conducted a systematic review on all available studies (randomized or not) of pre-hospital cooling in comparison to normothermia or later induction of cooling (Cabanas 2011). Among the studies from our systematic review they only included Kim 2007. They did not pool the data but concluded that cooling can

be efficiently induced in the pre-hospital environment. Further, it was stated that more research would be needed to understand the effectiveness and optimal timing of early cooling.

AUTHORS' CONCLUSIONS

Implications for practice

Currently, the overall quality of the available evidence on the effects of pre-hospital cooling on survival and neuroprotection is very low. There is no convincing evidence to clearly delineate beneficial or harmful effects of pre-hospital induction of cooling in comparison to in-hospital cooling. The currently available studies suggest that pre-hospital cooling induction is feasible to lower body temperature on hospital admission, but it may also increase the risk of cardiac re-arrest. We do not have sufficient evidence to determine the effects of other determinants (duration, target temperature, etc.) of cooling from this review.

Implications for research

Previous trials have focused on the feasibility and safety of pre-hospital cooling, or have not rigorously applied nor reliably controlled either form of targeted temperature management. This resulted in considerable heterogeneity in the pre-hospital phase but even more in the in-hospital phase, which made it impossible to pool the outcome data from the single studies. Future trials should tackle these shortcomings, include the full initial 36 hours in the treatment protocols, and evaluate at least neurological outcome and survival at six months. One ongoing trial that compares pre-hospital intra-arrest transnasal evaporative cooling with standard targeted temperature management in the hospital, with neurological intact survival as the primary outcome parameter, would fit our inclusion criteria and is currently recruiting participants (Nordberg 2013).

ACKNOWLEDGEMENTS

We would like to thank Nicola Petrucci (content editor), Vibeke E Horstmann (statistical editor), Shu Ming Pan, Charles D Deakin, Jasmeet Soar (peer reviewers), and Roy Buffery (consumer referee) for their help and editorial advice during the preparation of this systematic review.

REFERENCES

References to studies included in this review

Bernard 2010 {published data only}

Bernard SA, Smith K, Cameron P, Masci K, Taylor DM, Cooper DJ, et al. Rapid Infusion of Cold Hartmanns (RICH) Investigators. Induction of therapeutic hypothermia by paramedics after resuscitation from out-of-hospital ventricular fibrillation cardiac arrest: a randomized controlled trial. *Circulation* 2010;**122**(7):737-42. [PUBMED: 20679551]

Bernard 2012 {published data only}

Bernard SA, Smith K, Cameron P, Masci K, Taylor DM, Cooper DJ, et al. Rapid Infusion of Cold Hartmanns Investigators. Induction of prehospital therapeutic hypothermia after resuscitation from nonventricular fibrillation cardiac arrest*. *Critical Care Medicine* 2012;**40**(3):747-53. [PUBMED: 22020244]

Castren 2010 {published data only}

Busch HJ, Eichwede F, Födisch M, Taccone FS, Wöbker G, Schwab T, et al. Safety and feasibility of nasopharyngeal evaporative cooling in the emergency department setting in survivors of cardiac arrest. *Resuscitation* 2010;**81**(8):943-9. [PUBMED: 20627524]

* Castrén M, Nordberg P, Svensson L, Taccone F, Vincent JL, Desruelles D, et al. Intra-arrest transnasal evaporative cooling: a randomized, prehospital, multicenter study (PRINCE: Pre-ROSC IntraNasal cooling effectiveness). *Circulation* 2010;**122**(7):729-36. [PUBMED: 20679548]

Taccone FS, Vincent JL. Trans-nasal cooling during CPR: a single-center experience. *Critical Care* 2010;**14**(Suppl 1):S107-8. [DOI: [10.1186/cc8551](https://doi.org/10.1186/cc8551)]

Debaty 2014 {published data only}

Debaty G, Maignan M, Savary D, Koch F-X, Ruckly S, Durand M, et al. Impact of intra-arrest therapeutic hypothermia in outcomes of prehospital cardiac arrest: a randomized controlled trial. *Intensive Care Medicine* 2014;**40**(12):1832-42. [PUBMED: 25348858]

Kämäräinen 2009 {published data only}

Kämäräinen A, Virkkunen I, Tenhunen J, Yli-Hankala A, Silfvast T. Prehospital therapeutic hypothermia for comatose survivors of cardiac arrest: a randomized controlled trial. *Acta Anaesthesiologica Scandinavica* 2009;**53**(7):900-7. [PUBMED: 19496762]

Kim 2007 {published data only}

Kim F, Olsufka M, Longstreth WT Jr, Maynard C, Carlbom D, Deem S, et al. Pilot randomized clinical trial of prehospital induction of mild hypothermia in out-of-hospital cardiac arrest patients with a rapid infusion of 4 degrees C normal saline. *Circulation* 2007;**115**(24):3064-70. [PUBMED: 17548731]

Kim 2014 {published data only}

Kim F, Nichol G, Maynard C, Hallstrom A, Kudenchuk PJ, Rea T, et al. Effect of prehospital induction of mild hypothermia on survival and neurological status among adults with cardiac

arrest: a randomized clinical trial. *JAMA* 2014;**311**(1):45-52. [PUBMED: 24240712]

References to studies excluded from this review

Belohlavek 2013 {published data only}

Belohlavek J, Franek O, Pokorna M, Danda J, Kandrnal V, Balik M, et al. Hyperinvasive approach to out-of hospital cardiac arrest using mechanical chest compression device, prehospital intraarrest cooling, extracorporeal life support and early invasive assessment compared to standard of care. A randomized parallel groups comparative study. *JACC. Cardiovascular Interventions* 2013;**6**(2 Suppl 1):S41. [DOI: [10.1016/j.jcin.2013.01.050](https://doi.org/10.1016/j.jcin.2013.01.050)]

Busch 2010 {published data only}

Busch HJ, Eichwede F, Födisch M, Taccone FS, Wöbker G, Schwab T, et al. Safety and feasibility of nasopharyngeal evaporative cooling in the emergency department setting in survivors of cardiac arrest. *Resuscitation* 2010;**81**(8):943-9. [PUBMED: 20627524]

Taccone 2010 {published data only}

Taccone FS, Vincent JL. Trans-nasal cooling during CPR: a single-center experience. *Critical Care* 2010;**14**(1):P319. [DOI: [10.1186/cc8551](https://doi.org/10.1186/cc8551)]

References to ongoing studies

Nordberg 2013 {published data only}

Nordberg P, Taccone FS, Castren M, Truhlar A, Desruelles D, Forsberg S, et al. Design of the PRINCESS trial: pre-hospital resuscitation intra-nasal cooling effectiveness survival study (PRINCESS). *BMC Emergency Medicine* 2013;**13**(1):21.

Additional references

Abella 2004

Abella BS, Zhao D, Alvarado J, Hamann K, Vanden Hoek TL, Becker LB. Intra-arrest cooling improves outcomes in a murine cardiac arrest model. *Circulation* 2004;**109**(22):2786-91. [PUBMED: 15159295]

Abella 2008

Abella BS, Zhao D, Alvarado J, Hamann K, Vanden Hoek TL, Becker LB. Intra-arrest cooling improves outcomes in a murine cardiac arrest model. *Circulation* 2008;**109**(22):2786-91. [PUBMED: 15159295]

Arrich 2016

Arrich J, Holzer M, Havel C, Mullner M, Herkner H. Hypothermia for neuroprotection in adults after cardiopulmonary resuscitation. *The Cochrane database of systematic reviews* 2016;**2**:CD004128. [PUBMED: 26878327]

Berdowski 2010

Berdowski J, Berg RA, Tijssen JG, Koster RW. Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation* 2010;**81**(11):1479-87. [PUBMED: 20828914]

Bernard 2002

Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *New England Journal of Medicine* 2002;**346**(8):557-63. [MEDLINE: 11856794]

Cabanas 2011

Cabanas JG, Brice JH, De Maio VJ, Myers B, Hinchey PR. Field-induced therapeutic hypothermia for neuroprotection after out-of-hospital cardiac arrest: a systematic review of the literature. *Journal of Emergency Medicine* 2011;**40**(4):400-9. [PUBMED: 20850254]

Callaway 2002

Callaway CW, Tadler SC, Katz LM, Lipinski CL, Brader E. Feasibility of external cranial cooling during out-of-hospital cardiac arrest. *Resuscitation* 2002;**52**:159-65. [PUBMED: 11841883]

Callaway 2015

Callaway CW, Soar J, Aibiki M, Bottiger BW, Brooks SC, Deakin CD, et al. Part 4: Advanced Life Support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2015;**132**(16 Suppl 1):S84-S145. [PUBMED: 26472860]

Che 2011

Che D, Li L, Kopil CM, Liu Z, Guo W, Neumar RW. Impact of therapeutic hypothermia onset and duration on survival, neurologic function, and neurodegeneration after cardiac arrest. *Critical Care Medicine* 2011;**39**(6):1423-30. [PUBMED: 21610611]

Colbourne 1995

Colbourne F, Corbett D. Delayed postischemic hypothermia: a six month survival study using behavioral and histological assessments of neuroprotection. *Journal of Neuroscience* 1995;**15**(11):7250-60. [PUBMED: 7472479]

Cullen 2011

Cullen D, Augenstine D, Kaper L, Tinkham S, Utz D. Therapeutic hypothermia initiated in the pre-hospital setting: a meta-analysis. *Advanced Emergency Nursing Journal* 2011;**33**(4):314-21. [PUBMED: 22075682]

Cummins 1991

Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation* 1991;**84**:960-75. [PUBMED: 1860248]

Diao 2013

Diao M, Huang F, Guan J, Zhang Z, Xiao Y, Shan Y, et al. Prehospital therapeutic hypothermia after cardiac arrest: a systematic review and meta-analysis of randomized controlled trials. *Resuscitation* 2013;**84**(8):1021-8. [PUBMED: 23454259]

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**:629-34. [MEDLINE: 9310563]

Guyatt 2008

Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ. What is "quality of evidence" and why is it important to clinicians?. *BMJ (Clinical research ed.)* 2008;**336**(7651):995-8. [PUBMED: 18456631]

HACA 2002

Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *New England Journal of Medicine* 2002;**346**(8):549-56. [MEDLINE: 11856793]

Hachimi-Idrissi 2001

Hachimi-Idrissi S, Corne L, Ebinger G, Michotte Y, Huyghens L. Mild hypothermia induced by a helmet device: a clinical feasibility study. *Resuscitation* 2001;**51**(3):275-81. [MEDLINE: 11738778]

Haugk 2011

Haugk M, Testori C, Sterz F, Uranitsch M, Holzer M, Behringer W, et al. Time to Target Temperature Study Group. Relationship between time to target temperature and outcome in patients treated with therapeutic hypothermia after cardiac arrest. *Critical Care (London, England)* 2011;**15**(2):R101. [PUBMED: 21439038]

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557-60. [MEDLINE: 12958120]

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Holzer 2010

Holzer M. Targeted temperature management for comatose survivors of cardiac arrest. *New England Journal of Medicine* 2010;**363**(13):1256-64. [PUBMED: 20860507]

Hopewell 2009

Hopewell S, Loudon K, Clarke M, Oxman AD, Dickersin K. Publication bias in clinical trials due to statistical significance or direction of trial results. *Cochrane Database of Systematic Reviews* 2009; Vol. 1. [DOI: [10.1002/14651858.MR000006.pub3](https://doi.org/10.1002/14651858.MR000006.pub3)]

Janata 2010

Janata A, Weihs W, Schratter A, Bayegan K, Holzer M, Frossard M, et al. Cold aortic flush and chest compressions enable good neurologic outcome after 15 mins of ventricular fibrillation in cardiac arrest in pigs. *Critical Care Medicine* 2010;**38**(8):1637-43. [PUBMED: 20543671]

Jennett 1975

Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet* 1975;**1**:480-4. [MEDLINE: 46957]

Krizanac 2013

Krizanac D, Stratil P, Hoerburger D, Testori C, Wallmueller C, Schober A, et al. Femoro-iliac artery versus pulmonary artery core temperature measurement during therapeutic hypothermia: an observational study. *Resuscitation* 2013;**84**(6):805-9. [PUBMED: 23200998]

Kuboyama 1993

Kuboyama K, Safar P, Radovsky A, Tisherman SA, Stezoski SW, Alexander H. Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: a prospective, randomized study. *Critical Care Medicine* 1993;**21**(9):1348-58. [PUBMED: 8370299]

Nichol 2008

Nichol G, Thomas E, Callaway CW, Hedges J, Powell JL, Aufderheide TP, et al. Resuscitation Outcomes Consortium Investigators. Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA* 2008;**300**(12):1423-31. [PUBMED: 18812533]

Nielsen 2009

Nielsen N, Hovdenes J, Nilsson F, Rubertsson S, Stammet P, Sunde K, et al. Hypothermia Network. Outcome, timing and adverse events in therapeutic hypothermia after out-of-hospital cardiac arrest. *Acta Anaesthesiologica Scandinavica* 2009;**53**(7):926-34. [PUBMED: 19549271]

Nielsen 2013

Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, et al. TTM Trial Investigators. Targeted temperature management at 33°C versus 36°C after cardiac arrest. *New England Journal of Medicine* 2013;**369**(23):2197-206. [PUBMED: 24237006]

Nolan 2008

Nolan JP, Neumar RW, Adrie C, Aibiki M, Berg RA, Böttiger BW, et al. Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A Scientific Statement from the International Liaison Committee on Resuscitation; the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; the Council on Stroke. *Resuscitation* 2008;**79**(3):350-79. [PUBMED: 18963350]

RevMan 5.3 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Scolletta 2012

Scolletta S, Taccone FS, Nordberg P, Donadello K, Vincent JL, Castren M. Intra-arrest hypothermia during cardiac arrest: a systematic review. *Critical Care (London, England)* 2012;**16**(2):R41. [PUBMED: 22397519]

Sendelbach 2012

Sendelbach S, Hearst MO, Johnson PJ, Unger BT, Mooney MR. Effects of variation in temperature management on cerebral performance category scores in patients who received therapeutic hypothermia post cardiac arrest. *Resuscitation* 2012;**83**(7):829-34. [PUBMED: 22230942]

Soar 2015

Soar J, Callaway CW, Aibiki M, Bottiger BW, Brooks SC, Deakin CD, et al. Part 4: Advanced life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2015;**95**:e71-e120. [PUBMED: 26477429]

Sterz 1991

Sterz F, Safar P, Tisherman S, Radovsky A, Kuboyama K, Oku K. Mild hypothermic cardiopulmonary resuscitation improves outcome after prolonged cardiac arrest in dogs. *Critical Care Medicine* 1991;**19**(3):379-89. [PUBMED: 1999100]

Suffoletto 2009

Suffoletto B, Peberdy MA, van der Hoek T, Callaway C. Body temperature changes are associated with outcomes following in-hospital cardiac arrest and return of spontaneous circulation. *Resuscitation* 2009;**80**(12):1365-70. [PUBMED: 19804929]

Testori 2012

Testori C, Sterz F, Holzer M, Losert H, Arrich J, Herkner H, et al. The beneficial effect of mild therapeutic hypothermia depends on the time of complete circulatory standstill in patients with cardiac arrest. *Resuscitation* 2012;**83**(5):596-601. [PUBMED: 22138057]

Wolff 2009

Wolff B, Machill K, Schumacher D, Schulzki I, Werner D. Early achievement of mild therapeutic hypothermia and the neurologic outcome after cardiac arrest. *International Journal of Cardiology* 2009;**133**(2):223-8. [PUBMED: 18353458]

Yannopoulos 2009

Yannopoulos D, Zviman M, Castro V, Koldaivelu A, Ranjan R, Wilson RF, et al. Intra-cardiopulmonary resuscitation hypothermia with and without volume loading in an ischemic model of cardiac arrest. *Circulation* 2009;**120**(14):1426-35. [PUBMED: 19770397]

Zhao 2008

Zhao D, Abella BS, Beiser DG, Alvarado JP, Wang H, Hamann KJ, et al. Intra-arrest cooling with delayed reperfusion yields

higher survival than earlier normothermic resuscitation in a mouse model of cardiac arrest. *Resuscitation* 2008;**77**(2):242-9. [PUBMED: 18096292]

and neuroprotection after out-of-hospital cardiac arrest. *Cochrane Database of Systematic Reviews* 2013, Issue 6. [DOI: [10.1002/14651858.CD010570](https://doi.org/10.1002/14651858.CD010570)]

References to other published versions of this review

* Indicates the major publication for the study

Arrich 2013

Arrich J, Havel C, Holzer M, Herkner H. Prehospital versus in-hospital initiation of mild therapeutic hypothermia for survival

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bernard 2010

Methods	Parallel-group, randomized trial, October 2005 to November 2007, multicentre study
Participants	<p>Total number of participants 234, mean age 63 years, 15% female, patients' temperature on arrival of resuscitation team: pre-hospital cooling group 35.9°C, hospital cooling group: 35.8°C</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • out-of-hospital cardiac arrest • cause of cardiac arrest: any • primary rhythm: ventricular fibrillation • return of spontaneous circulation with a systolic blood pressure 90 mmHg (with epinephrine infusion if needed) • total cardiac arrest time > 10 minutes • age older than 14 years • intravenous access available <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • dependent on others for activities of daily living • already hypothermic (temperature < 34°C) • females who were obviously pregnant • cardiac arrest after trauma
Interventions	<ul style="list-style-type: none"> • Pre-hospital cooling: up to 2000 mL ice-cold lactated Ringer's solution, given after resuscitation, n = 118 • Control: in-hospital cooling, n = 116
Outcomes	<ul style="list-style-type: none"> • Primary: favourable outcome defined as discharge either to home or to a rehabilitation facility • Secondary: core (bladder or oesophageal) temperatures at hospital arrival, the development of pre-hospital pulmonary oedema, and recurrent pre-hospital cardiac arrest after enrolment
Funding	Funding: funding for the study was provided by grants from the Australian National Health and Medical Research Council (Drs Bernard, Cameron, Taylor, Cooper, Kelly, and Silvester) and the National Heart Foundation of Australia (Drs Bernard and Smith).
Declarations of interest	None
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Bernard 2010 (Continued)

Random sequence generation (selection bias)	Low risk	The envelope allocation was computer-randomized and allocated in blocks of 10 to each intensive care paramedic unit
Allocation concealment (selection bias)	Low risk	The treating intensive care paramedics randomized eligible participants by opening an opaque, sealed envelope that indicated treatment allocation
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating ambulance and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	Low risk	Before hospital discharge, conscious participants were evaluated by a rehabilitation physician who was unaware of the study allocation
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomized participants analysed for the primary outcome
Selective reporting (reporting bias)	Low risk	All expected outcomes included
Other bias	High risk	<ul style="list-style-type: none"> From 398 eligible participants 164 were not enrolled; no reasons could be determined. Of the 118 participants allocated to pre-hospital cooling only 48% received the full 2000 ml, 11 received 1500 to 2000 ml, 37 received 1000 to 1500 ml, 5 received 500 to 1000 ml, and 8 (7%) received no ice-cold fluid. Participants in the intervention group rewarmed after admission to hospital, which reduced the actual temperature difference between the 2 participants groups to around 1°C for up to 1 hour. No reasons or discussion were given by authors. The prespecified target temperature of 33°C could not be reached for a relevant number of participants. The mean target temperature presented for both groups was around 34°C, which implies that 50% of all participants did not reach the 32°C to 34°C recommended by the then current guidelines.

Bernard 2012

Methods	Parallel-group, randomized trial, October 2005 to November 2007, multicentre study
Participants	<p>Total number of participants 163, mean age 62 years, 36% female, patients' average temperature on arrival of resuscitation team: pre-hospital cooling group 35.9°C, hospital cooling group: 35.8°C</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> out-of-hospital cardiac arrest cause of cardiac arrest: any

Bernard 2012 (Continued)

- primary rhythm: asystole or PEA
- return of spontaneous circulation with a systolic blood pressure 90 mmHg (with epinephrine infusion if needed)
- total cardiac arrest time > 10 mins
- age older than 14 years
- intravenous access available

Exclusion criteria:

- dependent on others for activities of daily living
- already hypothermic (temperature < 34°C)
- females who were obviously pregnant
- cardiac arrest after trauma

Interventions	Pre-hospital cooling: 40 mL/kg and up to 2000 mL ice-cold lactated Hartmann solution, given after re-suscitation, n = 82 Control: in-hospital cooling, n = 81
Outcomes	<ul style="list-style-type: none"> • Favourable outcome defined as discharge either to home or to a rehabilitation facility • Core (bladder or oesophageal) temperatures at hospital arrival, the development of pre-hospital pulmonary oedema, and recurrent pre-hospital cardiac arrest after enrolment
Funding	Supported by the Australian National Health and Medical Research Committee
Declarations of interest	None
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-randomized and allocated in blocks of 10 (5 to pre-hospital cooling and 5 to hospital cooling) to each intensive care paramedic unit rating
Allocation concealment (selection bias)	Low risk	Opaque, sealed envelopes. The envelope allocation was computer-randomized and allocated in blocks of 10 to each intensive care paramedic unit.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating ambulance and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	Low risk	Before hospital discharge, conscious participants were evaluated by a rehabilitation physician who was unaware of the study allocation
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated

Bernard 2012 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomized participants analysed for primary outcome
Selective reporting (reporting bias)	Low risk	All expected outcomes included
Other bias	High risk	<ul style="list-style-type: none"> From 309 eligible participants 146 were not enrolled; no reasons could be determined. Of 82 participants assigned to pre-hospital cooling, only 36 (44%) of all participants in the intervention group received the full intervention of 2 L; 13 participants (16%) received no intervention at all. Participants in the intervention group rewarmed after admission to hospital, which reduced the actual temperature difference between the 2 groups. The prespecified target temperature of 33°C could not be reached for a relevant number of participants. The mean target temperature presented for both groups was around 34°C, which implies that 50% of all participants did not reach the 32°C to 34°C recommended by the then current guidelines.

Castren 2010

Methods	Parallel-group, randomized trial, November 2008 to June 2009, multicentre study
Participants	<p>Total number of participants 200, mean age: intervention 66, control 64 years, 27% female, patients' average temperature at ROSC: pre-hospital cooling group 35.5°C, hospital cooling group: 35.8°C</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> out-of-hospital cardiac arrest cause of cardiac arrest: any primary rhythm: any witnessed cardiac arrest CPR initiated within 20 minutes of collapse age older than 17 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> very old participants and/or "do-not-attempt-resuscitation" order intranasal obstruction cardiac arrest after trauma and drug overdose participants with cerebrovascular accidents known coagulopathy asphyxia or requirement for oxygen support electrocution
Interventions	<p>Intra-arrest cooling with RhinoChill device, continued in-hospital according to institutional standards</p> <p>Control: in-hospital cooling according to institutional standards</p>
Outcomes	<ul style="list-style-type: none"> Cooling rates (i.e. temperature at ROSC and on hospital arrival) Time to target temperature of 34°C) ROSC rate Survival to discharge Survival with good neurological outcome (CPC 1 or 2) at hospital discharge

Castren 2010 (Continued)

- Adverse events

Funding	This work was supported by BeneChill, Inc, San Diego, California
Declarations of interest	Dr Barbut is the founder and Chairman of BeneChill. She participated in study design, data analysis, and writing of the manuscript.
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The randomization assignments were generated under a randomized permuted-block design, with block sizes of 8, in a 1:1 allocation
Allocation concealment (selection bias)	Low risk	Sequentially numbered, sealed envelopes that contained single randomization assignments
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating ambulance and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	High risk	The discharge assessment may not always have been performed by an individual blinded to the treatment group
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	7 of 200 participants lost to follow-up (4 from intervention, 3 from control group), due to misclassification of eligibility criteria and transport to non-participating hospital. Reasons and numbers are balanced and the proportion of missing outcomes compared with the observed event risk is not enough to have a clinically relevant impact on the intervention effect estimate.
Selective reporting (reporting bias)	Unclear risk	Presentation of data included only participants with acquired ROSC (after randomization)
Other bias	High risk	<ul style="list-style-type: none"> • Target temperature only 34°C (guidelines recommend 33°C to 34°C), which may have reduced overall treatment effect • No information on the conduct of temperature management in the hospital

Debaty 2014

Methods	Parallel-group, randomized trial, February 2009 to August 2012, multicentre study
---------	---

Debaty 2014 (Continued)

Participants	<p>Total number of participants 245, mean age 67 years, 29% female, patients' average temperature at randomization: pre-hospital cooling group 35.2°C, hospital cooling group: 35.4°C (estimated from graph)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> OHCA participants over 18 years of age eligible for advanced life support were included irrespective of rhythm <p>Exclusion criteria:</p> <ul style="list-style-type: none"> participants with trauma participants with haemorrhage participants with asphyxia participants already hypothermic (temperature < 34°C) women who were obviously pregnant participants who had achieved ROSC before randomization participants with a do-not-attempt resuscitation order
Interventions	Intra-arrest cooling, infusion of up to 2000 mL of ice-cold 0.9% saline solution at 100 mL/min during cardiac arrest by use of a standard infusion set and a pressure bag inflated to 300 mmHg. Surface cooling was also induced using gel pads.
Outcomes	<p>Types of outcome measures:</p> <p>Primary outcomes</p> <ul style="list-style-type: none"> neuron specific enolase (NSE) at 24 hours <p>Secondary outcomes</p> <ul style="list-style-type: none"> IL-6 concentrations during the first 72 h IL-8 concentrations during the first 72 h IL-10 concentrations during the first 72 h cooling rates ROSC rate length of stay in intensive care survival (discharge, 30 days and 1 year) neurological outcome comparing individual CPC scores (hospital discharge and 30 days)
Funding	French Society of Emergency Medicine
Declarations of interest	None
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization assignments were generated under a randomized permuted-block design, with block sizes of 4, in a 1:1 allocation
Allocation concealment (selection bias)	Low risk	Each mobile intensive care unit was given sequentially numbered, sealed envelopes containing single randomization assignments

Debaty 2014 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) Good neurological outcome	Low risk	Neurological outcome at hospital discharge and 30 days was assessed by a physician blinded to the study allocation
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	Analysis of clinical outcomes was according to intention-to-treat
Selective reporting (reporting bias)	Low risk	All expected outcomes included
Other bias	Unclear risk	No information on application of pre-hospital cold fluids (which proportion of patients received how much) Over 24 hours of cooling the body temperature of the intra-arrest cooling group seemed to be higher than the body temperature of the hospital cooling group According to the authors the study was not powered to show a clinical difference

Kim 2007

Methods	Parallel-group, randomized trial, November 2004 to February 2006, multicentre study
Participants	Total number of participants 125, mean age 66 years, 30% female, patients' average temperature at randomization: pre-hospital cooling group 35.8°C, hospital cooling group: 35.5°C Inclusion criteria: <ul style="list-style-type: none"> • out-of-hospital cardiac arrest • cause of cardiac arrest: any • primary rhythm: any • unresponsive • return of spontaneous circulation with palpable pulses • age older than 17 years • intubated and intravenous access available • oesophageal temperature probe Exclusion criteria: <ul style="list-style-type: none"> • already hypothermic (temperature < 34°C)

Kim 2007 (Continued)

- cardiac arrest after trauma

Interventions	Pre-hospital induction of cooling (32°C to 34°C); up to 2 L of 4°C normal saline after resuscitation Control: standard care with and without hypothermia
Outcomes	Primary: <ul style="list-style-type: none"> • temperature differences field-hospital Secondary: <ul style="list-style-type: none"> • re-arrests after randomization • deaths before hospital admission • in-hospital death • neurological outcome (severe neurological deficit) • time to awakening, discharge, death • safety data
Funding	This work was supported by a grant from the Medic One Foundation and National Institutes of Health grant HL04346 (F.K.)
Declarations of interest	None
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"...balanced blocks of 4"
Allocation concealment (selection bias)	Low risk	Paramedics called the emergency department physician at Harborview Medical Center to verify eligibility and to learn treatment assignment. The emergency room physician opened sequentially numbered envelopes that randomized participants to either receive or not receive cooling.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating ambulance and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	High risk	Study personnel during data collection and analysis could not be entirely unaware of treatment assignment
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias)	Low risk	All randomized participants analysed for primary outcome

Kim 2007 (Continued)

All outcomes

Selective reporting (reporting bias)	Unclear risk	Of participants discharged alive, only severe neurological deficit was reported, but not in detail (neurological scores missing)
Other bias	High risk	<ul style="list-style-type: none"> Of all 63 participants assigned to pre-hospital cooling only 12 (20%) received full 2 L, 37 participants received between 500 mL and 2 L, 6 participants received 500 mL, 8 participants (13%) did not receive any fluid Primary endpoint of study was temperature differences on admission and not mortality Only 61% of all participants received in-hospital cooling; it is unclear how many in which group No information on the conduct of temperature management in the hospital

Kim 2014

Methods	Parallel-group, randomized trial, December 2007 to December 2012, multicentre study	
Participants	<p>Total number of participants 1359, mean age VF: 62; non-VF: 68, 36% female, baseline average temperature of cooling groups not reported</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> out-of-hospital cardiac arrest cause of cardiac arrest: any primary rhythm: any unresponsive return of spontaneous circulation with palpable pulses age older than 17 years intubated and intravenous access available oesophageal temperature probe <p>Exclusion criteria:</p> <ul style="list-style-type: none"> already hypothermic (temperature < 34°C) cardiac arrest after trauma 	
Interventions	<p>Pre-hospital cooling: up to 2L of 4°C saline after resuscitation</p> <p>Control: standard care alone (control) or standard care plus induction of cooling(intervention)</p>	
Outcomes	<ul style="list-style-type: none"> Survival to discharge, full recovery, mildly to moderately impaired, severely impaired, comatose, or dead Neurological status at discharge Safety data Number of days to death without awakening and to awakening 	
Funding	National Heart, Lung, and Blood Institute and with additional support from the Medic One Foundation (Seattle, Washington)	
Declarations of interest	Dr Nichol reported receiving institutional grant funding from the Asmund S. Laerdal Foundation for Acute Medicine, the National Heart, Lung, and Blood Institute, the National Institutes of Health, Medtronic Foundation, Velomedix Inc, Philips Healthcare Inc, Physio-Control Inc, HealthSine Technologies Inc, and Zoll Inc; serving on the board of Medic One Foundation; being part of a patent assigned to the University of Washington; and receiving travel reimbursement from the American Heart Association. Dr Hallstrom reported receiving grants, support for travel to meetings, fees for participating in	

Kim 2014 (Continued)

review activities, payment for writing or reviewing a manuscript, and provision for writing assistance, medicines, equipment, or administrative support from the National Heart, Lung, and Blood Institute; and serving as a consultant to Amarin and St Jude Medical for data and safety monitoring board activity on several trials. Dr Rea reported receiving grant support for community-based resuscitation from Medtronic Foundation. Dr Deem reported receiving institutional grant funding from the National Institutes of Health and Medic One Foundation. No other author reported disclosures.

Notes

Only 77% of all participants with VF received in-hospital cooling (equally with field cooling or not, 224 in each group).

It is unknown how many participants with non-VF rhythms received in-hospital cooling (only one hospital cooled participants with non-VF cardiac arrests)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization was stratified by first recorded rhythm (VF or without VF) and destination hospital and by using randomly permuted blocks of concealed size to ensure temporal equality of assignment in each stratum
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating ambulance and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	Low risk	Study personnel who abstracted the medical records for the primary outcome were unaware of the study allocation
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding of survival data considered 'low risk'
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 of 1364 participants withdrawn because in prison; the proportion of missing outcomes compared with the observed event risk is not enough to have a clinically relevant impact on the intervention effect estimate
Selective reporting (reporting bias)	Low risk	
Other bias	High risk	<ul style="list-style-type: none"> Of all 688 participants assigned to pre-hospital cooling only 50% did receive the full 2 L of fluids, 50 participants (7%) did not receive the intervention Only 77% of all participants with VF received in-hospital cooling (equally with field cooling or not, 224 in each group). It is unknown how many participants with non-VF rhythms received in-hospital cooling (only one hospital cooled participants with non-VF cardiac arrest). No information on the conduct of temperature management in the hospital

Kämäräinen 2009

Methods	Parallel-group, randomized trial, May 2005 to December 2008, multicentre study
Participants	<p>Total number of participants 43, mean age intervention: 59, control: 63 years, 2% female, baseline average temperature pre-hospital cooling group: 35.5°C, hospital cooling group 35.3°C</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • out-of-hospital cardiac arrest treated by the physician-staffed Helsinki Area Helicopter Emergency Medical Air Service • ROSC with palpable pulses • cause of cardiac arrest: any • primary rhythm: any • unconscious (GCS < 6) • time to ROSC exceeds 9 minutes • age older than 17 years • intubated and intravenous access available • oesophageal temperature probe <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • pregnancy • cardiac arrest after trauma or intoxication • persistent initial hypotension after ROSC (systolic blood pressure < 100 mmHg) not responding to fluid challenge or medication
Interventions	<p>Pre-hospital cooling: a rapid infusion of 4°C Ringer's acetate, 100 mL/min, target nasopharyngeal temperature was set at 33°C or alternatively the maximum volume of cold fluid was 30 mL/kg</p> <p>Control: no fluid cooling/conventional therapy; the use of in-hospital cooling was left at the discretion of the hospital's physicians</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> • nasopharyngeal temperature on arrival at the emergency department • first blood gas analyses after hospital admission including electrolyte, creatinine and lactate measurements <p>Secondary:</p> <ul style="list-style-type: none"> • hospital mortality • neurological outcome, CPC 1, CPC 2 at discharge
Funding	None
Declarations of interest	None
Notes	The authors emphasize that the focus of this study was on the pre-hospital phase of the participants

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomization
Allocation concealment (selection bias)	Low risk	Unmarked, sealed and opaque envelopes opened by helicopter staff upon inclusion

Kämäräinen 2009 *(Continued)*

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Treating helicopter and in-hospital staff were not blinded
Blinding of outcome assessment (detection bias) Good neurological outcome	High risk	Treating personnel and investigators could not be blinded to the treatment
Blinding of outcome assessment (detection bias) Survival	Low risk	Lack of blinding considered low risk
Blinding of outcome assessment (detection bias) Adverse events	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	Of 43 participants, 4 participants in the intervention group and 2 participants in the control group excluded from outcome analysis; 2 in the intervention group died between randomization and administration of interventions; the proportion of missing outcomes compared with the observed event risk is not enough to have a clinically relevant impact on the intervention effect estimate
Selective reporting (reporting bias)	Low risk	All expected outcomes included
Other bias	High risk	Only 10 out of 19 participants (52%) in the intervention group and 13 out of 18 (72%) participants in the control group received in-hospital cooling; no detailed information on the conduct of temperature management in the hospital was given

CPC = cerebral performance categories

CPR = cardiopulmonary resuscitation

GCS = Glasgow Coma Scale

h = hours

N = number of participants

OHCA = out-of-hospital cardiac arrest

PEA = pulseless electric activity

ROSC = return of spontaneous circulation

VF = ventricular fibrillation

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Belohlavek 2013	Mixed intervention/ongoing study
Busch 2010	Substudy to Castren 2010
Taccone 2010	Substudy to Castren 2010

Characteristics of ongoing studies *[ordered by study ID]*

Nordberg 2013

Trial name or title	Pre-hospital Resuscitation Intra Nasal Cooling Effectiveness Survival Study (PRINCESS)
Methods	Randomized, controlled, open-label, parallel design, phase 2 study
Participants	<p>Participants with out-of-hospital cardiac arrest</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age $\geq$ 18 years • Collapse was witnessed (heard or seen) • Do not have a pulse • Are unresponsive to external stimuli <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Age $\geq$ 80 years • Have an aetiology of cardiac arrest due to trauma, severe bleeding, drug overdose, cerebrovascular accident, drowning, smoke inhalation, electrocution, hanging • Already hypothermic (e.g. avalanche victim; found in the snow) • Have an obvious barrier to placing intra nasal catheters (e.g. intranasal obstruction) • Do Not Attempt to Resuscitate (DNAR) orders • Have a terminal disease • Known or clinically apparent pregnancy • Have a known coagulopathy (except therapeutically induced) • Are known to have a need for supplemental oxygen • Achieve ROSC prior to randomization • Response time (call to arrival) of the ambulance > 15 minutes
Interventions	<ul style="list-style-type: none"> • No intervention: control participants in the control group standard advanced cardiac life support care. Participants that achieve return of spontaneous circulation will be treated with cooling according to current guidelines upon arrival at the intensive care unit. • Experimental intervention: intra-arrest transnasal cooling with RhinoChill will be initiated during advanced cardiac life support. In participants achieving return of spontaneous circulation, transnasal cooling will continue until systemic cooling is started at the intensive care unit. Intervention: Device: pre-hospital intranasal cooling with RhinoChill
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> • Neurologically intact survival (CPC - cerebral performance categories scale 1 to 2) (time frame: 90 days after cardiac arrest) <p>Secondary:</p> <ul style="list-style-type: none"> • Total survival (time frame: 90 days) (designated as safety issue: no) • Proportion of participants achieving return of spontaneous circulation (ROSC) (time frame: 1 hour) (designated as safety issue: no) • Time to target temperature of 32°C to 34°C (time frame: 8 to 10 hours) (designated as safety issue: no) • Admitted alive to hospital (time frame: 2 to 4 hours) (designated as safety issue: no); proportion of participants that are admitted alive to hospital
Starting date	June 2010
Contact information	<p>Leif Svensson, MD, PhD, leif.svensson@sodersjukhuset.se</p> <p>Maaret Castrén, MD, PhD, maaret.castren@sodersjukhuset.se</p>

Nordberg 2013 (Continued)

Notes

This study is currently recruiting participants

CPC = cerebral performance categories

DNAR = do not attempt resuscitation

ROSC = return of spontaneous circulation

DATA AND ANALYSES

Comparison 1. Survival: pre-hospital cooling versus in-hospital cooling

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Survival	7		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Analysis 1.1. Comparison 1 Survival: pre-hospital cooling versus in-hospital cooling, Outcome 1 Survival.

Study or subgroup	pre-hospital n/N	in-hospital n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
Bernard 2010	56/118	62/116		0.89[0.69,1.15]
Kämäräinen 2009	8/19	8/18		0.95[0.45,1.98]
Kim 2014	259/688	249/676		1.02[0.89,1.17]
Kim 2007	21/63	18/62		1.15[0.68,1.94]
Castren 2010	14/93	13/101		1.17[0.58,2.36]
Debaty 2014	7/123	5/122		1.39[0.45,4.26]
Bernard 2012	11/82	7/81		1.55[0.63,3.8]
			Favours in-hospital 0.2 0.5 1 2 5 Favours pre-hospital	

Comparison 2. Neurological outcome: pre-hospital cooling versus in-hospital cooling

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Good neurological outcome	7		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Analysis 2.1. Comparison 2 Neurological outcome: pre-hospital cooling versus in-hospital cooling, Outcome 1 Good neurological outcome.

Study or subgroup	pre-hospital n/N	in-hospital n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
Bernard 2010	56/118	61/116		0.9[0.7,1.17]
Kämäräinen 2009	8/19	8/18		0.95[0.45,1.98]
Kim 2014	225/688	221/671		0.99[0.85,1.16]
			Favours in-hospital 0.5 0.7 1 1.5 2 Favours pre-hospital

ADDITIONAL TABLES

Table 1. Adverse events: any

Adverse event	Studies	Participants	Statistical method	Effect estimate
Pulmonary oedema first evaluation after ROSC	4	1457	Risk ratio (M-H, random, 95% CI)	1.02 (0.67 to 1.57)
Pulmonary congestions	1	103	Risk ratio (M-H, fixed, 95% CI)	1.81 (0.17 to 19.40)
Cardiomegaly	1	103	Risk ratio (M-H, fixed, 95% CI)	0.53 (0.28 to 0.99)
Pleural effusions	1	103	Risk ratio (M-H, fixed, 95% CI)	0.91 (0.19 to 4.29)
Re-arrest after random-ization	4	1713	Risk ratio (M-H, fixed, 95% CI)	1.23 (1.02 to 1.48)
Acidosis	1	194	Odds ratio (M-H, fixed, 95% CI)	0.21 (0.01 to 4.49)
Acute myocardial in-farction	1	194	Odds ratio (M-H, fixed, 95% CI)	0.36 (0.01 to 8.90)
Bleed	2	271	Odds ratio (M-H, fixed, 95% CI)	0.92 (0.22 to 3.85)
Convulsions	2	271	Odds ratio (M-H, fixed, 95% CI)	3.04 (0.78 to 11.81)
Lethal/long-lasting ar-rhythmia	2	271	Odds ratio (M-H, fixed, 95% CI)	0.57 (0.19 to 1.72)
Renal failure	1	194	Odds ratio (M-H, fixed, 95% CI)	0.54 (0.05 to 6.03)
Sepsis/multiorgan fail-ure	1	194	Odds ratio (M-H, fixed, 95% CI)	0.36 (0.04 to 3.47)
Hyperglycaemia	1	1322	Odds ratio (M-H, fixed, 95% CI)	0.70 (0.55 to 0.89)
Hyperthermia	1	77	Odds ratio (M-H, fixed, 95% CI)	1.74 (0.53 to 5.79)
Pneumonia	1	77	Odds ratio (M-H, fixed, 95% CI)	2.26 (0.54 to 9.51)
Bacteraemia	1	77	Odds ratio (M-H, fixed, 95% CI)	2.70 (0.11 to 68.47)
Adverse events any to-tal	1	194	Odds ratio (M-H, fixed, 95% CI)	1.81 (0.86 to 3.82)

Table 1. Adverse events: any (Continued)

Adverse events serious total	1	194	Odds ratio (M-H, fixed, 95% CI)	0.51 (0.19 to 1.31)
------------------------------	---	-----	---------------------------------	---------------------

CI = confidence interval

ROSC = return of spontaneous circulation

Table 2. Adverse events: device-related

Adverse event	Studies	Participants	Statistical method	Effect estimate
Epistaxis	1	194	Odds ratio (M-H, fixed, 95% CI)	7.85 (0.40 to 154.06)
Periorbital emphysema	1	194	Odds ratio (M-H, fixed, 95% CI)	3.29 (0.13 to 81.81)
Nasal whitening	1	194	Risk ratio (M-H, fixed, 95% CI)	29.30 (1.77 to 486.02)

CI = confidence interval

APPENDICES

Appendix 1. Search strategy: CENTRAL, The Cochrane Library

```
#1 MeSH descriptor Resuscitation explode all trees
#2 MeSH descriptor Cardiopulmonary Resuscitation explode all trees
#3 MeSH descriptor Resuscitation Orders explode all trees
#4 MeSH descriptor Heart Arrest explode all trees
#5 MeSH descriptor Heart Massage explode all trees
#6 ((cardio?pulmonary or order*) near2 resuscitation):ti,ab
#7 reanimation:ti,ab
#8 ((circulatory or circulation or cardiac) near arrest):ti,ab or heart standstill:ti,ab
#9 (#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8)
#10 MeSH descriptor Cryotherapy explode all trees
#11 MeSH descriptor Hypothermia explode all trees
#12 MeSH descriptor Hypothermia, Induced explode all trees
#13 ((resuscitative or therapeutic or artificial or induced or extracorporeal) near hypothermia)
#14 artificial hibernation or body cooling or refrigeration anesthesia or body temperature:ti,ab or refrigeration:ti,ab
#15 (#10 OR #11 OR #12 OR #13 OR #14)
#16 (#9 AND #15)
```

Appendix 2. Search strategy: MEDLINE (OvidSP)

```
1. Resuscitation/ or Cardiopulmonary Resuscitation/ or Resuscitation Orders/ or Heart Arrest/ or Heart Massage/ or advanced cardiac life support.mp. or ((cardio?pulmonary or order*) adj2 resuscitation).ti,ab. or reanimation.ti,ab. or ((circulatory or circulation or cardiac) adj3 arrest).ti,ab. or heart standstill.ti,ab.
2. Cryotherapy/ or Hypothermia/ or Circulatory Arrest, Deep Hypothermia Induced/ or Hypothermia, Induced/ or ((resuscitative or therapeutic or artificial or induced or extracorporeal) adj3 hypothermia).mp. or artificial hibernation.mp. or body cooling.mp. or chilling.mp. or refrigeration anesthesia.mp. or body temperature.ti,ab. or refrigeration.ti,ab.
3. 1 and 2
4. ((randomised controlled trial or controlled clinical trial).pt. or randomised.ab. or placebo.ab. or clinical trials as topic.sh. or randomly.ab. or trial.ti.) not (animals not (humans and animals)).sh.
5. 3 and 4
```

Appendix 3. Search strategy: EMBASE (OvidSP)

```
1. resuscitation/ or heart arrest/ or heart massage/ or advanced cardiac life support.mp. or ((cardio?pulmonary or order*) adj2 resuscitation).ti,ab. or reanimation.ti,ab. or ((circulatory or circulation or cardiac) adj3 arrest).ti,ab. or heart standstill.ti,ab.
```

2. cryotherapy/ or hypothermia/ or ((resuscitative or therapeutic or artificial or induced or extracorporeal) adj3 hypothermia).mp. or artificial hibernation.mp. or body cooling.mp. or chilling.mp. or refrigeration anesthesia.mp. or body temperature.ti,ab. or refrigeration.ti,ab.
3. 1 and 2
4. (placebo.sh. or controlled study.ab. or random*.ti,ab. or trial*.ti,ab. or ((singl* or doub* or trebl* or tripl*) adj3 (blind* or mask*)).ti,ab.) not (animals not (humans and animals)).sh.
5. 3 and 4

Appendix 4. Search strategy: CINAHL (EBSCOhost)

S1 ((MH "Resuscitation") OR (MH "Resuscitation Orders") OR (MH "Resuscitation, Cardiopulmonary") OR (MH "Heart Arrest") OR (MH "Heart Massage")) OR AB (((cardio?pulmonary or order*) and resuscitation)) OR AB reanimation OR ((circulatory or circulation or cardiac) and arrest) OR heart standstill

S2 ((MH "Cryotherapy") OR (MH "Hypothermia") OR (MH "Hypothermia, Induced")) OR (((resuscitative or therapeutic or artificial or induced or extracorporeal) and hypothermia)) OR artificial hibernation OR body cooling OR refrigeration anesthesia

S3 ((MH "randomised Controlled Trials") OR (MH "Random Assignment") OR (MH "Prospective Studies") OR (MH "Multicenter Studies") OR (MH "Clinical Trials") OR (MH "Clinical Trial Registry") OR (MH "Double-Blind Studies") OR (MH "Single-Blind Studies") OR (MH "Triple-Blind Studies") OR (MH "Placebos")) OR (random* or controlled clinical trial or placebo)

S4 S1 and S2 and S3

Appendix 5. Search strategy: BIOSIS (OvidSP)

1. advanced cardiac life support.mp. or ((cardio?pulmonary or order*) adj2 resuscitation).ti,ab. or reanimation.ti,ab. or ((circulatory or circulation or cardiac) adj3 arrest).ti,ab. or heart standstill.ti,ab.
2. (((resuscitative or therapeutic or artificial or induced or extracorporeal) adj3 hypothermia) or artificial hibernation or body cooling or chilling or refrigeration anesthesia).mp. or body temperature.ti,ab. or refrigeration.ti,ab.
3. 1 and 2

Appendix 6. Data extraction form

Data extraction form

Pre-hospital cooling versus in-hospital cooling for patients with cardiac arrest

Reviewer:

Date:

Decision:

- Inclusion
- Exclusion

Reasons for exclusion:

Study characteristics
Publication type:

Language:

Randomization:

- RCT
- Quasi-RCT
- Cluster-randomized

Setting
Multicentre:

- yes
- no

(Continued)

Participants	Total number of patients:
	Mean age:
	Percent female:
	Cause of cardiac arrest <ul style="list-style-type: none"> • cardiac • non-cardiac
	Primary cardiac rhythm <ul style="list-style-type: none"> • ventricular fibrillation • ventricular tachycardia • asystole • pulseless electrical activity
Quality	Allocation concealment <ul style="list-style-type: none"> A. adequate B. unclear <ul style="list-style-type: none"> 1. inadequate
	Outcome assessor blind <ul style="list-style-type: none"> • yes • no • if unclear, please explain
	Intention-to-treat: <ul style="list-style-type: none"> • yes • no • if unclear, please explain
	Selective reporting: <ul style="list-style-type: none"> • yes • no • if unclear, please explain
	Relevant amount of missing outcome data: <ul style="list-style-type: none"> • yes • no • if not, please specify
	Baseline characteristics comparable: <ul style="list-style-type: none"> • yes • no • if not, please specify
	Type of intervention:

(Continued)

Time point of intervention:

- intra-arrest
- post-arrest

Controls:

Duration of cardiac arrest intervention:

Duration of cardiac arrest control:

Target temperature intervention:

Target temperature control:

Cooling rate intervention:

Cooling rate control:

Temperature of patient at admission intervention:

Temperature of patient at admission control:

Total duration of cooling intervention:

Total duration of cooling control:

Rewarming rate intervention:

Rewarming rate control:

Multiple treatment groups:

- yes
- no

Outcomes

Types of outcome measures:

- ..
- ..
- ..
- ..
- ..
- ..

Time point of assessment of outcome measures:

- ..
- ..

(Continued)

- ..
- ..
- ..
- ..

Funding

Notes

Primary outcomes:

Type of outcome:

Intervention		Control	
Events (n)	Total (N)	Events (n)	Total (N)

Type of outcome:

Intervention		Control	
Events (n)	Total (N)	Events (n)	Total (N)

Secondary outcomes (dichotomous):

Type of outcome:

Intervention		Control	
Events (n)	Total (N)	Events (n)	Total (N)

Secondary outcomes (continuous):

Type of outcome:

Intervention
Control

Mean

SD

Total

Mean

SD

Total

WHAT'S NEW

Date	Event	Description
3 January 2019	Amended	Editorial team changed to Cochrane Emergency and Critical Care

CONTRIBUTIONS OF AUTHORS

Jasmin Arrich: (JA), Christof Havel (CH), Michael Holzer (MH), Alexandra-Maria Warenits (AMW), Harald Herkner (HH)

Conceiving the review: JA, CH, MH, and HH

Co-ordinating the review: HH and JA

Undertaking manual searches: JA, HH, MH, AMW, and CH

Screening search results: JA, HH, MH, AMW, and CH

Organizing retrieval of papers: JA

Screening retrieved papers against inclusion criteria: JA, HH, MH, AMW, and CH

Appraising quality of papers: JA, HH, MH, AMW, and CH

Abstracting data from papers: JA, HH, MH, AMW, and CH

Writing to authors of papers for additional information: JA

Providing additional data about papers: JA

Obtaining and screening data on unpublished studies: JA, HH, MH, AMW, and CH

Data management for the review: HH and JA

Entering data into Review Manager ([RevMan 5.3](#)): JA and HH

RevMan statistical data: HH and JA

Other statistical analysis not using RevMan: HH

Interpretation of data: JA, HH, MH, AMW, and CH

Statistical inferences: HH and JA

Writing the review: JA, HH, MH, AMW, and CH

Securing funding for the review: not applicable

Performing previous work that was the foundation of the present study: JA, CH, MH, and HH

Guarantor for the review (one author): JA

Person responsible for reading and checking review before submission: JA

DECLARATIONS OF INTEREST

Jasmin Arrich has no conflict of interest.

Michael Holzer received travel grants for scientific conferences and honoraria for lectures from Bard Medical, EmCools, Polimed Sp. z o.o. and Zoll Medical Österreich. He also received honoraria for consulting from Zoll Medical Österreich and was responsible for studies where the Department of Emergency Medicine received study grants from Velomedix and Philips.

Michael Holzer and Christof Havel were involved in the design, conduct, and publication of the [HACA 2002](#) trial.

Alexandra-Maria Warenits has no conflict of interest

Harald Herkner has no conflict of interest.

SOURCES OF SUPPORT

Internal sources

- Medical University of Vienna, Austria.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We made the following changes to the protocol ([Arrich 2013](#)):

1. We added one additional author (Alexandra-Maria Warenits) and slightly changed the order. We did not search the PASCAL database as it was no longer available at our institution.
2. We changed the title of the published protocol from 'Prehospital versus in-hospital initiation of mild therapeutic hypothermia for survival and neuroprotection after out-of-hospital cardiac arrest' to 'Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest' following suggestions in the latest resuscitation guidelines. 'Cooling' seemed the most fitting and simple term especially when the relatively short pre-hospital period is described. We tried to use the term 'cooling' throughout the manuscript to avoid unnecessary confusion. Occasionally we used 'targeted temperature management' (in the context of the whole cooling period).
3. For the same reason stated in 2, we changed the wording from 'hypothermia' to 'cooling' or 'targeted temperature management' in the [Objectives](#), [Types of participants](#), and [Types of interventions](#).
4. We changed the structure of the [Objectives](#) in order to comply with Cochrane standards (Methodological Expectations of Cochrane Intervention Reviews).
5. We have additionally searched three trials registers (EudraCT, ClinicalTrials.gov, and the International Clinical Trials Registry Platform).
6. As we did not pool the results we did not carry out sensitivity or subgroup analyses.
7. We did not include cluster-RCTs and studies with multiple treatment groups. If we include them when we update the review we will handle them as described in the [Methods](#) section.

NOTES

1. When we extracted the outcome data from [Kim 2014](#) (in January 2015) we noticed a discrepancy in the presented outcome table. After contacting the authors we were sent a corrected table of the outcome data and accordingly adjusted the events in the control group for the outcome "Good neurologic outcome" from 231 as presented in the paper to 221 in the tables sent to us.
2. We would like to thank Dr Nicola Petrucci (content editor), Dr Cathal Walsh (statistical editor), Dr Kjetil Sunde, Dr Clifton Callaway, Dr Jasmeet Soar (peer reviewers), and Jane Cracknell (Cochrane Anaesthesia Review Group Managing Editor) for their help and editorial advice during the preparation of the protocol for the systematic review ([Arrich 2013](#)).

INDEX TERMS

Medical Subject Headings (MeSH)

*Death, Sudden, Cardiac; *Neuroprotection; Emergency Medical Services [*methods]; Hypothermia, Induced [adverse effects] [*methods]; Hypoxia, Brain [*prevention & control]; Out-of-Hospital Cardiac Arrest [*mortality] [*therapy]; Randomized Controlled Trials as Topic; Recurrence; Risk

MeSH check words

Adult; Aged; Humans; Middle Aged