Published in final edited form as:

Curr Psychiatry Rep. ; 20(6): 39. doi:10.1007/s11920-018-0907-x.

What Does Sex Have to Do with It? The Role of Sex as a Biological Variable in the Development of Posttraumatic Stress Disorder

Sara L. Kornfield¹, Liisa Hantsoo¹, and C. Neill Epperson^{1,2,3}

¹Department of Psychiatry, Perelman School of Medicine at the University of Pennsylvania, 3535 Market Street, Room 3005, Philadelphia, PA 19104, USA

²Department of Obstetrics and Gynecology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA

³Penn PROMOTES Research on Sex and Gender in Health, University of Pennsylvania, Philadelphia, PA, USA

Abstract

Purpose of Review—This review highlights the neurobiological aspects of sex differences in posttraumatic stress disorder (PTSD), specifically focusing on the physiological responses to trauma and presents evidence supporting hormone and neurosteroid/peptide differences from both preclinical and clinical research.

Recent Findings—While others have suggested that trauma type or acute emotional reaction are responsible for women's disproportionate risk to PTSD, neither of these explanations fully accounts for the sex differences in PTSD. Sex differences in brain neurocircuitry, anatomy, and neurobiological processes, such as those involved in learning and memory, are discussed as they have been implicated in risk and resilience for the development of PTSD. Gonadal and stress hormones have been found to modulate sex differences in the neurocircuitry and neurochemistry underlying fear learning and extinction.

Summary—Preclinical research has not consistently controlled for hormonal and reproductive status of rodents nor have clinical studies consistently examined these factors as potential moderators of risk for PTSD. Sex as a biological variable (SABV) should be considered, in addition to the endocrine and reproductive status of participants, in all stress physiology and PTSD research.

Keywords

Posttraumatic stress disorder (PTSD); Sex as a biological variable (SABV); Neurobiology; Stress physiology; Neurocircuitry; Gonadal hormones

Correspondence to: Sara L. Kornfield.

Conflict of Interest Sara L. Kornfield declares no potential conflicts of interest.

Liisa Hantsoo and C. Neill Epperson are section editors for *Current Psychiatry Reports*.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

Introduction

Initially codified by the American Psychiatric Association in the Diagnostic and Statistical Manual of Mental Disorders (DSM-III) in 1980 [1], posttraumatic stress disorder (PTSD) described the effects of combat-related trauma on male military Veterans. However, it has now been well established that PTSD disproportionately affects women and girls [2–5]. PTSD is codified most recently in the DSM-5 as a trauma or stressor-related disorder [6] and requires the following functionally impairing symptoms to be present for a month at minimum for diagnosis: (a) direct or indirect exposure to a traumatic event which includes actual or threatened death, serious injury, or sexual violence; (b) persistent reexperiencing of the event; (c) persistent effortful avoidance of trauma-related stimuli after the event; (d) negative alterations in mood or affect that began or worsened after the event; and (e) alterations in arousal or reactivity. Women in the general population are less likely to experience traumatic events than men; however, epidemiologic studies suggest that women are approximately twice as likely as men to meet diagnostic criteria for PTSD [e.g., 3, 4] and are over four times as likely to experience chronic PTSD (lasting more than 6 months) [2]. This review aims to address the question of why females may be more vulnerable to the development of PTSD and/or trauma-related symptoms through consideration of sex as a biological variable (SABV) that influences risk and resilience to PTSD after trauma exposure.

Before considering sex differences in the underlying biology that contributes to PTSD susceptibility, it is important to consider structural (types of traumas) and socialization (peritraumatic reactions/characteristics) theories as they address differential exposures and reactions to trauma that may differ between men and women because of their *gender more than their sex*. Though structural and socialization factors contribute to greater rates and prevalence of PTSD in women compared to men [7], it is unlikely that these factors act in isolation. Instead, individual reaction to and processing of a traumatic experience may depend as much on the type and duration of trauma and the environment or context in which the trauma is enacted as the underlying biology of the individual's stress reaction, genotype, or hormonal state.

The structural theory attempts to explain gender differences in PTSD vulnerability, severity, and chronicity as a result of the types of traumas to which men and women may be differentially exposed. A meta-analysis of research published over a 25-year span (1980–2005) found that men are more likely to experience accidents, non-sexual assaults, combat or war, disasters, illness, unspecified injuries, and witnessing the death or injury of others. Women, in comparison, are more likely to experience sexual assault and childhood sexual abuse (CSA). Despite these differences, however, the research is inconclusive in suggesting that trauma type predicts PTSD diagnosis; when researchers control for trauma type, gender differences in PTSD prevalence rates are still evident. Similarly, among both men and women with a history of childhood abuse and neglect, women were more likely to be diagnosed with PTSD and this gender difference remained even after controlling for subsequent rape and multiple stressor exposures [8]. Furthermore, that women exposed to more typically male-oriented traumas remain more likely to meet criteria for PTSD than

men experiencing those same traumas lends validity to the hypothesis that the greater prevalence of interpersonal or sexual trauma among women is not the sole driver of sex or gender differences in PTSD prevalence and severity [5, 9].

According to the socialization theory of PTSD development, men and women have been shown to display different peritraumatic cognitive, emotional, and behavioral responses [10–13]. To what degree cognitive appraisal of traumatic events is due to biological differences between sexes or socio-cultural norms that lead to gender differences in response to trauma is not clear. After puberty, women in general are at greater risk than men for depression and anxiety disorders, which may increase their risk for PTSD development after trauma exposure [14, 15]. While greater exposure to more types of toxic trauma and a higher prevalence of preexisting psychiatric disorders [16] have been purported to contribute to gender or sex differences, respectively, in prevalence of PTSD, studies controlling for trait anxiety [17] and pre-existing mood or anxiety disorder were not able to fully explain the elevated PTSD rates among women [18, 19].

Women and men's immediate post-trauma reactions are markedly different and may be a contributing factor to the later development of PTSD. For example, women's reliance on emotion-focused coping styles has been associated with increased PTSD severity [10, 11]. These types of reactions include the following: a greater likelihood to report self-blame and a greater belief in their own incompetence, damaged state, or that the world is a dangerous place [12, 13]; wishful thinking, mental disengagement, and suppression or avoidance of trauma memories, all cognitive strategies which have been shown to enhance PTSD symptoms [20]; and trauma-related dissociation which is a predictor for the development of PTSD [21–25]. These female reactions contrast with a more typically male problem-focused coping style, which has been shown to be protective against PTSD [10, 11, 26]. Again, these factors may be better characterized as gender differences as they are routed in the socio-cultural context in large. As they explain a relatively small portion of the observed differences in prevalence, natural history, and treatment of PTSD, it is critical to consider sex as a biological variable (SABV) with respect to risk and resilience for PTSD among men and women.

Sex Differences in the Physiologic Response to Trauma

The female-specific risk factors related to trauma and its aftermath suggest that the physiologic response to trauma differs in women compared to men, perhaps due to the changing hormonal environment necessary for female reproductive function. Recent evidence in clinical and preclinical studies have also pointed to sex differences in hypothalamic pituitary adrenal (HPA) axis and autonomic response to threat, contributing to risk for PTSD and post-traumatic stress symptoms in females compared to males [27•, 28••, 29••, 30].

Sex Differences in Corticotropin Releasing Factor

There are well-described differences between males and females with respect to their HPA axis response to stress [29••]. Corticotropin-releasing factor (CRF) is implicated in the pathophysiology of disorders that are more common in women; specifically, hypersecretion

of this peptide is associated with both major depressive disorder (MDD) and PTSD [31, 32]. Given this association, it was hypothesized that sex differences in CRF reactivity could contribute to disease expression and the increased prevalence in women [33•]. For example, neuronal differences in response to CRF likely contribute to these differences as noradrenergic neurons in the locus coeruleus (LC) arousal system showed a higher sensitivity to CRF in female compared to male rodents. This phenomenon was linked to sex differences in CRF receptor coupling and signaling [34]. Similarly, in female rats, CRF receptor binding in the amygdala and certain regions of the cortex are higher than in male rats. And lastly, sex differences have been found in CRF receptor co-localization with GABAergic neurons in the dorsal raphe and delta opioid receptor-containing neurons in the hippocampus [35•, 36, 37]. Combined with earlier research showing that central administration of CRF can induce stress-related behaviors in male rats (e.g., head-shaking, burying, self-grooming) [38], and anatomical research which showed differences in CRF-induced activity in cortical, limbic, and hindbrains regions in males, the findings that CRF receptors differ by sex even at the cellular/neuronal level suggest that CRF may regulate brain circuits and behavior differently across the sexes.

More recently, this literature was tied together with a study comparing the effect of central administration of CRF in both male *and* female rats, in order to see how these cellular and anatomical differences in CRF reactivity impacted anxietylike behaviors. Consistent with the current hypotheses that CRF mediates the anxiety response differently in males and females, this study showed that higher doses of CRF resulted in increased anxious behaviors for female rats compared to male rats [29••]. Interestingly, there seems to be an interaction with female gonadal hormones as the female rats in their proestrous phase (higher ovarian hormones) showed even more anxious behaviors than females in the diestrous phase. Furthermore, brain regions associated with the anxiety-like behaviors were identified; the infralimbic region, responsible for regulating anxiety, showed a positive association between CRF-induced neuronal activation and anxious behaviors in proestrous females. However, this association was negative for diestrous females and males. Similarly, hormonal status altered the association between CRF-induced neuronal activation and brain region activation, especially in the prefrontal cortex, and other forebrain regions. These findings highlight the role of CRF in the expression of anxiety, not only from a cellular and anatomical perspective, but also with respect to how ovarian hormones alter brain response to CRF infusion. While this observation in rodents underscores the potential for ovarian hormones to sensitize the female to stress, this relationship is complicated in humans with studies suggesting modification by reproductive stage [39, 40].

Sex Differences in Learning and Memory Processes

Sex differences in neurobiological processes, such as those involved in learning and memory, have also been implicated in the sex differences in PTSD prevalence following a traumatic experience. Specifically, deficits in fear extinction are implicated in PTSD as patients experience exaggerated fear/startle response to events reminiscent of the original traumatic event. Sub-optimal fear extinction is central to the development of PTSD as it contributes to the hyperarousal and hypervigilance that are considered hallmark symptoms of the disorder. While the sexual dimorphisms found in the brain structures (i.e., medial

prefrontal cortex, amygdala, and hippocampus) that respond to stress and fear conditioning are thought to contribute to the sex differences seen in poor or adequate fear learning and extinction, results of preclinical and clinical studies have been mixed [27•]. Some, but not all [41], studies show male rats have superior fear conditioning and/or fear extinction, [e.g., 42, 43, 44, 45, 46] compared to female rats [47].

Importantly, anxiety-like response in fear conditioning and extinction experiments may be skewed towards more maletypical behaviors (i.e., freezing) given early studies focused solely on male rodents. Female rodents may show a different set of anxiety behaviors (i.e., darting, rearing, or scanning), which, when overlooked, contributed to the seemingly lower fear acquisition and increased extinction in females in earlier studies [48, 49, 50••, 51]. In an elegant experiment meant to examine this potential sex difference in anxiety-like behaviors, no sex differences were evident over the course of fear conditioning, fear extinction, and extinction retrieval (i.e., long-term memory of the extinction). Rats of both sexes displayed variable responses to an aversive tone (conditioned stimulus). In female rats, however, the behavioral distinction between resilience and susceptibility to anxiety-like behaviors in response to the tone emerged earlier, during fear conditioning, compared to male rats who only showed differentiating responses at fear extinction. Among both male and female rats who showed the susceptible response (high freezing to the tone), extinction retrieval did not differ. This suggests that despite showing the same behavioral response, the underlying neural mechanism may differ as a result of the different point of susceptible/resilience distinction [50••].

Inconsistencies regarding potential sex bias in fear conditioning and extinction are not limited to preclinical models. Men performed better than women in the conditioned fear response during fear acquisition in one study [52], but women showed greater fear conditioning in another [53]. Neuroimaging studies in humans support preclinical findings suggesting sex differences in fear conditioning and extinction. Functional and structural irregularities in the ventromedial prefrontal cortex (vmPFC) and hippocampus result in reduced activation in these areas in individuals with PTSD compared to healthy controls during fear conditioning and extinction paradigms [54]. Others have shown that women and men display distinct patterns of neural activation when under acute stress [55] and during fear learning tasks [56, 57]. Specifically, in healthy individuals, right rostral anterior cingulate (rACC) activity was found to be greater in women than men during extinction recall [58], while among PTSD patients, there is a significantly greater acquisition of conditioned fear (higher skin conductance response (SCR)) in women compared to men [59]. In an fMRI study that compared women and men with and without PTSD after exposure to a traumatic event, men with PTSD were found to show increased activity in the left rostral dACC during extinction recall compared to women with PTSD [28••]. While this work corroborated other studies' findings that dACC activation is implicated in PTSD, it failed to provide an explanation of the mechanism responsible for the greater prevalence of PTSD in women. Notably, menstrual cycle phase was not considered in the above studies. Given the prominent effects of gonadal steroids on brain neurochemistry, structure and function [60•, 61, 62, 63•, 64, 65] across various stages of development and preclinical models indicating estrous cycle effects on stress neurocircuitry, we would argue for controlling for hormonal and reproductive status in all stress- and PTSD-focused research.

Contribution of Gonadal Hormones and Neurosteroids-

The emergence of sex differences in anxiety at puberty [66, 67] and the increased risk of anxiety disorder development or symptom exacerbation that occurs during the luteal phase of the menstrual cycle and under conditions of suppressed endogenous hormone production such as that occurring with steroid contraceptive use [68] highlight the importance of considering neuroactive steroid levels and reproductive status of the individual when evaluating the response to stress and trauma [69•, 70•]. Reproductive hormones and neurosteroids such as the progesterone metabolite and GABA_A receptor agonist allopregnanolone (ALLO) not only impact stress neurocircuitry [54, 70•], they act through receptor and ion channel pathways to regulate numerous neurotransmitter systems involved in the regulation of affect and learning and memory [60•, 71]. Lowered progesterone and estradiol levels characteristic of hormonal contraceptive use are associated with decreased ALLO in women [72], a possible risk factor for contraceptive-induced depression and menstrual cycle-related negative affective states such as premenstrual dysphoric disorder [73, 74].

However, high progesterone levels characteristic of the mid-luteal phase of the menstrual cycle may contribute to the development of more salient and intrusive memories following an aversive event [69•, 75–78]. Elevated exogenous progesterone levels have been implicated in increased amygdala activity and increased functional coupling of the amygdala and the dorsal anterior cingulate cortex, both of which are associated with the acquisition of learned fear [79]. That ALLO is a metabolite of progesterone suggests that the so-called progesterone effects may actually be a result of associated elevations of this neurosteroid [80]. Manipulations of the ALLO activity within the bed nucleus of the stria terminalis (BNST) have been shown to modulate the expression of conditional contextual fear in rats [81]. This extended previous research on contextual fear behaviors [e.g., 82, 83] which together suggests that the reduced levels of context-dependent fear in female rodents with elevated levels of progesterone may actually be more closely related to ALLO's actions within the BNST. However, this preclinical data is inconsistent with human data. Females, compared to males, have higher levels of ALLO naturally, but are *more* susceptible to development of PTSD [84, 85] perhaps due to its fluctuation across the cycle or secondary to organizational differences between males and females at birth that are unmasked by such fluctuations after puberty [86]. Studies of ALLO effects in rodent models have largely been conducted with contextual fear paradigms, not fear extinction paradigms, and it is the deficits in fear extinction that are widely believed to be responsible for PTSD development and maintenance [87, 88]. Pregnant women with PTSD had greater fear-potentiated startle to a safety signal than non-pregnant women with PTSD, which correlated with PTSD hyperarousal symptoms [89]. This suggests that discrimination between the danger signal (the reinforced conditioned stimulus) and safety signal (non-reinforced conditioned) was impaired among the pregnant women. Although ovarian hormones such as progesterone were not assessed, they may play a role in this impaired inhibition of conditioned fear. At this point, it is unknown how progesterone and its metabolites contribute to fear extinction [81]. Work by Milad and colleagues suggests that in fear extinction paradigms, estrogen may be more strongly associated with deficits in extinction recall than progesterone [90]. Women with higher levels of estradiol during an extinction learning task had enhanced extinction

recall, however, estradiol had no impact on fear acquisition or extinction learning. In a fear-potentiated startle study, fear acquisition was similar among women with PTSD and trauma-exposed controls, regardless of gonadal hormone levels [56]. However, fear extinction differed by diagnosis and estradiol level; women with PTSD and low estradiol exhibit heightened fear-potentiated startle during extinction training compared with trauma-exposed controls, which was not exhibited by women with higher estradiol. This suggests that estrogen influences ability to inhibit learned fear.

That higher levels of estradiol are associated with increased activation of brain structures responsible for learning and memory (different sub-regions of the insular and cingulate cortices, amygdala, hippocampus, and hypothalamus) during fear conditioning suggests that the sex differences in the activity of these structures may be partially attributed to female gonadal hormones [91•]. Relevant to the development of PTSD, elevated activation of the circuitry responsible for arousal (hypothalamus, amygdala, and cingulate cortex) is also associated with elevated estradiol in response to emotional stimuli [55]. Notably, women using oral contraceptives, who have prolonged, non-cycling levels of lower endogenous estradiol, but higher levels of synthetic estrogen show reduced extinction memory as compared to women in a high estradiol menstrual phase such as peri-ovulation [92, 93]. Women using oral contraceptives also have enhanced acquisition of a conditioned eyeblink response compared with naturally cycling women, particularly mid-cycle [94]. These findings were consistent with those of rodents under similar hormonal conditions [92].

Given that the experience of intrusive memories and physiological arousal are common symptoms of PTSD, the relationship between sex hormones, emotional memory formation, and fear association seems to be implicated in women's vulnerability to the disorder.

Sex Differences in Stress Neurocircuitry

Another area of research into the discrepancy in PTSD prevalence between men and women is potential differences in brain anatomy and function responsible for fear conditioning. For example, the amygdala plays a key role in conditioned fear and has been shown to activate differently in healthy women and men when exposed to threatening conditions [37, 95]. Other brain structures are at play in these differences, too; women displayed greater activation than men in the dorsal anterior cingulate cortex (ACC), hippocampus, periaqueductal gray region of the brainstem, and cerebellum in response to viewing fearful faces [37]. These patterns of enhanced activity compared to men were sustained in healthy women when individuals were expecting a mild shock [96]. These findings were also confirmed in a metaanalysis of 65 studies showing that women experienced greater activation in the subcallosal ACC, thalamus, midbrain, and brainstem when exposed to affective stimuli [97]. Interestingly, the studies included in the meta-analysis did not consider female hormonal state, indicating that gonadal hormones are not solely responsible for sex differences. Rather, we must consider the organization of the male versus female brain in utero as being responsible for some of the differences noted here.

Among women exposed to trauma and those with PTSD, there is comparatively more activity in the dorsal brainstem (including the superior colliculi, reticular activating system, and midbrain periaqueductal gray region) compared to men with any level of trauma

exposure or PTSD, suggesting that brainstem activity is enhanced in women following trauma exposure but may not be as relevant in the pathophysiology of PTSD among males. These are the same regions which are activated more readily in women than men in healthy samples being exposed to aversive stimuli suggesting baseline sex differences regardless of trauma exposure or PTSD status [37, 96, 97]. Greater brainstem arousal in women is also associated with autonomic arousal, orienting to threat, and an automatic alarm mechanism. Prospective studies have found that heightened autonomic arousal (i.e., tachycardia, hyperventilation) in the immediate aftermath of trauma is associated with later development of PTSD [23]. While in these studies, women with PTSD showed increased activity in regions that may be responsible for PTSD acquisition, men showed an increase in amygdala inhibition via unexpected heightened hippocampal activity. Because the hippocampus works with the ventral medial prefrontal cortex to inhibit amygdala activity, this greater hippocampal activity seen in men may suggest the greater contextual processing and activation of inhibitory networks over fear and arousal networks, contributing to the lower prevalence of PTSD in men [98].

Conclusions

Sex differences in PTSD prevalence are numerous and often attributed to one particular area of research investigation. This review attempts to shed light on the multiple types of differences that exist between the sexes, in order to explain women's vulnerability to the development of this disorder after traumatic exposures. While overall, men may be the more visible recipients of care for PTSD due to the high-profile nature of war and combat trauma and their higher lifetime prevalence of trauma exposure, women are more vulnerable to the aftermath of traumatic events. This review highlights the unique contribution of neurobiology, hormonal shifts, and anatomy that may lead to risk or resilience for PTSD among females. Future research should take into consideration the endocrine conditions, specifically menstrual phase and oral contraceptive use, under which women experience traumatic events as reproductive hormones have pronounced impact on neurochemistry and stress neurocircuitry underlying the development of PTSD.

Acknowledgements

The editors would like to thank Dr. Susan Kornstein for taking the time to review this article.

Funding Sources Kornfield: NIMH K23MH102360; Hantsoo: NIMH K23MH107831, Brain & Behavior Research Foundation NARSAD Young Investigator Award; Epperson: NIMH and Office of Research on Women's Health P50MH099910, NICHD K12HD085848.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (3rd ed.). Washington, DC: Author; 1980.

2. Breslau N, Davis GC. Posttraumatic stress disorder in an urban population of young adults: risk factors for chronicity. *Am J Psychiatry*. 1992;149(5):671–5. [PubMed: 1575259]
3. Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P. Trauma and posttraumatic stress disorder in the community: the 1996 Detroit area survey of trauma. *Arch Gen Psychiatry*. 1998;55(7):626–32. [PubMed: 9672053]
4. Davidson JR, Hughes D, Blazer DG, George L. Post-traumatic stress disorder in the community: an epidemiological study. *Psychol Med*. 1991;21(3):713–21. [PubMed: 1946860]
5. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1995;52:1048–60. 10.1001/archpsyc.1995.03950240066012. [PubMed: 7492257]
6. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th ed.). Washington, DC: Author; 2013.
7. Christiansen DM, Elklit A. Sex differences in PTSD In: Lazinica A, Ovuga E, editors. Posttraumatic stress disorder in a global context. Rijeka: InTech–Open Access Book; 2012 p. 113–42.
8. Koenen KC, Widom CS. A prospective study of sex differences in the lifetime risk of posttraumatic stress disorder among abused and neglected children grown up. *J Trauma Stress*. 2009;22(6):566–74. 10.1002/jts.20478. [PubMed: 19937646]
9. Tolin DF, Foa EB. Sex differences in trauma and posttraumatic stress disorder: a quantitative review of 25 years of research. *Psychol Bull*. 2006;132(6):959–92. [PubMed: 17073529]
10. Bödvarsdóttir I, Elklit A. Psychological reactions in Icelandic earthquake survivors. *Scand J Psychol*. 2004;45(1):3–13. [PubMed: 15016274]
11. Gil S, Caspi Y, Zilberman Ben-Ari I, Koren D, Klein E. Does memory of a traumatic event increase the risk for posttraumatic stress disorder in patients with traumatic brain injury? A prospective study. *Am J of Psychiatry*. 2005;162(5):963–9. [PubMed: 15863799]
12. Tolin DF, Gender FEB. PTSD: a cognitive model In: Kimerling R, Ouimette P, Wolfe J, editors. Gender and PTSD. New York: The Guilford Press; 2002 p. 66–97.
13. Cromer LD, Smyth JM. Making meaning of trauma: trauma exposure doesn't tell the whole story. *J Contemp Psychother*. 2010;40: 65–72.
14. Ahern J, Galea S, Fernandez WG, Koci B, Waldman R, Vlahov D. Gender, social support, and posttraumatic stress in postwar Kosovo. *J Nerv Ment Dis*. 2004;192(11):762–70. [PubMed: 15505520]
15. Zeidner M Gender group differences in coping with chronic terror: the Israeli scene. *Sex Roles*. 2006;54:297–310. 10.1007/s11199-006-9346-y.
16. Hapke U, Schumann A, Rumpf HJ, John U, Meyer C. Posttraumatic stress disorder: the role of trauma, pre-existing psychiatric disorders, and gender. *Eur Arch Psychiatry Clin Neurosci*. 2006;256(5):299–306. [PubMed: 16783498]
17. Spindler H, Elklit A, Christiansen D. Risk factors for posttraumatic stress disorder following an industrial disaster in a residential area: a note on the origin of observed gender differences. *Gend Med*. 2010;7(2):156–65. [PubMed: 20435278]
18. Fullerton CS, Ursano RJ, Epstein RS, Crowley B, Vance K, Kao TC, et al. Gender differences in posttraumatic stress disorder after motor vehicle accidents. *Am J Psychiatry*. 2001 ;158(9): 1486–91. [PubMed: 11532736]
19. Breslau N, Davis GC, Andreski P, Peterson EL, Schultz LR. Sex differences in posttraumatic stress disorder. *Arch Gen Psychiatry*. 1997;54(11):1044–8. [PubMed: 9366662]
20. Clohessy S, Ehlers A. PTSD symptoms, response to intrusive memories and coping in ambulance service workers. *Br Clin Psychol*. 1999;38(Pt 3):251–65.
21. Ehring T, Ehlers A, Cleare AJ, Glucksman E. Do acute psychological and psychobiological responses to trauma predict subsequent symptoms severities of PTSD and depression? *Psychiatry Res*. 2006;161 (1):67–75.
22. Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull*. 2003;129(1):52–73. [PubMed: 12555794]
23. Bryant RA, Creamer M, O'Donnell M, Silove D, McFarlane AC. A multisite study of initial respiration rate and heart rate as predictors of posttraumatic stress disorder. *J Clin Psychiatry*. 2008;69(11): 1694–701. [PubMed: 19014750]

24. Bryant RA. Acute stress reactions: can biological responses predict posttraumatic stress disorder? *CNS Spectr*. 2003;8(9):668–74. [PubMed: 15079140]
25. Irish LA, Fischer B, Fallon W, Spoonster E, Sledjeski EM, Delahanty DL. Gender differences in PTSD symptoms: an exploration of peritraumatic mechanisms. *Journal of Anxiety Disorders*. 2011;25(2):209–16. [PubMed: 20956066]
26. Tamres LK, Janicki D, Helgeson VS. Sex differences in coping behavior: a meta-analytic review and an examination of relative coping. *Personal Soc Psychol Rev*. 2002;6(1):2–30.
- 27•. Maeng LY, Milad MR. Sex differences in anxiety disorders: Interactions between fear, stress, and gonadal hormones. *Horm Behav*. 2015;76:106–17. [PubMed: 25888456] This review paper discusses the intersection between stress and fear mechanisms and their modulation by gonadal hormones relevant to sex differences in anxiety and fear-based disorders.
- 28••. Shvil E, Sullivan GM, Schafer S, Markowitz JC, Campeas M, Wager TD, et al. Sex differences in extinction recall in posttraumatic stress disorder: a pilot fMRI study. *Neurobiol Learn Mem*. 2014;113:101–8. [PubMed: 24560771] Shvil and colleagues explored the biological markers of sex differences in relation to trauma by using a fear learning and extinction paradigm and fMRI in PTSD vs trauma exposed non-PTSD individuals.
- 29••. Wiersielis KR, Wicks B, Simko H, Cohen SR, Khantsis S, Baksh N, et al. Sex differences in corticotropin releasing factor-evoked behavior and activated networks. *Psychoneuroendocrinology*. 2016;73:204–16. [PubMed: 27521739] This paper found that ovarian hormones alter the way brain regions work in response to corticotropin releasing factor, which may be responsible for the different coping strategies displayed by males and females. It also suggests an explanation for female vulnerability to psychiatric illness.
30. Puralewski R, Vasilakis G, Seney ML. Sex-related factors influence expression of mood-related genes in the basolateral amygdala differentially depending on age and stress exposure. *Biol Sex Differ*. 2016 9 17;7:50. [PubMed: 27660699]
31. Bremner JD, Licino J, Darnell A, Krystal JH, Owens MJ, Southwick SM, et al. Elevated CSF corticotropin-releasing factor concentrations in posttraumatic stress disorder. *Am J Psychiatry*. 1997;154(5):624–9. [PubMed: 9137116]
32. Nemeroff CB, Widerlov E, Bissette G, Walleus H, Karlsson I, Eklund K, et al. Elevated concentrations of CSF corticotropin-releasing factor-like immunoreactivity in depressed patients. *Science*. 1984;226(4680):1342–4. [PubMed: 6334362]
- 33•. Bangasser DA, Valentino RJ. Sex difference in stress-related psychiatric disorders: neurobiological perspectives. *Front Neuroendocrinol*. 2014;35(3):303–19. [PubMed: 24726661] This review describes clinical and preclinical studies that identify sex differences in the underlying circuitry of the neuroendocrine and the corticolimbic systems, and the HPA axis, and demonstrate cellular and molecular sex differences in stress response systems.
34. Bangasser DA, Curtis A, Reyes BA, Bethea TT, Parastatidis I, Ischiropoulos H, Van Bockstaele EJ, Valentino RJ. Sex differences in corticotropin-releasing factor receptor signaling and trafficking: potential role in female vulnerability to stress related psychopathology. *Mol Psychiatry*. 2010; 15(9): 877, 896–904.
- 35•. Howerton AR, Roland AV, Fluharty JM, Marshall A, Chen A, Daniels D, et al. Sex differences in corticotropin-releasing factor receptor-1 action within the dorsal raphe nucleus in stress responsivity. *Biol Psychiatry*. 2014;75(11):873–83. [PubMed: 24289884] These authors found that male mice were more receptive to the effects of CRF antagonist infusions compared to females which showed a blunted response. These sex differences in the CRF receptor-1 throughout the dorsal raphe suggest unique stress modulation of the dorsal raphe among females and may underlie women's vulnerability to affective disorder development and maintenance.
36. Williams TJ, Akama KT, Knudsen MG, McEwen BS, Milner TA. Ovarian hormones influence corticotropin releasing factor receptor colocalization with delta opioid receptors in CA1 pyramidal cell dendrites. *Exp Neurol*. 2011;230(2):186–96. 10.1016/j.expneurol.2011.04.012. [PubMed: 21549703]
37. Williams MA, McGlone F, Abbott DF, Mattingley JB. Differential amygdala responses to happy and fearful facial expressions depend on selective attention. *NeuroImage*. 2005;24(2):417–25. [PubMed: 15627583]

38. Howard O, Carr GV, Hill TE, Valentino RJ, Lucki I. Differential blockade of CRF evoked behaviors by depletion of norepinephrine and serotonin in rats. *Psychopharmacology*. 2008;199(4):569–82. [PubMed: 18516596]
39. Albert K, Pruessner J, Newhouse P. Estradiol levels modulate brain activity and negative responses to psychosocial stress across the menstrual cycle. *Psychoneuroendocrinology*. 2015;59:14–24. [PubMed: 26123902]
40. Dumas JA, Albert KM, Naylor MR, Sites CK, Benkelfat C, Newhouse PA. The effects of age and estrogen on stress responsivity in older women. *Am J Geriatr Psychiatry*. 2012;20(9):734–43. [PubMed: 22832417]
41. Daviu N, Andero R, Armario A, Nadal R. Sex differences in the behavioural and hypothalamic-pituitary-adrenal response to contextual fear conditioning in rats. *Horm Behav*. 2014;66:713–23. [PubMed: 25311689]
42. Aguilar R, Gil L, Gray JA, Driscoll P, Flint J, Dawson GR, et al. Fearfulness and sex in F2 roman rats: males display more fear though both sexes share the same fearfulness traits. *Physiol Behav*. 2003;78(4-5):723–32. [PubMed: 12782229]
43. Baker-Andresen D, Flavell CR, Li X, Bredy TW. Activation of BDNF signaling prevents the return of fear in female mice. *Learn Mem*. 2013;20(5):237–40. [PubMed: 23589089]
44. Baran SE, Armstrong CE, Niren DC, Conrad CD. Prefrontal cortex lesions and sex differences in fear extinction and perseveration. *Learn Mem*. 2010;17(5):267–78. [PubMed: 20445082]
45. Baran SE, Armstrong CE, Niren DC, Hanna JJ, Conrad CD. Chronic stress and sex differences on the recall of fear conditioning and extinction. *Neurobiol Learn Mem*. 2009;91(3):323–32. [PubMed: 19073269]
46. Fenton GE, Pollard AK, Halliday DM, Mason R, Bredy TW, Stevenson CW. Persistent prelimbic cortex activity contributes to enhanced learned fear expression in females. *Learn Mem*. 2014;21(2):55–60. [PubMed: 24429423]
47. Ribiero A, Barbosa FF, Godhino MR, Fernandes VS, Munguba H, Melo TG, et al. Sex differences in aversive memory in rats: possible role of extinction and reactive emotional factors. *Brain Cogn*. 2010;74(2):145–51. [PubMed: 20727653]
48. Gruene TM, Flick K, Stefano A, Shea SD, Shansky RM. Sexually divergent expression of active and passive conditioned fear responses in rats. *elife*. 2015;4:e11352. [PubMed: 26568307]
49. Gruene TM, Roberts E, Thomas V, Ronzio A, Shansky RM. Sexspecific neuroanatomical correlates of fear expression in prefrontalamygdala circuits. *Biol Psychiatry*. 2015;78(3):186–93. 10.1016/j.biopsych.2014.11.014. [PubMed: 25579850]
- 50••. Shansky RM, Woolley CS. Considering sex as a biological variable will be valuable for neuroscience research. *J Neurosci*. 2016;36(47):11817–22. [PubMed: 27881768] This paper is a call for research to include sex as a biological variable (SABV) and the authors present compelling evidence for why doing so will strengthen basic science research and improve health outcomes.
51. Shansky RM. Sex differences in PTSD resilience and susceptibility: challenges for animal models of fear learning. *Neurobiol of Stress*. 2015;1:60–5.
52. Milad MR, Rauch SL, Pitman RK, Quirk GJ. Fear extinction in rats: implications for human brain imaging and anxiety disorders. *Biol Psychol*. 2006;73:61–71. [PubMed: 16476517]
53. Kelly MM, Forsyth JP. Sex differences in response to an observational fear conditioning procedure. *Behav Ther*. 2007;38(4):340–9. [PubMed: 18021949]
54. Milad MR, Quirk GJ. Fear extinction as a model for translational neuroscience: ten years of progress. *Annu Rev Psychol*. 2012;63: 129–51. [PubMed: 22129456]
55. Goldstein JM, Jerram M, Abbs B, Whitfield-Gabrieli S, Makris N. Sex differences in stress response circuitry activation dependent on female hormonal cycle. *J Neurosci*. 2010;30(2):431–8. [PubMed: 20071507]
56. Glover EM, Jovanovic T, Mercer KB, Kerley K, Bradley B, Ressler KJ, et al. Estrogen levels are associated with extinction deficits in women with posttraumatic stress disorder. *Biol Psychiatry*. 2012;72(1):19–24. 10.1016/j.biopsych.2012.02.031. [PubMed: 22502987]

57. Zeidan MA, Igoe SA, Linnman C, Vitalo A, Levine JB, Klibanski A, et al. Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biol Psychiatry*. 2011;70:920–7. [PubMed: 21762880]
58. Lebron-Milad K, Graham BM, Milad MR. Low estradiol levels: a vulnerability factor for the development of posttraumatic stress disorder. *Biol Psychiatry*. 2012;72(1):6–7. [PubMed: 22682395]
59. Inslicht SS, Metzler TJ, Garcia NM, Pineles SL, Milad MR, Orr SP, et al. Sex differences in fear conditioning in posttraumatic stress disorder. *J Psychiatr Res*. 2013;47(1):64–71. [PubMed: 23107307]
- 60•. Shanmugan S, Epperson CN. Estrogen and the prefrontal cortex: towards a new understanding of estrogen's effects on executive functions in the menopause transition. *Hum Brain Mapp*. 2014;35(3):847–65. [PubMed: 23238908] This review presents evidence from preclinical rodent and non-human primate as well as human studies for consideration of the interactions of estrogen with neurotransmitter systems, stress, genotype, and individual life events related to cognitive changes associate with menopause.
61. Epperson CN, Haga K, Mason GF, Sellers E, Gueorguieva R, Zhang W, et al. Cortical gamma-aminobutyric acid levels across the menstrual cycle in healthy women and those with premenstrual dysphoric disorder: a proton magnetic resonance spectroscopy study. *Arch Gen Psychiatry*. 2002;59(9):851–8. [PubMed: 12215085]
62. Shanmugan S, Wolf DH, Calkins ME, Moore TM, Ruparel K, Hopson RD, et al. Common and dissociable mechanisms of executive system dysfunction across psychiatric disorders in youth. *Am J Psychiatry*. 2016;173(5):517–26. [PubMed: 26806874]
- 63•. Shanmugan S, Satterthwaite TD, Sammel MD, Cao W, Ruparel K, Gur RC, et al. Impact of early life adversity and tryptophan depletion on functional connectivity in menopausal women: A doubleblind, placebo - controlled crossover study. *Psychoneuroendocrinology*. 2017;84:197–205. [PubMed: 28755550] Results from this study of menopausal women suggest that early life adversity has lasting impacts on large-scale functional networks underlying executive function.
64. Amin Z, Mason GF, Cavus I, Krystal JH, Rothman DL, Epperson CN. The interaction of neuroactive steroids and GABA in the development of neuropsychiatric disorders in women. *Pharmacol Biochem Behav*. 2006;84(4):635–43. [PubMed: 16860856]
65. Hantsoo L, Epperson CN. Premenstrual dysphoric disorder: epidemiology and treatment. *Curr Psychiatry Rep*. 2015;17(11):87. [PubMed: 26377947]
66. McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. *J Psychiatr Res*. 2011;45(8):1027–35. [PubMed: 21439576]
67. Paus T, Keshavan M, Giedd JN. Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci*. 2008;9(12): 947–57. [PubMed: 19002191]
- 68•. Skovlund CW, Mørch LS, Kessing LV, Lidegaard O. Association of hormonal contraception with depression. *JAMA Psychiatry*. 2016;73(11):1154–62. 10.1001/jamapsychiatry.2016.2387. [PubMed: 27680324] These authors found that among women who used hormonal contraceptive, initiation of anti-depressant medications was more common than in women who did not use hormonal contraception. This suggests a connection between neuroactive steroid levels/ reproductive status and development or exacerbation of trauma related psychopathology.
- 69••. Li SH, Graham BM. Why are women so vulnerable to anxiety, trauma-related and stress-related disorders? The potential role of sex hormones. *Lancet Psychiatry*. 2017;4(1):73–82. [PubMed: 27856395] This review paper highlights two broad mechanisms by which estradiol and progesterone influence sex differences in anxiety disorders: augmentation of vulnerability factors associated with the development of anxiety disorders; and facilitation of the maintenance of anxious symptoms.
- 70•. Bale TL, Epperson CN. Sex as a biological variable: who, what, when, why, and how. *Neuropsychopharmacology*. 2017;42(2): 386–96. [PubMed: 27658485] The authors of this review discuss examples of both animal models and clinical studies and provide guidance and references for the inclusion of sex as a biological variable in neuropsychopharmacological research

71. Gordon JL, Girdler SS, Meltzer-Brody SE, Stika CS, Thurston RC, Clark CT, et al. Ovarian hormone fluctuation, Neurosteroids and HPA Axis dysregulation in Perimenopausal depression: a novel heuristic model. *Am J Psychiatry*. 2015;172(3):227–36. 10.1176/appi.ajp.2014.14070918. [PubMed: 25585035]
72. Andreen L, Nyberg S, Turkmen S, van Wingen G, Fernandez G, Backstrom T. Sex steroid induced negative mood may be explained by the paradoxical effect mediated by GABA_A modulators. *Psychoneuroendocrinology*. 2009;34(8):1121–32.
73. Zorumski CF, Paul SM, Izumi Y, Covey DF, Mennerick S. Neurosteroids, stress and depression: potential therapeutic opportunities. *Neurosci Biobehav Rev*. 2013;37(1):109–22. [PubMed: 23085210]
74. Uhde TW, Singareddy R. Psychiatry as a neuroscience In: Juan Jose L-I, Wolfgang G, Mario M, Norman S, editors. *Biological research in anxiety disorders*. Chichester: John Wiley & Sons, Ltd; 2002 p. 237–86.
75. Ertman N, Andreano JM, Cahill L. Progesterone at encoding predicts subsequent emotional memory. *Learn Mem*. 2011;18(12): 759–63. [PubMed: 22101178]
76. Ferree NK, Cahill L. Post-event spontaneous intrusive recollections and strength of memory for emotional events in men and women. *Conscious Cogn*. 2009;18:126–34. [PubMed: 19131257]
77. Ferree NK, Kamat R, Cahill L. Influences of menstrual cycle position and sex hormone levels on spontaneous intrusive recollections following emotional stimuli. *Conscious Cogn*. 2011 ;20(4): 1154–62. [PubMed: 21353599]
78. Soni M, Curran VH, Kamboj SK. Identification of a narrow postovulatory window of vulnerability to distressing involuntary memories in healthy women. *Neurobiol Learn Mem*. 2013;104:32–28. [PubMed: 23611942]
79. Van Wingen GA, van Broekhoven F, Verkes RJ, Petersson KM, Backstrom T, Buitelaar JK, et al. Progesterone selectively increases amygdala reactivity in women. *Mol Psych*. 2008;13:325–33.
80. Schumacher M, Mattern C, Ghomari A, Oudinet JP, Liere P, Lambombarda F, et al. Revisiting the roles of progesterone and allopregnanolone in the nervous system: resurgence of the progesterone receptors. *Prog Neurobiol*. 2014;113:6–39. [PubMed: 24172649]
81. Nagaya N, Acca GM, Maren S. Allopregnanolone in the bed nucleus of the stria terminalis modulates contextual fear in rats. *Front Behav Neurosci*. 2015;9:205 10.3389/fnbeh.2015.00205. [PubMed: 26300750]
82. Markus EJ, Zecevic M. Sex differences and estrous cycle changes in hippocampus-dependent fear conditioning. *Psychobiology*. 1997;25:246–52.
83. Toufexis DJ, Davis C, Hammond A, Davis M. Progesterone attenuates corticotropin-releasing factor-enhanced but not fearpotentiated startle via the activity of its neuroactive metabolite, allopregnanolone. *J Neurosci*. 2004;24(45):10280–7. [PubMed: 15537900]
84. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62:617–27. [PubMed: 15939839]
85. Chang Y-J, Yang C-H, Liang Y-C, Yeh C-M, Huang C-C, Hsu K-S. Estrogen modulates sexually dimorphic contextual fear extinction in rats through estrogen receptor β . *Hippocampus*. 2009;19:1142–50. [PubMed: 19338017]
86. Bale TL, Epperson CN. Sex differences and stress across the lifespan. *Nat Neurosci*. 2015;18(10): 1413–20. [PubMed: 26404716]
87. Pitman RK, Rasmusson AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, et al. Biological studies of post-traumatic stress disorder. *Nat Rev Neurosci*. 2012;13:769–87. [PubMed: 23047775]
88. Maren S, Phan KL, Liberzon I. The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nat Rev Neurosci*. 2013;14:417–28. [PubMed: 23635870]
89. Michopoulos V, Rothbaum AO, Corwin E, Bradley B, Ressler KJ, Jovanovic T. Psychophysiology and posttraumatic stress disorder symptom profile in pregnant African-American women with trauma exposure. *Arch Women's Ment Health*. 2015;18(4):639–48. 10.1007/s00737-014-0467-y. [PubMed: 25278341]

90. Milad MR, Zeidan MA, Contero A, Pitman RK, Klibanski A, Rauch SL, et al. The influence of gonadal hormones on conditioned fear extinction in healthy humans. *Neuroscience*. 2010;168(3):652–8. 10.1016/j.neuroscience.2010.04.030. [PubMed: 20412837]
91. Hwang MJ, Zsido RG, Song H, Pace-Schott EF, Miller KK, Lebron-Milad K, et al. Contribution of estradiol levels and hormonal contraceptives to sex differences within the fear network during fear conditioning and extinction. *BMC Psychiatry*. 2015;15:295. [PubMed: 26581193] This study found that women with higher levels of estradiol have greater activations of sub regions of the insular and cingulate cortices, amygdala, hippocampus, and hypothalamus during conditioning, extinction, and recall. These results suggest the need to consider gonadal hormone levels and reproductive status/phase in women in order to avoid overlooking sex differences when examining the neurobiology of emotional regulation.
92. Graham BM, Milad MR. Blockade of estrogen by hormonal contraceptives impairs fear extinction in female rats and women. *Biol Psychiatry*. 2013;73(4):371–8. 10.1016/j.biopsych.2012.09.01893. [PubMed: 23158459]
93. Wegerer M, Kerschbaum H, Blechert J, Wilhelm FH. Low levels of estradiol are associated with elevated conditioned responding during fear extinction and with intrusive memories in daily life. *Neurobiol Learn Mem*. 2014;116:145–54. [PubMed: 25463649]
94. Beck KD, McLaughlin J, Bergen MT, Cominski TP, Moldow RL, Servatius RJ. Facilitated acquisition of the classically conditioned eyeblink response in women taking oral contraceptives. *Behav Pharmacol*. 2008;19(8):821–8. 10.1097/FBP.0b013e32831c3b8295. [PubMed: 19020417]
95. McClure EB, Monk CS, Nelson ES, Zarahn E, Leibenluft E, Bilder R (2004). A developmental examination of gender differences in brain engagement during evaluation threat. *Biol Psychiatry*, 55(11): 1047–1055. 96. Fear-related activity in subgenual anterior cingulate differs between men and women. [PubMed: 15158422]
96. Butler T, Pan H, Epsiein J, Protopopescu X, Tuescher O, Goldstein M, et al. Fear-related activity in subgenual anterior cingulate differs between men and women. *Neuroreport*. 2005;16(11):1233–6. [PubMed: 16012355]
97. Wager TD, Phan KL, Liberzon I, Taylor SF. Valence, gender, and lateralization of functional brain anatomy in emotion: a metaanalysis of findings from neuroimaging. *NeuroImage*. 2003;19(3):513–31. [PubMed: 12880784]
98. Felmingham K, Williams LM, Kemp AH, Liddell B, Falconer E, Peduto A, et al. Neural responses to masked fear faces: sex differences and trauma exposure in posttraumatic stress disorder. *J Abnorm Psychol*. 2010;119(1):241–7. [PubMed: 20141261]