The on-site differential diagnosis of decompression sickness from endogenous cerebral ischaemia in an elderly Ama diver using ultrasound

Youichi Yanagawa, Kazuhiko Omori, Ikuto Takeuchi, Kei Jitsuiki, Hiromichi Ohsaka, Kouhei Ishikawa

Department of Acute Critical Care Medicine, Juntendo Shizuoka Hospital, Izunokuni City, Shizuoka Prefecture, Japan

Corresponding author: Youichi Yanagawa, 1129 Nagaoka, Izunokuni City, Shizuoka, Japan 410-2295
yyanaga@juntendo.ac.jp

Key words

Breath-hold diving; Indigenous divers; Stroke; Doppler; Case reports

Abstract

(Yanagawa Y, Omori K, Takeuchi I, Jitsuiki K, Ohsaka H, Ishikawa K. The on-site differential diagnosis of decompression sickness from endogenous cerebral ischaemia in an elderly Ama diver using ultrasound. *Diving and Hyperbaric Medicine*. 2018 December 24;48(4):262–263. doi: 10.28920/dhm48.4.262-263. PMID: 30517960.)

Commercial or occupational breath-hold (BH) harvest divers along the coast and islands of Japan are collectively called Ama divers. Repetitive BH diving by Ama divers may place them at risk of developing neurological decompression sickness (DCS). We report a 74-year-old Ama diver who demonstrated right hemiparesis during an ascent after free diving at a depth of 5 metres' sea water. This report suggests the usefulness of on-site ultrasound for making a differential diagnosis of DCS from endogenous cerebral ischaemia. Further clinical studies of this management approach are warranted.

Introduction

It was previously believed that a diver must be exposed to a certain minimum depth before bubbles could form and that a diver could spend an unlimited amount of time at shallow depths (< 10 metres' sea water, msw).¹ However, it is now known that bubble formation can occur even after shallow dives and that it is inappropriate to exclude the diagnosis of decompression sickness (DCS) based on a perceived minimum depth.¹ Even when a diver follows the decompression tables, excess nitrogen remains in the blood for hours after a dive. Doppler studies have shown that up to 10% of remaining nitrogen may be released as bubbles after a normal dive.²

Commercial or occupational breath-hold (BH) harvest divers along the coast and islands of Japan are collectively known as Ama divers.³ Repetitive BH diving can lead to the accumulation of nitrogen (N₂) in blood and tissues, which may give rise to DCS,⁴ and previous studies have suggested that Ama divers are at risk of neurological DCS.^{3–5} We report a patient with endogenous transient ischaemia who demonstrated right hemiparesis during her ascent after free diving to a depth of 5 metres' sea water (msw).

Case report

A 74-year-old Japanese female Ama diver was rescued by colleagues after she developed right hemiparesis during ascent while free diving to a depth of 5 msw. She had made

multiple dives over three hours to similar depths collecting turban shells and abalone. The exact times of the dives could not be determined. She had known hypertension on medication. She was taken ashore, and an emergency call resulted in an ambulance being dispatched. A physician-staffed helicopter from Eastern Shizuoka was also requested, as divers with DCS are transported to a hospital located further away for recompression since there are no suitable hospitals on the Izu Peninsula.

When examined on site by a physician an hour post dive, she was asymptomatic. As portable ultrasound (Vscan®; General Electric, USA) using the longitudinal sub-xiphoid window did not reveal any bubbles in her inferior vena cava, she was diagnosed with endogenous cerebral ischaemia not induced by DCS or air embolism from barotrauma. She was transported to our hospital (which does not have recompression facilities) by helicopter whilst receiving intravenous Ringer's solution and on oxygen.

On arrival, she was fully conscious and orientated, with isocoric, reactive pupils. Neurological examination was unremarkable. Whole-body computed tomography (CT) revealed no gas in her body, and there were no blebs, small bullae or air trapping in the lungs. Magnetic resonance imaging (MRI) of her head two hours after the incident, including diffusion-weighted imaging and angiography, were negative. Biochemical analyses of blood revealed no abnormalities, including for D-dimer. Right-to-left shunt, including a persistent foramen ovale (PFO), was excluded on

ultrasound. A diagnosis of a transient endogenous ischaemic attack and hypertension was made, and she received anti-platelet therapy without recompression treatment. Her post-admission course was uneventful and she was discharged on hospital day five without sequelae.

Discussion

This report suggests the potential usefulness of on-site ultrasound for the differential diagnosis of a central neurological event. The different diagnosis of DCS, air embolism from pulmonary barotrauma or endogenous stroke was necessary in the present case. Free diving as performed by Japanese Ama is a rare cause of DCS but still one that should be considered.³ The mechanism of stroke induced by DCS or air embolism is intravascular gas produced by abrupt decompression or barotrauma. Also there is a strong correlation between DCS and PFO, where venous bubbles can become arterialized via the right-to-left shunt and, thus, result in cerebral symptoms.⁶ Repetitive deep breath-hold diving, long dive times and short surface intervals result in N₂ loading predisposing divers to DCS, which characteristically manifests as cerebral stroke.^{3,7–9} This elderly Ama had no air bubbles on ultrasound at the scene or on subsequent CT, a negative D-dimer level and no right-to-left shunt, so the possibility of DCS was unlikely.

Even in asymptomatic divers, gas bubbles can be detected by ultrasound immediately after diving.¹⁰ Loss of consciousness during diving may be from decompression illness (DCI), hypoxia induced by drowning, immersion pulmonary oedema or a cardiac event from immersion and exercise.¹¹ In such cases, physicians on board the Eastern Shizuoka helicopter routinely use ultrasound to make a differential diagnosis at the scene in order to decide to which hospital the diver will be transported, dependent on the availability of recompression facilities. However, the use of this device in making a differential diagnosis has not been validated in the literature. There may be criticism that the absence of bubbles in the inferior vena cava, even in the acute phase, cannot exclude the presence of bubbles in the whole body of a diver, so divers with focal neurological signs should always be transported to a hospital with a recompression chamber as a precaution.

Conclusion

This report suggests the usefulness of on-site ultrasound for the differential diagnosis of DCS from endogenous cerebral ischaemia in an elderly Ama diver. Further clinical experience with ultrasound for the acute differential diagnosis of divers with neurological symptoms and signs is merited.

References

- 1 Moon RE, de Lisle Dear G, Stolp BW. Treatment of decompression illness and iatrogenic gas embolism. *Resp Care Clin N Am*. 1999;5:93–135. PMID: 10205814.
- 2 Hardy KR. Diving-related emergencies. *Emerg Med Clin North Am*. 1997;15:223–39. PMID: 9056577.
- 3 Tamaki H, Kohshi K, Ishitake T, Wong RM. A survey of neurological decompression illness in commercial breath-hold divers (Ama) of Japan. *Undersea Hyperb Med*. 2010;37:209–17. PMID: 20737928.
- 4 Kohshi K, Wong RM, Abe H, Katoh T, Okudera T, Mano Y. Neurological manifestations in Japanese Ama divers. *Undersea Hyperb Med*. 2005;32:11–20. PMID: 15796310.
- 5 Tamaki H, Kohshi K, Ishitake T, Wong RM. A survey of neurological decompression illness in commercial breath-hold divers (Ama) of Japan. *Undersea Hyperb Med*. 2010;37:209–17. PMID: 20737928.
- 6 Kohshi K, Wong RM, Abe H, Katoh T, Okudera T, Mano Y. Neurological manifestations in Japanese Ama divers. *Undersea Hyperb Med*. 2005;32:11–20. PMID: 15796310.
- 7 Kohshi K, Tamaki H, Lemaître F, Okudera T, Ishitake T, Denoble PJ. Brain damage in commercial breath-hold divers. *PLoS One*. 2014;9(8):e105006. doi: 10.1371/journal.pone.0105006. PMID: 25115903. PMCID: PMC4130625.
- 8 Gempp E, Blatteau JE. Neurological disorders after repetitive breath-hold diving. *Aviat Space Environ Med*. 2006;77:971–3. PMID: 16964749.
- 9 Goldman S, Solano-Altamirano JM. Decompression sickness in breath-hold diving and its probable connection to the growth and dissolution of small arterial gas emboli. *Math Biosci*. 2015;262:1–9. doi: 10.1016/j.mbs.2015.01.001. PMID: 25598211.
- 10 Blogg SL, Gennser M, Møllerlækken A, Brubakk AO. Ultrasound detection of vascular decompression bubbles: the influence of new technology and considerations on bubble load. *Diving Hyperb Med*. 2014;44:35–44. PMID: 24687484.
- 11 Gempp E, Morin J, Louge P, Blatteau JE. Reliability of plasma D-dimers for predicting severe neurological decompression sickness in scuba divers. *Aviat Space Environ Med*. 2012;83:771–5. PMID: 22872991.

Funding

This work received funds from the Ministry of Education, Culture, Sports, Science and Technology (MEXT - Supported Program for the Strategic Research Foundation at Private Universities, 2015–2019).

Conflicts of interest: nil

Submitted: 28 February 2018; revised 28 May and 12 July 2018

Accepted: 07 November 2018

Copyright: This article is the copyright of the authors who grant *Diving and Hyperbaric Medicine* a non-exclusive licence to publish the article in electronic and other forms.