Corrigendum

Effect of phase feeding and valine-to-lysine ratio during lactation on sow and piglet performance¹

A. Craig,^{*,†,2} W. Henry,[‡] and E. Magowan[†]

*Agri-Food and Bioscience Institute, Hillsborough, Northern Ireland, BT26 6DR; [†]Queens University Belfast, Belfast, Northern Ireland, BT1 7NN; and [‡]Rekify Ltd, Co. Armagh, Northern, Ireland

© The Author(s) 2019. Published by Oxford University Press on behalf of the American Society of Animal Science. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

J. Anim. Sci. 2019.97:981

doi: 10.1093/jas/sky444

In “Effect of phase feeding and valine-to-lysine ratio during lactation on sow and piglet performance” by A. Craig et al [*Journal of Animal Science*, Volume 94, Issue 9, 1 September 2016,

Pages 3835–3843, <https://doi.org/10.2527/jas.2016-0648>], values for SID in Table 2 had been calculated on the formulated instead of actual. The updated table now contains the correct values.

Table 2: Formulated and actual analysis of experimental diets on an ‘as fed’ (wet) basis

	Control	Flat		Phased	
		Low Valine	Norm Valine	Low Valine	Norm Valine
Formulated Composition					
DE ¹ , MJ/kg	13.5	14.4	14.4	15.0	15.0
CP, g/kg	170	198	198	199	200
Crude Fiber, g/kg	359	266	259	250	251
NDF, g/kg	142	112	109	106	106
Lysine, g/kg	10.0	12.5	12.5	14.0	14.0
Valine, g/kg	8.2	9.5	15.0	11.0	16.8
Valine:Lysine	0.82:1	0.76:1	1.2:1	0.79:1	1.2:1
Actual composition					
DE ¹ , MJ/kg	13.6	14.1	14.4	14.9	15.1
CP, g/kg	144	159	174	156	162
NDF, g/kg	106	91	91	79	91
Lysine, g/kg	8.8	9.2	10.0	11.1	9.6
Valine, g/kg	6.7	6.2	12.0	7.7	10.3
Valine:Lysine	0.76:1	0.67:1	1.2:1	0.69:1	1.07:1
SID ² Lysine, g/kg	7.2	7.5	8.2	9.1	7.9
SID ² Methionine, g/kg	2.2	2.3	2.5	2.7	2.4
SID ² M+C, g/kg	4.2	4.4	4.8	5.4	4.7
SID ² Tryptophan, g/kg	1.4	1.4	1.6	1.7	1.5
SID ² Threonine, g/kg	4.7	4.9	5.3	5.9	5.1
SID ² Valine, g/kg	5	5.3	5.7	6.4	5.5