Risk factors and distribution of MDROs among patients with healthcare associated burn wound infection

Mariam ALfadli¹, Eman M. ELsehsah², Moustapha Ahmed-Maher Ramadan^{3,*}

Abstract

Introduction Patients with burn wounds are more susceptible to develop healthcare associated (HA) infection. Bacterial isolates from HA burn wound infection vary from one place to the other and also differ in terms of antimicrobial susceptibility pattern. The aim of our study was to assess the incidence of HA burn wound infection, risk factors and to determine the microbiological profile and antimicrobial susceptibility pattern of isolated pathogens.

Methods This was a one-year retrospective descriptive study conducted between January 2017 and December 2017, in Al-Babtain Burn and Plastic Surgery center, Kuwait. The identification of bacterial isolates was conducted by conventional biochemical methods according to standard microbiological techniques and commercially available kits using analytical profile index procedure.

Results Total body surface area >35% affected and length of stay more than 14 days were statistically significant risk factors for HA burn infection (RR of 10.057 and 5.912, respectively). Analysis of microbiological profile of these positive cultures indicated that the most common isolated organism was *Acinetobacter baumannii* followed by *Klebsiella pneumoniae*. Multidrug resistant organisms composed 65.85% of the positive isolates. Vancomycin, tigecycline, teicoplanin and linezolid showed 100% effectiveness for all Gram positive isolates. For Gram negative organisms, imipenem and meropenem showed 38.71% and 41.93% efficacy, respectively.

Conclusion Identification of antimicrobial susceptibility patterns helps to tailor the required antibiotic policy to minimize the acquired infections among these vulnerable patients.

Keywords Epidemiology, antimicrobial resistance, health care associated infection, burn, infection control, surveillance.

Introduction

Burn injuries are part of the most common and serious types of trauma, representing a major public health concern all over the world.¹ Hospitalized patients with burn wounds are more vulnerable to infection due to a variety of factors such as the loss of the normal protective skin barrier, impaired immune system and the

presence of dead and necrotic tissues that provide an environment conducive to the growth and proliferation of pathogens.² About 75% of all deaths after burn injuries are related to infections, with a high rate of healthcare associated infection in burn units.³

As bacterial isolates from burn patients change with time, it is essential to update the

Received: 17 September 2018; revised: 07 November 2018; accepted: 15 November 2018.

¹PhD, Director of Infection Control Directorate, Ministry of Health, Sabbah health region, P.O.Box: 12414 Alshamiya, Kuwait; ²PhD, Lecturer at Microbiology and Immunology Department, Faculty of medicine, Mansoura University, Egypt, Specialist of Infection Control at Infection Control Directorate, Ministry of Health, Sabbah health region, P.O.Box: 12414 Alshamiya, Kuwait; ³PhD, Fellow at Community Medicine Department, Faculty of Medicine, Alexandria University, Egypt, Specialist of Infection Control at Infection Control Directorate, Ministry of Health, Sabbah health region, P.O.Box: 12414 Alshamiya, Kuwait.

*Corresponding author: Moustapha Ahmed-Maher Ramadan, PhD, Fellow at Community Medicine Department, Faculty of Medicine, Alexandria University, Egypt, Specialist of Infection Control at Infection Control Directorate, Ministry of Health, Sabbah health region, P.O.Box: 12414 Alshamiya, Kuwait. ficas_alex@yahoo.fr

Article downloaded from www.germs.ro

Published December 2018

© GERMS 2018

ISSN 2248 - 2997

ISSN - L = 2248 - 2997

microbiological profile of burn wound infection periodically with identification of the most common isolated pathogens and their antimicrobial sensitivity pattern in order to improve management and outcome.⁴

Antibiotic resistance is an emerging concern among patients with healthcare associated infection (HAI) in this era, increasing rapidly and becoming a public health threat; this resistance is in part due to misuse and over use of antibiotic prescription.⁵ The emergence of highly drug resistant bacteria in HAI of burn patients, with limited therapeutic options, is an alarming new concern with subsequent morbidity and mortality. Therefore, in depth knowledge of the microbiological profile and antibiotic susceptibility pattern of HAI in patients with burn injuries is essential because many patients with wound infection need to receive antibiotic therapy before availability of the results of microbiological cultures as this is particularly important to reduce morbidity and mortality related to infection.⁶

The aim of our study was to assess the incidence of health care associated burn wound infection, risk factors and determination of microbiological profile and antimicrobial susceptibility pattern of isolated pathogens focusing on the distribution of multidrug resistant organisms (MDRO) that can be helpful in the design of empirical antibiotic policy for this category of patients in our setting.

Methods

Study design

This was a one-year prospective study conducted between January 2017 and December 2017, in Al-Babtain Burn and Plastic surgery center, Kuwait.

The study protocol was reviewed and approved by the research committee of the Ministry of Health Kuwait (887/30 Oct 2018).

Study setting

Al-Babtain Burn and Plastic Surgery center is the only referral center for burn patients in Kuwait City, State of Kuwait. Data were retrieved from patients' medical record and infection control surveillance sheets.

Inclusion criteria

All patients presenting with burn injury and requiring admission were included.

Exclusion criteria

Patients with minor burn injury treated on outpatient department basis.

Patients transferred from plastic surgery ward to burn ward and not presenting with burn injury.

Identification of healthcare associated burn wound infection was based on the Centers for Disease Control and Prevention (CDC) criteria for year 2017, which is "a change in burn wound appearance or character, such as rapid eschar separation, or dark brown, black, or violaceous discoloration of the eschar", with "organism(s) identified from blood by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment".⁷

Microbiological analysis; sample collection and inoculation

Blood cultures were processed using manual standard method (Vitek, Marcy-l'Étoile, France). All blood specimens were put into thioglycolate broth and subcultured after overnight incubation on 5% blood agar, Mac Conkey agar and Chocolate agar plates and incubated overnight at 37°C aerobically.

The identification of bacterial isolates was conducted by conventional biochemical methods according to standard microbiological techniques and commercially available kits, using the analytical profile index procedure (API; Vitek, bioMérieux, Hazelwood, MO, USA).⁸

Antibiotic susceptibility was determined by using classical Kirby Bauer disc diffusion method on Mueller-Hinton agar according to Clinical and Laboratory Standards Institute (CLSI) guidelines.⁹ Interpretation of antimicrobial resistance and susceptibility was determined using CLSI 2015 recommendations.¹⁰

For identification of MDRO we used the definitions outlined by Magiorakos et al. to define multidrug-resistant (MDR) as "acquired

non-susceptibility to at least one agent in three or more antimicrobial categories”.¹¹

Statistical analysis

Different risk factors were assessed to identify the risk of developing healthcare associated burn wound infection. For total body surface area (TBSA) patients were divided according to affected TBSA percentage into two groups: the first group with TBSA% >35% and the second group with TBSA% ≤35%. Regarding the length of hospital stay (LOS), patients were divided into two groups: patients above the mean hospital stay for the studied sample and patients who stayed equally or below the mean calculated length of stay.

Relative risks were calculated to show significance in developing healthcare associated burn wound infection and acquiring multidrug resistant organism. Fischer's exact test was used to identify significance at p value <0.05. Statistical analysis was performed using SPSS version 18 (IBM, Chicago, IL, USA).

Results

The study population was composed of 201 patients. Male patients represented 61.19% of the study sample. The flame burn was the most common type of injury among our studied population (62.69%), while scald came second common (27.86%) and all other injuries represented only 9.45% (Table 1).

For the 201 patients who constituted 2834 patient days, the affected TBSA ranged from 1%-98% with a mean of 19.7% ± 17.65%. The average length of hospital stay was 14 days ± 18.14 days, while for patients who developed healthcare associated burn infection the average length of stay was 31 days ± 25.48 days.

During the period of study, 17 patients developed healthcare associated burn wound infection with an incidence rate of 6/1000 patient days. Eleven male patients developed healthcare associated burn wound infection, and six patients were female. This distribution showed no statistical significance (p=0.756). All patients who developed healthcare associated burn infection had a flame injury, and this was statistically significant (p<0.001). Patients with

Table 1. Distribution of studied risk factors among patients with burn injuries in Al Babbain center during 2017

Risk factor category	N	%
Gender		
Male	123	61.19
Female	78	38.81
Type of burn		
Flame	126	62.69
Scald	56	27.86
Contact	10	4.98
Electric	8	3.98
Chemical	1	0.50
TBSA		
>35%	25	12.44
≤35%	176	87.56
Length of hospital stay		
>14 days	58	28.86
≤14 days	143	71.14

TBSA – total body surface area.

burns covering more than 35% of TBSA were found to be statistically more likely to develop healthcare associated burn infection (p<0.001). Patients who stayed more than the mean LOS of the study (14 days) were more prone to develop healthcare associated infection (p<0.001). These data are presented in Table 2. Further statistical analysis for the TBSA and LOS by linear regression showed their statistical significance as risk factors to develop healthcare burn associated infection (p<0.000).

The 17 patients who developed healthcare associated burn wound infection revealed 41 positive blood cultures. These were repeated blood cultures collected at different time points for each patient within the infection period. On the other hand, TBSA of burn wound and LOS did not show any statistical significance among patients with healthcare associated burn infection regarding acquisition of multidrug resistant organisms in blood cultures. These data are presented in Table 3.

Analysis of the microbiological profile of these positive cultures indicated that the most commonly isolated organism was *Acinetobacter*

Table 2. Risk of acquiring healthcare associated burn wound infection in relation to studied risk factors in patients with burn injuries in Al Babbain center during 2017

Risk factor	Patient with healthcare associated burn infection	Patient without healthcare associated burn infection	Relative risk	Confidence interval
Male	11	112	1.163	0.448 – 3.017
Female	6	72		
Flame burn	17	109	20.945	1.278 – 343.3
Other burn injuries	0	75		
Patient with >35 TBSA burn	10	15	10.057	4.212 – 24.013
Patient with ≤35 TBSA burn	7	169		
LOS >14 days	12	46	5.917	2.182 – 16.047
LOS ≤14	5	138		

LOS – length of hospital stay; TBSA – total body surface area.

Table 3. Risk of acquiring MDROs for patients with healthcare associated burn wound infection in relation to studied risk factors in patients with burn injuries in Al Babbain center during 2017

Risk factor	Blood culture with MDRO in patients with healthcare associated burn infection	Blood culture without MDRO in patients with healthcare associated burn infection	Relative risk	Confidence interval
Male	15	12	0.648	0.435 – 0.966
Female	12	2		
Patient with >35 TBSA burn	20	9	1.182	0.691 – 2.025
Patient with ≤35 TBSA burn	7	5		
LOS >14 days	18	11	0.827	0.536 – 1.276
LOS ≤14	9	3		

HAI – healthcare associated infection; LOS – length of hospital stay; MDRO – multidrug resistant organisms; TBSA – total body surface area.

baumannii 17 (41.46%) followed by eight *Klebsiella pneumoniae* (19.51%). Multidrug resistant organisms composed 27 (65.85%) of the positive isolates. All *Acinetobacter baumannii* isolates were MDR (100%). These data are shown in Table 4.

Regarding the antimicrobial sensitivity pattern among Gram positive organisms we found that (66.67%) of the isolates of *Staphylococcus aureus* were MRSA and that both *Enterococcus gallinarum* and *Enterococcus faecalis* were sensitive to penicillins. Vancomycin,

tigecycline, teicoplanin and linezolid showed 100% susceptibility in all Gram positive isolates. This is presented in Figure 1.

For Gram negative organisms, tigecycline was 100% active on tested isolates, while imipenem and meropenem were active on 38.71% and 41.93% of the isolates, respectively. In our study we found one isolate (*Acinetobacter baumannii*) which showed resistance to colistin. This is shown in Figure 2.

Table 4. Microbiological profile and distribution of MDRO among patients with healthcare associated burn infection in Al Babtain center in 2017

Microorganism	No. of isolates	Percentage	Number of MDR isolates	Percentage of MDR isolates*
<i>Acinetobacter baumannii</i>	17	41.46	17	100
<i>Klebsiella pneumoniae</i>	8	19.51	4	50
<i>Staphylococcus aureus</i>	6	14.63	4	66.67
<i>Pseudomonas aeruginosa</i>	6	14.63	1	16.66
<i>Enterococcus faecalis</i>	2	4.88	0	0
<i>Serratia marcescens</i>	1	2.44	1	100
<i>Enterococcus gallinarum</i>	1	2.44	0	0
TOTAL	41	100	27	65.85

*The percentage is calculated from the total isolates number of the same microorganism
MDR – multidrug resistant.

Figure 1. Antimicrobial susceptibility pattern for Gram positive isolates in patients with healthcare associated burn wound infections in Al Babtain center during 2017

Discussion

Healthcare associated infections represent a serious issue for health services all over the world, impacting the survival of high-risk patients, such as burn patients. Burn site infections can influence both the survival of the patient and the outcome of reconstructive treatment.¹²

In our study, we revealed that the incidence rate of healthcare associated burn infection was 6/1000 patient days. This rate is higher than that of the study of Van Duin and his colleagues who identified a rate of 1.78/1000 patient days.¹³

However, another comprehensive study of the epidemiology of burn infection showed that the

rate among patients with TBSA >30% exceeds 5/1000 patient days.¹⁴ It is worth to mention that our setting is the only referral center for burn patients in the state of Kuwait, so any age group or degree of burn, type of burn injury or TBSA will be receiving management in our setting.

Our findings showed that flame burn is the most common (62.69%) cause of burn injury. This is similar to a systematic review of epidemiology of burn injuries in the east Mediterranean Region and reports from India and Nepal.¹⁵⁻¹⁷ Several factors may contribute to the predominance of flame burn injury. Diverse intentions for flame injuries have been reported,

Figure 2. Antimicrobial susceptibility pattern for Gram negative isolates in patients with healthcare associated burn infections in Al Babbain center during 2017

marital problems, family problems, psychological and psychiatric disorders as well as poverty. Low socioeconomic status with poorly secured liquefied petroleum gas cylinders, or cooking equipment, or using open fire for cooking are usually implicated in flame burn injuries.^{15,17}

A total body surface area affected above 35% was shown to be a significant risk factor for developing healthcare associated burn infection in our study. This is similar to different studies conducted in the USA, Turkey and Egypt, which showed that a TBSA above 30% is an independent risk factor for burn wound infection.^{14,18,19} Increase in TBSA is usually associated with loss of natural physical defense, decrease in patient immune system, and these patients are usually in need of invasive devices. All these factors will lead to an increased risk of HAIs.

Patients who spend more time in the hospital are more vulnerable to develop HAIs. Our patients spent an average of 14 days in the hospital, this is in accordance with an Iranian study and a systematic review of different countries in the eastern Mediterranean Region that showed similar duration of hospital stay.^{15,20} Van Duin and his colleagues reported that half of the patients who spend 80 days from admission will experience at least one HAI.¹³ Also Coetzee et al. reported that the risk of wound infection

increases by time.²¹ Burn wounds are known to be good media for bacterial growth, and colonization occurs shortly after the injury as virulent microorganism strains invade the skin and soft tissue; as the host defense is weak, an increase in the number of microorganisms will be able to reach the venous system to cause infection.²²

In the current study, the most commonly isolated microorganisms from healthcare associated burn wound infection were *Acinetobacter baumannii* (41.46%) followed by *Klebsiella pneumoniae* (19.51%) then *Staphylococcus aureus* and *Pseudomonas aeruginosa* (14.63% each). This result is in agreement with a Turkish study which revealed *Acinetobacter baumannii* as the most common pathogen isolated (23.6%); in the same study the percentage of *S. aureus* was near to our value (11.2%).²³

The microbiological profile in our study is somewhat different from that shown in a related study done by Rathod et al., which showed *Pseudomonas aeruginosa* as the most frequent isolate (34.93%).²⁴

Possible explanations to the difference in the microbiological profile may be due to differences in geographic location or patient age (children and adults), differences in hygienic measures applied in healthcare facilities and associated

cross-infection by the hands of healthcare personnel.

In the current study, we found that MDROs represented 65.85% of the positive cultures; 100% of *Acinetobacter baumannii* strains were resistant, resistance in *S. aureus* strains was 66.67%, while for *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* the percentage of resistant strains was 50% and 11.11%, respectively. These findings are in agreement with a related Egyptian study conducted in 2016 which showed that multidrug resistant Gram negative organisms represented about 60% of the isolates.²⁵

Regarding the antibiotic sensitivity pattern of the isolates from healthcare associated burn infections, Gram positive organisms were 100% sensitive to vancomycin, teicoplanin, tigecycline and linezolid, and to a lower extent to other antimicrobials such as clindamycin (88%) and gentamicin (66%). Similar results are shown in the study of Rathod et al. that revealed 100% sensitivity to vancomycin and linezolid followed by clindamycin (64.55%).²⁴

In our study, sensitivity to imipenem was only 38.71% in Gram negative isolates; this may be explained by overuse of imipenem in our setting to improve the clinical outcome of such critically ill patients. This result is similar to that found by Mohammed in which sensitivity to imipenem was 35%.²⁵ In another study by Rathod VS et al., in which *Pseudomonas aeruginosa* was the most common isolate, the organism showed 75% resistance to ceftazidime.²⁴

Lack of compliance to the infection control measures, absence of an antimicrobial stewardship program, and patients' characteristics, especially for this vulnerable category, all can lead to emergence of resistant microorganisms.

Among study limitations, the absence of an electronic file system has limited the collection of other potential risk factors such as degree of burn, time of injury and different source of flame burn. Also, the establishment of the MDRO profile depended on the disc diffusion method not on PCR for detection of resistance genes due to unavailability of these resources in the hospital lab.

Conclusion

Identification of the microbiological profile of HAI in burn wounds is of great importance both epidemiologically and to improve the clinical outcome, as this data may guide effective empirical antibiotic therapy. In our setting the most prevalent microorganism identified was *Acinetobacter baumannii* and all its isolates were found to be MDR. Proper compliance with infection control measures and development of an antibiotic stewardship program will help in reducing the number of infections and the emergence of resistant strains for the critically ill patient with burn wound injury.

Authors' contributions statement: MA, EE, MAMR performed the literature review. MAMR designed the study, collected the data and performed the statistical analysis. EE and MAMR performed the data interpretation. All authors prepared the first draft, read and approved the final version of the manuscript.

Conflicts of interest: All authors – none to declare.

Funding: None to declare.

References

1. Mock C, Peck M, Peden M, et al. A WHO plan for burn prevention and care. Geneva: World Health Organization; 2008.
2. Bhat VG, Vasaikar SD. Bacteriological profile and antibiogram of aerobic burn wound isolates in Mthatha, Eastern Cape, South Africa. South Afr J Epidemiol Infect 2010;25:16-9. [[Crossref](#)]
3. Fouzia B, Damle AS, Maher G. Changing patterns of burn infections IOSR J Dent Med Sci 2013;5:114.
4. Kumarasamy KK, Toleman MA, Walsh TR, et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. Lancet Infect Dis 2010;10:597-602. [[Crossref](#)]
5. Lushniak BD. Antibiotic resistance: a public health crisis. Public Health Rep 2014;129:314-6. [[Crossref](#)]
6. Srinivasan S, Vartak AM, Patil A, Saldanha J. Bacteriology of the burn wound at the Bai Jerbai Wadia Hospital for children, Mumbai, India – A 13 year study, Part I-Bacteriological profile. Indian J Plast Surg 2009;42:213-8.
7. Centers for Disease Control and Prevention. CDC/NHSN surveillance definitions for specific types of infections. Accessed on 12 August 2018. Available at: https://www.cdc.gov/nhsn/PDFs/pscManual/17pscNosInfDef_current.pdf.
8. Immunochemical methods used for organism detection. In: Forbes BA, Sham DF, Weissfeld AS. AS Bailey &

- Scott's Diagnostic Microbiology, 12th Edition. St Louis, Missouri: Mosby; 2007, p 147-158.
9. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial disk susceptibility tests; approved standard - twelfth edition. CLSI document M02-A12. Wayne, PA: Clinical and Laboratory Standards Institute; 2015.
 10. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; twenty-fifth informational supplement. CLSI document M100-S25. Wayne, PA: Clinical and Laboratory Standards Institute; 2015.
 11. Magiorakos AP, Srinivasan A, Carey RB, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012;18:268-81. [\[Crossref\]](#)
 12. Cosgrove SE. The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clin Infect Dis* 2006;42 Suppl 2:S82-9. [\[Crossref\]](#)
 13. van Duin D, Strassle PD, DiBiase LM, et al. Timeline of healthcare-associated infections and pathogens after burn injuries. *Am J Infect Control* 2016;44:1511-6. [\[Crossref\]](#)
 14. Weber J, McManus A, Nursing Committee of the International Society for Burn Injuries. Infection control in burn patients. *Burns* 2004;30:A16-24. [\[Crossref\]](#)
 15. Othman N, Kendrick D. Epidemiology of burn injuries in the East Mediterranean Region: a systematic review. *BMC Public Health* 2010;10:83. [\[Crossref\]](#)
 16. Maske AN, Deshmukh SN. Clinico-epidemiological study of burns: our experience with 500 patients. *Int Surg J* 2016;3:1234-9. [\[Crossref\]](#)
 17. Tripathi S, Basnet SJ. Epidemiology of burn injuries in Nepal: a systemic review. *Burns Trauma* 2017;5:10. [\[Crossref\]](#)
 18. GÜldoğan CE, Kendirci M, Tikici D, Gündoğdu E, Yasti A. Clinical infection in burn patients and its consequences. *Ulus Travma Acil Cerrahi Derg* 2017;23:466-71. [\[Crossref\]](#)
 19. El-Maghawry HAMM, El Nem W, Sherif N, Hagag SA. An interventional study to decrease healthcare associated burn wound infections in the burn unit of Al Ahrar Hospital in Zagazig city, Sharkia Governorate. *Int J Curr Microbiol Appp Sci* 2016;5:566-78. [\[Crossref\]](#)
 20. Emami SA, Karimi H, Alijanpour A. Epidemiology of burn wound infection and its bacterial resistance, burn registry program. *Merit Res J Med Med Sci* 2015;3:135-9.
 21. Coetzee E, Rode H, Kahn D. *Pseudomonas aeruginosa* burn wound infection in a dedicated paediatric burns unit. *S Afr J Surg* 2013;51:50-3.
 22. Essayagh M, Essayagh T, Essayagh S, El Hamzaoui S. [Epidemiology of burn wound infection in Rabat, Morocco: Three-year review]. *Med Sante Trop* 2014;24:157-64. [\[Crossref\]](#)
 23. Bayram Y, Parlak M, Aypak C, Bayram I. Three-year review of bacteriological profile and antibiogram of burn wound isolates in Van, Turkey. *Int J Med Sci* 2013;10:19-23. [\[Crossref\]](#)
 24. Rathod VS, Kasturi, Raut SS. Emergence of multi-drug resistant strains among bacterial isolates in burn wound swabs in a tertiary care centre, Nanded, Maharashtra, India. *Int J Res Med Sci* 2017;5:973-7. [\[Crossref\]](#)
 25. Mohamed H. One year prevalence of critically ill burn wound bacterial infections in surgical ICU in Egypt: Retrospective study. *Egypt J Anesth* 2016;32:431-4. [\[Crossref\]](#)

Please cite this article as:

ALfadli M, EL-sehsah EM, Ramadan MAM. Risk factors and distribution of MDROs among patients with healthcare associated burn wound infection. *GERMS* 2018;8(4):199-206. doi: 10.18683/germs.2018.1147.