Published in final edited form as:

Neuromuscul Disord. 2018 October ; 28(10): 868–877. doi:10.1016/j.nmd.2018.07.005.

GsMTx4-D provides protection to the D2.*mdx* mouse

Christopher W. Ward^a, Frederick Sachs^b, Ernest D. Bush^c, and Thomas M. Suchyna^{b,*}

^aDepartment of Orthopedics, University of Maryland School of Medicine, 100 Penn Street, Room 515C, Baltimore, MD 21201, USA

^bDepartment of Physiology and Biophysics, University of Buffalo, 301 Cary Hall, Buffalo, NY 14214, USA

^cAkashi Therapeutics, 245 First Street, 18th Floor, Cambridge, MA 02142, USA

Abstract

Duchenne muscular dystrophy is a life-limiting muscle disease that has no current effective therapy. Despite mounting evidence that dysregulation of mechanosensitive ion channels is a significant contributor to dystrophy pathogenesis, effective pharmacologic strategies targeting these channels are lacking. GsMTx4, and its enantiomer GsMTx4-D, are peptide inhibitors of mechanosensitive channels with identical activity. In previous studies, acute *in vitro* application of GsMTx4 to dystrophic murine muscle effectively reduced the excess MSC dependent calcium influx linked to contraction-induced muscle damage. Here we sought to determine if *in vivo* treatment with GsMTx4-D proffered benefit in the D2.*mdx* mouse. GsMTx4-D showed a 1-week half-life when administered by subcutaneous injection over four weeks. Informed by these results, D2.*mdx* mice were then treated by a subcutaneous injection regimen of GsMTx4-D for six weeks followed by determination of muscle mass, muscle susceptibility to eccentric contraction injury and multiple histological indicators of disease progression. The mice showed a reduction in the loss of muscle mass and a decrease in susceptibility to contraction induced injury. These protective effects were realized without reduction in fibrosis, supporting a model where GsMTx4-D acts directly on muscle cells. We propose GsMTx4-D represents a promising new therapy to slow disease progression and may complement other therapies such as anti-inflammatory agents and gene-replacement strategies.

Keywords

GsMTx4; Dystrophy; Treatment; D2.*mdx* mouse; Eccentric contraction injury

1. Background

DMD is a muscle disease caused by the genetic depletion of dystrophin, a cytoskeletal protein whose functions include the organization of sarcolemmal and cytoskeletal components [1] and regulation of signaling proteins. In muscle from *mdx* mice (model of human DMD) the absence of dystrophin is linked to an increase in the permeability of the

*Corresponding author. suchyna@buffalo.edu (T.M. Suchyna).

sarcolemma to extracellular calcium (Ca^{2+}) [2,3]. Given firm evidence linking excess Ca^{2+} permeability to the cellular dysfunction that drives the dystrophic phenotype, pharmacologic approaches targeting dysregulated Ca^{2+} influx have been a major focus towards a strategy to halt or slow disease progression in DMD.

While studies have identified multiple channel candidates that may contribute to the excess Ca^{2+} permeability in DMD muscle, their targeting has generated mixed results. Preclinical efficacy studies targeting voltage-activated Ca^{2+} channels in *mdx* mice have shown inconsistent results [1] and disease progression appears to be unaffected by treatment in most human trials [4,5]. Antimicrobial aminoglycosides, nonselective cationic channel inhibitors, have shown promise in short-term efficacy trials (6 week) in *mdx* mice; [6], however longer-term treatment (up to 6 months) was associated with an increase in degeneration of the heart and diaphragm. These deleterious effects may result from the non-specific pore blocking of aminoglycosides [7], and their impact on protein expression [8] and the inositol signaling system [9].

In addition to its scaffold function, dystrophin participates in mechanotransduction by linking the dystrophin–glycoprotein complex (DGC) to the underlying cytoskeleton. Recent evidence shows that the loss of dystrophin leads to increased mechanosensitive ion channels (MSCs) activity [10,11], underscored by the increased sarcolemmal fragility [12–14]. GsMTx4 is a 34 amino acid peptide representing a new class of drugs that selectively inhibits nonselective cationic MSCs [15]. GsMTx4 does not bind directly to the channels, but inhibits MSC gating by association with the cell membrane and buffering tension transmission to the channels [16]. *In vitro* studies have demonstrated that GsMTx4 is effective at reducing excess stretch-dependent sarcolemma Ca^{2+} influx through these channels [10,12,14,17–21]. GsMTx4-D is a synthetic enantiomeric form of GsMTx4 made from D amino acids and it has the same potency as the WT peptide [22]. Given evidence that GsMTx4 acts as a cardioprotective in ischemic reperfusion injury [23] and mechanically induced atrial fibrillation [24], we reasoned that GsMTx4-D dependent suppression of MSC activity may proffer benefit as treatment in DMD. The goal of this study was to assess the short-term efficacy of GsMTx4-D in a disease-relevant murine model of DMD. If successful, this study would serve as a foundation for exploring GsMTx4-D as a chronic therapy for DMD.

2. Methods

2.1. Animal model

We chose the D2.*mdx* murine model of DMD because the *mdx* crossed into the DBA/2 J background (i.e., D2.*mdx*) yields a severe dystrophic phenotype that models the human pathogenesis; a severity not observed in other commonly used murine dystrophic models [25,26]. Clearance pharmacokinetics and efficacy testing were conducted in male DBA/2 J and D2.*mdx* mice at 6–9 weeks of age. All mice were obtained from Jackson Laboratories (Bar Harbor, ME, USA). Animal care and procedures were approved and performed in accordance with the standards set by the University of Maryland Institutional Animal Care and Use Committee and by the National Institutes of Health (NIH).

2.2. Pharmacokinetics

We chose 0.1–5.0 μM in skeletal muscle as our target range based on previous studies [14,18,19,24,27]. We previously demonstrated that a single dose of 50 mg/kg subcutaneous dosing achieved $\sim 1 \mu\text{M}$ concentration in the heart [23]. Here we conducted accumulation and clearance testing in five key tissues, plasma and urine by a continuous injection regimen. Cohorts of five D2.*mdx* mice were subcutaneously injected into the back with 10 mg/kg every other day. Dosage times and sacrifice times were as follows:

- 5 mice injected with 10 mg/kg GsMTx4, 3 times over 7 days – sacrifice on day 7.
- 5 mice injected with 10 mg/kg GsMTx4, 6 times over 14 days – sacrifice on day 14.
- 5 mice injected with 10 mg/kg GsMTx4, 6 times over 14 days – sacrifice on day 21.
- 5 mice injected with 10 mg/kg GsMTx4, 6 times over 14 days – sacrifice on day 28.

Two cohorts were sacrificed at 7 and 14 days during GsMTx4-D accumulation. The peptide injections were ceased at 14 days for the remaining two cohorts which were then sacrificed at 21 and 28 days to determine clearance rate. The kidney, liver, heart, gastrocnemius, diaphragm, blood plasma and urine samples were harvested, flash frozen and shipped to Custom Biologics (Mississauga, Canada) for liquid chromatography coupled with tandem mass spectrometry analysis. Each tissue sample was individually analyzed and the mean concentrations determined.

2.2.1. In vivo muscle strength testing and eccentric contraction-induced injury

Muscle performance was measured *in situ* with a 305C-FP muscle lever system (Aurora Scientific Inc., Aurora, Canada). Mice were anesthetized (isoflurane; to effect), and placed on a thermostatically controlled table. The knee was stabilized and the foot firmly fixed to the footplate of a torque sensor (Fig. 1). Isometric contractions were elicited from the gastrocnemius muscle with needle electrodes inserted behind the knee in proximity to the sciatic nerve. Optimal isometric twitch torque was determined by increasing the current with a minimum of 30 s between each contraction to avoid fatigue.

- *Peak isometric torque:* Peak isometric tetanic torque contractions (0.2 ms pulse, 250 ms train, 150 Hz) were determined by taking the peak of three successive contractions elicited 3 min apart. The muscle was then allowed to rest for 5 min.
- *Injury susceptibility:* Mice were injected with Evans Blue dye (EBD) (1 mg/kg i.p.) 24 h prior to the contraction assay. The susceptibility to contraction induced was examined with 20 tetanic contractions (100 Hz, 0.2 ms pulse width, 250 ms duration) elicited where at 200 ms, the muscle was stretched at a rate of $40^\circ/\text{s}$ through a 30° angle [12,13]. The muscle was then passively returned to resting length. Maximal force of the isometric plateau (prior to the eccentric stretch) was measured and used to normalize the force decrease.

2.2.2. Histology—The gastrocnemius muscle from unstimulated (resting) and stimulated (exercised) legs were snap frozen in isopentane cooled in liquid N₂ in Cryomatrix. Tissues were sectioned from the muscle midbelly using a cryostat, air dried to a glass slide, and fixed with 4% paraformaldehyde. In exercised legs, muscle permeability post-exercise was assessed by dye incorporation within each muscle as the cross sectional area of EBD containing muscle fibers expressed as a % of total muscle area. To determine the extent of fibrotic inclusions, and central located nuclei, unstimulated sections of muscle were labeled with Texas Red-conjugated wheat germ agglutinin (WGA) to decorate the extracellular matrix and nuclei as described [28]. The mean pixel area of the WGA label, the cross-sectional areas of the muscle fibers, and percentage of those myofibers with central nuclei, were determined in two regions of interest (ROI) per muscle (two ROI's; one from each head of the gastrocnemius muscle) using Nikon Elements Software analysis tools.

3. Results

3.1. GsMTx4-D tissue accumulation and clearance

Our target concentration of 0.1–5.0 μM in the skeletal muscle was based on multiple studies showing (1) A concentration of 1 μM GsMTx4 inhibited MSC current by > 50% in patch-clamp studies [10,14,15], and (2) reports of *in vitro* experiments showing 0.1–5.0 μM effective in suppressing MSC activity in various disease models [14,18,19,24,27].

Our previous pharmacokinetic study, aimed at determining rapid tissue exposure to GsMTx4-D in C57BL/6 J mice [23] demonstrated that a single subcutaneous dose of 50 mg/kg achieved $\sim 1 \mu\text{M}$ concentration in the cardiac muscle within 24 h. Here, we wanted to develop a dosing regimen that could be used for extended periods; so, we performed subcutaneous injections of 10 mg/kg of GsMTx4-D into D2.*mdx* mice every other day to determine the accumulation (7 or 14 days) and clearance (7 or 14 days) profile of the peptide (Fig. 2). Five tissues and two body fluids (plasma and urine) were followed over both phases. We show that gastrocnemius and cardiac muscle achieved a $\sim 2 \mu\text{M}$ concentration of GsMTx4-D while diaphragm approached 8 μM . Liver and kidney, two highly vascularized organs, yielded > 10 fold higher concentrations than any of the muscle groups. During the clearance phase all tissues showed a > 50% decrease in concentration after one week that dropped to between 12–19% of the peak concentration after two weeks. Blood concentration dropped rapidly to near 0 after injections ceased, as was previously observed with higher temporal resolution [23]. This suggests that tissue binding sites have higher affinity than plasma binding sites. Urine analysis was limited since samples were not attainable from all animals, though we observed a trend of increased secretion during the accumulation period that decreased after ceasing injections.

3.2. Six week efficacy study

Based on our published pharmacokinetic analyses [23] and the clearance study conducted here (Fig. 2), we designed a six-week *in vivo* efficacy study with muscle force and susceptibility to eccentric injury as the primary outcome measure. There were four study groups ($n = 10$ mice/group): D2.*mdx* treated with vehicle (saline), or GsMTx4-D at the target concentration (10 mg/kg) or at a 10-fold lower concentration (1 mg/kg) to determine if

the target approximates the least effective dose. A group of DBA2 mice treated with vehicle served as a non-diseased control. The clearance data showed that muscle concentrations remained in the target range two weeks after ceasing injections suggesting we could reduce the time between doses following a period of accumulation over the six-week test. Therefore, dosing began at ~8 weeks of age with subcutaneous injections on alternate days for 2 weeks followed by dosing every 4th day for the remaining 4 weeks.

3.2.1. Bodyweight and muscle mass—The D2.*mdx* exhibited significant losses in bodyweight and muscle weight compared to DBA2 controls (Fig. 3A). In response to GsMTx4-D treatment, the D2.*mdx* exhibited a non-significant trend in protection from loss of bodyweight. In contrast to the impact on bodyweight, the 10 mg/kg treatment was shown to effectively protect the disease driven loss in gastrocnemius muscle weight compared to the untreated D2.*mdx* (Fig. 3B).

3.2.2. Muscle function in vivo—Nerve-evoked isometric torque generated by the plantarflexor group (i.e., gastrocnemius, soleus) was assessed in deeply anesthetized mice as described [12,13]. Consistent with the impact of the dystrophic pathology, we observed a significant deficit in maximal muscle torque in the D2.*mdx* compared to the DBA2 control (Fig. 4A). Treatment of the D2.*mdx* with 10 mg/kg of GsMTx4-D proffered benefit in isometric torque compared to either the 1 mg/kg or vehicle treated D2.*mdx*. Given that normalizing the isometric torque to the mass of the muscle accounted for the identified differences in the total torque output (Fig. 4B), we conclude that protection from the loss of muscle mass was responsible for the treatment induced increase in muscle torque.

3.2.3. Susceptibility to eccentric injury—Following the determination of maximal muscle force, we examined the susceptibility to eccentric contraction injury using 20 eccentric contractions (Fig. 5) as described [12,13]. Consistent with dystrophic disease increasing injury susceptibility, the D2.*mdx* exhibited a ~70% drop in isometric force after 20 eccentric contractions (Fig. 5A) In response to short term treatment with 10 or 1 mg/kg GsMTx4-D, we found a significant protection (~43% and 30%, respectively) from the eccentric contraction induced injury.

Prior to the contractility testing, mice were injected (24 h prior; $n = 5$ per group; i.p. injected) with Evans Blue dye (EBD). Following the eccentric contraction protocol (15 min post), the gastrocnemius was flash frozen and transversely sectioned to assess EBD accumulation in the muscle fibers. Consistent with the increased susceptibility to eccentric injury seen in the D2.*mdx*, we identified a significant increase in the percentage of EBD positive muscle fibers in the D2.*mdx* vs. the DBA. However, GsMTx4 treatment had no significant impact on preventing the eccentric contraction induced EBD dye accumulation (Fig. 5B).

3.2.4. Histopathology—Reports have aligned the enhanced dystrophic phenotype in the D2.*mdx* to an increase in inflammatory signaling of the DBA2 background. In this strain a mutation in Latent TGF Binding Protein 4 (LTBP4) [25,29] underscores a dramatic increase in fibrosis in the D2.*mdx* [29]. In muscle cryosections we used fluorescent conjugated wheat germ agglutinin (Texas red) to quantify the abundance of interstitial fibrosis and to identify

the presence of centrally nucleated muscle fibers (CNF). As previously reported [25,29], we found significantly greater levels of interstitial fibrosis and CNF's in the D2.*mdx* compared to DBA2 (Fig. 6A). Treatment with GsMTx4-D had no impact on the level of fibrosis or CNF's in the D2.*mdx* muscle (Fig. 6B and C).

The muscle fiber cross-sectional area (CSA) was quantified to further elucidate the effect of GsMTx4-D on gastrocnemius size. Population histograms of CSA (Fig. 7A) revealed a relative absence of hypertrophic muscle fibers, and an increased percentage of smaller diameter myofibers in the D2.*mdx*, an effect further highlighted when results were visualized as a plot of CSA cumulative percentage (Fig. 7C). This result aligned with a previous report in the D2.*mdx* [25]. In response to GsMTx4-D treatment we revealed that myofibers from the [1 μ M] group had a larger overall diameter than the non-treated D2.*mdx* with the [10 μ M] treatment group having an even larger impact when compared to the low-dose treated or the non-treated D2.*mdx*. Taken together, we conclude that GsMTx4-D treatment impacted muscle function by sparing the disease-dependent decrease in muscle fiber CSA.

4. Discussion

4.1. Pharmacokinetics

Our previous [23] and current pharmacokinetic studies demonstrated our ability to achieve a [0.1–5 μ M] of GsMTx4-D in skeletal muscle and heart by subcutaneous injections. Most pharmaceutical peptides of less than 70 kDa (kidney filtration cutoff size) suffer from short half-lives due to filtration by the kidney and susceptibility to rapid enzymatic digestion [30] (e.g. 1–2 min for serum half-life of glucagon [31]). However D amino acid peptides are less susceptible to enzymatic digestion which affords longer half-lives. The wash-out study revealed that concentrations in gastrocnemius, heart and diaphragm muscle were still within ~20% of the peak concentration a week after ceasing injections; concentrations still within the expected therapeutic range. The long half-life of GsMTx4-D in tissues, along with its rapid depletion from the blood, suggests that GsMTx4-D has a higher affinity for tissue than for serum proteins, and that GsMTx4-D is resistant to enzymatic digestion or instability.

4.2. Molecular targets

The specific target(s) of GsMTx4 in skeletal muscle need to be rigorously determined. However, both the L and D enantiomers have been shown to inhibit mechanosensitive Piezo channels [32] and multiple TRP channel family members [20,33–35] expressed in heterologous systems. GsMTx4 inhibits endogenous cationic MSCs and exogenously expressed Piezo 1 channels in outside-out patches with a K_D of ~0.5 μ M [15,32]. The D enantiomer shows strong inhibition of endogenous MSCs in muscle cell types such as chick cardiac cells [22] and murine skeletal muscle myotubes [10,14].

We have shown that GsMTx4 associates superficially with membranes under low tension, and penetrates deeper when membrane tension increases acting as an area/tension clamp for the outer monolayer [16,36]. Electron micrographic studies of the Piezo1 channels place the

mechanosensing domains for these channels on the outer monolayer [37], supporting GsMTx4 inhibition of these channels. However, relief of tension in the outer monolayer would transfer tension to the cytoplasmic monolayer. Interestingly, GsMTx4 appears to potentiate K^+ selective 2P domain channels [16] which are predicted to have gating elements sensitive to tension changes in the inner monolayer [38,39]. Potentiation of K^+ flux would tend to repolarize cells decreasing the activity of voltage gated cationic fluxes.

Functioning as a tension clamp, GsMTx4 may also affect enzymatic mechanosensitive signaling molecules like Nox2 (gp91^{phox}) and gp22^{phox} subunits, that are part of a transmembrane protein complex that generates reactive oxygen species (ROS) as signaling molecules [40]. The active complex responds to mechanical stress in both cardiac and skeletal muscle [41], and we have shown that Nox2 activity is sensitive to sarcolemmal mechanical integrity and microtubule assembly [13]. Indeed, we have shown that GsMTx4-L inhibits the Ca^{2+} influx pathways triggered by Nox2-ROS production [12]; however, it is unclear whether mechanically sensitive ROS production is inhibited directly by GsMTx4 inhibiting Nox2. Never-the-less, this work and others [12,13,42] support a model in which eccentric contraction-induced force loss is driven by an increase in sarcolemmal Ca^{2+} influx, a pathway inhibited by GsMTx4.

4.3. Toxicity

There is growing evidence that MSC's contribute to many physiological functions including arterial pressure regulation [43–46], the skeletal muscle pressor reflex [47], cardiac stretch induced slow force response [20], red blood cell volume regulation [48,49], neuronal touch and pain sensation [50–52], mechanotransduction of sound within the inner ear [53,54], and renal epithelial luminal pressure detection [55]. While we showed that the brain receives almost no GsMTx4-D, a finding consistent with previous studies [23], we showed elevated levels in the kidneys and liver compared to the intended muscular target tissues. Despite the pharmacokinetics analyses suggesting there would be a minimum of 20–60 μM in these organs throughout the six-week study, we observed no overt signs of toxicity. This lack of toxicity aligns with our recently proposed model for GsMTx4-D's action [16] in which ideal peptide concentrations act to suppress outer monolayer tension that is beyond the bounds of physiological stress, acting solely during conditions of pathological stress. It is important to note that the majority of cell and tissue-based experiments describing GsMTx4 effects on MSC function (i.e. single channel gating in pipette patches, cancer cell migration, atrial fibrillation, blood pressure regulation, peripheral pain sensation, etc.) were conducted under non-physiological stress conditions. While we have some idea of mechanical stress cells experience *in vitro*, our understanding of the actual stresses and thresholds for cellular mechanosensitivity *in situ* or *in vivo* are largely unknown. Nonetheless, it will be important to determine if the peptide elicits adverse effects over longer time frames, and particularly during growth and development, and in mechanically sensitive tissues like the kidney where high levels of GsMTx4-D accumulates.

4.4. GsMTx4-D proffers benefit in D2.mdx mice

Here we show that GsMTx4-D treatment blunts the decline in muscle mass and improves the functional capacity (i.e., the isometric torque) of D2.*mdx* skeletal muscle. As normalizing

isometric torque to the mass of the muscle accounted for the drop in the total torque output, we conclude that GsMTx4-D's effect was primarily on slowing the disease-dependent decline in muscle mass. Given that this decline is driven by an enhanced calcium-dependent proteolysis and necrosis in dystrophic muscle fibers [42], the GsMTx4-induced suppression of MSC-dependent cation/ Ca^{2+} influx is a parsimonious explanation for these results.

Reduction in contraction injury is a well-established preclinical endpoint in murine DMD [56]. GsMTx4-D also decreased susceptibility of skeletal muscle to EC injury by ~50%, a degree of protection similar to our previous findings with microtubule targeted therapeutics [12,13] or inhibition of Nox2 activity [12] and other signaling pathways linked to excessive MSC activity [57]. Furthermore, these findings are aligned with our recent study demonstrating GsMTx4-D's function as a cardio-protectant in ischemic reperfusion injury [23]. Given that DMD patients succumb to cardiovascular and pulmonary pathologies, we speculate that GsMTx4-D's benefit may extend from skeletal muscle to include heart.

Even though potent protection from force loss was observed in GsMTx4-treated D2.*mdx* mice, we found no difference in EBD incorporation, a classic measure of membrane fragility [58]. While the EBD result is in apparent contrast to the reduced contraction-induced dye accumulation in GsMTx4 treated *mdx* muscles *in vitro* [59], the concentration used *in vitro* (10 μM) was significantly above those reached with chronic *in vivo* dosing in this study (0.5–2 μM). Furthermore, given that GsMTx4-D had no impact on the level of interstitial fibrosis, and fibrosis changes the mechanical stress to muscle fibers during contraction, it is reasonable to hypothesize that altered mechanics may be a mechanism that drives the incorporation of EBD into the muscle. To this end, EBD has been shown to pass through connexin hemichannels [60]. Given reports of increased connexin permeability in dystrophic muscle [61], a disconnection between the protection of contraction injury and the permeability of EBD may lie within the dysregulated function of connexins.

4.5. GsMTx4-D therapeutic potential

GsMTx4-D is the first member in a class of compounds targeting MSCs. The therapeutic potential for GsMTx4-D is not as a cure, rather as a drug that can act by slowing disease progression. Given growing evidence that dysregulated Ca^{2+} signals from sarcolemmal MSC's driving is pathognomonic across several dystrophies, GsMTx4-D may prove effective in other forms of dystrophy as well.

Given the genetic basis of DMD, gene editing and exon skipping therapies that can restore dystrophin expression have the potential to cure the disease. While dystrophin expression to ~10% of normal levels is thought necessary to realize clinical benefits, these levels may be difficult to achieve based on current technology. In fact, phase III trials with eteplirsen, the FDA approved morpholino exon skipping drug, has shown ~1% of normal dystrophin levels over 180 weeks [62]. New evidence that normalizing dysregulated Ca^{2+} signaling improves exon skipping efficiency [63] raises the intriguing possibility that compounds like GsMTx4-D may be necessary adjuvants to realize effective genetic cures.

5. Conclusion

GsMTx4-D treatment partially protects D2.*mdx* mice from the disease associated loss of muscle mass and protects from contraction induced injury. These findings, in combination with a recent report of GsMTx4-D's cardioprotective effects, support further development of GsMTx4-D for treatment of muscular pathology.

Acknowledgments

This work was supported by funding from Tonus Therapeutics and Akashi Therapeutics (where GsMTx4 has been renamed AT-300). We would also like to thank Mr. Jeff Harvey (CSO of Tonus Therapeutics) for his generous financial support for these studies.

References

- [1]. Blake DJ, Weir A, Newey SE, Davies KE. Function and genetics of dystrophin and dystrophin-related proteins in muscle. *Physiol Rev.* 2002;82(2):291–329. [PubMed: 11917091]
- [2]. Mallouk N, Jacquemond V, Allard B. Elevated subsarcolemmal Ca²⁺ in mdx mouse skeletal muscle fibers detected with Ca²⁺-activated K⁺ channels. *Proc Natl Acad Sci USA* 2000;97(9):4950–5. [PubMed: 10781103]
- [3]. De Backer F, Vandebrouck C, Gailly P, Gillis JM. Long-term study of Ca(2+) homeostasis and of survival in collagenase-isolated muscle fibres from normal and mdx mice. *J Physiol.* 2002;542(Pt 3):855–65. [PubMed: 12154184]
- [4]. Phillips MF, Quinlivan R. Calcium antagonists for Duchenne muscular dystrophy. *The Cochrane Library*; 2008.
- [5]. Ruegg UT, Nicolas-Metral V, Challet C, Bernard-Helary K, Dorchies OM, Wagner S, et al. Pharmacological control of cellular calcium handling in dystrophic skeletal muscle. *Neuromuscul Disord.* 2002(Suppl 1):S155–61. [PubMed: 12206810]
- [6]. Jørgensen LH, Blain A, Grealley E, Laval SH, Blamire AM, Davison BJ, et al. Long-term blocking of calcium channels in mdx mice results in differential effects on heart and skeletal muscle. *Am J Pathol* 2011;178(1):273–83. [PubMed: 21224064]
- [7]. Haws CM, Winegar BD, Lansman JB. Block of single L-type Ca²⁺ channels in skeletal muscle fibers by aminoglycoside antibiotics. *J Gen Physiol* 1996;107:421–32. [PubMed: 8868052]
- [8]. Malik V, Rodino-Klapac LR, Viollet L, Mendell JR. Aminoglycoside-induced mutation suppression (stop codon readthrough) as a therapeutic strategy for Duchenne muscular dystrophy. *Therap Adv Neurol Disord* 2010;3(6):379–89. [PubMed: 21179598]
- [9]. Du X-J, Anderson KE, Jacobsen A, Woodcock EA, Dart AM. Suppression of ventricular arrhythmias during ischemia-reperfusion by agents inhibiting Ins (1, 4, 5) P 3 release. *Circulation* 1995;91(11):2712–16. [PubMed: 7758174]
- [10]. Suchyna TM, Sachs F. Mechanosensitive channel properties and membrane mechanics in mouse dystrophic myotubes. *J Physiol* 2007;581 (Pt 1):369–87. [PubMed: 17255168]
- [11]. Franco-Obregon A, Jr, Lansman JB. Mechanosensitive ion channels in skeletal muscle from normal and dystrophic mice. *J Physiol* 1994;481:299–309. [PubMed: 7537813]
- [12]. Khairallah RJ, Shi G, Sbrana F, Prosser BL, Borroto C, Mazaitis MJ, et al. Microtubules underlie dysfunction in duchenne muscular dystrophy. *Sci Signal* 2012;5(236):ra56. [PubMed: 22871609]
- [13]. Kerr JP, Robison P, Shi G, Bogush AI, Kempema AM, Hexum JK, et al. Detyrosinated microtubules modulate mechanotransduction in heart and skeletal muscle. *Nat Commun* 2015;6.
- [14]. Yeung EW, Whitehead NP, Suchyna TM, Gottlieb PA, Sachs F, Allen DG. Effects of stretch-activated channel blockers on [Ca²⁺]_i and muscle damage in the mdx mouse. *J Physiol.* 2005;562(Pt 2):367–80. [PubMed: 15528244]
- [15]. Suchyna TM, Johnson JH, Clemo HF, Huang ZH, Gage DA, Baumgarten CM, et al. Identification of a peptide toxin from *Grammostola spatulata* spider venom that blocks stretch activated channels. *J Gen Physiol* 2000;115:583–98. [PubMed: 10779316]

- [16]. Gnanasambandam R, Ghatak C, Yasmann A, Nishizawa K, Sachs F, Ladokhin AS, et al. GsMTx4: mechanism of inhibiting mechanosensitive ion channels. *Biophys J* 2017;112(1):31–45. [PubMed: 28076814]
- [17]. Ducret T, Vandebrout C, Cao ML, Lebacq J, Gailly P. Functional role of store-operated and stretch-activated channels in murine adult skeletal muscle fibres. *J Physiol* 2006;575(Pt 3):913–24. [PubMed: 16825296]
- [18]. Gailly P, De Backer F, Van Schoor M, Gillis JM. In situ measurements of calpain activity in isolated muscle fibres from normal and dystrophin-lacking mdx mice. *J Physiol* 2007;582(Pt 3):1261–75. [PubMed: 17510188]
- [19]. Stiber JA, Zhang ZS, Burch J, Eu JP, Zhang S, Truskey GA, et al. Mice lacking Homer 1 exhibit a skeletal myopathy characterized by abnormal transient receptor potential channel activity. *Mol Cell Biol* 2008;28(8):2637–47. [PubMed: 18268005]
- [20]. Ward ML, Williams IA, Chu Y, Cooper PJ, Ju YK, Allen DG. Stretch-activated channels in the heart: contributions to length-dependence and to cardiomyopathy. *Prog Biophys Mol Biol* 2008;97(2–3):232–49. [PubMed: 18367238]
- [21]. Williams IA, Allen DG. Intracellular calcium handling in ventricular myocytes from mdx mice. *Am J Physiol Heart Circ Physiol* 2007;292(2):H846–55. [PubMed: 17012353]
- [22]. Suchyna TM, Tape SE, Koeppe RE, Andersen OS, Sachs F, Gottlieb PA. Bilayer-dependent inhibition of mechanosensitive channels by neuroactive peptide enantiomers. *Nature* 2004;430(6996):235–40. [PubMed: 15241420]
- [23]. Wang J, Ma Y, Sachs F, Li J, Suchyna TM. GsMTx4-D is a cardioprotectant against myocardial infarction during ischemia and reperfusion. *J Mol Cell Cardiol* 2016;98:83–94. [PubMed: 27423272]
- [24]. Bode F, Sachs F, Franz MR. Tarantula peptide inhibits atrial fibrillation – a peptide from spider venom can prevent the heartbeat from losing its rhythm. *Nature* 2001;409(6816):35–6. [PubMed: 11343101]
- [25]. Coley WD, Bogdanik L, Vila MC, Yu Q, Van Der Meulen JH, Rayavarapu S, et al. Effect of genetic background on the dystrophic phenotype in mdx mice. *Hum Mol Gen* 2016;25(1):130–45. [PubMed: 26566673]
- [26]. Fukada S-i, Morikawa D, Yamamoto Y, Yoshida T, Sumie N, Yam-aguchi M, et al. Genetic background affects properties of satellite cells and mdx phenotypes. *Am J Pathol* 2010;176(5):2414–24. [PubMed: 20304955]
- [27]. Teichmann MD, Wegner FV, Fink RH, Chamberlain JS, Launikonis BS, Martinac B, et al. Inhibitory control over Ca(2+) sparks via mechanosensitive channels is disrupted in dystrophin deficient muscle but restored by mini-dystrophin expression. *PLoS One* 2008;3(11):e3644. [PubMed: 18982068]
- [28]. Wood LK, Kayupov E, Gumucio JP, Mendias CL, Claflin DR, Brooks SV. Intrinsic stiffness of extracellular matrix increases with age in skeletal muscles of mice. *J Appl Physiol* 2014;117(4):363–9. [PubMed: 24994884]
- [29]. Heydemann A, Ceco E, Lim JE, Hadhazy M, Ryder P, Moran JL, et al. Latent TGF- β -binding protein 4 modifies muscular dystrophy in mice. *J Clin Invest* 2010;120(2):645.
- [30]. Diao L, Meibohm B. Pharmacokinetics and pharmacokinetic–pharmacodynamic correlations of therapeutic peptides. *Clin Pharmacokinet* 2013;52(10):855–68. [PubMed: 23719681]
- [31]. Strohl WR. Fusion proteins for half-life extension of biologics as a strategy to make biobetters. *BioDrugs* 2015;29(4):215–39. [PubMed: 26177629]
- [32]. Bae C, Sachs F, Gottlieb PA. The mechanosensitive ion channel Piezo1 is inhibited by the peptide GsMTx4. *Biochemistry* 2011;50(29):6295–300. [PubMed: 21696149]
- [33]. Dyachenko V, Husse B, Rueckschloss U, Isenberg G. Mechanical deformation of ventricular myocytes modulates both TRPC6 and Kir2.3 channels. *Cell Calc* 2009;45(1):38–54.
- [34]. Patel A, Sharif-Naeini R, Folgering JR, Bichet D, Duprat F, Honoré E. Canonical TRP channels and mechanotransduction: from physiology to disease states. *Pflügers Arch Eur J Physiol* 2010;460(3):571–81. [PubMed: 20490539]

- [35]. Parpaite T, Cardouat G, Mauroux M, Gillibert-Duplantier J, Robillard P, Quignard J-F, et al. Effect of hypoxia on TRPV1 and TRPV4 channels in rat pulmonary arterial smooth muscle cells. *Pflügers Arch Eur J Physiol* 2016;468(1):111–30. [PubMed: 25799977]
- [36]. Nishizawa K, Nishizawa M, Gnanasambandam R, Sachs F, Sukharev SI, Suchyna TM. Effects of Lys to Glu mutations in GsMTx4 on membrane binding, peptide orientation, and self-association propensity, as analyzed by molecular dynamics simulations. *Biochim Biophys Acta (BBA) Biomembr* 2015;1848(11):2767–78.
- [37]. Zhao Q, Zhou H, Chi S, Wang Y, Wang J, Geng J, et al. Structure and mechanogating mechanism of the Piezo1 channel. *Nature* 2018;554(7693):487. [PubMed: 29469092]
- [38]. Patel AJ, Honore E, Maingret F, Lesage F, Fink M, Duprat F, et al. A mammalian two pore domain mechano-gated S-like K⁺ channel. *EMBO J* 1998;17(15):4283–90. [PubMed: 9687497]
- [39]. Brohawn SG, del Mármol J, MacKinnon R. Crystal structure of the human K2P TRAAK, a lipid- and mechano-sensitive K⁺ ion channel. *Science* 2012;335(6067):436–41. [PubMed: 22282805]
- [40]. Lambeth JD, Neish AS. Nox enzymes and new thinking on reactive oxygen: a double-edged sword revisited. *Ann Rev Pathol Mech Dis* 2014;9:119–45.
- [41]. Prosser BL, Ward CW, Lederer WJ. X-ROS signaling: rapid mechano-chemo transduction in heart. *Science* 2011;333(6048):1440–5. [PubMed: 21903813]
- [42]. Whitehead NP, Yeung EW, Allen DG. Muscle damage in mdx (dystrophic) mice: role of calcium and reactive oxygen species. *Clin Exp Pharmacol Physiol* 2006;33(7):657–62. [PubMed: 16789936]
- [43]. Spassova MA, Hewavitharana T, Xu W, Soboloff J, Gill DL. A common mechanism underlies stretch activation and receptor activation of TRPC6 channels. *Proc Natl Acad Sci USA* 2006;103(44):16586–91. [PubMed: 17056714]
- [44]. Fanchaouy M, Bychkov R, Meister JJ, Beny JL. Stretch-elicited calcium responses in the intact mouse thoracic aorta. *Cell Calc* 2007;41(1):41–50.
- [45]. Gilbert G, Ducret T, Marthan R, Savineau J-P, Quignard J-F. Stretch-induced Ca²⁺ signaling in vascular smooth muscle cells depend on Ca²⁺ store segregation. *Cardiovasc Res* 2014;cvu069.
- [46]. Gonzales AL, Yang Y, Sullivan MN, Sanders L, Dabertrand F, Hill-Eubanks DC, et al. A PLC (gamma) 1-dependent, force-sensitive signaling network in the myogenic constriction of cerebral arteries. *Sci Signal* 2014;7(327):ra49. [PubMed: 24866019]
- [47]. Copp SW, Kim JS, Ruiz-Velasco V, Kaufman MP. The mechano-gated channel inhibitor GsMTx4 reduces the exercise pressor reflex in decerebrate rats. *J Physiol* 2016;594(3):641–55. [PubMed: 26608396]
- [48]. Vadorpe DH, Xu C, Shmukler BE, Otterbein LE, Trudel M, Sachs F, et al. Hypoxia activates a Ca²⁺-permeable cation conductance sensitive to carbon monoxide and to GsMTx-4 in human and mouse sickle erythrocytes. *PLoS One* 2010;5(1):e8732. [PubMed: 20090940]
- [49]. Stewart AK, Vadorpe DH, Heneghan JF, Chebib F, Stolpe K, Akhavein A, et al. The GPA-dependent, spherostomatocytosis mutant AE1 E758K induces GPA-independent, endogenous cation transport in amphibian oocytes. *Am J Physiol Cell Physiol* 2010;298(2):C283–97. [PubMed: 19907019]
- [50]. Woo S-H, Ranade S, Weyer AD, Dubin AE, Baba Y, Qiu Z, et al. Piezo2 is required for Merkel-cell mechanotransduction. *Nature* 2014;509(7502):622–6. [PubMed: 24717433]
- [51]. Park SP, Kim BM, Koo JY, Cho H, Lee CH, Kim M, et al. A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain. *Pain* 2008;137(1):208–17. [PubMed: 18359568]
- [52]. Alessandri-Haber N, Dina OA, Chen X, Levine JD. TRPC1 and TRPC6 channels cooperate with TRPV4 to mediate mechanical hyperalgesia and nociceptor sensitization. *J Neurosci* 2009;29(19):6217–28. [PubMed: 19439599]
- [53]. Beurg M, Kim KX, Fettiplace R. Conductance and block of hair-cell mechanotransducer channels in transmembrane channel-like protein mutants. *J Gen Physiol* 2014;144(1):55–69. [PubMed: 24981230]
- [54]. Wu Z, Grillet N, Zhao B, Cunningham C, Harkins-Perry S, Coste B, et al. Mechanosensory hair cells express two molecularly distinct mechanotransduction channels. *Nat Neurosci* 2017;20(1):24–33. [PubMed: 27893727]

- [55]. Peyronnet R, Martins JR, Duprat F, Demolombe S, Arhatte M, Jodar M, et al. Piezo1-dependent stretch-activated channels are inhibited by Polycystin-2 in renal tubular epithelial cells. *EMBO Rep* 2013;14(12):1143–8. [PubMed: 24157948]
- [56]. Spurney CF, Gordish-Dressman H, Guerron AD, Sali A, Pandey GS, Rawat R, et al. Preclinical drug trials in the mdx mouse: assessment of reliable and sensitive outcome measures. *Muscle Nerve* 2009;39(5):591–602.
- [57]. Ismail HM, Scapozza L, Ruegg UT, Dorchies OM. Diapocynin, a dimer of the NADPH oxidase inhibitor apocynin, reduces ROS production and prevents force loss in eccentrically contracting dystrophic muscle. *PloS One* 2014;9(10):e110708. [PubMed: 25329652]
- [58]. Allen DG, Whitehead NP, Froehner SC. Absence of dystrophin disrupts skeletal muscle signaling: roles of Ca²⁺, reactive oxygen species, and nitric oxide in the development of muscular dystrophy. *Physiol Rev* 2016;96(1):253–305. [PubMed: 26676145]
- [59]. Whitehead NP, Streamer M, Lusambili LI, Sachs F, Allen DG. Streptomycin reduces stretch-induced membrane permeability in muscles from mdx mice. *Neuromusc Disord* 2006;16(12):845–54. [PubMed: 17005404]
- [60]. Cisterna BA, Vargas AA, Puebla C, Sáez JC. Connexin hemichannels explain the ionic imbalance and lead to atrophy in denervated skeletal muscles. *Biochim Biophys Acta (BBA) Mol Basis Dis* 2016;1862(11):2168–76.
- [61]. Cea LA, Puebla C, Cisterna BA, Escamilla R, Vargas AA, Frank M, et al. Fast skeletal myofibers of mdx mouse, model of Duchenne muscular dystrophy, express connexin hemichannels that lead to apoptosis. *Cell Mol Life Sci* 2016;73(13):2583–99. [PubMed: 26803842]
- [62]. Lim KRQ, Maruyama R, Yokota T. Eteplirsen in the treatment of Duchenne muscular dystrophy. *Drug Des Dev Therapy* 2017;11:533.
- [63]. Kendall GC, Mokhonova EI, Moran M, Sejbuk NE, Wang DW, Silva O, et al. Dantrolene enhances antisense-mediated exon skipping in human and mouse models of Duchenne muscular dystrophy. *Sci Trans Med* 2012;4(164):164ra160.

Fig. 1. Experimental setup showing the anesthetized mouse with gastrocnemius muscle exposed and knee and foot in restraints in place for force measurements. The stimulating electrodes are embedded in the muscle and foot restraint is attached to a force torque sensor.

Fig. 2.

Pharmacokinetic measurement of GsMTx4-D accumulation and clearance from DBA2-*mdx* mice. Mice were injected subcutaneously every other day for either 7 or 14 days with 10 mg/kg GsMTx4-D (top panel). Tissues were harvested and fluids were collected at 7, 14, 21 and 28 days. Tissue and body fluid concentrations of GsMTx4-D are shown at the different time points ($n = 5$ mice/time point). Urine samples were not available from all mice at each time point. Values are means $\pm$ SE.

Fig. 3.

Body weight (**A**) and muscle weight (**B**) measured at the conclusion of six weeks of high and low dose treatments. ($n = 10$ mice/cohort, with weights showing means $\pm$ SE). * $p < 0.05$ ANOVA.

Fig. 4.

(A) Isometric torque observed in DBA2 and D2.*mdx* mice. The decline in force generation observed in the D2-*mdx* was partially rescued by treatment with 1 or 10 mg/kg GsMTx4-D. (B) Normalizing the isometric torque to gastrocnemius weight ablated the disease and treatment effect consistent with the impact of muscle mass on muscle torque output. ($n = 10$ mice/cohort, with forces showing means $\pm$ SE). $* = p < 0.05$ ANOVA.

Fig. 5.

(A) Following 20 EC contractions D2-*mdx* (red) show a significant force decline when compared to DBA2 (black; 1st contraction in solid, 20th in dashed) Vertical bar = 0.1 N/cm, and Horizontal bar = 100 ms. (B) Treatment with either the target or low concentration of GsMTx4-D protected the D2-*mdx* against contraction induced decline in force. (C) The percentage of muscle fibers that accumulated EBD post EC contraction was not impacted by GsMTx4-D treatment. ($n = 10$ mice/cohort, with forces and dye incorporation showing means $\pm$ SE). $*$ = $p < 0.05$ ANOVA.

Fig. 6.

WGA staining denotes interstitial fibrotic accumulation and central nuclei. (A) Shows representative cryosections from WGA stained DBA2 and D2.*mdx* muscle after six weeks of disease progression. Expanded regions of interest (ROI) below the main images highlight the accelerated fibrosis and fiber irregularity in the DBA2-mdx muscle. Central nuclei are readily distinguished in these images and are indicated by arrows. (B) Quantitatively shows the significant increase in fibrosis in DBA2-mdx over DBA2 WT, and that GsMTx4-D treatment has no significant effect on the level of fibrosis. (C) The percent of myofibers with

central nucleated fibers's (CNF) revealed no difference following treatment. ($n = 10$ mice/cohort, with WGA staining and central nucleation values showing means $\pm$ SE).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 7.

(A) Population histogram of myofibers cross-sectional area (CSA) with mean and median values summarized in (B) Results visualized as plot of CSA cumulative percentage. Non-parametric K-sample analysis (w/ Kruskal Wallis ANOVA; $p < 0.05$) was used to determine the impact of disease and treatment on muscle fiber CSA. * $p < 0.05$ vs. DBA2, # $p < 0.05$ vs. vehicle, [^] $p < 0.05$ vs. 1 μ M (GsMTx4).