Published in final edited form as:

Eur J Med Chem. 2019 March 01; 165: 250–257. doi:10.1016/j.ejmech.2019.01.019.

Benzimidazolone-based selective σ_2 receptor ligands: synthesis and pharmacological evaluation

Sebastiano Intagliata^a, Walid F. Alsharif^b, Christophe Mesangeau^b, Nicola Fazio^b, Michael Seminerio^c, Yan-Tong Xu^c, Rae R. Matsumoto^{c,d}, and Christopher R. McCurdy^{a,b,*}

^aDepartment of Medicinal Chemistry, College of Pharmacy, University of Florida, Gainesville, Florida 32610, USA

^bDepartment of Medicinal Chemistry, School of Pharmacy, The University of Mississippi, University, Mississippi 38677, USA

^cDepartment of Basic Pharmaceutical Sciences, West Virginia University, Morgantown, West Virginia 26506, USA

^dCollege of Pharmacy, Touro University California, Vallejo, CA 94592, USA

Abstract

Sigma receptors (σ Rs) are considered to be a significant and valid target for developing new medications to address several diseases. Their potential involvement in numerous central nervous system disorders, neuropathic pain, addiction, and cancer has been extensively reported. In particular, the σ_2 R has been identified as potential target for the development of pharmaceutical agents intended to treat the negative effects associated with drugs of abuse. As a continuation of our previous efforts to develop new selective σ_2 R ligands, a series of benzimidazolone derivatives were designed, synthesized, and characterized. The newly synthesized ligands were evaluated through *in vitro* radioligand binding assays to determine their affinity and selectivity towards both σ_1 and σ_2 receptors. Several derivatives displayed high affinity for the σ_2 R (K_i = 0.66–68.5 nM) and varied from preferring to selective, compared to σ_1 R (σ_1/σ_2 = 5.8–1139). Among them, compound 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-propyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (**14**) displayed the ability to produce a dose-dependent reduction in the convulsive effects of cocaine in a rodent model after injecting 10 mg/kg (i.p.). These preliminary results support the use of selective σ_2 R ligands in the development of useful pharmacological tools or potential pharmacotherapies for cocaine toxicity.

Graphical Abstract

*Corresponding author. Tel.: +1 (352) 294-8691; fax: +1 (352) 392-9455. CMcCurdy@cop.ufl.edu (C.R. McCurdy).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Sigma receptors; Sigma-2 receptors; Sigma-2 ligands; Benzimidazolone; Structure-affinity relationships; Cocaine-induced convulsions; Cocaine's toxicity

1. Introduction

Sigma receptors (σ Rs) were originally classified as members of the opioid receptor family. This was based on the behavioral actions observed with the opiate benzomorphan, ($\pm$)-*N*-allylnormetazocine, and its analogs [1]. The effects of ($\pm$)-*N*-allylnormetazocine were reported to be partially antagonized by the opioid antagonist naloxone, thus resulting in σ Rs being classified as a new opioid receptor subtype. Through subsequent studies using purified enantiomers, it was shown that the “psychotomimetic” effect produced by (–)-SKF-10047 was instigated through kappa opioid receptors and the “psychotomimetic” effect of (+)-SKF-10047 was not blocked by naltrexone, a potent analog of naloxone, and must be mediated through non-opioid receptors. (+)-SKF-10047 was eventually shown to interact with at least two distinct binding sites: 1) the phencyclidine (PCP) site on N-methyl-D-aspartate (NMDA) receptors, and it was due to this interaction that the term sigma-PCP was prevalent in the early 1980s, and 2) other binding sites, distinct from opioid and NMDA receptors, which today retains the designation of σ Rs [2, 3].

Later, several studies proved that σ Rs are unique proteins highly preserved across species, cell types, and organelles [4–7]. Additional studies identified two subtypes of the receptor, σ_1 R and σ_2 R. Although historical sigma ligands interacted with both σ_1 and σ_2 R, the subtypes are distinguishable from each other by their amino acid sequences and pharmacological preferences [8–10]. The σ_1 R was first cloned in 1996 from guinea pig liver and later from other sources, representing different species and tissues such as human placental choriocarcinoma cells [4], human brain [5], rat brain [6], and mouse brain [8]. The σ_1 R is highly conserved across species with an amino acid sequence that is distinct from other known mammalian receptors [4]. The σ_2 R has been more difficult to understand and clone. In 2011, the σ_2 R was suggested to be part of the progesterone receptor membrane component 1 (PGRMC1) protein complex [11]. However, further investigations showed no correlation between PGRMC1 expression and σ_2 R binding levels in knockout or over-expression studies [12]. More recently, the σ_2 R was purified from calf liver tissue by affinity

chromatography and, subsequently, the gene that codes for the σ_2 R was identified as identical to Transmembrane Protein 97 (TMEM97), which is an endoplasmic reticulum-resident transmembrane protein that acts as a binding partner of the lysosomal cholesterol transporter NPC1. Indeed, this latest finding provided further evidence that the σ_2 R and PGRMC1 are two distinct proteins [13]. Consensus in the field now accepts the TMEM97 protein as the σ_2 R. Over-expression of σ_2 R in proliferative cancer cells has been reported, suggesting that selective radiolabeled σ_2 R ligands may be useful to pinpoint the malignant tissues in SPECT (single-photon emission computed tomography) and PET (positron emission tomography) imaging [14–16]. Moreover, literature-based reports have shown that σ_2 R agonists are able to induce apoptosis and reduce cancer cell proliferation, thus, there is a rising interest in the development of σ_2 R agents as therapeutics or adjuvants to treat cancer [17–19].

The involvement of σ Rs in several CNS disorders such as depression, schizophrenia, and dementia, as well as their potential role in other neurological and psychological disorders, drove much of the early interest in these receptors [20–24]. This interest was due to the majority of typical antipsychotics, such as haloperidol, having affinity for σ Rs. Also, agents that instigate psychotomimetic activity, such as ($\pm$)-*N*-allylnormetazocine and its analogs, shared affinity for σ Rs [25]. Moreover, evidence-based literature demonstrated that some σ R antagonists were able to attenuate the effects of psychostimulant drugs such as cocaine [26–28]. Conversely, σ R agonists aggravated some of these effects, including the convulsive effects of cocaine, resulting in a leftward shift in the cocaine convulsion dose–response curve [28, 29]. There still remains a need to develop new medications to attenuate the convulsions associated with an overdose of cocaine, which cannot be controlled with existing antiepileptic drugs. However, unlike the σ_1 R the involvement of the σ_2 R in the toxicity of cocaine is still unclear, and improved, selective σ_2 R ligands are needed [30].

Previously, we reported a series of 1,3-benzothiazole-2(3*H*)-thione with high mixed affinity for σ_1 Rs and σ_2 Rs [31]. Among them, 3-[4-(4-cyclohexylpiperazin-1-yl)butyl]-2(3*H*)-benzothiazolethione compound **1** (CM156, Fig. 1) produced both acute and subchronic effects against cocaine in behavioral models in rodents [32]. Further studies showed the ability of **1** to mitigate the effects of methamphetamine in mice [33]. Furthermore, we developed 6-acetyl-3-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-2(3*H*)-benzoxazolone compound **2** (SN79, Fig. 1) which reduced the toxic and stimulant effects of cocaine, and also caused sedation and incoordination with acute treatment [34]. More studies have suggested that pretreatment with compound **2** afforded protection against methamphetamine-induced hyperthermia, and striatal dopaminergic and serotonergic neurotoxicity in mice [35].

Within the frame of our efforts to develop novel selective σ_2 R ligands [36], we have developed several σ ligands with high affinity for σ_2 Rs and preference over σ_1 Rs. Among them, the most selective σ_2 R ligand was selected in order to study its effect against cocaine-induced convulsions in a preclinical rodent model, and to assess the selectivity requirement of both receptor subtypes. As a strategy in the current study, we replaced the 1,3-benzothiazole-2(3*H*)-thione, and the 1,3-benzoxazol-2(3*H*)-one of **1** and **2** with the bioisostere 1,3-dihydrobenzimidazol-2-one. In particular, the new versatile core allowed us

to perform 1,3-bis-functionalization along with the introduction of an acetyl group in the 5-position of the heterocyclic ring. Similarly to **1** and **2**, the cyclic amine fragment of the new ligands were both 1-cyclohexylpiperazine and 1-(4-fluorophenyl)piperazine. Finally, based on our previous data, a four-carbon chain was selected as a linker as it results in greater σ_2R preference [31].

2. Results and discussion

2.1. Chemistry

The synthesis of final compounds **5** and **6**, was very straightforward as outlined in Scheme 1. Treatment of commercially available compound **3** with 1,4-dibromobutane in the presence of potassium carbonate in DMF, gave the 4-bromobutyl derivative (**4**) which was then coupled with 1-cyclohexylpiperazine, or 1-(4-fluorophenyl)piperazine in the presence of potassium carbonate in DMF to afford target compounds **5** and **6**, respectively (Scheme 1).

The synthesis of final compounds **12-20** is depicted in Scheme 2. *N*-alkylation of commercially available 2-(4-bromobutyl)isoindole-1,3-dione (**7**) with 1-(4-fluorophenyl)piperazine provided compound **8**. The desired primary amine was then liberated from the phthalimide by hydrazinolysis to afford compound **9** which was subsequently reacted, through a nucleophilic aromatic substitution, with 1-fluoro-2-nitrobenzene to give **10** or, with 1-(4-chloro-3-nitrophenyl)ethanone to give **11**. The nitro groups of **10** and **11** were reduced in the presence of Pd/C and the corresponding 1,2-diamine derivatives which were reacted with 1,1'-carbonyldiimidazole to give benzimidazolones **12** and **13**. The final compounds **14-20**, were obtained by *N*-alkylation with the appropriate alkyl halides.

2.2. Binding studies and extended non- σ binding profile

All final compounds were tested in radioligand binding studies at the σ_1R and σ_2R in rat brain membranes using well-established and previously described assay conditions [31]. The σ_1Rs were labeled with [3H](+)-pentazocine. The σ_2Rs were labeled with [3H]di-o-tolylguanidine (DTG) in the presence of (+)-pentazocine to block σ_1Rs . Nonspecific binding was determined in the presence of haloperidol. The binding affinities (K_i in nM), and selectivity ratios ($K_i \sigma_1 / K_i \sigma_2$) of the new compounds at σRs are summarized in Table 1.

Initially, we synthesized compounds **5** and **6** to investigate the effect of replacement of the heterocyclic scaffolds in **1** and **2** with a benzimidazolone moiety. This structural modification led to an increase of the σ_1/σ_2 selectivity ratio for both analogs (**5** vs **1**, **6** vs **2**). In particular, compound **6** resulted in 46-fold more selective preference than **2** for the σ_2R over the σ_1R . This result prompted us to maintain the 1-(4-fluorophenyl)piperazine ring as a fixed portion of the new series. Subsequently, the introduction of different alkyl substituents on the *N*-3 (such as nC_3H_7 , nC_5H_{11} , and $nC_{10}H_{21}$), as well as an acetyl group at the 5-position of the benzimidazolone scaffold, gave compounds **12-20** which resulted in affinity values in the low or subnanomolar range for the σ_2R ($K_i = 0.66-32.5$ nM).

Based on the binding data, reported in Table 1, the main structure-affinity relationships (SARs) were as follows. In general, the new benzimidazolone derivatives retained high σ_2R

affinity and showed increased selectivity over the σ_1 R (selectivity 5.8-1139 fold). Replacement of the 1-cyclohexylpiperazine with 1-(4-fluorophenyl)piperazine, greatly improved the selectivity for the σ_2 R subtype (**5** vs **6**). Functionalization at the *N*-3 of the benzimidazolone ring generally increased affinity for the σ_2 R. Introduction of an acetyl group at the 5-position of the heterocyclic ring led to a loss of the σ_1/σ_2 selectivity. Thus, the presence of both a bulky substituent as R^2 and an acetyl group as R^1 was detrimental for σ_2 R affinity. Regarding the set of compounds with no substituent on the benzo-fused ring (**6**, **12**, and **14-16**), the introduction of R^2 groups were ranked based on their affinity (K_i values) at σ_2 R as follows: $nC_3H_7 > CH_3 > nC_5H_{11} > nC_{10}H_{21} > H$. A different ranking was observed for the set of 5-acetyl-benzimidazolone analogs **13**, and **17-20**: $nC_3H_7 > nC_5H_{11} > CH_3 > H > nC_{10}H_{21}$. Among both sets, *n*-propyl and *n*-pentyl analogs (**14**, **18**, and **15**, **19**) displayed better results in terms of affinity and selectivity for the σ_2 R than other alkyl-substituted derivatives.

Based on these results, we selected compounds **14** and **15** which had the highest ratios of σ_1/σ_2 (>1000 fold and >500 fold, respectively) for further evaluation in extended non- σ Rs binding experiments (Table 2), using previously established and published methods [28]. Substantially, compound **14** did not exhibit significant affinities for the non- σ proteins tested, except for serotonin (5-HT₂) receptors ($K_i = 54.70$ nM, 5-HT₂/ σ_2 R = 83 fold preference for σ_2 Rs). Compound **15** showed similar results at serotonin (5-HT₂) receptors ($K_i = 249.6$ nM, 5-HT₂/ σ_2 R = 58 fold selectivity for σ_2 Rs), although possessed very low affinity for serotonin transporters ($K_i = 445.9$ nM, 5-HTT/ σ_2 R = 104 fold selectivity for σ_2 Rs).

2.3. In vivo convulsion studies

Seizure is a common result of cocaine overdose and earlier σ R antagonists attenuated the convulsive effects of cocaine in mice [27, 28]. Therefore, we tested whether pretreatment of mice with 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-propyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (**14**), similarly attenuated cocaine-induced convulsions. Compound **14** was selected because it possessed a very good extended non- σ binding profile, thus acting as a relatively selective σ_2 R ligand. As can be seen from Fig. 2, compound **14** produced a dose-dependent reduction in the convulsive effects of cocaine, with the protective effects being statistically significant at the 10 mg/kg pretreatment dose ($P < 0.005$). These results suggest that σ_2 R-selective ligands can mitigate cocaine-induced convulsions, and support the hypothesis of an active role for this target in cocaine's toxicity.

3. Conclusion

We successfully synthesized a series of benzimidazolone derivatives, and evaluated their binding affinity for σ Rs. In this work, a versatile 1,3-dihydrobenzimidazol-2-one scaffold was used, along with a 1-(4-fluorophenyl)piperazine as a cyclic amine moiety. Based on the SARs, both structural elements were identified as preferential for optimizing affinity and selectivity for the σ_2 R. Derivatives **14**, and **15**, which gave the best results in terms of affinity for the σ_2 R ($K_i = 0.66$ and 4.27 nM) and selectivity over the σ_1 R ($\sigma_1/\sigma_2 = 1139$ and 532), were selected for additional binding evaluation *versus* different non- σ sites. In

particular, only moderate affinity for serotonin (5-HT₂) receptors (**14** and **15**, $K_i = 54.70$ and 249.60 nM respectively) and very low affinity for serotonin transporters (**15**, $K_i = 445.9$ nM) were observed. Noteworthy, *in vivo* convulsion studies in mice showed that compound **14** was able to dose-dependently attenuate cocaine-induced convulsions. Thus, the data reported in this paper are indicative of the potential utility of targeting σ_2 Rs to mitigate the convulsive effects of cocaine. The synthesis of new derivatives and further *in vivo* studies are currently under development and will be reported elsewhere.

4. Experimental protocols

4.1. Chemistry

Reagents and starting materials were obtained from commercial suppliers and were used without purification. Precoated silica gel GF Uniplates from Analtech were used for thin-layer chromatography (TLC). Column chromatography was performed on silica gel 60 (Sorbent Technologies). ¹H and ¹³C NMR spectra were recorded with a Bruker APX400 in CDCl₃ and DMSO-*d*₆ solution. Chemical shift (δ) values are given in parts per million (ppm) downfield to tetramethylsilane (TMS, $\delta = 0.00$ ppm); coupling constants (J values) are given in hertz (Hz). For signal multiplicities, the following abbreviations are used as follows: s (singlet), d (doublet), dd (doublets of doublet), t (triplet), q (quartet), quint (quintet), br s (broad singlet) and m (multiplet). Except where otherwise noted, ¹H and ¹³C NMR data for final compounds are given for materials in their salt form. The high resolution mass spectra (HRMS) were recorded on a Waters Micromass Q-ToF Micro mass spectrometer with a lock spray source. The mass spectra (MS) were recorded on a WATERS ACQUITY Ultra Performance LC with ZQ detector in ESI mode. Elemental analysis (C, H, N) was recorded on an elemental analyzer, Perkin-Elmer CHN/SO Series II Analyzer.

4.1.1. 1-(4-bromobutyl)-3-methyl-1,3-dihydrobenzimidazol-2-one (4).—K₂CO₃ (0.56 g, 4.05 mmol) and 1,4-dibromobutane (1.12 mL, 9.45 mmol) were added, under mechanical stirring, to a solution of 3-methyl-1*H*-benzimidazol-2-one (**3**) (0.20 g, 1.35 mmol) in anhydrous DMF (8 mL). The reaction mixture was heated at 60 °C for 3 h. After cooling, the mixture was poured into 100 mL of water, extracted with ethyl acetate (3 × 40 mL), washed with saturated aqueous NaCl and dried. The solvent was removed *in vacuo*, and the residue was purified by chromatography on a silica gel column using petroleum ether/ethyl acetate (7:3) as the eluent to give **4** (0.27 g, 70%), as a colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 7.06-7.03 (m, 2H), 6.96-6.91 (m, 2H), 3.88-3.86 (m, 2H), 3.41-3.38 (m, 2H), 3.63 (s, 3H), 1.89-1.85 (m, 4H). ¹³C NMR (100 MHz, CDCl₃): δ 154.41, 130.10, 129.17, 121.26, 121.23, 107.53, 107.50, 40.08, 33.13, 29.66, 27.15, 26.98. MS (ESI) m/z 305 [M + Na]⁺ for ⁷⁹Br, m/z 307 [M + Na]⁺ for ⁸¹Br.

4.1.2. 1-[4-(4-cyclohexylpiperazin-1-yl)butyl]-3-methyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (5).—K₂CO₃ (0.06 g, 0.44 mmol) and 1-cyclohexylpiperazine (0.04 g, 0.21 mmol) were added, under mechanical stirring, to a solution of **4** (0.05 g, 0.18 mmol) in anhydrous DMF (3 mL). The reaction mixture was heated at 60 °C for 2 h. After cooling, the mixture was poured into 20 mL of water, extracted with ethyl acetate (3 × 30 mL), and the combined organic layers were washed with saturated

aqueous NaCl and dried. The solvent was removed *in vacuo*, and the residue was chromatographed on a silica gel column using methylene chloride/methanol (95:5) as the eluent. The free base was converted into the hydrochloride salt by addition of HCl/dioxane to give **5** (0.05 g, 62%), as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.01-11.80 (m, 2H), 7.21-7.19 (m, 1H), 7.15-7.12 (m, 1H), 7.08-7.06 (m, 2H), 3.84-3.02 (m, 17H), 1.80-1.77 (m, 2H), 1.66-1.05 (m, 11H). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 154.01, 129.80, 128.89, 121.33, 121.30, 108.09, 64.32, 55.37, 48.60, 45.39, 39.96, 27.13, 26.37, 25.28, 24.75, 24.55, 21.14. HRMS (ESI) calcd for C₂₂H₃₅N₄O [M + H]⁺ 371.2811, found 371.2828. Anal. calcd for C₂₂H₃₆Cl₂N₄O. H₂O: C, 57.26; H, 8.30; N, 12.14. Found: C, 57.46; H, 7.92; N, 12.11.

4.1.3. 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-methyl-1,3-

dihydrobenzimidazol-2-one dihydrochloride (6).—This compound was prepared from **4** and 1-(4-fluorophenyl)piperazine as described for **5**. Yield 56%, as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 11.40 (br s, 1H), 7.30-7.00 (m, 9H), 3.88 (t, *J* = 6.8 Hz, 2H), 3.70-3.67 (m, 2H), 3.50-3.47 (m, 2H), 3.32 (s, 3H), 3.23-3.08 (m, 6H), 1.81-1.68 (m, 4H). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 154.81 (d, *J* = 258.0 Hz), 149.09, 129.59, 128.69, 120.78, 120.75, 117.84 (d, *J* = 7.0 Hz), 115.25 (d, *J* = 22.0 Hz), 107.65, 107.63, 54.68, 50.33, 45.95, 39.64, 26.80, 25.16, 20.18. HRMS (ESI) calcd for C₂₂H₂₈N₄OF [M + H]⁺ 383.2247, found 383.2237. Anal. calcd for C₂₂H₂₉Cl₂FN₄O: C, 58.02; H, 6.42; N, 12.30. Found: C, 58.31; H, 6.14; N, 12.23.

4.1.4. 2-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-1H-isoindole-1,3(2H)-dione (8).

—K₂CO₃ (8.45 g, 61.2 mmol) and 1-(4-fluorophenyl)piperazine (4.77 g, 26.50 mmol) were added, under mechanical stirring, to a solution of 2-(4-bromobutyl)isoindole-1,3-dione (**7**) (3.75 g, 20.00 mmol) in anhydrous DMF (30 mL). The reaction mixture was stirred at 60 °C for 1 h and at room temperature for 15 h. The reaction mixture was then poured into 40 mL of water, extracted with ethyl acetate (3 × 40 mL), and the combined organic layers were washed with saturated aqueous NaCl and dried. The solvent was removed *in vacuo*, and the residue was purified by chromatography on a silica gel column using a gradient of methylene chloride and methanol (10:0 to 9.5:0.5) as the eluent to give **8** (4.70 g, 60%), as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.86-7.80 (m, 4H), 7.00 (t, *J* = 9.0 Hz, 2H), 6.90-6.87 (m, 2H), 3.58 (t, *J* = 7.0 Hz, 2H), 3.01-2.99 (m, 4H), 2.43-2.41 (m, 4H), 2.30 (t, *J* = 7.0 Hz, 2H), 1.63-1.57 (m, 2H), 1.48-1.42 (m, 2H). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 167.88, 155.87 (d, *J* = 234.0 Hz), 147.89 (d, *J* = 1.8 Hz), 134.27, 131.56, 122.91, 116.91 (d, *J* = 7.5 Hz), 115.13 (d, *J* = 21.6 Hz), 57.14, 52.65, 48.92, 37.31, 25.90, 23.58.

4.1.5. 4-[4-(4-fluorophenyl)piperazin-1-yl]butan-1-amine (9).

—A solution of **8** (4.25 g, 11.10 mmol), and hydrazine monohydrate (2.7 mL, 55.70 mmol) in ethanol (220 mL) was heated at reflux for 16 h. The reaction mixture was then cooled with an ice bath. The precipitate was eliminated by filtration, and the filtrate was evaporated. The residue was purified by chromatography on a silica gel column using a gradient of methylene chloride and methanol saturated with ammonia (10:0 to 8:2) as the eluent to give **9** (2.33 g, 83%), as a low melting point pale yellow solid. ¹H NMR (400 MHz, CDCl₃): δ 6.91-6.89 (m, 2H), 6.83-6.79 (m, 2H), 3.06 (t, *J* = 4.8 Hz, 4H), 2.67 (t, *J* = 6.8 Hz, 2H), 2.54 (t, *J* = 5.0 Hz, 4H),

2.34 (t, $J = 7.0$ Hz, 2H), 2.00 (br s, 2H), 1.55-1.40 (m, 4H). ^{13}C NMR (100 MHz, CDCl_3): δ 157.24 (d, $J = 237.1$ Hz), 148.11 (d, $J = 2.1$ Hz), 117.88 (d, $J = 7.5$ Hz), 115.58 (d, $J = 21.9$ Hz), 58.52, 53.35, 50.23, 42.07, 31.57, 24.42.

4.1.6. *N*-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-2-nitroaniline (10).—A mixture of 1-fluoro-2-nitrobenzene (2.51 g, 17.80 mmol), compound **9** (2.24 g, 8.91 mmol), *N,N*-diisopropylethylamine (7.8 mL, 44.56 mmol) in DMF (30 mL) was heated at 120 °C for 2 h. After cooling, the mixture was poured into 50 mL of water, extracted with ethyl acetate (3 × 50 mL), and the combined organic layers were washed with brine and dried. The solvent was removed *in vacuo*, and the residue was purified by chromatography on a silica gel column using ethyl acetate/hexanes (6:4) as the eluent to give **10** (3.09 g, 93%), as an orange solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.14-8.11 (m, 1H), 8.04 (d, $J = 8.6$ Hz, 1H), 7.50 (t, $J = 7.6$ Hz, 1H), 7.06 (d, $J = 8.8$ Hz, 1H), 7.01 (t, $J = 8.8$ Hz, 2H), 6.92-6.88 (m, 2H), 6.65 (t, $J = 7.6$ Hz, 1H), 3.36 (q, $J = 6.3$ Hz, 2H), 3.04-3.02 (m, 4H), 2.47-2.45 (m, 4H), 2.34 (t, $J = 6.9$ Hz, 2H), 1.68-1.61 (m, 2H), 1.57-1.50 (m, 2H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 155.90 (d, $J = 234.0$ Hz), 147.91 (d, $J = 1.8$ Hz), 145.21, 136.52, 130.79, 126.19, 116.93 (d, $J = 7.5$ Hz), 115.16 (d, $J = 21.6$ Hz), 115.00, 114.51, 57.09, 52.65, 48.96, 42.00, 26.07, 23.50.

4.1.7. 1-[4-({4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}amino)-3-nitrophenyl]ethan-1-one (11).—A mixture of 1-(4-chloro-3-nitrophenyl)ethanone (3.89 g, 19.50 mmol), compound **9** (2.45 g, 9.75 mmol), *N,N*-diisopropylethylamine (6.32 g, 48.74 mmol) in DMF (50 mL) was heated at 100 °C for 6 h. After cooling, the mixture was poured into 80 mL of water, extracted with ethyl acetate (3 × 80 mL), and the combined organic layers were washed with brine and dried. The solvent was removed *in vacuo*, and the residue was purified by chromatography on a silica gel column using a gradient of ethyl acetate and hexanes (6:4 to 10:0) as the eluent to give **11** (3.35 g, 83%), as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.61-8.60 (m, 2H), 7.99 (d, $J = 5.2$ Hz, 1H), 7.16 (d, $J = 9.1$ Hz, 1H), 7.02 (t, $J = 9.0$ Hz, 2H), 6.93-6.90 (m, 2H), 3.45 (q, $J = 2.5$ Hz, 2H), 3.37 (s, 3H), 3.05 (t, $J = 4.0$ Hz, 4H), 2.48 (s, 4H), 2.37 (t, $J = 6.9$ Hz, 2H), 1.66 (quint, $J = 7.0$ Hz, 2H), 1.54 (quint, $J = 6.9$ Hz, 2H).

4.1.8. 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-1,3-dihydrobenzimidazol-2-one (12).—A solution of **10** (3.00 g, 8.06 mmol) in methanol (220 mL) was hydrogenated in the presence of 10% Pd/C (0.30 g) under pressure (40 psi) for 1 h. The catalyst was removed by filtration and the solvent was evaporated. The residue was dissolved in THF (80 mL), and 1,1'-carbonyldiimidazole (2.61 g, 16.10 mmol) was added. The reaction mixture was stirred under argon at room temperature. After 2 h, brine was added and the mixture was extracted with ethyl acetate (3 × 40 mL). The combined organic layers were washed with brine, dried and evaporated. The residue was purified by chromatography on a silica gel column using a gradient of methylene chloride and methanol (10:0 to 9.5:0.5) as the eluent to give **12** (2.03 g, 68%), as a white solid. A small portion was converted to the corresponding dihydrochloride salt for the biological testing. ^1H NMR (free amine, 400 MHz, $\text{DMSO}-d_6$): δ 10.81 (s, 1H), 7.12-7.10 (m, 1H), 7.03-6.97 (m, 5H), 6.90-6.87 (m, 2H), 3.79 (t, $J = 6.9$ Hz, 2H), 3.01-2.99 (m, 4H), 2.44-2.42 (m, 4H), 2.30 (t, $J = 7.0$ Hz, 2H),

1.68-1.62 (m, 2H), 1.48-1.41 (m, 2H). ^{13}C NMR (free amine, 100 MHz, $\text{DMSO}-d_6$): δ 155.90 (d, $J = 234.0$ Hz), 154.73, 147.91 (d, $J = 1.8$ Hz), 130.19, 128.22, 120.62, 120.39, 116.93 (d, $J = 7.5$ Hz), 115.15 (d, $J = 21.7$ Hz), 108.64, 107.67, 57.04, 52.60, 48.93, 39.59, 25.61, 23.33. Anal. calcd for $\text{C}_{21}\text{H}_{27}\text{Cl}_2\text{FN}_4\text{O} \cdot 1/2 \text{H}_2\text{O}$: C, 56.00; H, 6.27; N, 12.44. Found: C, 56.44; H, 6.02; N, 12.44.

4.1.9. 5-acetyl-1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-1,3-dihydrobenzimidazol-2-one (13).—This compound was prepared from **11** as described for **12**. Yield 51%, as a white solid. A small portion was converted to the corresponding dihydrochloride salt for the biological testing. ^1H NMR (free amine, 400 MHz, $\text{DMSO}-d_6$): δ 11.16 (br s, 1H), 7.71 (d, $J = 6.5$ Hz, 1H), 7.51 (s, 1H), 7.24 (d, $J = 6.5$ Hz, 1H), 7.00 (t, $J = 7.0$ Hz, 2H), 6.89-6.87 (m, 2H), 3.84 (t, $J = 5.4$ Hz, 2H), 2.99 (s, 4H), 2.53 (s, 3H), 2.41 (s, 4H), 2.30 (t, $J = 5.6$ Hz, 2H), 1.66 (quint, $J = 5.3$ Hz, 2H), 1.44 (quint, $J = 5.4$ Hz, 2H). ^{13}C NMR (free amine, 100 MHz, $\text{DMSO}-d_6$): δ 196.59, 155.93 (d, $J = 237.3$ Hz), 154.54, 147.93, 134.34, 130.18, 128.26, 122.59, 116.95 (d, $J = 5.9$ Hz), 115.19 (d, $J = 17.3$ Hz), 108.03, 107.22, 57.05, 52.64, 48.96, 40.01, 26.51, 25.66, 23.32. Anal. calcd for $\text{C}_{23}\text{H}_{29}\text{Cl}_2\text{FN}_4\text{O}_2 \cdot \text{H}_2\text{O}$: C, 55.09; H, 6.23; N, 11.17. Found: C, 55.25; H, 5.81; N, 10.67.

4.1.10. 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-propyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (14).—Sodium hydride (60% in mineral oil, 0.02 g, 0.38 mmol) and tetrabutylammonium iodide (0.01 g, 0.03 mmol) were added to a solution of **12** (0.10 g, 0.27 mmol) in anhydrous DMF (4 mL) at room temperature under argon. After stirring for 15 min, iodopropane (0.03 mL, 0.32 mmol) in DMF (2 mL) was added. The reaction mixture was stirred for 4 h at room temperature, poured into brine (20 mL) and extracted with ethyl acetate (3×20 mL). The organic layers were combined, washed with water, dried and evaporated. The residue was purified by chromatography on a silica gel column using ethyl acetate as the eluent. The free base was converted into the hydrochloride salt by the addition of HCl/dioxane to give **14** (0.10 g, 80%), as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 11.41 (s, 1H), 7.23-7.22 (m, 1H), 7.19-7.18 (m, 1H), 7.11-7.01 (m, 6H), 6.39 (br s, 1H), 3.86 (t, $J = 6.6$ Hz, 2H), 3.79 (t, $J = 6.9$ Hz, 2H), 3.69 (d, $J = 12.3$ Hz, 2H), 3.48 (d, $J = 11.3$ Hz, 2H), 3.23-3.08 (m, 6H), 1.78-1.62 (m, 6H), 0.85 (t, $J = 7.0$ Hz, 3H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 156.75 (d, $J = 235.8$ Hz), 153.50, 146.20, 129.09, 128.78, 120.89, 120.80, 118.03 (d, $J = 7.7$ Hz), 115.56 (d, $J = 21.9$ Hz), 107.96, 107.91, 54.76, 50.42, 46.10, 41.90, 39.69, 25.26, 21.22, 20.30, 11.08. Anal. calcd for $\text{C}_{24}\text{H}_{33}\text{Cl}_2\text{FN}_4\text{O}$: C, 59.63; H, 6.88; N, 11.59. Found: C, 59.38; H, 6.66; N, 11.10.

4.1.11. 1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-pentyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (15).—This compound was prepared from **12** as described for **14**. Yield 68%, as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 11.42 (s, 1H), 7.24-7.21 (m, 1H), 7.18-7.16 (m, 1H), 7.11-7.01 (m, 6H), 3.86 (t, $J = 6.8$ Hz, 2H), 3.81 (t, $J = 7.0$ Hz, 2H), 3.68 (d, $J = 12.7$ Hz, 2H), 3.49 (d, $J = 11.2$ Hz, 2H), 3.23-3.08 (m, 6H), 1.78-1.62 (m, 6H), 1.31-1.21 (m, 4H), 0.82 (t, $J = 6.6$ Hz, 3H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 156.93 (d, $J = 236.0$ Hz), 153.61, 146.23, 129.11, 128.88, 121.11, 121.00, 118.23 (d, $J = 7.6$ Hz), 115.70 (d, $J = 21.9$ Hz), 108.09, 108.08, 54.92, 50.60, 46.32,

40.44, 39.81, 28.42, 27.63, 25.31, 21.86, 20.47, 13.97. Anal. calcd for $C_{26}H_{37}Cl_2FN_4O$: C, 61.05; H, 7.29; N, 10.95. Found: C, 60.74; H, 7.25; N, 10.65.

4.1.12. 1-decyl-3-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-1,3-

dihydrobenzimidazol-2-one dihydrochloride (16).—This compound was prepared from **12** as described for **14**. Yield 33%, as a white solid. 1H NMR (400 MHz, DMSO- d_6): δ 11.15 (s, 1H), 7.24-7.22 (m, 1H), 7.19-7.16 (m, 1H), 7.09-7.01 (m, 6H), 5.85 (br s, 1H), 3.86 (t, J = 6.7 Hz, 2H), 3.81 (t, J = 7.0 Hz, 2H), 3.68 (d, J = 11.9 Hz, 2H), 3.48 (d, J = 11.2 Hz, 2H), 3.17-3.06 (m, 6H), 1.76-1.62 (m, 6H), 1.15-1.20 (m, 14H), 0.83 (t, J = 6.2 Hz, 3H). ^{13}C NMR (125 MHz, DMSO- d_6): δ 156.61 (d, J = 235.7 Hz), 153.41, 146.37, 129.00, 128.77, 120.86, 120.76, 117.87 (d, J = 7.7 Hz), 115.48 (d, J = 21.9 Hz), 107.88, 107.87, 54.77, 50.49, 46.02, 40.30, 39.75, 31.23, 28.88, 28.87, 28.61, 28.59, 27.76, 26.09, 25.20, 22.03, 20.29, 13.91. Anal. calcd for $C_{31}H_{47}Cl_2FN_4O$: C, 64.01; H, 8.14; N, 9.63. Found: C, 63.94; H, 8.04; N, 9.59.

4.1.13. 5-acetyl-1-(4-(4-(4-fluorophenyl)piperazin-1-yl)butyl)-3-methyl-1H-benzo[d]imidazol-2(3H)-one dihydrochloride (17)—A solution of **13** (0.11 g, 0.27

mmol), potassium carbonate (0.11 g, 0.80 mmol), tetrabutylammonium iodide (0.30 g, 0.08 mmol) in anhydrous DMF (10 mL) was heated at 60 °C under argon. After 30 min, a solution of iodomethane (16.7 μ L, 0.27 mmol) in anhydrous DMF (8 mL) was added dropwise over 1 h. After 3 h, a solution of iodomethane (5 μ L, 0.08 mmol) in anhydrous DMF (3 mL) was added dropwise and the reaction was stirred for another 3 h. The mixture was then poured into 80 mL of water, extracted with ethyl acetate (3 $\times$ 50 mL) and the solvent was evaporated. The residue was purified by chromatography on a silica gel column using a gradient of ethyl acetate and methanol saturated with ammonia (10:0 to 9:1) as the eluent. The free base was converted into the hydrochloride salt by addition of HCl/dioxane to give **17** (0.11 g, 89%), as a white solid. 1H NMR (400 MHz, DMSO- d_6): δ 11.17 (br s, 1H), 7.78 (d, J = 8.2 Hz, 1H), 7.73 (s, 1H), 7.36 (d, J = 8.2 Hz, 1H), 7.11-6.99 (m, 4H), 5.20 (br s, 1H), 3.92 (t, J = 6.7 Hz, 2H), 3.69 (d, J = 11.7 Hz, 2H), 3.49 (d, J = 11.0 Hz, 2H), 3.40 (s, 3H), 3.18-3.06 (m, 6H), 2.58 (s, 3H), 1.74-1.71 (m, 4H). ^{13}C NMR (100 MHz, DMSO- d_6): δ 196.80, 156.77 (d, J = 236.0 Hz), 154.01, 146.17, 132.92, 130.42, 129.76, 122.84, 118.05 (d, J = 8.5 Hz), 115.56 (d, J = 21.8 Hz), 107.42, 107.39, 54.82, 50.48, 46.15, 40.10, 27.14, 26.50, 20.29. Anal. calcd for $C_{24}H_{31}Cl_2FN_4O_2 \cdot H_2O$: C, 55.92; H, 6.45; N, 10.87. Found: C, 54.88; H, 6.05; N, 10.58.

4.1.14. 5-acetyl-1-(4-(4-(4-fluorophenyl)piperazin-1-yl)butyl)-3-propyl-1,3-dihydrobenzimidazol-2-one dihydrochloride (18).—This compound was prepared

from **13** as described for **17**. Yield 89%, as a white solid. 1H NMR (400 MHz, DMSO- d_6): δ 11.48 (br s, 1H), 9.69 (br s, 1H), 7.76 (d, J = 9.0 Hz, 1H), 7.74 (s, 1H), 7.36 (d, J = 8.1 Hz, 1H), 7.11-7.04 (m, 4H), 3.93-3.84 (m, 4H), 3.69 (d, J = 12.1 Hz, 2H), 3.49 (d, J = 11.3 Hz, 2H), 3.22 (t, J = 12.1 Hz, 2H), 3.19-3.09 (m, 4H), 2.57 (s, 3H), 1.79-1.64 (m, 6H), 0.86 (t, J = 7.3 Hz, 3H). ^{13}C NMR (100 MHz, DMSO- d_6): δ 196.83, 156.79 (d, J = 235.7 Hz), 153.85, 146.12, 132.90, 130.47, 129.13, 122.78, 118.07 (d, J = 7.6 Hz), 115.57 (d, J = 21.9 Hz), 107.52, 107.44, 54.74, 50.41, 46.13, 42.04, 40.45, 26.69, 25.28, 21.23, 20.26, 11.06.

Anal. calcd for $C_{26}H_{35}Cl_2FN_4O_2 \cdot 1/2H_2O$: C, 58.43; H, 6.79; N, 10.48. Found: C, 58.85; H, 6.46; N, 10.46.

4.1.15. 5-acetyl-1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-3-dihydrobenzimidazol-2-one dihydrochloride (19).—This compound was prepared from **13** as described for **17**. Yield 84%, as a white solid. 1H NMR (400 MHz, DMSO- d_6): δ 11.53 (br s, 1H), 8.75 (br s, 1H), 7.76 (d, J = 8.2 Hz, 1H), 7.71 (s, 1H), 7.36 (d, J = 8.2 Hz, 1H), 7.11-7.02 (m, 4H), 3.93-3.86 (m, 4H), 3.69 (d, J = 12.4 Hz, 2H), 3.51 (d, J = 11.4 Hz, 2H), 3.23 (t, J = 11.8 Hz, 2H), 3.20-3.09 (m, 4H), 2.57 (s, 3H), 1.79-1.62 (m, 6H), 1.31-1.22 (m, 4H), 0.81 (t, J = 6.7 Hz, 3H). ^{13}C NMR (100 MHz, DMSO- d_6): δ 196.82, 156.85 (d, J = 235.7 Hz), 153.79, 146.07, 132.93, 130.47, 129.08, 122.84, 118.13 (d, J = 7.7 Hz), 115.59 (d, J = 21.9 Hz), 107.53, 107.32, 54.74, 50.39, 46.19, 40.87, 40.45, 28.28, 27.53, 26.69, 25.28, 21.75, 20.27, 13.86. Anal. calcd for $C_{28}H_{39}Cl_2FN_4O_2 \cdot 1/2 H_2O$: C, 59.78; H, 7.17; N, 9.96. Found: C, 59.71; H, 7.04; N, 9.93.

4.1.16. 5-acetyl-3-decyl-1-{4-[4-(4-fluorophenyl)piperazin-1-yl]butyl}-1,3-dihydrobenzimidazol-2-one dihydrochloride (20).—This compound was prepared from **13** as described for **17**. Yield 87%, as a white solid. 1H NMR (free amine, 400 MHz, $CDCl_3$): δ 7.71 (dd, J = 6.7, 1.5 Hz, 1H), 7.61 (d, J = 1.3 Hz, 1H), 7.00 (d, J = 8.2 Hz, 1H), 6.93-6.89 (m, 2H), 6.84-6.81 (m, 2H), 3.92 (t, J = 7.2 Hz, 2H), 3.87 (t, J = 7.4 Hz, 2H), 3.09 (t, J = 5.0 Hz, 4H), 2.61 (t, J = 4.9 Hz, 4H), 2.59 (s, 3H), 2.45 (t, J = 7.4 Hz, 2H), 1.79 (quint, J = 7.3 Hz, 2H), 1.72 (quint, J = 7.2 Hz, 2H), 1.59 (quint, J = 7.8, 2H), 1.30-1.21 (m, 16H), 0.83 (t, J = 7.0 Hz, 3H). ^{13}C NMR (free amine, 100 MHz, $CDCl_3$): δ 197.22, 157.44 (d, J = 237.6 Hz), 154.64, 147.98, 133.67, 131.13, 129.91, 123.47, 118.08 (d, J = 8.0 Hz), 115.74 (d, J = 22.0 Hz), 107.32, 106.86, 57.73, 53.12, 50.01, 41.57, 41.23, 32.04, 29.70, 29.45, 29.43, 28.60, 27.00, 26.65, 26.41, 23.73, 22.83, 21.67, 14.28. Anal. calcd for $C_{33}H_{49}Cl_2FN_4O_2$: C, 63.55; H, 7.92; N, 8.98. Found: C, 63.49; H, 7.67; N, 8.81.

4.2. Competition binding assays for the σ_1R and σ_2R

Preparation of rat brain membranes and *in vitro* binding assays for the σ_1R and σ_2R were performed using methods published previously in detail [31]. In brief, homogenates of whole rat brain excluding the cerebellum (400-500 μ g) were incubated with 5 nM [3H](+)-pentazocine to label σ_1Rs , or 3 nM [3H]-DTG in the presence of 300 nM (+)-pentazocine to label σ_2Rs . Non-specific binding was determined in the presence of 10 μ M haloperidol. Ten concentrations (0.1-1000 nM) of each test compound were incubated for 120 min at 25 $^{\circ}C$ in 50 mM Tris-HCl, pH 8.0 to measure their ability to displace each radioligand from its binding sites. The total reaction volume was 500 μ l. All of the assays were terminated with the addition of ice-cold buffer and rapid vacuum filtration over glass fiber filters. Counts were extracted from the filters using Ecoscint (National Diagnostics, Manville, NJ) for at least 8 h prior to counting. K_i values were calculated using the Cheng-Prusoff equation [37] and K_d values determined in separate saturation assays.

4.3. Extended non- σ binding profile

The two compounds with the highest σ_1/σ_2 ratios (**14** and **15**) were tested for interactions with other binding sites known to interact with many historic σ compounds and

psychostimulants: opioid receptors, PCP binding site on NMDA receptors, dopamine (D₂) receptors, serotonin (5-HT₂) receptors, monoamine transporters (dopamine, serotonin, norepinephrine). Well-established, previously published methods were used [28]. Briefly, dopamine transporters were labeled using 0.5 nM [³H]WIN 35,428 in membranes prepared from rat striata; nonspecific binding was determined in the presence of 50 μM cocaine. Serotonin transporters were labeled using 0.2 nM [³H]paroxetine in membranes prepared from rat brainstem; nonspecific binding was determined in the presence of 1.5 μM imipramine. Norepinephrine transporters were labeled using 0.5 nM [³H]nisoxetine in membranes prepared from rat cerebral cortex; nonspecific binding was determined in the presence of 4 μM desipramine. All of the receptor assays were conducted using rat brain membranes minus cerebellum, similarly to those for σ R_s. Dopamine (D₂) receptors were labeled using 5 nM [³H](–)-sulpiride and nonspecific binding was determined with 1 μM haloperidol. Serotonin (5-HT₂) receptors were labeled using 2 nM [³H]ketanserin and nonspecific binding was determined with 1 μM mianserin. The PCP site on NMDA receptors were labeled using 5 nM [³H]1-[1-(2-thienyl)cyclohexyl]piperidine (TCP) and nonspecific binding was determined with 10 μM cyclazocine. Opioid receptors were labeled using 0.5 nM [³H]bremazocine and nonspecific binding was determined with 10 μM levallorphan. All of the assays were terminated with the addition of ice-cold buffer and vacuum filtration through glass fiber filters. Counts were extracted and K_i values determined as described for the σ R assays.

4.4. In vivo convulsion studies

Male, Swiss Webster mice (21–30 g; Harlan, USA) were pretreated (i.p.) with either saline or compound **14** (0.1, 1 or 10 mg/kg), then challenged 15 min later with a convulsive dose of cocaine (70 mg/kg, i.p.). Mice were observed for the next 30 min for convulsions, which were defined as a loss of righting reflexes for at least 5 s, combined with the presence of clonic limb movements or popcorn jumping. The data were analyzed using Fisher's exact tests. Animals were housed 1–5 per cage with a 12:12H light/dark cycle (lights on 06:00 hours) and *ad libitum* food and water. They were allowed to climatize following their arrival for one week prior to being utilized in any experiment. All procedures were approved by the Institutional Animal Care and Use Committee at West Virginia University.

Acknowledgments

We appreciate the technical assistance of Matthew Robson, Colleen Beatty, and Bahar Noorbakhsh. This project was supported by grants from NIDA R01 DA013978 and R01 DA023205.

References

- [1]. Martin WR, Eades CG, Thompson JA, Huppler RE, Gilbert PE, The effects of morphine- and nalorphine-like drugs in the nondependent and morphine-dependent chronic spinal dog, *The Journal of pharmacology and experimental therapeutics*, 197 (1976) 517–532. [PubMed: 945347]
- [2]. Vaupel DB, Naltrexone fails to antagonize the sigma effects of PCP and SKF 10,047 in the dog, *European journal of pharmacology*, 92 (1983) 269–274. [PubMed: 6313397]
- [3]. Su TP, Hayashi T, Vaupel DB, When the endogenous hallucinogenic trace amine N,N-dimethyltryptamine meets the sigma-1 receptor, *Science signaling*, 2 (2009) pe12. [PubMed: 19278957]

- [4]. Hanner M, Moebius FF, Flandorfer A, Knaus HG, Striessnig J, Kempner E, Glossmann H, Purification, molecular cloning, and expression of the mammalian sigma1-binding site, *Proceedings of the National Academy of Sciences of the United States of America*, 93 (1996) 8072–8077. [PubMed: 8755605]
- [5]. Kekuda R, Prasad PD, Fei YJ, Leibach FH, Ganapathy V, Cloning and functional expression of the human type 1 sigma receptor (hSigmaR1), *Biochemical and biophysical research communications*, 229 (1996) 553–558. [PubMed: 8954936]
- [6]. Seth P, Fei YJ, Li HW, Huang W, Leibach FH, Ganapathy V, Cloning and functional characterization of a sigma receptor from rat brain, *Journal of neurochemistry*, 70 (1998) 922–931. [PubMed: 9489711]
- [7]. Mei J, Pasternak GW, Molecular cloning and pharmacological characterization of the rat sigma1 receptor, *Biochemical pharmacology*, 62 (2001) 349–355. [PubMed: 11434908]
- [8]. Hellewell SB, Bruce A, Feinstein G, Orringer J, Williams W, Bowen WD, Rat liver and kidney contain high densities of sigma 1 and sigma 2 receptors: characterization by ligand binding and photoaffinity labeling, *European journal of pharmacology*, 268 (1994) 9–18. [PubMed: 7925616]
- [9]. Quirion R, Bowen WD, Itzhak Y, Junien JL, Musacchio JM, Rothman RB, Su TP, Tam SW, Taylor DP, A proposal for the classification of sigma binding sites, *Trends in pharmacological sciences*, 13 (1992) 85–86. [PubMed: 1315463]
- [10]. Walker JM, Bowen WD, Walker FO, Matsumoto RR, De Costa B, Rice KC, Sigma receptors: biology and function, *Pharmacological reviews*, 42 (1990) 355–402. [PubMed: 1964225]
- [11]. Xu J, Zeng C, Chu W, Pan F, Rothfuss JM, Zhang F, Tu Z, Zhou D, Zeng D, Vangveravong S, Johnston F, Spitzer D, Chang KC, Hotchkiss RS, Hawkins WG, Wheeler KT, Mach RH, Identification of the PGRMC1 protein complex as the putative sigma-2 receptor binding site, *Nature communications*, 2 (2011) 380.
- [12]. Chu UB, Mavlyutov TA, Chu ML, Yang H, Schulman A, Mesangeau C, McCurdy CR, Guo LW, Ruoho AE, The Sigma-2 Receptor and Progesterone Receptor Membrane Component 1 are Different Binding Sites Derived From Independent Genes, *EBioMedicine*, 2 (2015) 1806–1813. [PubMed: 26870805]
- [13]. Alon A, Schmidt HR, Wood MD, Sahn JJ, Martin SF, Kruse AC, Identification of the gene that codes for the sigma2 receptor, *Proceedings of the National Academy of Sciences of the United States of America*, 114 (2017) 7160–7165. [PubMed: 28559337]
- [14]. van Waarde A, Rybczynska AA, Ramakrishnan N, Ishiwata K, Elsinga PH, Dierckx RA, Sigma receptors in oncology: therapeutic and diagnostic applications of sigma ligands, *Current pharmaceutical design*, 16 (2010) 3519–3537. [PubMed: 21050178]
- [15]. Vilner BJ, John CS, Bowen WD, Sigma-1 and sigma-2 receptors are expressed in a wide variety of human and rodent tumor cell lines, *Cancer research*, 55 (1995) 408–413. [PubMed: 7812973]
- [16]. Wheeler KT, Wang LM, Wallen CA, Childers SR, Cline JM, Keng PC, Mach RH, Sigma-2 receptors as a biomarker of proliferation in solid tumours, *British journal of cancer*, 82 (2000) 1223–1232. [PubMed: 10735510]
- [17]. Georgiadis MO, Karoutzou O, Foscolos AS, Papanastasiou I, Sigma Receptor (sigmaR) Ligands with Antiproliferative and Anticancer Activity, *Molecules (Basel, Switzerland)*, 22 (2017).
- [18]. Crawford KW, Bowen WD, Sigma-2 receptor agonists activate a novel apoptotic pathway and potentiate antineoplastic drugs in breast tumor cell lines, *Cancer research*, 62 (2002) 313–322. [PubMed: 11782394]
- [19]. Nicholson HE, Alsharif WF, Comeau AB, Mesangeau C, Intagliata S, Mottinelli M, McCurdy CR, Bowen WD, Divergent Cytotoxic and Metabolically Stimulative Functions of Sigma-2 Receptors: Structure-activity Relationships of 6-acetyl-3-(4-(4-(4-fluorophenyl)piperazin-1-yl)butyl)benzo[d]oxazol-2(3H)-one (SN79) Derivatives, *The Journal of pharmacology and experimental therapeutics*, (2018).
- [20]. Smith SB, Introduction to Sigma Receptors: Their Role in Disease and as Therapeutic Targets, *Advances in experimental medicine and biology*, 964 (2017) 1–4. [PubMed: 28315260]
- [21]. Nguyen L, Lucke-Wold BP, Mookerjee S, Kaushal N, Matsumoto RR, Sigma-1 Receptors and Neurodegenerative Diseases: Towards a Hypothesis of Sigma-1 Receptors as Amplifiers of

Neurodegeneration and Neuroprotection, *Advances in experimental medicine and biology*, 964 (2017) 133–152. [PubMed: 28315269]

- [22]. Albayrak Y, Hashimoto K, Sigma-1 Receptor Agonists and Their Clinical Implications in Neuropsychiatric Disorders, *Advances in experimental medicine and biology*, 964 (2017) 153–161. [PubMed: 28315270]
- [23]. Fukunaga K, Moriguchi S, Stimulation of the Sigma-1 Receptor and the Effects on Neurogenesis and Depressive Behaviors in Mice, *Advances in experimental medicine and biology*, 964 (2017) 201–211. [PubMed: 28315273]
- [24]. Maurice T, Gogvadze N, Role of sigma1 Receptors in Learning and Memory and Alzheimer's Disease-Type Dementia, *Advances in experimental medicine and biology*, 964 (2017) 213–233. [PubMed: 28315274]
- [25]. Bowen WD, Sigma receptors: recent advances and new clinical potentials, *Pharmaceutica acta Helveticae*, 74 (2000) 211–218. [PubMed: 10812960]
- [26]. Guitart X, Codony X, Monroy X, Sigma receptors: biology and therapeutic potential, *Psychopharmacology*, 174 (2004) 301–319. [PubMed: 15197533]
- [27]. Matsumoto RR, McCracken KA, Friedman MJ, Pouw B, De Costa BR, Bowen WD, Conformationally restricted analogs of BD1008 and an antisense oligodeoxynucleotide targeting sigma1 receptors produce anti-cocaine effects in mice, *European journal of pharmacology*, 419 (2001) 163–174. [PubMed: 11426838]
- [28]. Matsumoto RR, McCracken KA, Pouw B, Zhang Y, Bowen WD, Involvement of sigma receptors in the behavioral effects of cocaine: evidence from novel ligands and antisense oligodeoxynucleotides, *Neuropharmacology*, 42 (2002) 1043–1055. [PubMed: 12128006]
- [29]. Matsumoto RR, McCracken KA, Pouw B, Miller J, Bowen WD, Williams W, De Costa BR, N-alkyl substituted analogs of the sigma receptor ligand BD1008 and traditional sigma receptor ligands affect cocaine-induced convulsions and lethality in mice, *European journal of pharmacology*, 411 (2001) 261–273. [PubMed: 11164383]
- [30]. Nastasi G, Miceli C, Pittala V, Modica MN, Prezzavento O, Romeo G, Rescifina A, Marrazzo A, Amata E, S2RSLDB: a comprehensive manually curated, internet-accessible database of the sigma-2 receptor selective ligands, *Journal of cheminformatics*, 9 (2017) 3. [PubMed: 28123452]
- [31]. Mesangeau C, Narayanan S, Green AM, Shaikh J, Kaushal N, Viard E, Xu YT, Fishback JA, Poupaert JH, Matsumoto RR, McCurdy CR, Conversion of a highly selective sigma-1 receptor-ligand to sigma-2 receptor preferring ligands with anticocaine activity, *Journal of medicinal chemistry*, 51 (2008) 1482–1486. [PubMed: 18278854]
- [32]. Xu YT, Kaushal N, Shaikh J, Wilson LL, Mesangeau C, McCurdy CR, Matsumoto RR, A novel substituted piperazine, CM156, attenuates the stimulant and toxic effects of cocaine in mice, *The Journal of pharmacology and experimental therapeutics*, 333 (2010) 491–500. [PubMed: 20100904]
- [33]. Kaushal N, Seminerio MJ, Shaikh J, Medina MA, Mesangeau C, Wilson LL, McCurdy CR, Matsumoto RR, CM156, a high affinity sigma ligand, attenuates the stimulant and neurotoxic effects of methamphetamine in mice, *Neuropharmacology*, 61 (2011) 992–1000. [PubMed: 21762711]
- [34]. Kaushal N, Robson MJ, Vinnakota H, Narayanan S, Avery BA, McCurdy CR, Matsumoto RR, Synthesis and Pharmacological Evaluation of 6-Acetyl-3-(4-(4-(4-fluorophenyl)piperazin-1-yl)butyl)benzo[d]oxazol-2(3H)-one (SN79), a Cocaine Antagonist, in Rodents, *The AAPS Journal*, 13 (2011) 336–346. [PubMed: 21494909]
- [35]. Kaushal N, Seminerio MJ, Robson MJ, McCurdy CR, Matsumoto RR, Pharmacological evaluation of SN79, a sigma (σ) receptor ligand, against methamphetamine-induced neurotoxicity in vivo, *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*, 23 (2013) 960–971. [PubMed: 22921523]
- [36]. Mesangeau C, Amata E, Alsharif W, Seminerio MJ, Robson MJ, Matsumoto RR, Poupaert JH, McCurdy CR, Synthesis and pharmacological evaluation of indole-based sigma receptor ligands, *European journal of medicinal chemistry*, 46 (2011) 5154–5161. [PubMed: 21899931]

- [37]. Cheng Y, Prusoff WH, Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction, *Biochemical pharmacology*, 22 (1973) 3099–3108. [PubMed: 4202581]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- A series of benzimidazolone derivatives as highly selective σ_2 R ligands were prepared and characterized.
- Replacement of the heterocyclic scaffolds of the lead compounds with a benzimidazolone moiety increased the σ_2 selectivity.
- The cyclic amine fragment 1-(4-fluorophenyl)piperazine is preferred rather than 1-cyclohexylpiperazine for the σ_2 selectivity.
- Introduction of n-propyl and n-pentyl group at *N*-3 position of the benzimidazolone, resulted in better affinity and selectivity for the σ_2 R.
- Selective σ_2 R ligands might attenuate cocaine-induced convulsions.

1 (CM156) $\sigma_1 K_i = 1.3 \text{ nM}$ $\sigma_2 K_i = 0.6 \text{ nM}$ $K_i \sigma_1 / K_i \sigma_2 = 2.17$ **2 (SN79)** $\sigma_1 K_i = 27 \text{ nM}$ $\sigma_2 K_i = 7 \text{ nM}$ $K_i \sigma_1 / K_i \sigma_2 = 3.86$

Fig. 1.
Chemical structures and affinity values of compounds **1** and **2**.

Fig. 2.

In *vivo* cocaine-induced convulsion studies on compound **14**. ** $P < 0.005$ vs vehicle response; $n = 6-8$ per group.

Scheme 1.

Reagents and conditions: (a) 1,4-dibromobutane, K₂CO₃, DMF, 60 °C, 70%; (b) cyclic amine, K₂CO₃, DMF, 60 °C, 56-62%.

Scheme 2.

Reagents and conditions: (a) 1-(4-fluorophenyl)piperazine, K_2CO_3 , DMF, 60 °C, 60%; (b) NH_2-NH_2 , EtOH, reflux, 83%; (c) 1-fluoro-2-nitrobenzene or 1-(4-chloro-3-nitrophenyl)ethanone, DIPEA, DMF, 100-120 °C, 83-93%; (d) H_2 (40 psi), 10% Pd/C, EtOH then CDI, THF, 20 °C, 51-68%; (e) NaH, TBAI, alkyl halide, DMF, 20 °C, 33-80% or K_2CO_3 , TBAI, alkyl halide, DMF, 60 °C, 84-89%.

Table 1.

σ Rs binding affinities and selectivity ratios of compounds **5**, **6**, **12-20**, and haloperidol.

Compd.	R ¹	R ²	K_i (nM $\pm$ SEM) ^a		$K_i \sigma_1 / K_i \sigma_2$
			σ_1 R	σ_2 R	
5	H	CH ₃	50.22 $\pm$ 7.59	2.57 $\pm$ 0.47	19.5
6	H	CH ₃	366.23 $\pm$ 101	2.05 $\pm$ 0.33	178.6
12	H	H	144 $\pm$ 6.6	18.4 $\pm$ 1.4	7.8
13	COCH ₃	H	166 $\pm$ 12	28.5 $\pm$ 8.6	5.8
14	H	nC ₃ H ₇	752 $\pm$ 63	0.66 $\pm$ 0.01	1139
15	H	nC ₅ H ₁₁	2274 $\pm$ 187	4.27 $\pm$ 0.29	532.5
16	H	nC ₁₀ H ₂₁	724 $\pm$ 13.6	4.55 $\pm$ 0.21	159.1
17	COCH ₃	CH ₃	72.3 $\pm$ 2.6	7.6 $\pm$ 0.31	9.5
18	COCH ₃	nC ₃ H ₇	273.5 $\pm$ 1.33	1.81 $\pm$ 0.02	151.1
19	COCH ₃	nC ₅ H ₁₁	299 $\pm$ 32	2.15 $\pm$ 0.94	139
20	COCH ₃	nC ₁₀ H ₂₁	988 $\pm$ 53	32.5 $\pm$ 5.3	30.4
Haloperidol			3.35 $\pm$ 0.80	80.60 $\pm$ 14.10	0.042

^aThe values represent the mean $\pm$ SEM from triplicate assays.

Table 2.Binding profile of **14** and **15** for other non- σ proteins.

Compd.	K_i (nM $\pm$ SEM) ^a						
	Monoamine transporters			Other receptors			
	Dopamine	Serotonin	Norepinephrine	Dopamine (D ₂)	Serotonin (5-HT ₂)	NMDA	Opioid
14	>1000	>1000	>10000	>10000	54.70 $\pm$ 0.27	>10000	>10000
15	>1000	445.9 $\pm$ 3.2	>10000	>1000	249.6 $\pm$ 4.8	>10000	>10000

^aAffinities (K_i values in nanomolar) were determined in rat brain homogenates. The values represent the mean $\pm$ SEM from triplicate assays. Values of >10000 nM represent less than 30% displacement of the radioligand at that concentration.