Published in final edited form as:

J Emerg Med. 2019 February ; 56(2): 153–165. doi:10.1016/j.jemermed.2018.10.032.

Blunt Thoracolumbar-Spine Trauma Evaluation in the Emergency Department: A Meta-Analysis of Diagnostic Accuracy for History, Physical Examination, and Imaging

James VandenBerg, MD, MSC^{a,b}, Kevin Cullison, MD^{b,c}, Susan A. Fowler, MLIS^d, Matthew S. Parsons, MD^e, Christopher M. McAndrew, MD, MSC^f, and Christopher R. Carpenter, MD, MSC^b

^aDepartment of Emergency Medicine, Detroit Receiving Hospital and University Health Center, Detroit, Michigan

^bDivision of Emergency Medicine, Washington University School of Medicine, St. Louis, Missouri

^cDepartment of Emergency Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania

^dBernard Becker Medical Library, Washington University School of Medicine, St. Louis, Missouri

^eDepartment of Radiology, Washington University School of Medicine, St. Louis, Missouri

^fDepartment of Orthopedic Surgery, Washington University School of Medicine, St. Louis, Missouri

Abstract

Background: Delayed diagnoses of unstable thoracolumbar-spine (TL-spine) fractures can result in neurological deficits and avoidable pain, so it is important for clinicians to reach prompt diagnostic decisions. There are no validated decision aids for determining which trauma patients warrant TL-spine imaging.

Objectives: Quantifying the diagnostic accuracy of the injury mechanism, physical examination, associated injuries, clinical decision aids, and imaging for evaluating blunt TL-spine trauma patients.

Methods: A search strategy for studies including adult blunt TL-spine trauma using PubMed, Embase, Scopus, CENTRAL, Cochrane Database of Systematic Reviews, and [ClinicalTrials.gov](https://www.clinicaltrials.gov) was performed. Excluded studies lacked data to construct 2×2 tables, were duplicates, not primary research, did not focus on blunt trauma, examined associated injuries without any utility in identifying TL-spine injuries, only studied cervical-spine fractures, were non-English, had a pediatric setting, or were cadaver/autopsy reports. Risk of bias was assessed using the Quality Assessment Tool for Diagnostic Accuracy Studies. Diagnostic predictors were analyzed with a meta-analysis of sensitivity, specificity, and likelihood ratios.

Results: In blunt trauma patients in the emergency department, the weighted pretest probability of a TL-spine fracture was 15%. The estimates for the detection of TL-spine fractures with plain

film was positive likelihood ratio (+LR)=25.0 (95% CI=4.1-152.2, I²=94%, p<0.001) and negative likelihood ratio (-LR)=0.43 (95% CI=0.32-0.59, I²=84%, p<0.001), while for CT were +LR=81.1 (95% CI=14.1-467.9, I²=87%, p<0.001) and -LR=0.04 (95% CI=0.02-0.08, I²=23%, p=0.26).

Conclusions: CT is more accurate than plain films for detecting TL-spine fractures. Injury mechanism, physical exam, and associated injuries alone are not accurate to rule-in or -out TL-spine fractures.

Keywords

Blunt trauma; spine trauma; diagnostic imaging; radiography; computed tomography

Introduction

Thoracolumbar-spine (TL-spine) injuries occur in approximately 7% of blunt trauma injuries, accounting for approximately half of vertebral injuries.¹⁻³ Complications of misdiagnosis include radiculopathy or spinal cord injuries, with a 19-50% occurrence.⁴⁻⁶ Neurological deterioration is up to eight-fold more likely with delayed or missed diagnosis.^{4,7,8} Plain films can miss injuries or assessments for instability. As depicted in Figure 1, computed tomography (CT) of the spine more accurately identifies fracture comminution.

Unlike imaging of the cervical spine (C-spine), there are no validated decision aids for determining which trauma patients warrant TL-spine imaging. This lack of evidence may explain why diagnostic guidelines vary. The Eastern Association for the Surgery of Trauma and the American College of Radiology recommend CT as the modality of choice, whereas the Advanced Trauma Life Support (ATLS) and the National Institute for Health and Care Excellence recommend plain films initially.^{6,9-11}

CT is increasingly used to screen the torso for visceral trauma, raising the issue of whether dedicated TL-spine reformations are required to accurately identify clinically significant TL-spine injuries. Furthermore, it is unclear if patients with minor trauma (who don't require initial CT screening) may have their TL-spine cleared by physical examination alone. The disadvantages of CT imaging include increased radiation exposure, incidental findings prompting consideration of or the obtaining of further testing, prolonged ED lengths of stay, and higher costs.¹²⁻¹⁵ Expert panels within the specialty of emergency medicine and other organizations and specialty societies, and larger guideline panels have identified the radiographic clearance of blunt TL-spine trauma patients as a priority for further investigation.^{3,16}

The primary objective of this systematic review was to quantify parameters of diagnostic accuracy of commonly used imaging modalities for blunt TL spine trauma in patients 14 years or older, with a focus upon the mechanism of injury, the physical examination findings, and commonly-employed imaging modalities.

Materials and Methods

Study Design

This manuscript adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and the Meta-Analysis of Observational Studies in Epidemiology (MOOSE) reporting guidelines.^{17,18} Eligible studies included patients 14 years or older in an Emergency Department (ED) following blunt trauma in whom evaluations for TL-spine injuries was reported in sufficient detail to reconstruct 2×2 tables. “True Positive” (TP), “False Positive” (FP), “True Negative” (TN), and “False Negative” (FN) were defined *a priori*. “Disease” indicated TL-spine injury and “non-disease” the absence of TL-spine injury as defined by the original study’s criterion standard.

Study Setting and Protocol

A medical librarian created the electronic search strategy, which was completed on January 30, 2017, limited to English, and exported to EndNote. Full search strategies are available in Appendix A. Two authors independently reviewed the titles and abstracts to identify relevant articles. Studies were excluded if they lacked data to construct 2×2 tables, were duplicates, were case reports/narrative reviews, lacked focus on blunt trauma, examined associated injuries without using them as predictive factors for identifying TL-spine injuries (such as using spine injuries to predict traumatic aortic injuries), only studied C-spine fractures, were not printed in the English language, were performed at a pediatric hospital, or were cadaver/autopsy reports. Authors were contacted if their manuscripts lacked sufficient detail to reconstruct 2×2 tables. Two investigators independently hand-searched abstracts from scientific assemblies, including: Academic Emergency Medicine, Annals of Emergency Medicine, American Journal of Roentgenology, Skeletal Radiology, The Spine Journal, Spine, Journal of Neurological Surgery: Spine, Journal of Trauma-injury, Infection, and Critical Care, Journal of Trauma and Acute Care Surgery, Journal of Orthopaedic Trauma, Journal of Bone and Joint Surgery (American and British editions), and society presentations from Orthopedic Trauma Association meetings and American Association of Neurological Surgeons meetings.

Two authors independently evaluated study’s risk of bias and applicability using the revised Quality Assessment Tool for Diagnostic Accuracy Studies (QUADAS-2).¹⁹ The QUADAS-2 evaluation was performed multiple times for studies examining multiple index tests. Responses were compared, disagreements recorded, and discussed. A third author adjudicated disagreements. The authors established several *a priori* considerations for the QUADAS-2 assessments:

- Increased risk of inappropriate exclusions occurred when studies excluded patients with difficult to diagnose TL-injuries; those who were not admitted or underwent spine surgery; or when certain patients did not receive the index/criterion test but should have.
- A level I or II trauma center represented the most relevant, least biased setting. Acute blunt trauma patients represented the patients of direct relevance. Studies in other settings or different populations were “low applicability”.

- If there was no mention of blinding index test interpreters to the criterion standard or vice versa, then prospective studies were “uncertain” and retrospective studies were “high risk”.
- If plain film radiography lacked antero-posterior (AP) and lateral views, or if CT test protocols used less than 64 slices, the index test conduct or interpretation were “low applicability”.
- Studies in which all patients received either CT, magnetic resonance imaging (MRI), surgical confirmation, or clinical follow-up were low risk of bias.
- An appropriate interval between the index test and the criterion standard was when both tests occurred during the patient’s ED visit or initial hospitalization.

Measures

Two authors independently abstracted study data: clinical setting, location, age, inclusion criteria, criterion standard, diagnostic modality specifications, TL-spine injury prevalence, study design, index tests, sensitivity (SN), specificity (SP), positive likelihood ratio (+LR), and negative likelihood ratio (–LR). Index tests or data included injury mechanism, history or physical examination findings of presence or absence of “associated injuries”, results of plain film studies, and results of CT for diagnosing TL-spine fractures. CT studies had two groups: routine Chest-Abdomen-Pelvis (CAP) CT scans, and reformatted CT images of the TL-spine (which required a specified axial magnification and bone algorithm reconstruction). We referred to these as “CAP CT” and “reformatted TL-spine CT”, respectively. We also evaluated the accuracy of ultrasound (US) and MRI to detect injuries to the TL-spine posterior ligamentous complex (PLC). When possible, we distinguished all fractures from clinically significant fractures as the latter required surgical stabilization, orthotics, or were deemed “unstable”.

Data Analysis

Abstracted data was used to recreate 2×2 tables and calculate SN, SP, +LR, and –LR. If studies used 3×2 tables, including a third “equivocal” result, 2×2 tables were created by conservatively adding the equivocal results with a true injury to the FN cell and those without injury into the FP cell.²⁰ We also analyzed thoracic or lumbar injury subgroups. Data was preferentially abstracted based on the presence of injuries using individual patients as the population. Results presenting individual vertebrae as the population (e.g. fractured vs. unfractured vertebrae), were analyzed separately. If studies used multiple observers and calculated SN and SP based on mean averages, these values were used to recreate 2×2 tables. When greater than one study assessed the same index test, we computed meta-analysis estimates using MetaDiSc (Hospital Universitario Ramon y Cajal, Madrid, Spain) and forest tree plots for SN, SP, +LR, and –LR using a random-effects model.²¹ Pretest probability (prevalence) was estimated using a weighted average. The DerSimonian-Laird random effects model was used to calculate interstudy heterogeneity amongst pooled SN and SP using the index of inconsistency (I^2).^{22,23} QUADAS-2 between-rater agreement was quantified using Cohen’s kappa (κ) with IBM SPSS Statistic for Windows, Version 22.0 (Armonk, NY).

Results

The primary search yielded 6,420 articles, with 224 selected for review and 14 abstracts identified via hand search (Figure 2). The 48 included studies are summarized in Appendix B. The mean age ranged from 35-44 and the prevalence of TL-spine fractures ranged from 4-72%, thoracic-spine fractures 2-61%, lumbar-spine fractures 10-25%, and PLC injuries amongst those with TL-spine fractures was 28-67%. For studies with a criterion standard likely to correctly classify the presence of TL-spine fractures, the TL-spine fracture prevalence was 8-41%.²⁴⁻³⁵ Based on these studies, the estimated pretest probability (prevalence) for TL-spine fracture in blunt trauma was 15%.

Quality Assessment:

The QUADAS-2 assessment inter-rater kappa ranged from 0.49 – 1.00. Appendix C shows the final adjudicated assessment for each study. The included studies were of variable quality. For patient selection, there were only two case-control studies.^{36,37} Eight studies only included “sicker” trauma patients, such as patients admitted to the hospital or intubated, which increases the risk of spectrum bias when applying these results to less ill patients.^{24,29,32,33,38-41} Aside from the studies utilizing US or MRI, most occurred in the ED. Another source of bias involved the un-blinded interpretation of index tests and criterion standards. Most studies were retrospective, and we assumed that index tests occurred with awareness of the criterion standard results unless otherwise specified. Lastly, since most criterion standards included the index test, incorporation bias likely falsely raised the SN and SP of the index.⁴²

Injury Mechanism

Ten studies examined injury mechanisms as predictors of TL-spine fractures.^{1,2,27,28,31,43-47} Injury incidents included falls, motor vehicle crashes (MVC), pedestrian struck, bicyclist accidents, motorcycle accidents, assaults, and “high risk mechanisms” as defined by individual studies. Injury mechanism pooled +LR ranged from 0.5-1.7 and –LR ranged from 0.63-1.25 (see Appendix D1). Thus, the mechanism of injury did not serve as a tool for highly confident decision-making.

History and Physical Examination

Eighteen studies examined the diagnostic accuracy of history or physical examination, including complaints of or findings of spine pain or tenderness upon palpation, lateral spine pain upon palpation, distracting injury, abnormal neurologic examination, altered consciousness, intoxication, palpable spine deformity, back bruising, skin abrasion/laceration, physician gestalt, and the whole physical examination.^{25,27,31,32,34,41,43-52} The most useful physical examination to rule-in a TL-spine fracture was a palpable spine deformity with +LR=15.3 (95% CI=7.1-33.0, $I^2=0\%$, $p=0.79$).^{46,47} No single negative physical examination finding (such as lack of spine tenderness) significantly reduced the probability of a TL-spine fracture (see Appendix D2).

One study examined the diagnostic accuracy of history and physical examination for thoracic-spine fractures.⁵³ Spine tenderness with palpation showed +LR=3.4 (95%

CI=2.4-4.8) and $-LR=0.57$ (95% CI=0.41-0.81); altered consciousness $+LR=2.8$ (95% CI=2.0-4.0) and $-LR=0.70$ (95% CI=0.57-0.87); distracting injury $+LR=1.9$ (95% CI=1.6-2.3) and $-LR=0.52$ (95% CI=0.37-0.75); and the whole physical examination was $+LR=1.9$ (95% CI=1.8-2.0) and $-LR=0.07$ (95% CI=0.02-0.27).

In studies assessing clinician's gestalt for TL-spine fractures, a physician estimate of "greater than low pretest probability" (from a scale of very low, low, intermediate, high, and very high) had a $+LR=1.8$ (95% CI=1.4-2.3) and $-LR=0.50$ (95% CI=0.29-0.84). In comparison, a "greater than high pretest probability" had $+LR=12.6$ (95% CI=4.2-37.3) and $-LR=0.80$ (95% CI=0.67-0.96).³² Reynolds et al.⁵⁴ studied gestalt for lumbar-spine fractures, and showed $+LR=2.5$ (95% CI=1.7-3.7) and $-LR=0.47$ (95% CI=0.26-0.87) for a greater than low pretest probability.⁵⁴ In comparison to gestalt, the whole physical examination for TL-spine fractures showed a pooled $+LR=1.7$ (95% CI=1.0-2.8, $I^2=100\%$, $p<0.001$) and a $-LR=0.40$ (95% CI=0.27-0.59, $I^2=76\%$, $p<0.001$).

One study involved a diagnostic pathway requiring imaging (plain film, CT, or MRI) if patients had complaints of spine pain, tenderness upon palpation (of midline), local signs of TL-spine injury, neurologic deficits, or a C-spine fracture. In the absence of these, patients were imaged if Glasgow Coma Scale was less than 15. If it equaled 15, they received imaging in the presence of distracting injury or alcohol/drug intoxication. With this algorithm 92% of patients would have received imaging, with a reported SN=100% (95% CI=88-100%) and SP=11% (95% CI=5-21%) for TL-spine fractures.⁴⁶ A second study used the criteria of spine pain, midline tenderness on palpation, spine deformity, neurologic deficit, age ≥ 60 , or high risk mechanism to detect clinically significant TL-spine fractures with a reported SN=99% (95% CI=97-100%), SP=29% (95% CI=27-31%), and SN=100% for clinically significant fractures. In this study, the addition of age ≥ 60 appeared to help improve the SN of the history and physical examination findings, though SP was reduced.⁴⁷

Associated Injuries

Seven studies examined the accuracy of associated injuries for diagnosing TL-spine fractures.^{1,29,39,43,45,46,55-57} Associated injuries included C-spine injury, major injuries (defined as an Abbreviated Injury Scale greater than or equal to three), injuries divided anatomically (cardiac, pulmonary/thoracic, abdominal, orthopedic, ophthalmic/maxillofacial, or a combination of any of these injuries), and pelvic fractures. Appendix D3 shows that the $+LR$ ranged from 0.4-2.4, while $-LR$ ranged from 0.83-2.49.

One study evaluated the accuracy of rib fractures to diagnose thoracic-spine fractures, and showed $+LR=1.6$ (95% CI=1.2-2.0) and $-LR=0.76$ (95% CI=0.63-0.93).³⁹ Another investigated the accuracy of the presence of abdominal aortic calcification and low bone mineral density with lumbar-spine fractures in blunt trauma. Two board-certified fellowship-trained radiologists interpreted the studies for the presence of fractures.⁵⁵ Abdominal aortic calcification was studied as its severity is associated with bone fragility and vertebral fractures.^{55,58} Abdominal aortic calcification resulted in $+LR=2.1$ (95% CI=1.5-2.9) and $-LR=0.52$ (95% CI=0.33-0.83), while low bone mineral density was $+LR=2.9$ (95% CI=2.2-3.9) and $-LR=0.33$ (95% CI=0.18-0.60).

Plain Film

Eight studies investigated the accuracy of plain films, all diagnostic accuracy results for TL-spine, thoracic-spine, and lumbar-spine studies are detailed in Appendix D4. Five studies evaluated the diagnostic accuracy of TL-spine plain films.^{24,29,33,59,60} As shown in the forest plots of Figure 3, plain films had a +LR=25.0 (95% CI=4.1-152.2, $I^2=94\%$, $p<0.001$), and -LR=0.43 (95% CI=0.32-0.59, $I^2=84\%$, $p<0.001$). Table 1 details the index tests and criterion standards for each study in this analysis. Three studies examined the diagnostic accuracy of plain films to detect thoracic-spine fractures, with a +LR=2.2 (95% CI=0.7-7.2, $I^2=89\%$, $p<0.001$) and -LR=0.62 (95% CI=0.30-1.27, $I^2=87\%$, $p=0.005$).^{33,61,62} Two studies investigated lumbar-spine fractures: +LR=15.7 (95% CI=7.6-32.4, $I^2=0\%$, $p=0.78$) and -LR= 0.60 (95% CI=0.35-1.03, $I^2=68\%$, $p=0.08$).^{33,38}

Barrios et al.⁶³ evaluated the accuracy of chest x-ray for thoracic-spine fractures in blunt trauma patients. The prevalence of thoracic-spine fractures was 2% (four patients), with SN=0% (95% CI=0-60%) and SP=100% (95% CI=98-100%). Chen et al.²⁶ and Deyle et al.⁶⁴ reported the accuracy of the Lodox/Statscan, a digital x-ray device that can generate full-body anterior and lateral views. With this device, the set of plain radiographs called for in the primary survey can be replaced. As of 2012, it is installed in 50 sites throughout the world.⁶⁵ Chen et al.²⁶ examined TL-spine fractures with a +LR=16.7 (95% CI=8.3-33.5) and -LR=0.18 (95% CI=0.07-0.43). Deyle et al.⁶⁴ evaluated thoracic-spine and lumbar-spine fractures separately, with +LR=175.1 (95% CI=10.8-2,846.0) and -LR=0.57 (95% CI=0.44-0.73) for thoracic-spine fractures and +LR=282.8 (95% CI=17.7-4,524.8) and -LR=0.26 (95% CI=0.17-0.41) for lumbar-spine fractures.

CAP CT

Eight studies investigated the accuracy of CT, and all diagnostic accuracy results are detailed in Appendix D5. Six studies evaluated the diagnostic accuracy of CAP CT for diagnosing TL-spine fractures.^{24,28,30,33,59,60} As shown in the forest plots of Figure 4, CAP CT had +LR=81.1 (95% CI=14.1-467.9, $I^2=87\%$, $p<0.001$) and -LR=0.04 (95% CI=0.02-0.08, $I^2=23\%$, $p=0.26$). The index tests and criterion standards for each study in this analysis are shown in Table 2. Additional studies that were not included in the meta-analysis (2x2 tables could not be recreated) showed inter-radiologist κ range of 0.80-0.98 for detecting TL-spine fractures.^{37,66}

Smith MW et al.³³ examined thoracic-spine fractures amongst individual patients, showing a +LR=77.3 (95% CI=4.8-1,241.2) and -LR=0.23 (95% CI=0.08-0.68). Overall, 3 mm slices are superior to 5 mm slices, and multiplanar reformations improved accuracy. Using 3 mm slices with or without multiplanar reformations demonstrate a +LR=45.0 (95% CI=2.9-697.8) and -LR=0.02 (95% CI=0.0-0.4); 5 mm slices with multiplanar reformations demonstrate +LR=42.9 (95% CI=2.8-666.4) and -LR=0.07 (95% CI=0.01-0.33); and 5 mm slices without showed +LR=38.7 (95% CI=2.5-603.6) and -LR=0.16 (95% CI=0.06-0.43).⁴⁰

Three studies investigated the diagnostic accuracy of CAP CT for diagnosing lumbar-spine fractures.^{33,38,67} The pooled estimate for +LR=202.5 (95% CI=41.1-998.7, $I^2=0\%$, $p=0.46$) and -LR=0.02 (95% CI=0.00-0.11, $I^2=0\%$, $p=0.63$).

Reformatted TL-Spine CT and Other CT Studies

Rozenberg et al.³⁰ investigated the diagnostic accuracy of reformatted TL-spine CT with +LR=40.2 (95% CI=16.8-95.8) and -LR=0.05 (95% CI=0.01-0.21). Carter et al.⁶⁸ also examined reformatted TL-spine CT but was excluded (2×2 tables could not be recreated). They showed reformatted TL-spine CT had a SN=97% (95% CI=89-100%). No included study examined inter-radiologist agreement, but a previous study showed $\kappa=0.98$ using a reformatted TL-spine CT protocol.⁶⁶

Sroka et al.³⁴ demonstrated that the scout AP and lateral CT images of the TL-spine from a four-slice scanner had a +LR and -LR of 187.5 (95% CI=11.7-2,999.5) and 0.12 (95% CI=0.03-0.52). A study by Karaca et al.⁶⁹ compared the diagnostic accuracy of 256-slice dual-source CT scans (DSCT) to MRI in detecting bone marrow edema. Bone marrow edema served as a predictive factor for vertebral compression fractures. The study showed that DSCT had a +LR=72.4 (95% CI=18.2-288.0) and -LR=0.11 (95% CI=0.05-0.25). Inter-observer agreement for the determination of bone marrow edema had a $k=0.82$ and intra-observer agreement $k=0.80$.

Clinically Significant Fractures for Plain Film and CT Studies

The summary data for clinically significant fractures is shown in Table 3. Wintermark et al.³⁵ evaluated the accuracy of plain films for clinically significant TL-spine fractures, reporting a +LR=61.6 (95% CI=3.5-1,079.4) and -LR=0.66 (95% CI=0.44-0.98).

Plain films missed 66.7% clinically significant fractures.³⁵ In comparison, Hauser et al.⁵⁹ showed that no clinically significant fractures were missed and Berry et al.²⁴ showed that 7.7% of clinically significant fractures were missed (one patient who required an orthotic brace). Mejia et al.⁶⁰ provided no abstractable data, but noted that 12 clinically significant fractures were missed on plain film (one requiring surgical stabilization and 11 orthoses), whereas CT missed none. For clinically significant thoracic-spine fractures, SN for plain films reportedly ranged from 75-100%.^{33,62} Huddleston et al.⁶¹ also reported on plain film's ability to detect "severe" thoracic-spine fractures based on the Thoracolumbar Injury Classification and Severity Score.⁷⁰ They found that the SN of plain films did not improve with increasing injury severity: "least severe" SN=55%, "moderate" SN=67%, and "most severe" injuries SN=59%. For clinically significant lumbar-spine fractures the SN=63%.³³

Mancini et al.²⁸ showed a +LR=138.9 (95% CI=8.8-2,200.8) and -LR=0.02 (95% CI=0.00-0.34) for CAP CT, while Wintermark et al.³⁵ demonstrated a +LR=166.6 (95% CI=10.5-2,651.8) and -LR=0.06 (95% CI=0.01-0.51). The remaining studies reported that CAP CT did not miss any clinically significant TL-spine fractures.^{24,30,33,59,60} Gestring et al.⁶⁷ provided sufficient data for clinically significant lumbar-spine fractures and showed +LR=122.4 (95% CI=7.6-1,967.1) and -LR=0.10 (95% CI=0.01-1.40). Deunk et al.³⁸ showed that CT led to a change in management in 21% of patients with lumbar-spine fractures when compared to plain films.

Rozenberg et al.³⁰ was the only included study that compared both reformatted TL-spine CT and CAP CT, showing that neither missed any clinically significant TL-spine fractures. Carter et al.⁶⁸ was excluded because of inadequate detail to reconstruct 2×2 tables, but

reported that reformatted TL-spine CT missed no clinically significant TL-spine fractures, while CAP CT missed 29% (all of which required orthotic bracing).

Ligamentous Spinal Injuries (US, CT, MRI)

Three studies used US to examine injuries to the PLC.^{36,71,72} These studies also examined individual components of the PLC, including the supraspinous ligament and interspinous ligament. Appendix D6 shows all diagnostic accuracy summary data. US was most accurate for detecting interspinous ligament injuries, with a +LR of 4.8 (95% CI=0.1-379.9, I²=89%, p=0.002) and -LR of 0.20 (95% CI=0.03-1.36, I²=0%, p=0.32).

Haba et al.⁷³ used MRI to detect supraspinous or interspinous ligament injury.⁷³ The diagnostic accuracy for detecting supraspinous ligament injuries was +LR=11.6 (95% CI=1.6-82.8) and -LR=0.11 (95% CI=0.03-0.38), and interspinous ligament injuries +LR=7.7 (95% CI=1.8-33.7) and -LR=0.02 (95% CI=0.00-0.48). Ghanem et al.⁷⁴ evaluated MRI's accuracy in detecting adjoining intervertebral disc injury (PLC and anterior/posterior longitudinal ligaments) in TL-spine fractures with a +LR=3.7 (95% CI=1.4-10.0) and -LR=0.09 (95% CI=0.01-0.61).

Discussion

History and physical examination findings are generally inaccurate to rule-in or rule-out TL-spine fractures. The findings were based on mostly Retrospective studies were the data source for most of the findings reported herein. Retrospective studies are inherently prone to diagnostic bias.⁴² The criterion standard for many included the composite of all clinical/radiologic information (including the index test), introducing incorporation bias that raises both the observed SN and SP of the index test.⁴² Though history and physical examination had a lower accuracy than CT imaging, decision aids to guide TL-spine imaging exist and await prospective validation.^{46,47,75}

A +LR over 10 and -LR below 0.1 are regarded as diagnostic tests with sufficient accuracy to rule-in or rule-out disease, respectively.⁷⁶ Pooled LR of CT's accuracy were within these thresholds. The CT studies with the lowest SN were due to clinically insignificant fractures, and missed fractures may be attributable to imaging technique.^{30,35,40} In contrast, plain film's diagnostic accuracy for TL-spine fractures had pooled +LR=25.0 (95% CI=4.1-152.2). The +LR point estimate exceeds 10, but the lower margin of the confidence interval is 4. While thoracic-spine plain films were inadequate for ruling in fractures, lumbar-spine plain films may be an adequate alternative to CT as the +LR=15.7 (95% CI=7.6-32.4).

For clinically significant fractures, plain film had a wide SN range: 33-100% (Table 3).^{24,35,59} While the CT studies had a consistently high SN, one excluded study compared the diagnostic accuracy of CAP CT to reformatted TL-spine CT, and reported that CAP CT had a SN=64% (95% CI=51-75%) for all TL-spine fractures and missed 28.6% of clinically significant fractures.⁶⁸ Therefore, CAP CT can also miss both clinically significant and insignificant TL-spine fractures.

Reformatted TL-spine CT is not associated with improvements in management when compared to CAP CT.^{30,66} The missed fractures on CAP CT generally required only conservative management.^{30,68} The American College of Radiology deemed the use of reformatted images as appropriate, because it may enhance the visualization of spinal alignment at no additional cost or radiation. However, it should be noted that a 3-megapixel diagnostic display is used in most instances within the reading rooms of radiology departments. This is in contrast to the usual 1-megapixel display that can be found within the clinical areas of an emergency department. Given the difference pixel density, subtle findings may be more apparent to radiologists.

The current data is sparse, and observational studies providing quantitative estimates of SN and SP represent the lowest tier of diagnostic evidence, evidence that does not equate to patient-centric benefit.⁷⁷ Diagnostic randomized controlled trials are often non-existent and fail to evaluate patient-oriented outcomes.⁷⁸ In lieu of randomized controlled trials, researchers substitute trauma registries and highlight missed spine fractures when advanced imaging is not obtained,⁷⁹ but usually fail to measure the potential harms of reformatted imaging including incidental findings and downstream consequences of overdiagnosis and overtreatment.⁸⁰ However, as reformatted TL-spine CT has no additional cost or radiation to the patient, it would be beneficial to investigate larger scale prospective studies to delineate their impact on immediate management and downstream outcomes.¹⁰

Accelerating patterns of CT utilization over the past two decades, and overlying controversies of the management for neurologically intact TL-spine injury patients shape the contemporary research landscape for blunt trauma victims. This meta-analysis includes CT studies from 2002-2017, a period when advancing CT technology fueled the role of advanced imaging in the first-line ED evaluation of trauma patients.^{12,81,82} As a result, the need for imaging in blunt trauma patients is debatably limited to patients who don't require CAP CT for visceral torso injury. None of the studies included in our meta-analysis focused on this population, and therefore more research is needed in this area. Additionally, certain injuries, including TL-spine burst fractures without neurological deficit, can often be managed non-operatively.^{83,84} Moreover, a recent multicenter prospective randomized study in Canada revealed no benefit in short- or long-term outcomes in terms of pain relief or disability assessment in patients with incomplete TL-spine burst fractures treated with orthotic bracing versus no bracing.^{85,86} Surgical intervention or orthotics traditionally define what constitutes a "clinically significant" injury. Further investigations into which patients warrant TL-spine imaging after blunt trauma should incorporate rigid definitions for clinically significant injuries based upon patient-centered outcomes of available interventions, with input from spine surgeons, rather than being based solely upon whether or not a medical intervention (such as an orthotic) or a surgical intervention ensued.

Limitations

The overall quantity and quality of the available evidence is imperfect and far from ideal for deriving a clinical decision rule. The applicability of our findings to older populations is limited, since the majority of our studies had a mean age between 30-50 years, and advanced age may have an influence upon the accuracy of combined physical examination data.

Incorporating advanced age (60 years and older) has been shown to improve the accuracy of TL-spine fracture decision aids.⁴⁷

Temporal bias exists in radiology as advancing technology improves image quality and accuracy.⁸⁷ While scan quality may be similar between different CT scanners, 64-slice scanners have increased image quality, scan speed, and advanced image post-processing applications.⁸⁸ If an older CT scanner was used as a criterion standard and misclassified patients as having no fracture, imperfect gold standard bias is possible, which could decrease the observed SN and SP of the index test (assuming the errors of the index test and criterion standard are independent). Conversely, if errors of the index test and criterion standard are correlated, then this increases the observed SN and SP.⁴²

Numerous studies were excluded, usually because of an inability to create 2×2 tables. We contacted authors when appropriate to clarify reported data, but some did not respond or could not supply the required data. Adherence to the Standards for Reporting Diagnostic Accuracy (STARD) reporting guidelines improves the quality of reported data, and would improve the pool of available evidence for future meta-analyses.⁸⁹

Conclusion

High-quality research to guide evidence-based diagnostic decision-making for TL-spine fractures in blunt trauma victims is lacking. In acute blunt trauma patients with suspected TL-spine injuries, the prevalence of a TL-spine fracture is approximately 15%. No injury mechanism, physical examination, or associated injury accurately ruled in the presence of a TL-spine fracture. CT is superior to plain film both in the detection of all TL-spine fractures and clinically significant fractures. While reformatted TL-spine CT is more accurate than CAP CT, it is unknown whether these differences impact management or patient-centric outcomes. Of course, individual clinicians may occasionally obtain plain films when they have a low but non-zero percent suspicion of significant injury, or when an injury that could be clearly demonstrated with a plain film series is anticipated. This review does not advocate that plain films be abandoned or that CT is the new "standard of care" for blunt spine trauma. However, this manuscript has delineated certain advantages and disadvantages of various diagnostic techniques, as applied to patients with blunt trauma and possible TL-spine injury.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Cooper C, Dunham CM, Rodriguez A. Falls and major injuries are risk factors for thoracolumbar fractures: cognitive impairment and multiple injuries impede the detection of back pain and tenderness. *J Trauma* 1995;38:692–6. [PubMed: 7760394]
2. Holmes JF, Miller PQ, Panacek EA, Lin S, Horne NS, Mower WR. Epidemiology of thoracolumbar spine injury in blunt trauma. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2001;8:866–72. [PubMed: 11535478]

3. Katsuura Y, Osborn JM, Cason GW. The epidemiology of thoracolumbar trauma: A meta-analysis. *J Orthop* 2016;13:383–8. [PubMed: 27504058]
4. Levi AD, Hurlbert RJ, Anderson P, et al. Neurologic deterioration secondary to unrecognized spinal instability following trauma--a multicenter study. *Spine* 2006;31:451–8. [PubMed: 16481957]
5. O'Connor E, Walsham J. Review article: indications for thoracolumbar imaging in blunt trauma patients: a review of current literature. *Emergency medicine Australasia : EMA* 2009;21:94–101. [PubMed: 19422405]
6. Sixta S, Moore FO, Ditillo MF, et al. Screening for thoracolumbar spinal injuries in blunt trauma: an Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg* 2012;73:S326–32. [PubMed: 23114489]
7. Poonnoose PM, Ravichandran G, McClelland MR. Missed and mismanaged injuries of the spinal cord. *J Trauma* 2002;53:314–20. [PubMed: 12169940]
8. Reid DC, Henderson R, Saboe L, Miller JD. Etiology and clinical course of missed spine fractures. *J Trauma* 1987;27:980–6. [PubMed: 3656481]
9. Advanced trauma life support (ATLS(R)): the ninth edition. *J Trauma Acute Care Surg* 2013;74:1363–6. [PubMed: 23609291]
10. Daffner RH, Weissman BN, Wippold FJ, et al. Expert Panels on Musculoskeletal and Neurologic Imaging. ACR Appropriateness Criteria® suspected spine trauma. American College of Radiology (ACR) 2012.
11. National Clinical Guideline C. National Institute for Health and Care Excellence: Clinical Guidelines Spinal Injury: Assessment and Initial Management. London: National Institute for Health and Care Excellence (UK) Copyright (c) National Clinical Guideline Centre, 2016.; 2016.
12. Korley FK, Pham JC, Kirsch TD. Use of advanced radiology during visits to US emergency departments for injury-related conditions, 1998–2007. *Jama* 2010;304:1465–71. [PubMed: 20924012]
13. Smith-Bindman R, Lipson J, Marcus R, et al. Radiation dose associated with common computed tomography examinations and the associated lifetime attributable risk of cancer. *Archives of internal medicine* 2009;169:2078–86. [PubMed: 20008690]
14. Rodriguez RM, Baumann BM, Raja AS, et al. Diagnostic yields, charges, and radiation dose of chest imaging in blunt trauma evaluations. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2014;21:644–50. [PubMed: 25039548]
15. Kelly ME, Heeney A, Redmond CE, et al. Incidental findings detected on emergency abdominal CT scans: a 1-year review. *Abdominal imaging* 2015;40:1853–7. [PubMed: 25576049]
16. Moore CL, Broder J, Gunn ML, et al. Comparative Effectiveness Research: Alternatives to "Traditional" Computed Tomography Use in the Acute Care Setting. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2015;22:1465–73. [PubMed: 26576033]
17. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *Bmj* 2009;339:b2700. [PubMed: 19622552]
18. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama* 2000;283:2008–12. [PubMed: 10789670]
19. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Annals of internal medicine* 2011;155:529–36. [PubMed: 22007046]
20. Simel DL, Feussner JR, DeLong ER, Matchar DB. Intermediate, indeterminate, and uninterpretable diagnostic test results. *Medical decision making : an international journal of the Society for Medical Decision Making* 1987;7:107–14. [PubMed: 3574020]
21. Zamora J, Abaira V, Muriel A, Khan K, Coomarasamy A. Meta-DiSc: a software for meta-analysis of test accuracy data. *BMC medical research methodology* 2006;6:31. [PubMed: 16836745]
22. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials* 1986;7:177–88. [PubMed: 3802833]

23. DerSimonian R, Kacker R. Random-effects model for meta-analysis of clinical trials: an update. *Contemporary clinical trials* 2007;28:105–14. [PubMed: 16807131]
24. Berry GE, Adams S, Harris MB, et al. Are plain radiographs of the spine necessary during evaluation after blunt trauma? Accuracy of screening torso computed tomography in thoracic/lumbar spine fracture diagnosis. *J Trauma* 2005;59:1410–3; discussion 3. [PubMed: 16394914]
25. Cason B, Rostas J, Simmons J, Frotan MA, Brevard SB, Gonzalez RP. Thoracolumbar spine clearance: Clinical examination for patients with distracting injuries. *J Trauma Acute Care Surg* 2016;80:125–30. [PubMed: 26491795]
26. Chen RJ, Fu CY, Wu SC, et al. Diagnostic accuracy, biohazard safety, and cost effectiveness-the Lodox/Statscan provides a beneficial alternative for the primary evaluation of patients with multiple injuries. *J Trauma* 2010;69:826–30. [PubMed: 20938269]
27. Inaba K, DuBose JJ, Barmparas G, et al. Clinical examination is insufficient to rule out thoracolumbar spine injuries. *J Trauma* 2011;70:174–9. [PubMed: 20489662]
28. Mancini DJ, Burchard KW, Pekala JS. Optimal thoracic and lumbar spine imaging for trauma: are thoracic and lumbar spine reformats always indicated? *J Trauma* 2010;69:119–21. [PubMed: 20622586]
29. Pouw MH, Deunk J, Brink M, et al. Is a pelvic fracture a predictor for thoracolumbar spine fractures after blunt trauma? *J Trauma* 2009;67:1027–32. [PubMed: 19901664]
30. Rozenberg A, Weinstein JC, Flanders AE, Sharma P. Imaging of the thoracic and lumbar spine in a high volume level 1 trauma center: are reformatted images of the spine essential for screening in blunt trauma? *Emergency radiology* 2017;24:55–9. [PubMed: 27663571]
31. Shahrami A, Shojaei M, Tabatabaee SM, Mianehsaz E. Diagnostic Value of Clinical Findings in Evaluation of Thoracolumbar Blunt Traumas. *Emergency (Tehran, Iran)* 2016;4:127–31.
32. Smith CB, Barrett TW, Berger CL, Zhou C, Thurman RJ, Wrenn KD. Prediction of blunt traumatic injury in high-acuity patients: bedside examination vs computed tomography. *The American journal of emergency medicine* 2011;29:1–10. [PubMed: 20825767]
33. Smith MW, Reed JD, Facco R, et al. The reliability of nonreconstructed computerized tomographic scans of the abdomen and pelvis in detecting thoracolumbar spine injuries in blunt trauma patients with altered mental status. *J Bone Joint Surg Am* 2009;91:2342–9. [PubMed: 19797568]
34. Sroka NL, Combs J, Mood R, Henderson V. Scout anteroposterior and lateral CT scans as a screening test for thoracolumbar spine injury in blunt trauma. *The American surgeon* 2007;73:780–5; discussion 5–6. [PubMed: 17879684]
35. Wintermark M, Mouhsine E, Theumann N, et al. Thoracolumbar spine fractures in patients who have sustained severe trauma: depiction with multi-detector row CT. *Radiology* 2003;227:681–9. [PubMed: 12702827]
36. Moon SH, Park MS, Suk KS, et al. Feasibility of ultrasound examination in posterior ligament complex injury of thoracolumbar spine fracture. *Spine* 2002;27:2154–8. [PubMed: 12394931]
37. Roos JE, Hilfiker P, Platz A, et al. MDCT in emergency radiology: is a standardized chest or abdominal protocol sufficient for evaluation of thoracic and lumbar spine trauma? *AJR American journal of roentgenology* 2004;183:959–68. [PubMed: 15385287]
38. Deunk J, Brink M, Dekker HM, et al. Routine versus selective computed tomography of the abdomen, pelvis, and lumbar spine in blunt trauma: a prospective evaluation. *J Trauma* 2009;66:1108–17. [PubMed: 19359922]
39. Dunham CM, Hileman BM, Ransom KJ, Malik RJ. Trauma patient adverse outcomes are independently associated with rib cage fracture burden and severity of lung, head, and abdominal injuries. *Int J Burns Trauma* 2015;5:46–55. [PubMed: 26064801]
40. Herzog C, Ahle H, Mack MG, et al. Traumatic injuries of the pelvis and thoracic and lumbar spine: does thin-slice multidetector-row CT increase diagnostic accuracy? *European radiology* 2004;14:1751–60. [PubMed: 15300395]
41. Stanislas MJ, Latham JM, Porter KM, Alpar EK, Stirling AJ. A high risk group for thoracolumbar fractures. *Injury* 1998;29:15–8. [PubMed: 9659474]
42. Kohn MA, Carpenter CR, Newman TB. Understanding the direction of bias in studies of diagnostic test accuracy. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2013;20:1194–206. [PubMed: 24238322]

43. Samuels LE, Kerstein MD. 'Routine' radiologic evaluation of the thoracolumbar spine in blunt trauma patients: a reappraisal. *J Trauma* 1993;34:85–9. [PubMed: 8437202]
44. Terregino CA, Ross SE, Lipinski MF, Foreman J, Hughes R. Selective indications for thoracic and lumbar radiography in blunt trauma. *Ann Emerg Med* 1995;26:126–9. [PubMed: 7618772]
45. Holmes JF, Panacek EA, Miller PQ, Lapidis AD, Mower WR. Prospective evaluation of criteria for obtaining thoracolumbar radiographs in trauma patients. *The Journal of emergency medicine* 2003;24:1–7. [PubMed: 12554032]
46. Hsu JM, Joseph T, Ellis AM. Thoracolumbar fracture in blunt trauma patients: guidelines for diagnosis and imaging. *Injury* 2003;34:426–33. [PubMed: 12767788]
47. Inaba K, Nosanov L, Menaker J, et al. Prospective derivation of a clinical decision rule for thoracolumbar spine evaluation after blunt trauma: An American Association for the Surgery of Trauma Multi-Institutional Trials Group Study. *J Trauma Acute Care Surg* 2015;78:459–65; discussion 65–7. [PubMed: 25710414]
48. Liberman M, Farooki N, Lavoie A, Mulder DS, Sampalis JS. Clinical evaluation of the spine in the intoxicated blunt trauma patient. *Injury* 2005;36:519–25. [PubMed: 15755434]
49. Durham RM, Luchtfeld WB, Wibbenmeyer L, Maxwell P, Shapiro MJ, Mazuski JE. Evaluation of the thoracic and lumbar spine after blunt trauma. *Am J Surg* 1995;170:681–4; discussion 4–5. [PubMed: 7492026]
50. Sava J, Williams MD, Kennedy S, Wang D. Thoracolumbar fracture in blunt trauma: is clinical exam enough for awake patients? *J Trauma* 2006;61:168–71. [PubMed: 16832266]
51. Terregino CA, Nyce A, Killian AJ, Lipinski MF, Ross SE. Prospective evaluation of a protocol for selective thoracolumbar radiography. *Annals of Emergency Medicine* 1999;34:S45–S6.
52. Bulthuis DB, Vemulapalli P, Nyce AL, Killian AJ, Ross SE, Terregino CA. 2000 SAEM ANNUAL MEETING ABSTRACTS. *Academic Emergency Medicine* 2000;7:425–595.
53. Langdorf MI, Zuabi N, Khan NA, et al. Yield and clinical predictors of thoracic spine injury from chest computed tomography for blunt trauma. *The western journal of emergency medicine* 2014;15:465–70. [PubMed: 25035753]
54. Reynolds SL, Hernandez J, Hogg M, Runyon M. SAEM Annual Meeting Abstracts. *Academic Emergency Medicine* 2014;21:S241–S2.
55. Hanna TN, Zygmunt ME, Harmouche E, Salastekar N, Johnson JO, Khosa F. Association of lumbar fractures, abdominal aortic calcification, and osteopenia. *Clinical imaging* 2015;39:662–6. [PubMed: 25537431]
56. Winslow JE, 3rd, Hensberry R, Bozeman WP, Hill KD, Miller PR. Risk of thoracolumbar fractures doubled in victims of motor vehicle collisions with cervical spine fractures. *J Trauma* 2006;61:686–7. [PubMed: 16967008]
57. Mehta A, Byrne L, Moran V, Armbrrecht E. SAEM Abstracts, Plenary Session. *Academic Emergency Medicine* 2012;19:S4–S393.
58. Szulc P, Samelson EJ, Sornay-Rendu E, Chapurlat R, Kiel DP. Severity of aortic calcification is positively associated with vertebral fracture in older men—a densitometry study in the STRAMBO cohort. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2013;24:1177–84.
59. Hauser CJ, Visvikis G, Hinrichs C, et al. Prospective validation of computed tomographic screening of the thoracolumbar spine in trauma. *J Trauma* 2003;55:228–34; discussion 34–5. [PubMed: 12913630]
60. Mejia VA, Diaz JJJ, Guy J, et al. Plain Films Vs. Helical CT for Thoracolumbar Spine Clearance. *The Journal of Trauma Injury, Infection, and Critical Care* 2005;57:1376.
61. Huddleston J, Davis K, Blankenbaker D, Lee K, Anderson P, Haaland B. Are Radiographs of the Thoracic Spine Truly Inadequate in the Setting of Trauma? Accuracy of Thoracic Radiographs for Diagnosing Injuries of Varying Severity. *American Journal of Roentgenology* 2010;194:A65–A70.
62. Karul M, Bannas P, Schoennagel BP, et al. Fractures of the thoracic spine in patients with minor trauma: comparison of diagnostic accuracy and dose of biplane radiography and MDCT. *European journal of radiology* 2013;82:1273–7. [PubMed: 23422283]

63. Barrios C, Malinoski D, Dolich M, Lekawa M, Hoyt D, Cinat M. Utility of thoracic computed tomography after blunt trauma: when is chest radiograph enough? *The American surgeon* 2009;75:966–9. [PubMed: 19886146]
64. Deyle S, Wagner A, Benneker LM, et al. Could full-body digital X-ray (LODOX-Statscan) screening in trauma challenge conventional radiography? *J Trauma* 2009;66:418–22. [PubMed: 19204516]
65. Whiley SP, Mantokoudis G, Ott D, Zimmerman H, Exadaktylos AK. A review of full-body radiography in nontraumatic emergency medicine. *Emerg Med Int* 2012;2012:108129. [PubMed: 23243508]
66. Kim S, Yoon CS, Ryu JA, et al. A comparison of the diagnostic performances of visceral organ-targeted versus spine-targeted protocols for the evaluation of spinal fractures using sixteen-channel multidetector row computed tomography: is additional spine-targeted computed tomography necessary to evaluate thoracolumbar spinal fractures in blunt trauma victims? *J Trauma* 2010;69:437–46. [PubMed: 20699755]
67. Gestring ML, Gracias VH, Feliciano MA, et al. Evaluation of the lower spine after blunt trauma using abdominal computed tomographic scanning supplemented with lateral scanograms. *J Trauma* 2002;53:9–14. [PubMed: 12131382]
68. Carter B, Griffith B, Mossa-Basha F, et al. Reformatted images of the thoracic and lumbar spine following CT of chest, abdomen, and pelvis in the setting of blunt trauma: are they necessary? *Emergency radiology* 2015;22:373–8. [PubMed: 25666301]
69. Karaca L, Yuceler Z, Kantarci M, et al. The feasibility of dual-energy CT in differentiation of vertebral compression fractures. *The British journal of radiology* 2016;89:20150300. [PubMed: 26537691]
70. Lee JY, Vaccaro AR, Lim MR, et al. Thoracolumbar injury classification and severity score: a new paradigm for the treatment of thoracolumbar spine trauma. *Journal of orthopaedic science : official journal of the Japanese Orthopaedic Association* 2005;10:671–5. [PubMed: 16307197]
71. Vordemvenne T, Hartensuer R, Lohrer L, Vieth V, Fuchs T, Raschke MJ. Is there a way to diagnose spinal instability in acute burst fractures by performing ultrasound? *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society* 2009;18:964–71.
72. Zhao JW, Liu Y, Yin RF, Wang JC, Yang YH, Liu P. Ultrasound assessment of injury to the posterior ligamentous complex in patients with mild thoracolumbar fractures. *The Journal of international medical research* 2013;41:1252–7. [PubMed: 23760913]
73. Haba H, Taneichi H, Kotani Y, et al. Diagnostic accuracy of magnetic resonance imaging for detecting posterior ligamentous complex injury associated with thoracic and lumbar fractures. *Journal of neurosurgery* 2003;99:20–6. [PubMed: 12859054]
74. Ghanem N, Uhl M, Muller C, et al. MRI and discography in traumatic intervertebral disc lesions. *European radiology* 2006;16:2533–41. [PubMed: 16775692]
75. Hunter BRCC. The Development of Clinical Prediction Rules In: Wilson MP GK, Hayden SR (eds), ed. *Doing Research in Emergency and Acute Care: Making Order out of Chaos*: John Wiley & Sons; 2016:139–47.
76. Jaeschke R, Guyatt G, J L. *Diagnostic tests. Users' guides to the medical literature*. Chicago AMA Press; 2002:121–40.
77. Kanzaria HK, McCabe AM, Meisel ZM, et al. Advancing Patient-centered Outcomes in Emergency Diagnostic Imaging: A Research Agenda. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2015;22:1435–46. [PubMed: 26574729]
78. El Dib R, Tikkinen KAO, Akl EA, et al. Systematic survey of randomized trials evaluating the impact of alternative diagnostic strategies on patient-important outcomes. *J Clin Epidemiol* 2017;84:61–9. [PubMed: 28063913]
79. Jawa RS, Singer AJ, Rutigliano DN, et al. Spinal Fractures in Older Adult Patients Admitted After Low-Level Falls: 10-Year Incidence and Outcomes. *Journal of the American Geriatrics Society* 2017;65:909–15. [PubMed: 27910090]

80. Hoffman JR, Carpenter CR. Guarding Against Overtesting, Overdiagnosis, and Overtreatment of Older Adults: Thinking Beyond Imaging and Injuries to Weigh Harms and Benefits. 2017;65:903–5.
81. Arasu VA, Abujudeh HH, Biddinger PD, et al. Diagnostic emergency imaging utilization at an academic trauma center from 1996 to 2012. *Journal of the American College of Radiology : JACR* 2015;12:467–74. [PubMed: 25622765]
82. Broder J, Warshauer DM. Increasing utilization of computed tomography in the adult emergency department, 2000-2005. *Emergency radiology* 2006;13:25–30. [PubMed: 16900352]
83. Oner C, Rajasekaran S, Chapman JR, et al. Spine Trauma-What Are the Current Controversies? *Journal of orthopaedic trauma* 2017;31 Suppl 4:S1–s6.
84. Kepler CK, Vroome C, Goldfarb M, et al. Variation in the management of thoracolumbar trauma and postoperative infection. *Journal of spinal disorders & techniques* 2015;28:E212–8. [PubMed: 25393665]
85. Urquhart JC, Alrehaili OA, Fisher CG, et al. Treatment of thoracolumbar burst fractures: extended follow-up of a randomized clinical trial comparing orthosis versus no orthosis. *Journal of neurosurgery Spine* 2017;27:42–7. [PubMed: 28409669]
86. Bailey CS, Urquhart JC, Dvorak MF, et al. Orthosis versus no orthosis for the treatment of thoracolumbar burst fractures without neurologic injury: a multicenter prospective randomized equivalence trial. *The spine journal : official journal of the North American Spine Society* 2014;14:2557–64. [PubMed: 24184649]
87. Pines JM, Carpenter CR, Raja AS, Schuur JD. Understanding Bias in Diagnostic Research Evidence-Based Emergency Care: John Wiley & Sons, Ltd; 2012:54–64.
88. Goldman LW. Principles of CT: multislice CT. *J Nucl Med Technol* 2008;36:57–68; quiz 75-6. [PubMed: 18483143]
89. Carpenter CR, Meisel ZF. Overcoming the Tower of Babel in Medical Science by Finding the "EQUATOR": Research Reporting Guidelines. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2017.

Article Summary:

1. Why is this topic important? No validated decision aids for determining which blunt trauma patients warrant thoracolumbar-spine (TL-spine) imaging exists. Therefore, a high practice variability and the presence of inconsistent clinical practice guidelines or recommendations are associated with this topic. Since delayed recognition of significant spine injuries risks preventable neurologic complications, while overutilization of diagnostic imaging increases cost, waiting times, and radiation exposure, a quantitative understanding of the diagnostic accuracy of history, physical examination, and commonly ordered imaging tests may improve diagnostic efficiency.
2. What does this review attempt to show? This systematic review and meta-analysis quantifies the diagnostic accuracy of history, physical examination, and diagnostic imaging modalities for blunt trauma-related TL-spine injuries, and to identify any knowledge gaps in the existing research literature.
3. What are the key findings? No injury mechanism, physical examination, or associated injury accurately ruled in the presence of a TL-spine fracture. Computed tomography (CT) is superior to plain film both in the detection of all TL-spine fractures and clinically significant fractures. While reformatted TL-spine CT is more accurate than chest-abdomen-pelvis (CAP) CT, the impact of these differences on management decisions or patient-centric outcomes is unknown.
4. How is patient care impacted? CT is currently the most accurate imaging test to diagnose TL-spine fractures in blunt trauma patients. As compared with other imaging choices, CT misses less TL-spine fractures and reformatted TL-spine CT can be done without any additional cost or radiation to the patient, with added accuracy compared to CAP CT. Whether the additional fractures identified yield improved patient outcomes is uncertain, as are the potential harms of over-diagnosis that might be associated with more widespread CT imaging.

Figure 1:

Three sagittal images of the lumbar-spine in a 28-year-old male involved in a motor vehicle crash. A: Cross-table lateral lumbar-spine radiograph in the Emergency Department at the time of injury demonstrates a fracture of L4 with loss of vertebral height; B: Sagittal reconstruction of the lumbar-spine CT demonstrates more detail, including comminution of the fracture, retropulsion of fragments into the spinal canal, and a three-column injury; C: Six-month follow-up after surgical stabilization and healing, upright lateral lumbar-spine radiograph with maintained alignment and implants.

Figure 2:
Study Selection Process

Figure 3:
Forest Plots of Diagnostic Accuracy of plain films for TL-Spine fracture

Figure 4:
Forest Plots of Diagnostic Accuracy of CT for TL-Spine Fracture

Table 1:**Plain Film Diagnostic Accuracy Study's Index Tests and Criterion Standards**

Study	Index Test	Criterion Standard
Hauser 2003	AP/lateral TL-spine plain film	Reformatted TL-spine CT (with reconstructions of 1-2 mm), and/or subsequent clinical examination when patient fully alert
Mejia 2004	AP/lateral TL-spine plain film	n/a
Berry 2005	AP/lateral TL-spine plain film	All available clinical/radiologic data
Pouw 2009	AP/lateral TL-spine plain film	Reformatted TL-spine CT (16-slice MDCT, with 120 kV tube potential, 200 mAs effective tube current time product, detector configuration of 16 mm × 1.5 mm, reconstructions at 3 mm section thickness with 1.5 mm increments for bone kernel, and sagittal/coronal reformations of the spine)
Smith MW 2009	AP/lateral TL-spine plain film	Reformatted TL-spine CT (either 16-slice MDCT or 64-slice scanner, reconstruction with 2 mm collimation, "thin-slice" in axial/sagittal/coronal planes)

Table 2:**CT Diagnostic Accuracy Study's Index Tests and Criterion Standards**

Study	Index Test	Criterion Standard
Hauser 2003	Helical CAP CT with IV contrast (collimation 5 mm, pitch 1.6, reconstructions with 5 mm image spacing)	TL-spine CT (with reconstructions of 1-2 mm thick slices), and/or subsequent clinical examination when patient fully alert
Mejia 2004	Helical CAP CT	n/a
Berry 2005	Helical CAP CT (pitch 1, reconstructions at 5 mm thick slices)	All available clinical/radiologic data
Smith MW 2009	CAP CT (16-slice MDCT or 64-slice machines, volume acquisition post-processed at axial collimation 5 mm, pitch of 1.375-1.000, "thick-slab" coronal reconstructions)	Reformatted TL-spine CT (either 16-slice MDCT or 64-slice scanner, reconstruction with 2 mm collimation, "thin-slice" in axial/sagittal/coronal planes)
Mancini 2010	CAP CT (16-slice CT machine, reconstructions with 5 mm image spacing)	Reformatted TL-spine CT (16-slice CT machine, reformations with 2.5 mm image spacing, sagittal/coronal planes)
Rozenberg 2017	CAP CT with IV contrast (64-slice CT machine, reconstructions with 3 mm slice thickness)	Consensus amongst imaging modalities (using 64-slice MDCT w/IV contrast, CAP CT with reconstructions at 3 mm slice thickness, and reformatted TL-spine CT with axial slice thickness at 1.25 mm, with coronal/sagittal images reconstructed at 3 mm slice thickness); amongst indeterminate fractures discharge diagnosis of spinal fracture was used (determined by clinical symptoms or MR imaging in certain patients)

Table 3:

Plain Film and CT's Diagnostic Accuracy of Clinically Significant TL-Spine Fractures

Study	SN%	SP%
Plain Film:		
Hauser 2003	100	-
Wintermark 2004	33	100
Berry 2005	92	-
CAP CT:		
Hauser 2003	100	-
Mejia 2004	100	-
Wintermark 2004	97	100
Berry 2005	100	-
Smith MW 2009	100	-
Mancini 2010	100	100
Rozenberg 2017	100	-