Published in final edited form as:

Exp Neurol. 2019 March ; 313: 79–87. doi:10.1016/j.expneurol.2018.12.007.

Broadening the Definition of Brain Insulin Resistance in Aging and Alzheimer's Disease

Hilaree N. Frazier, Adam O. Ghoweri, Katie L. Anderson, Ruei-Lung Lin, Nada M. Porter, and Olivier Thibault

University of Kentucky Department of Pharmacology and Nutritional Sciences, 800 Rose St., Lexington, KY 40536, United States of America

Abstract

It has been more than 20 years since studies first revealed that the brain is insulin sensitive, highlighted by the expression of insulin receptors in neurons and glia, the presence of circulating brain insulin, and even localized insulin production. Following these discoveries, evidence of decreased brain insulin receptor number and function was reported in both clinical samples and animal models of aging and Alzheimer's disease, setting the stage for the hypothesis that neuronal insulin resistance may underlie memory loss in these conditions. The development of therapeutic insulin delivery to the brain using intranasal insulin administration has been shown to improve aspects of memory or learning in both humans and animal models. However, whether this approach functions by compensating for poorly signaling insulin receptors, for reduced insulin levels in the brain, or for reduced trafficking of insulin into the brain remains unclear. Direct measures of insulin's impact on cellular physiology and metabolism in the brain have been sparse in models of Alzheimer's disease, and even fewer studies have analyzed these processes in the aged brain. Nevertheless, recent evidence supports the role of brain insulin as a mediator of glucose metabolism through several means, including altering glucose transporters. Here, we provide a review of contemporary literature on brain insulin resistance, highlight the rationale for improving memory function using intranasal insulin, and describe initial results from experiments using a molecular approach to more directly measure the impact of insulin receptor activation and signaling on glucose uptake in neurons.

Keywords

glucose transporters; intranasal insulin; afterhyperpolarization; calcium; insulin receptor; insulin signaling; brain insulin resistance

*Corresponding author at: Department of Pharmacology and Nutritional Sciences, 800 Rose St., UKMC MS-313, Lexington, KY 40536, United States.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Statement of Competing Interests: The authors declare that no financial or personal competing interests exist.

Introduction

Despite significant knowledge of neuronal energy expenditure in the cortex, including estimates of the metabolic cost for eliciting single action potentials and excitatory post-synaptic potentials, maintaining resting membrane potential, and buffering calcium transients in spines (Attwell and Laughlin, 2001; Howarth et al., 2012), the sensitivity of these dynamic signaling processes to insulin remains unclear. To date, it is still common to read that the brain is insulin independent despite evidence to the contrary. Indeed, many studies support the notion that insulin-dependent processes are present and, in some cases, able to influence memory or recall (Banks et al., 2012; Benedict et al., 2004; Bruning et al., 2000; de la Monte et al., 2006; Dou et al., 2005; Grillo et al., 2009; Kamal et al., 2013; Lannert and Hoyer, 1998; McNay et al., 2010; Pancani et al., 2013; Salameh et al., 2015; Stanley et al., 2016; Wickelgren, 1998; Woods et al., 1979; Zhao et al., 1999). Recently, therapeutic intranasal insulin (INI) administration has been successfully employed to ameliorate several cognitive functions (declarative memory [short-term or delayed], verbal fluency, attention, functional status, spatial memory, and several others). It is not clear, however, whether this approach functions by compensating for poorly signaling insulin receptors (IRs), for reduced insulin levels in the brain, or for reduced trafficking of insulin into the brain. Potential pathways mediating the benefits of INI include increases in cerebral blood flow (CBF), reductions in inflammation, and elevations in glucose metabolism. Together, the current body of literature supports the theory of an insulin-sensitive brain and highlights the physiological importance of maintained insulin signaling, yet the exact mechanisms involved in this process appear to be highly complex and variable.

In aging and Alzheimer's disease (AD), conditions where both glucose utilization and memory recall are known to decline, it is tempting to infer causality between these two processes. However, it is only recently that region-specific brain glucose concentrations were measured to test for associations between glucose metabolism and severity of AD pathology (An et al., 2018). Interestingly, this study found that reductions in glucose metabolism were associated with elevated glucose levels and reduced numbers of glucose transporter (GLUT) type 3 in the brains of AD patients compared to aged-match controls. On initial assessment, elevated brain glucose levels together with cognitive decline may appear contradictory; however, it is possible that the increased glucose in the brains of AD patients may reflect reduced glucose utilization through either dysfunctional IR (Frolich et al., 1998), reduced GLUTs at the cell membrane (An et al., 2018), or diminished insulin levels in the central nervous system (CNS) (Craft et al., 1998; Frolich et al., 1998; Molina et al., 2002), all of which may lead to reduced insulin sensitivity.

Here, we review the current evidence supporting the theory that the brain is insulin-sensitive and address potential mechanisms underlying the age- and AD-related decrease in insulin and IR signaling, a phenomenon often referred to as brain insulin resistance. Additionally, we discuss cellular components that may participate in this process. Finally, we introduce a molecular technique designed to directly investigate the relationship between IR activity and glucose metabolism in neurons and present preliminary data supporting this as a viable technique.

What elements and processes inform insulin resistance in the brain?

While it is becoming clear that insulin and IR signaling play an important role in overall brain health and contribute to numerous physiological functions, the question as to how insulin and IR dysfunction lead to reductions in memory remains unanswered. A PubMed query with the terms “insulin resistance,” “brain,” and either “aged” or “Alzheimer’s disease” shows a steady increase in articles published between the years 1998 and 2018, highlighting the growing scientific interest in this topic (Figure 1). One relevant concept based on a broad phenotype characterized by reduced brain insulin levels, IR signaling, and activity of insulin-related processes, is the brain insulin resistance hypothesis (Hoyer, 1998). This hypothesis was formulated following observations of insulin resistance in AD (Frolich et al., 1998). This work showed that compared to age-matched control brains, AD brains displayed region-specific reductions in insulin concentrations together with elevations in IR densities (Frolich et al., 1998). While the elevation in IR density was surprising, this result has since been interpreted as a failed compensation, given the evidence of reduced brain glucose utilization in AD (An et al., 2018; Hoyer, 1986, 2004; Mielke et al., 1992; Mosconi et al., 2008a; Mosconi et al., 2004; Rivera et al., 2005). Together, however, these initial observations, along with critical work from other labs, secured insulin resistance as a potential therapeutic target in the brains of mild cognitive impairment (MCI), AD, and perhaps even Type-2 diabetes mellitus (T2DM) patients.

Evidence for insulin resistance in the brain has been mostly derived from data using molecular approaches to quantify levels of accessory proteins in the insulin signaling pathway, from IR itself to targets further downstream in the signaling pathway, including insulin receptor substrate-1 (IRS-1) and the resulting transport of GLUTs to the plasma membrane (Barone et al., 2018; Bell and Fadool, 2017; Benedict and Grillo, 2018; Biessels and Reagan, 2015; Bomfim et al., 2012; De Felice et al., 2014; de la Monte and Wands, 2008; Grillo et al., 2015; Mao et al., 2016; Pearson-Leary and McNay, 2012; Schubert et al., 2004; Talbot et al., 2012; van der Heide et al., 2006; Willette et al., 2013; Zhao et al., 2008). However, these links are almost always based on associations, as it is difficult to directly address mechanisms and causality using CNS exposure to the ligand in clinical samples. Further, the nature of these techniques often limits experiments to transient signaling snapshots, thus preventing investigations of the physiological impact of insulin across time. Other approaches that do not introduce the ligand to the brain, but instead use molecular techniques to knockout the IR (Bruning et al., 2000; Schubert et al., 2004) or introduce an IR antisense sequence (LV-IRAS) via a lentiviral delivery system (Grillo et al., 2015), have been very informative in defining the role of IR and downstream signaling in the CNS. Similarly, introducing an IR binding peptide (i.e., affibody) has been employed to reduce insulin actions at the IR (Luckett et al., 2013; Maimaiti et al., 2017; Paranjape et al., 2010). Yet, these loss-of-function approaches only report on the impact of reducing IR activity in the brain by attempting to mimic conditions of reduced brain insulin levels or IR function, as seen in aging or AD.

Furthermore, a large body of evidence supports the hypothesis that peripheral metabolic dysregulation contributes to cognitive decline in T2DM and obesity, suggesting the presence of parallel mechanisms between neuronal and chronic peripheral insulin resistance (Arnold

et al., 2018; Benedict and Grillo, 2018; Biessels and Reagan, 2015; Cholerton et al., 2011; Craft, 2005; De Felice, 2013; de la Monte, 2014; Fadel and Reagan, 2016; Gispen and Biessels, 2000; Luchsinger, 2008; Luchsinger et al., 2004). With respect to peripheral insulin levels and their potential impact on cognitive function, well-powered studies have provided evidence that elevated peripheral insulin levels are associated with poorer cognitive function in elderly patients with or without hypertension or diabetes, perhaps even more so in females than males (Kuusisto et al., 1993; Stolk et al., 1997). Indeed, higher *chronic* insulin levels, peripheral insulin resistance, and a pre-diabetic state (e.g., greater insulin area-under-the-curve (AUC) in response to an acute intravenous glucose tolerance test (GTT)) are associated with greater decline in performance over a 2-year period in healthy elderly subjects (Burns et al., 2012). Interestingly, the same group also showed that in AD patients, increased insulin AUC during GTT was associated with less brain atrophy and lower dementia severity across a similar time period (Burns et al., 2007; Burns et al., 2012). These later results align well with early work from Craft and others showing that *acute* elevations in peripheral insulin levels along with maintained euglycemia enhances memory recall in AD patients (Craft et al., 1996; Craft and Watson, 2004; Kern et al., 1999; Long et al., 1992). It remains to be determined whether the duration of elevated peripheral insulin is important in the development of brain insulin resistance. While work from several labs has demonstrated a form of reduced brain insulin sensitivity, peripheral insulin resistance status was not always provided; thus, it is difficult to directly test for associations between the periphery and the brain in these samples (Moloney et al., 2010; Steen et al., 2005; Talbot et al., 2012; Tramutola et al., 2017). Nevertheless, initial assessment suggests that elevations in peripheral insulin, whether chronic or acute, might be beneficial in individuals with AD.

More investigations are needed to determine if, and/or how closely, central insulin resistance and associated molecular processes mirror peripheral elements of insulin resistance in muscle or fat cells. However, at least one study using magnetoencephalography has characterized insulin-mediated changes in cortical activity in lean and obese subjects and shows reductions in obese humans, likely through alterations in IRS-1 signaling (Tschritter et al., 2006). Interestingly, these authors also conclude that transport of insulin across the blood-brain barrier (BBB) could have an impact on cortical insulin sensitivity. Several other labs have shown that chronic hyperinsulinemia or insulin resistance may reduce insulin transport into the CNS (reviewed in Banks et al., 2012; Baura et al., 1996; reviewed in Craft, 2005; Heni et al., 2014; Israel et al., 1993; Schwartz et al., 1990), and have even raised the concept of cell-selective insulin resistance in the brain (Vogt and Bruning, 2013). Together, these results suggest that brain insulin resistance can take place through mechanisms that may depend on insulin transport, insulin levels, IRS-1 pathway sensitivity, and other yet unknown processes.

Evidence for preserved insulin sensitivity in the aged and AD brain

While the bulk of the literature supports the hypothesis that overall brain insulin signaling is reduced in aging, AD and T2DM, work from several labs has provided evidence that insulin *sensitivity* may actually be preserved. An early study measuring insulin binding across multiple brain regions in aged rats found no reduction in specific binding in the frontal cortex, hippocampus, or hypothalamus compared to young adult animals (Tchilian et al.,

1990). A more recent report showed that in both younger and older APP/PS1 mice, insulin could still signal normally, provided it was delivered directly to the brain (Stanley et al., 2016). Using electrophysiological analyses of calcium-dependent processes, our lab showed that neurons of mid-age and aged rats responded robustly to exogenous insulin following direct application of the ligand to hippocampal slices (Figure 2 and Figure 3). Of note, the insulin response, which reduces the calcium-dependent afterhyperpolarization (AHP), appears greater in aged compared to young animals (Maimaiti et al., 2016; Pancani et al., 2013). Our group has also reported that measures of CBF in aged animals respond more significantly to direct administration of exogenous insulin in the brain compared to younger animals (Anderson et al., 2017). Together, these results suggest that a deficiency in insulin availability might govern some parts of brain insulin resistance.

Indeed, transport of insulin across the BBB has been implicated as a potential contributor to CNS insulin resistance. In a recent study, measures of brain IR signaling in response to subcutaneous insulin administration showed reductions in aged mice; yet, in these same animals, a robust response was detected provided insulin was delivered directly to the brain (Sartorius et al., 2015). Clinical studies corroborate these findings, with the same group of investigators also showing decreased insulin transport in the brains of human subjects with impaired peripheral glucose tolerance (Heni et al., 2014). This series of studies suggests insulin resistance in aging or AD may be mediated by mechanisms independent of decreased IR sensitivity. Indeed, previous work from the Banks lab has proposed that entry of insulin into the brain is a likely component of CNS insulin resistance associated with aging, AD, or T2DM (Banks et al., 2012). Note that a similar reduction in transport has also been seen in the context of leptin resistance in obese humans and rodents (Banks, 2001).

Irrespective of the mechanism mediating insulin resistance, the evidence for the reduction in insulin transport suggests the aged brain is starved of insulin. Additionally, these results clearly point to the importance of ligand transport into the CNS, and also bolster the rationale for the work of Craft and others, who demonstrated that INI administration may overcome the chronic hyperinsulinemia-induced downregulation of insulin transport at the BBB by raising the ligand concentration in the brain (reviewed in Banks et al., 2012; Baura et al., 1996; reviewed in Craft, 2005; Heni et al., 2014; Israel et al., 1993; Schwartz et al., 1990). It is likely that brain insulin resistance manifests through a combination of altered pathways, from IR signaling to the presence and transport of the ligand into the brain, and it is therefore not surprising that bypassing these reduced elements is a therapeutically relevant and often beneficial approach to offset memory decline in aging and AD.

Combating brain insulin resistance using INI

Early work using intracerebroventricular (ICV) insulin administration in baboons, dogs, rats, and mice framed a series of experiments to introduce the ligand into the brain. Since then, alternative approaches, including INI delivery, have continued to be investigated in different animal models and have moved the field forward with much success in clinical settings. Indeed, initial work (Benedict et al., 2004; Kern et al., 1999; Reger et al., 2006) provided some evidence that INI could alter cerebral function, and in some cases, improve word recall in younger male subjects or early apolipoprotein E (ApoE)- ϵ 4 positive AD patients. Since

then, a significant number of studies have corroborated these findings (Benedict and Grillo, 2018; Chapman et al., 2017; de la Monte, 2013; Freiherr et al., 2013; Hughes and Craft, 2015) and highlighted potential mechanisms governing these beneficial effects, including alterations in CBF flow (Akintola et al., 2017; Kullmann et al., 2017) and circulating β -amyloid (Reger et al., 2008), as well as changes in peripheral corticosteroid concentrations (Thienel et al., 2017).

To date, the effects of INI have only been investigated in relatively small, albeit numerous, trials and for limited periods of time. Only one study has reported that INI could rapidly raise insulin concentrations in the brain (Born et al., 2002). This study was conducted in healthy adults and similar measures have not yet been obtained in cognitively impaired individuals. Nevertheless, as these trials continue, notable sex differences are beginning to emerge, further highlighting the complexity of this therapeutic approach (Benedict et al., 2008; Claxton et al., 2013; Hallschmid et al., 2004; Kullmann et al., 2013). With respect to the impact on feeding behavior, women appeared to gain more weight during an 8-week trial compared to men (Hallschmid et al., 2004). This result was corroborated in a later study (Benedict et al., 2008). In another report, this time focusing on short-term memory function in AD patients, sex differences were only apparent in ApoE negative individuals (Claxton et al., 2013). Furthermore, the authors provided evidence that the beneficial effects of INI were dependent on at least three factors, including sex, ApoE genotype, and dose. Finally, it is important to consider the potential impact of the different formulations of insulin, the duration of treatment, or the regimen of the interventions (reviewed in Arnold et al., 2018).

With respect to the use of INI in animal models of AD, Banks and colleagues have shown that the presence of the ligand in the brain following coadministration of INI with serum albumin is independent of age and phenotype in the SAMP8 mouse model of AD, but did not investigate downstream consequences of this application (Rhea et al., 2017). A 2014 study in female 3xTG mice exposed to INI for 7 days showed reductions in amyloid beta ($A\beta$) levels, restoration of insulin signaling, and enhancement of synaptic proteins (synapsin-1 and PSD-95) (Chen et al., 2014). A more recent report from the same group, this time conducted in female APPswe/PS1dE9 mice exposed to INI for 6 weeks, revealed a significant improvement in cognitive performance, neurogenesis, and insulin signaling, as well as reductions in $A\beta$ production and amyloid plaque burden (Mao et al., 2016). The impact of INI in the 3xTg animal model again revealed an improvement in spatial memory recall, as well as increased insulin signaling (Barone et al., 2018). The authors also suggested potential mechanisms mediating this increase, specifically biliverdin reductase A enhancement, reductions in Tau phosphorylation, and increased serine phosphorylation of IRS-1.

In wildtype mice (C57Bl/6) fed a control diet, short-term INI exposure (5–7 days) was able to improve object recognition memory and olfactory discrimination while reducing anxiety and increasing the phosphorylation state of a potassium channel (Kv1.3). However, compared to older mice fed a control diet, older mice made prediabetic using a high fat diet failed to show increases in object memory following INI administration (Marks et al., 2009). This same group recently published evidence that longer INI treatment (~2 months) did not improve object recognition or olfactory memory in adult male mice. The authors attribute

these differences to the onset of a brain insulin-resistant state (Bell and Fadool, 2017). Similarly, we have also witnessed that long-term INI (~3 months) has minimal impact on spatial learning or memory performance in younger or older F344 rats (in preparation). In another study, we reported a positive impact of two formulations of insulin on spatial memory performance following intranasal delivery in aged F344 rats compared to young (Maimaiti et al., 2016). Furthermore, we have also recently shown that INI increases CBF (Anderson et al., 2017). The question still remains as to which cognitive processes and cellular functions are directly mediated by insulin signaling and targeted by therapeutic INI. One potential answer is likely to involve elements of glucose metabolism in the brain.

The impact of brain insulin on glucose utilization in humans

With respect to brain glucose metabolism in the context of insulin resistance, a significant number of advances have been made through measures of glucose utilization *in vivo*. 2[18F]fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET) imaging has been used to infer rates of brain glucose metabolism in patients expressing a wide range of disease phenotypes, including prediabetes and T2DM, MCI, AD, non-AD dementia types, and cognitively normal, as well as across age. An FDG-PET study comparing young and aged individuals showed that aging in the absence of cognitive impairment is associated with reduced CNS aerobic glycolysis with no apparent change in glucose uptake (Goyal et al., 2017). It is important to note that in this study, age-related reductions in overall brain glucose metabolism are observed only when measures of glucose uptake are combined with oxygen utilization rates, suggesting that cellular glucose utilization and metabolism could be modulated by processes independent of uptake alone. In another FDG-PET study of 548 individuals displaying a spectrum of cognitive impairment, patterns of hypometabolism were seen across various brain regions compared to age-matched controls (Mosconi et al., 2008b). Additional work in AD patients has emphasized the potential use of FDG-PET to predict the progression of AD severity, further refine the initial clinical diagnosis of AD, and even identify individuals at risk for developing the disease (Mosconi, 2013; Swerdlow, 2007).

Using similar techniques, Craft and colleagues showed that aged, cognitively normal individuals with a higher level of peripheral insulin resistance (i.e. prediabetes or T2DM) had a marked reduction in cerebral glucose metabolic rate across several brain regions compared to age-matched controls under baseline conditions (Baker et al., 2011). However, in a study using a euglycemic-hyperinsulinemic clamp to induce a transient increase in peripheral insulin, this time performed in middle-age, cognitively intact individuals, those with impaired peripheral glucose tolerance showed increased, not decreased, brain glucose utilization during peripheral hyperinsulinemia (Hirvonen et al., 2011). As FDG-PET signals in the brains of subjects with impaired peripheral glucose tolerance were found to respond more robustly to the hyperinsulinemic condition compared to healthy controls, these results suggest that peripheral insulin resistance does not necessarily translate to reductions in IR sensitivity. While contradictory, the results presented by Baker and Hirvonen nevertheless suggest that peripheral insulin is capable of influencing central glucose utilization, albeit differently across age and phenotype. Finally, a small study of 8 healthy, middle-aged volunteers revealed that decreasing basal peripheral insulin levels significantly reduced FDG-PET measures of global brain glucose metabolism (Bingham et al., 2002). This same

study also showed that a lowdose insulin infusion in the periphery restored both glucose uptake and rate of utilization in the brain.

Clearly, brain glucose metabolism has an important role in the development of cognitive impairment and is associated with CNS hypometabolism in several pathological conditions. Importantly, it is also highly sensitive to both exogenous and peripheral insulin. However, clinical studies are limited in their ability to characterize the cellular mechanisms targeted by the ligand. In the next section, we provide a review of techniques used in animal models to characterize the potential cellular targets, receptors, and pathways that mediate this process.

Insulin and glucose in animal models of aging and AD

Work performed in animal models of traumatic brain injury (TBI) and diabetes suggests a role of insulin in central glucose metabolism similar to that seen in the clinic. One manuscript reported a decrease in brain glucose metabolism measured by FDG-PET following TBI (Brabazon et al., 2017). In this same study, daily INI administration to adult male rats for 14 days post-TBI lead to increased glucose uptake in the hippocampus, reduced hippocampal lesion volume and microglial reactivity, and even significantly improved memory and motor function. A similar report recently highlighted INI-mediated increases in regional cerebral glucose metabolism in an ICV streptozotocin-induced model of T2DM (Chen et al., 2018). The processes underlying alterations in CNS glucose metabolism, as well as the physiological impact of insulin in the brain, depend, at least in part, on the availability of the ligands, the presence of functional receptors and their respective signaling cascades, and the transport of the ligands into the brain. Once fully characterized, these mechanisms may help provide a more complete definition of central insulin resistance. One such mechanism that has been studied extensively at the intersection of glucose metabolism and insulin resistance is the transport of glucose via GLUTs at the plasma membrane.

GLUT3 has long been established as the primary neuronal GLUT (Vannucci et al., 1997). Other GLUTs, such as GLUT1, GLUT2, and the insulin-sensitive GLUT4, have also been identified in various parts of the brain (Choeiri et al., 2002; Kobayashi et al., 1996; Leino et al., 1997; McEwen and Reagan, 2004; Pearson-Leary et al., 2017; Reagan, 2005). An early study found that in the rat brain, GLUT4 was primarily expressed in areas that also possess high levels of IR, suggesting a potential role of circulating brain insulin and IR signaling in the regulation of neuronal GLUT4 function (El Messari et al., 1998). This same study found that regions with the highest level of GLUT4 expression corresponded with areas primarily involved in motor function. However, while a more recent study corroborated these initial findings, these investigators also discovered novel locations for GLUT4-expression in different subfields of the hippocampus (Choeiri et al., 2002). As the hippocampus expresses high levels of IR (Zhao et al., 1999), the localization of GLUT4 in this structure further supports the hypothesis that insulin signaling mediates brain glucose metabolism, and that this relationship may play a role in cognitive function.

Evidence for a reduction in brain glucose utilization with age, together with evidence for decreased IR numbers or function in animal model of aging (Zaia and Piantanelli, 2000; Zhao et al., 2004), strengthens the notion that these altered processes may be somehow

connected. It has been reported that the permissive action of insulin during a spatial alternation task requires GLUT4 translocation and signaling, and that insulin increases glucose utilization in neurons via a GLUT4 mechanism (Grillo et al., 2009; McEwen and Reagan, 2004; PearsonLeary et al., 2018; Reagan, 2005). Unfortunately, very few studies have directly investigated the impact of insulin on glucose utilization in the brain of aged animals, although evidence is present that aged F344 rats running a cognitively demanding task display significantly reduced hippocampal glucose levels (McNay and Gold, 2001). Considering this, the authors have argued that the enhanced depletion of glucose during demanding tasks may be due to reduced glucose supply (McNay, 2005). Importantly, McNay and colleagues have since reported that acute delivery of insulin to the hippocampus 10 min prior to a cognitive task enhances performance, increases glucose uptake and glycolytic metabolism, and reduces extracellular glucose concentration (McNay et al., 2010), again providing evidence that the hippocampus, at least, is highly sensitive to insulin.

In animal models of AD, marked decreases in brain insulin signaling and sensitivity, together with deteriorations in cognitive or spatial memory performance, have been detected (reviewed in Griffith et al., 2018; Lee et al., 2018). Studies in multiple mouse models have characterized the impact of peripheral and central metabolic dysregulation with respect to the onset of AD pathology and cognitive decline (Ho et al., 2004; Macklin et al., 2017; Rodriguez-Rivera et al., 2011; Velazquez et al., 2017). Further evidence that glucose- and insulin-dependent cellular processes are dysregulated in other AD models have also been reported, particularly in relation to peripheral and central metabolism or the use of high fat diets or insulin-sensitizers (Lee et al., 2018). It is interesting to note that one study in 3xTG mice exposed to INI for 7 days did not show alterations in GLUT levels (Chen et al., 2014). While this has helped further solidify associations between peripheral insulin sensitivity and the AD phenotype, the direct connection between insulin and glucose in the brain of these models is better addressed by using exogenous insulin delivery methods. Together, this body of evidence strongly supports the proposal that utilizing insulin as means to improve glucose availability in neuronal cells could be a viable therapeutic approach. Specifically, the current literature highlights the need for continued explorations into novel therapeutic strategies to increase insulin signaling by either elevating ligand availability (i.e., INI), increasing IR trafficking to the plasma membrane, or even using molecular techniques to increase IR signaling in neurons.

Raising insulin signaling and glucose uptake in neurons through a molecular approach

Recently, we have designed a constitutively active form of the human IR comprised of an intact catalytic β -subunit and a heavily truncated α -subunit, referred to as IR β (Frazier et al., 2018; Lebwohl et al., 1991), to directly test the impact of elevated IR signaling on neuronal metabolism. A lentiviral delivery system was used to express this IR β construct in mixed primary hippocampal cultures and to test for alterations in voltage-gated calcium currents. Our first study showed no change in channel activity but did reveal robust expression and constitutive activity of the receptor (Frazier et al., 2018). To measure glucose uptake, we loaded hippocampal neurons with 2-(N-(7-nitrobenz-2-oxa-1,3-diazol-4-

yl)amino)-2deoxyglucose (2-NBDG), a fluorescent glucose analogue. This analogue is metabolized to a non-fluorescent derivative at a rate shown to be dependent on glycolysis, therefore providing an indirect measure of glucose metabolism (Pancani et al., 2011). For this reason, 2-NBDG provides a powerful tool to observe not only measures of initial glucose uptake, but also the rates of glucose utilization.

Preliminary data show a robust and statistically significant increase in 2-NBDG uptake in IR β -expressing cells compared to cells transfected with the empty vector control (Figure 4). In addition to elevated uptake, it also appears that measures of glucose utilization rates are slightly increased following IR β -expression (data not shown). If 2-NBDG uptake and utilization are elevated following sustained IR signaling through this molecular approach, this may provide a new tool to combat reductions in glucose uptake in the brain of aged animals. Ongoing studies in our lab are addressing which GLUT isoform is responsible for this increased glucose uptake. Studies on GLUT trafficking recently showed that treatment with insulin induced GLUT4 translocation to the plasma membrane in both a human neuronal cell line (Benomar et al., 2006) and in the rat hippocampus (Grillo et al., 2009) through a phosphatidylinositol 3-kinase-dependent pathway. Additionally, some work suggests that translocation of GLUT3, largely considered to be an insulin-insensitive isoform, may also be mediated through insulin (Uemura and Greenlee, 2006). Given the importance of neuronal glucose utilization and the impaired glycolysis as fundamental features of AD, it is imperative to identify which isoform might represent key features of the disease (An et al., 2018). Overall, the current body of evidence strongly supports the proposal that utilizing molecular strategies (i.e., adeno-associated viral delivery of IR β) to improve glucose availability in neurons could lead to the identification of disease-modifying interventions.

Conclusion

Although not all studies align with the notion that insulin insensitivity is present in the aged or AD brain, the bulk of the current literature does indeed support the proposal that dysfunctional IR may be responsible, at least in part, for the memory decline associated with these disease states. Events that help give rise to brain insulin resistance include a reduction in IR signaling and/or a reduction in the ligand or its transport at the BBB. In this review, we attempted to highlight alternative sources of insulin dysregulation that could mediate the well-characterized reduction in glucose utilization seen in aging or AD and presented evidence for new approaches, from INI to IR β , aimed at offsetting this decline. Future work is needed to continue investigations using these two approaches as well as research into other, less characterized cellular processes that could also be involved in insulin-mediated brain dysfunction.

Acknowledgments

Funding: This work was supported by the National Institutes of Health [R01AG033649, T32DK007778, T32AG057461]; the University of Kentucky College of Medicine [Fellowship for Excellence in Graduate Research awarded to H. N. Frazier]; and the University of Kentucky Department of Pharmacology and Nutritional Sciences [Department Reinvestment Fund Award].

Abbreviations:

AD	Alzheimer's disease
IR	insulin receptor
GLUT	glucose transporter
INI	intranasal insulin
MCI	mild cognitive impairment
T2DM	Type-2 diabetes mellitus
AUC	area-under-the-curve
GTT	glucose tolerance test
IRS1	insulin receptor substrate-1
CNS	central nervous system
APP	amyloid precursor protein
PS1	presenilin 1
AHP	afterhyperpolarization
CBF	cerebral blood flow
ICV	intracerebroventricular
BBB	bloodbrain barrier
ApoE	apolipoprotein E
Aβ	amyloid beta
2-NBDG	2-[N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino]-2-deoxyglucose
TBI	traumatic brain injury
FDG-PET	2-[18F]fluoro-2-deoxy-D-glucose positron emission tomography

References

- Akintola AA, van Opstal AM, Westendorp RG, Postmus I, van der Grond J, van Heemst D, 2017 Effect of intranasally administered insulin on cerebral blood flow and perfusion; a randomized experiment in young and older adults. *Aging (Albany NY)* 9, 790–802. [PubMed: 28291957]
- An Y, Varma VR, Varma S, Casanova R, Dammer E, Pletnikova O, Chia CW, Egan JM, Ferrucci L, Troncoso J, Levey AI, Lah J, Seyfried NT, Legido-Quigley C, O'Brien R, Thambisetty M, 2018 Evidence for brain glucose dysregulation in Alzheimer's disease. *Alzheimers Dement* 14, 318–329. [PubMed: 29055815]
- Anderson KL, Frazier HN, Maimaiti S, Bakshi VV, Majeed ZR, Brewer LD, Porter NM, Lin AL, Thibault O, 2017 Impact of Single or Repeated Dose Intranasal Zinc-free Insulin in Young and

- Aged F344 Rats on Cognition, Signaling, and Brain Metabolism. *J Gerontol A Biol Sci Med Sci* 72, 189–197. [PubMed: 27069097]
- Arnold SE, Arvanitakis Z, Macauley-Rambach SL, Koenig AM, Wang HY, Ahima RS, Craft S, Gandy S, Buettner C, Stoekel LE, Holtzman DM, Nathan DM, 2018 Brain insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums. *Nat Rev Neurol* 14, 168–181. [PubMed: 29377010]
- Attwell D, Laughlin SB, 2001 An energy budget for signaling in the grey matter of the brain. *J Cereb Blood Flow Metab* 21, 1133–1145. [PubMed: 11598490]
- Baker LD, Cross DJ, Minoshima S, Belongia D, Watson GS, Craft S, 2011 Insulin resistance and Alzheimer-like reductions in regional cerebral glucose metabolism for cognitively normal adults with prediabetes or early type 2 diabetes. *Arch Neurol* 68, 51–57. [PubMed: 20837822]
- Banks WA, 2001 Leptin transport across the blood-brain barrier: implications for the cause and treatment of obesity. *Curr Pharm Des* 7, 125–133. [PubMed: 11172704]
- Banks WA, Owen JB, Erickson MA, 2012 Insulin in the brain: there and back again. *Pharmacol Ther* 136, 82–93. [PubMed: 22820012]
- Barone E, Tramutola A, Triani F, Calcagnini S, Di Domenico F, Ripoli C, Gaetani S, Grassi C, Butterfield DA, Cassano T, Perluigi M, 2018 Biliverdin Reductase-A Mediates the Beneficial Effects of Intranasal Insulin in Alzheimer Disease. *Mol Neurobiol*.
- Baura GD, Foster DM, Kaiyala K, Porte D, Jr., Kahn SE, Schwartz MW, 1996 Insulin transport from plasma into the central nervous system is inhibited by dexamethasone in dogs. *Diabetes* 45, 86–90. [PubMed: 8522065]
- Bell GA, Fadool DA, 2017 Awake, long-term intranasal insulin treatment does not affect object memory, odor discrimination, or reversal learning in mice. *Physiol Behav* 174, 104–113. [PubMed: 28259806]
- Benedict C, Grillo CA, 2018 Insulin Resistance as a Therapeutic Target in the Treatment of Alzheimer's Disease: A State-of-the-Art Review. *Front Neurosci* 12, 215. [PubMed: 29743868]
- Benedict C, Hallschmid M, Hatke A, Schultes B, Fehm HL, Born J, Kern W, 2004 Intranasal insulin improves memory in humans. *Psychoneuroendocrinology* 29, 1326–1334. [PubMed: 15288712]
- Benedict C, Kern W, Schultes B, Born J, Hallschmid M, 2008 Differential sensitivity of men and women to anorexigenic and memory-improving effects of intranasal insulin. *J Clin Endocrinol Metab* 93, 1339–1344. [PubMed: 18230654]
- Benomar Y, Naour N, Aubourg A, Bailleux V, Gertler A, Djiane J, Guerre-Millo M, Taouis M, 2006 Insulin and leptin induce Glut4 plasma membrane translocation and glucose uptake in a human neuronal cell line by a phosphatidylinositol 3-kinase-dependent mechanism. *Endocrinology* 147, 2550–2556. [PubMed: 16497805]
- Biessels GJ, Reagan LP, 2015 Hippocampal insulin resistance and cognitive dysfunction. *Nat Rev Neurosci* 16, 660–671. [PubMed: 26462756]
- Bingham EM, Hopkins D, Smith D, Pernet A, Hallett W, Reed L, Marsden PK, Amiel SA, 2002 The role of insulin in human brain glucose metabolism: an 18fluoro-deoxyglucose positron emission tomography study. *Diabetes* 51, 3384–3390. [PubMed: 12453890]
- Bomfim TR, Forny-Germano L, Sathler LB, Brito-Moreira J, Houzel JC, Decker H, Silverman MA, Kazi H, Melo HM, McClean PL, Holscher C, Arnold SE, Talbot K, Klein WL, Munoz DP, Ferreira ST, De Felice FG, 2012 An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer's disease-associated Aβ oligomers. *J Clin Invest* 122, 1339–1353. [PubMed: 22476196]
- Born J, Lange T, Kern W, McGregor GP, Bickel U, Fehm HL, 2002 Sniffing neuropeptides: a transnasal approach to the human brain. *Nat Neurosci* 5, 514–516. [PubMed: 11992114]
- Brabazon F, Wilson CM, Jaiswal S, Reed J, Frey WHN, Byrnes KR, 2017 Intranasal insulin treatment of an experimental model of moderate traumatic brain injury. *J Cereb Blood Flow Metab* 37, 3203–3218. [PubMed: 28058996]
- Bruning JC, Gautam D, Burks DJ, Gillette J, Schubert M, Orban PC, Klein R, Krone W, Muller-Wieland D, Kahn CR, 2000 Role of brain insulin receptor in control of body weight and reproduction. *Science* 289, 2122–2125. [PubMed: 11000114]

- Burns JM, Donnelly JE, Anderson HS, Mayo MS, Spencer-Gardner L, Thomas G, Cronk BB, Haddad Z, Klima D, Hansen D, Brooks WM, 2007 Peripheral insulin and brain structure in early Alzheimer disease. *Neurology* 69, 1094–1104. [PubMed: 17846409]
- Burns JM, Honea RA, Vidoni ED, Hutfles LJ, Brooks WM, Swerdlow RH, 2012 Insulin is differentially related to cognitive decline and atrophy in Alzheimer's disease and aging. *Biochim Biophys Acta* 1822, 333–339. [PubMed: 21745566]
- Chapman CD, Schioth HB, Grillo CA, Benedict C, 2017 Intranasal insulin in Alzheimer's disease: Food for thought. *Neuropharmacology*.
- Chen Y, Guo Z, Mao YF, Zheng T, Zhang B, 2018 Intranasal Insulin Ameliorates Cerebral Hypometabolism, Neuronal Loss, and Astroglialosis in Streptozotocin-Induced Alzheimer's Rat Model. *Neurotox Res* 33, 716–724. [PubMed: 28929339]
- Chen Y, Zhao Y, Dai CL, Liang Z, Run X, Iqbal K, Liu F, Gong CX, 2014 Intranasal insulin restores insulin signaling, increases synaptic proteins, and reduces Abeta level and microglia activation in the brains of 3xTg-AD mice. *Exp Neurol* 261, 610–619. [PubMed: 24918340]
- Choeiri C, Staines W, Messier C, 2002 Immunohistochemical localization and quantification of glucose transporters in the mouse brain. *Neuroscience* 111, 19–34. [PubMed: 11955709]
- Cholerton B, Baker LD, Craft S, 2011 Insulin resistance and pathological brain ageing. *Diabet Med* 28, 1463–1475. [PubMed: 21974744]
- Claxton A, Baker LD, Wilkinson CW, Trittschuh EH, Chapman D, Watson GS, Cholerton B, Plymate SR, Arbuckle M, Craft S, 2013 Sex and ApoE genotype differences in treatment response to two doses of intranasal insulin in adults with mild cognitive impairment or Alzheimer's disease. *J Alzheimers Dis* 35, 789–797. [PubMed: 23507773]
- Craft S, 2005 Insulin resistance syndrome and Alzheimer's disease: age- and obesity-related effects on memory, amyloid, and inflammation. *Neurobiol Aging* 26 Suppl 1, 65–69. [PubMed: 16266773]
- Craft S, Newcomer J, Kanne S, Dagogo-Jack S, Cryer P, Sheline Y, Luby J, Dagogo-Jack A, Alderson A, 1996 Memory improvement following induced hyperinsulinemia in Alzheimer's disease. *Neurobiol Aging* 17, 123–130. [PubMed: 8786794]
- Craft S, Peskind E, Schwartz MW, Schellenberg GD, Raskind M, Porte D, Jr., 1998 Cerebrospinal fluid and plasma insulin levels in Alzheimer's disease: relationship to severity of dementia and apolipoprotein E genotype. *Neurology* 50, 164–168. [PubMed: 9443474]
- Craft S, Watson GS, 2004 Insulin and neurodegenerative disease: shared and specific mechanisms. *Lancet Neurol* 3, 169–178. [PubMed: 14980532]
- De Felice FG, 2013 Alzheimer's disease and insulin resistance: translating basic science into clinical applications. *J Clin Invest* 123, 531–539. [PubMed: 23485579]
- De Felice FG, Lourenco MV, Ferreira ST, 2014 How does brain insulin resistance develop in Alzheimer's disease? *Alzheimers Dement* 10, S26–32. [PubMed: 24529521]
- de la Monte SM, 2013 Intranasal insulin therapy for cognitive impairment and neurodegeneration: current state of the art. *Expert Opin Drug Deliv* 10, 1699–1709. [PubMed: 24215447]
- de la Monte SM, 2014 Type 3 diabetes is sporadic Alzheimer's disease: mini-review. *Eur Neuropsychopharmacol* 24, 1954–1960. [PubMed: 25088942]
- de la Monte SM, Tong M, Lester-Coll N, Plater M, Jr., Wands JR, 2006 Therapeutic rescue of neurodegeneration in experimental type 3 diabetes: relevance to Alzheimer's disease. *J Alzheimers Dis* 10, 89–109. [PubMed: 16988486]
- de la Monte SM, Wands JR, 2008 Alzheimer's disease is type 3 diabetes-evidence reviewed. *J Diabetes Sci Technol* 2, 1101–1113. [PubMed: 19885299]
- Dou JT, Chen M, Dufour F, Alkon DL, Zhao WQ, 2005 Insulin receptor signaling in long-term memory consolidation following spatial learning. *Learn Mem* 12, 646–655. [PubMed: 16287721]
- El Messari S, Leloup C, Quignon M, Brisorgueil MJ, Penicaud L, Arluison M, 1998 Immunocytochemical localization of the insulin-responsive glucose transporter 4 (Glut4) in the rat central nervous system. *J Comp Neurol* 399, 492–512. [PubMed: 9741479]
- Fadel JR, Reagan LP, 2016 Stop signs in hippocampal insulin signaling: the role of insulin resistance in structural, functional and behavioral deficits. *Curr Opin Behav Sci* 9, 47–54. [PubMed: 26955646]

- Frazier HN, Anderson KL, Maimaiti S, Ghoweri AO, Kraner SD, Popa GJ, Hampton KK, Mendenhall MD, Norris CM, Craven RJ, Thibault O, 2018 Expression of a Constitutively Active Human Insulin Receptor in Hippocampal Neurons Does Not Alter VGCC Currents. *Neurochem Res*.
- Freiherr J, Hallschmid M, Frey WH, 2nd, Brunner YF, Chapman CD, Holscher C, Craft S, De Felice FG, Benedict C, 2013 Intranasal insulin as a treatment for Alzheimer's disease: a review of basic research and clinical evidence. *CNS Drugs* 27, 505–514. [PubMed: 23719722]
- Frolich L, Blum-Degen D, Bernstein HG, Engelsberger S, Humrich J, Laufer S, Muschner D, Thalheimer A, Turk A, Hoyer S, Zochling R, Boissl KW, Jellinger K, Riederer P, 1998 Brain insulin and insulin receptors in aging and sporadic Alzheimer's disease. *J Neural Transm (Vienna)* 105, 423–438. [PubMed: 9720972]
- Gispén WH, Biessels GJ, 2000 Cognition and synaptic plasticity in diabetes mellitus. *Trends Neurosci* 23, 542–549. [PubMed: 11074263]
- Goyal MS, Vlassenko AG, Blazey TM, Su Y, Couture LE, Durbin TJ, Bateman RJ, Benzinger TL, Morris JC, Raichle ME, 2017 Loss of Brain Aerobic Glycolysis in Normal Human Aging. *Cell Metab* 26, 353–360 e353. [PubMed: 28768174]
- Griffith CM, Eid T, Rose GM, Patrylo PR, 2018 Evidence for altered insulin receptor signaling in Alzheimer's disease. *Neuropharmacology* 136, 202–215. [PubMed: 29353052]
- Grillo CA, Pirolì GG, Hendry RM, Reagan LP, 2009 Insulin-stimulated translocation of GLUT4 to the plasma membrane in rat hippocampus is PI3-kinase dependent. *Brain Res* 1296, 35–45. [PubMed: 19679110]
- Grillo CA, Pirolì GG, Lawrence RC, Wrihten SA, Green AJ, Wilson SP, Sakai RR, Kelly SJ, Wilson MA, Mott DD, Reagan LP, 2015 Hippocampal Insulin Resistance Impairs Spatial Learning and Synaptic Plasticity. *Diabetes* 64, 3927–3936. [PubMed: 26216852]
- Hallschmid M, Benedict C, Schultes B, Fehm HL, Born J, Kern W, 2004 Intranasal insulin reduces body fat in men but not in women. *Diabetes* 53, 3024–3029. [PubMed: 15504987]
- Heni M, Schopfer P, Peter A, Sartorius T, Fritsche A, Synofzik M, Haring HU, Maetzler W, Hennige AM, 2014 Evidence for altered transport of insulin across the bloodbrain barrier in insulin-resistant humans. *Acta Diabetol* 51, 679–681. [PubMed: 24370925]
- Hirvonen J, Virtanen KA, Nummenmaa L, Hannukainen JC, Honka MJ, Bucci M, Nesterov SV, Parkkola R, Rinne J, Iozzo P, Nuutila P, 2011 Effects of insulin on brain glucose metabolism in impaired glucose tolerance. *Diabetes* 60, 443–447. [PubMed: 21270256]
- Ho L, Qin W, Pompl PN, Xiang Z, Wang J, Zhao Z, Peng Y, Cambareri G, Rocher A, Mobbs CV, Hof PR, Pasinetti GM, 2004 Diet-induced insulin resistance promotes amyloidosis in a transgenic mouse model of Alzheimer's disease. *FASEB J* 18, 902–904. [PubMed: 15033922]
- Howarth C, Gleeson P, Attwell D, 2012 Updated energy budgets for neural computation in the neocortex and cerebellum. *J Cereb Blood Flow Metab* 32, 1222–1232. [PubMed: 22434069]
- Hoyer S, 1986 Senile dementia and Alzheimer's disease. Brain blood flow and metabolism. *Prog Neuropsychopharmacol Biol Psychiatry* 10, 447–478. [PubMed: 3541046]
- Hoyer S, 1998 Is sporadic Alzheimer disease the brain type of non-insulin dependent diabetes mellitus? A challenging hypothesis. *J Neural Transm (Vienna)* 105, 415–422. [PubMed: 9720971]
- Hoyer S, 2004 Glucose metabolism and insulin receptor signal transduction in Alzheimer disease. *Eur J Pharmacol* 490, 115–125. [PubMed: 15094078]
- Hughes TM, Craft S, 2015 The role of insulin in the vascular contributions to age-related dementia. *Biochim Biophys Acta*.
- Israel PA, Park CR, Schwartz MW, Green PK, Sipols AJ, Woods SC, Porte D, Jr., Figlewicz DP, 1993 Effect of diet-induced obesity and experimental hyperinsulinemia on insulin uptake into CSF of the rat. *Brain Res Bull* 30, 571–575. [PubMed: 8457906]
- Kamal A, Ramakers GM, Gispén WH, Biessels GJ, 2013 Hyperinsulinemia in rats causes impairment of spatial memory and learning with defects in hippocampal synaptic plasticity by involvement of postsynaptic mechanisms. *Exp Brain Res* 226, 45–51. [PubMed: 23371746]
- Kern W, Born J, Schreiber H, Fehm HL, 1999 Central nervous system effects of intranasally administered insulin during euglycemia in men. *Diabetes* 48, 557–563. [PubMed: 10078556]

- Kobayashi M, Nikami H, Morimatsu M, Saito M, 1996 Expression and localization of insulin-regulatable glucose transporter (GLUT4) in rat brain. *Neurosci Lett* 213, 103–106. [PubMed: 8858619]
- Kullmann S, Frank S, Heni M, Ketterer C, Veit R, Haring HU, Fritsche A, Preissl H, 2013 Intranasal insulin modulates intrinsic reward and prefrontal circuitry of the human brain in lean women. *Neuroendocrinology* 97, 176–182. [PubMed: 22922661]
- Kullmann S, Fritsche A, Wagner R, Schwab S, Haring HU, Preissl H, Heni M, 2017 Hypothalamic insulin responsiveness is associated with pancreatic insulin secretion in humans. *Physiol Behav* 176, 134–138. [PubMed: 28347723]
- Kuusisto J, Koivisto K, Mykkanen L, Helkala EL, Vanhanen M, Hanninen T, Pyorala K, Riekkinen P, Laakso M, 1993 Essential hypertension and cognitive function. The role of hyperinsulinemia. *Hypertension* 22, 771–779. [PubMed: 8225537]
- Lannert H, Hoyer S, 1998 Intracerebroventricular administration of streptozotocin causes long-term diminutions in learning and memory abilities and in cerebral energy metabolism in adult rats. *Behav Neurosci* 112, 1199–1208. [PubMed: 9829797]
- Lebwohl DE, Nunez I, Chan M, Rosen OM, 1991 Expression of inducible membraneanchored insulin receptor kinase enhances deoxyglucose uptake. *J Biol Chem* 266, 386–390. [PubMed: 1845969]
- Lee JH, Jahrling JB, Denner L, Dineley KT, 2018 Targeting Insulin for Alzheimer's Disease: Mechanisms, Status and Potential Directions. *J Alzheimers Dis*.
- Leino RL, Gerhart DZ, van Bueren AM, McCall AL, Drewes LR, 1997 Ultrastructural localization of GLUT 1 and GLUT 3 glucose transporters in rat brain. *J Neurosci Res* 49, 617–626. [PubMed: 9302083]
- Long JM, Davis BJ, Garofalo P, Spangler EL, Ingram DK, 1992 Complex maze performance in young and aged rats: response to glucose treatment and relationship to blood insulin and glucose. *Physiol Behav* 51, 411–418. [PubMed: 1313593]
- Luchsinger JA, 2008 Adiposity, hyperinsulinemia, diabetes and Alzheimer's disease: an epidemiological perspective. *Eur J Pharmacol* 585, 119–129. [PubMed: 18384771]
- Luchsinger JA, Tang MX, Shea S, Mayeux R, 2004 Hyperinsulinemia and risk of Alzheimer disease. *Neurology* 63, 1187–1192. [PubMed: 15477536]
- Luckett BS, Frielle JL, Wolfgang L, Stocker SD, 2013 Arcuate nucleus injection of an anti-insulin affibody prevents the sympathetic response to insulin. *Am J Physiol Heart Circ Physiol* 304, H1538–1546. [PubMed: 23542919]
- Macklin L, Griffith CM, Cai Y, Rose GM, Yan XX, Patrylo PR, 2017 Glucose tolerance and insulin sensitivity are impaired in APP/PS1 transgenic mice prior to amyloid plaque pathogenesis and cognitive decline. *Exp Gerontol* 88, 9–18. [PubMed: 28025127]
- Maimaiti S, Anderson KL, DeMoll C, Brewer LD, Rauh BA, Gant JC, Blalock EM, Porter NM, Thibault O, 2016 Intranasal Insulin Improves Age-Related Cognitive Deficits and Reverses Electrophysiological Correlates of Brain Aging. *J Gerontol A Biol Sci Med Sci* 71, 30–39. [PubMed: 25659889]
- Maimaiti S, Frazier HN, Anderson KL, Ghoweri AO, Brewer LD, Porter NM, Thibault O, 2017 Novel calcium-related targets of insulin in hippocampal neurons. *Neuroscience* 364, 130–142. [PubMed: 28939258]
- Mao YF, Guo Z, Zheng T, Jiang Y, Yan Y, Yin X, Chen Y, Zhang B, 2016 Intranasal insulin alleviates cognitive deficits and amyloid pathology in young adult APPswe/PS1dE9 mice. *Aging Cell* 15, 893–902. [PubMed: 27457264]
- Marks DR, Tucker K, Cavallin MA, Mast TG, Fadool DA, 2009 Awake intranasal insulin delivery modifies protein complexes and alters memory, anxiety, and olfactory behaviors. *J Neurosci* 29, 6734–6751. [PubMed: 19458242]
- McEwen BS, Reagan LP, 2004 Glucose transporter expression in the central nervous system: relationship to synaptic function. *Eur J Pharmacol* 490, 13–24. [PubMed: 15094070]
- McNay EC, 2005 The impact of recurrent hypoglycemia on cognitive function in aging. *Neurobiol Aging* 26 Suppl 1, 76–79. [PubMed: 16236383]

- McNay EC, Gold PE, 2001 Age-related differences in hippocampal extracellular fluid glucose concentration during behavioral testing and following systemic glucose administration. *J Gerontol A Biol Sci Med Sci* 56, B66–71. [PubMed: 11213269]
- McNay EC, Ong CT, McCrimmon RJ, Cresswell J, Bogan JS, Sherwin RS, 2010 Hippocampal memory processes are modulated by insulin and high-fat-induced insulin resistance. *Neurobiol Learn Mem* 93, 546–553. [PubMed: 20176121]
- Mielke R, Herholz K, Grond M, Kessler J, Heiss WD, 1992 Differences of regional cerebral glucose metabolism between presenile and senile dementia of Alzheimer type. *Neurobiol Aging* 13, 93–98. [PubMed: 1542386]
- Molina JA, Jimenez-Jimenez FJ, Vargas C, Gomez P, de Bustos F, Gomez-Escalonilla C, Zurdo M, Tallon A, Martinez-Salio A, Porta-Etessam J, Villanueva C, Arenas J, 2002 Cerebrospinal fluid levels of insulin in patients with Alzheimer's disease. *Acta Neurol Scand* 106, 347–350. [PubMed: 12460139]
- Moloney AM, Griffin RJ, Timmons S, O'Connor R, Ravid R, O'Neill C, 2010 Defects in IGF-1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance to IGF-1 and insulin signalling. *Neurobiol Aging* 31, 224–243. [PubMed: 18479783]
- Mosconi L, 2013 Glucose metabolism in normal aging and Alzheimer's disease: Methodological and physiological considerations for PET studies. *Clin Transl Imaging* 1.
- Mosconi L, De Santi S, Li J, Tsui WH, Li Y, Boppana M, Laska E, Rusinek H, de Leon MJ, 2008a Hippocampal hypometabolism predicts cognitive decline from normal aging. *Neurobiol Aging* 29, 676–692. [PubMed: 17222480]
- Mosconi L, Pupi A, De Cristofaro MT, Fayyaz M, Sorbi S, Herholz K, 2004 Functional interactions of the entorhinal cortex: an 18F-FDG PET study on normal aging and Alzheimer's disease. *J Nucl Med* 45, 382–392. [PubMed: 15001677]
- Mosconi L, Tsui WH, Herholz K, Pupi A, Drzezga A, Lucignani G, Reiman EM, Holthoff V, Kalbe E, Sorbi S, Diehl-Schmid J, Perneckzy R, Clerici F, Caselli R, Beuthien-Baumann B, Kurz A, Minoshima S, de Leon MJ, 2008b Multicenter standardized 18F-FDG PET diagnosis of mild cognitive impairment, Alzheimer's disease, and other dementias. *J Nucl Med* 49, 390–398. [PubMed: 18287270]
- Pancani T, Anderson KL, Brewer LD, Kadish I, DeMoll C, Landfield PW, Blalock EM, Porter NM, Thibault O, 2013 Effect of high-fat diet on metabolic indices, cognition, and neuronal physiology in aging F344 rats. *Neurobiol Aging* 34, 1977–1987. [PubMed: 23545425]
- Pancani T, Anderson KL, Porter NM, Thibault O, 2011 Imaging of a glucose analog, calcium and NADH in neurons and astrocytes: Dynamic responses to depolarization and sensitivity to pioglitazone. *Cell Calcium* 50, 548–558. [PubMed: 21978418]
- Paranjape SA, Chan O, Zhu W, Horblitt AM, McNay EC, Cresswell JA, Bogan JS, McCrimmon RJ, Sherwin RS, 2010 Influence of insulin in the ventromedial hypothalamus on pancreatic glucagon secretion in vivo. *Diabetes* 59, 1521–1527. [PubMed: 20299468]
- Pearson-Leary J, Jahagirdar V, Sage J, McNay EC, 2017 Insulin modulates hippocampally-mediated spatial working memory via glucose transporter-4. *Behav Brain Res*.
- Pearson-Leary J, Jahagirdar V, Sage J, McNay EC, 2018 Insulin modulates hippocampally-mediated spatial working memory via glucose transporter-4. *Behav Brain Res* 338, 32–39. [PubMed: 28943428]
- Pearson-Leary J, McNay EC, 2012 Intrahippocampal administration of amyloid-beta(1–42) oligomers acutely impairs spatial working memory, insulin signaling, and hippocampal metabolism. *J Alzheimers Dis* 30, 413–422. [PubMed: 22430529]
- Reagan LP, 2005 Neuronal insulin signal transduction mechanisms in diabetes phenotypes. *Neurobiol Aging* 26 Suppl 1, 56–59. [PubMed: 16225964]
- Reger MA, Watson GS, Frey WH, 2nd, Baker LD, Cholerton B, Keeling ML, Belongia DA, Fishel MA, Plymate SR, Schellenberg GD, Cherrier MM, Craft S, 2006 Effects of intranasal insulin on cognition in memory-impaired older adults: modulation by APOE genotype. *Neurobiol Aging* 27, 451–458. [PubMed: 15964100]

Reger MA, Watson GS, Green PS, Wilkinson CW, Baker LD, Cholerton B, Fishel MA, Plymate SR, Breitner JC, DeGroot DW, Mehta P, Craft S, 2008 Intranasal insulin improves cognition and modulates beta-amyloid in early AD. *Neurology* 70, 440–448. [PubMed: 17942819]

Rhea EM, Humann SR, Nirkhe S, Farr SA, Morley JE, Banks WA, 2017 Intranasal Insulin Transport is Preserved in Aged SAMP8 Mice and is Altered by Albumin and Insulin Receptor Inhibition. *J Alzheimers Dis* 57, 241–252. [PubMed: 28222522]

Rivera EJ, Goldin A, Fulmer N, Tavares R, Wands JR, de la Monte SM, 2005 Insulin and insulin-like growth factor expression and function deteriorate with progression of Alzheimer's disease: link to brain reductions in acetylcholine. *J Alzheimers Dis* 8, 247–268. [PubMed: 16340083]

Rodriguez-Rivera J, Denner L, Dineley KT, 2011 Rosiglitazone reversal of Tg2576 cognitive deficits is independent of peripheral gluco-regulatory status. *Behav Brain Res* 216, 255–261. [PubMed: 20709114]

Salameh TS, Bullock KM, Hujoel IA, Niehoff ML, Wolden-Hanson T, Kim J, Morley JE, Farr SA, Banks WA, 2015 Central Nervous System Delivery of Intranasal Insulin: Mechanisms of Uptake and Effects on Cognition. *J Alzheimers Dis* 47, 715–728. [PubMed: 26401706]

Sartorius T, Peter A, Heni M, Maetzler W, Fritsche A, Haring HU, Hennige AM, 2015 The brain response to peripheral insulin declines with age: a contribution of the blood-brain barrier? *PLoS One* 10, e0126804. [PubMed: 25965336]

Schubert M, Gautam D, Surjo D, Ueki K, Baudier S, Schubert D, Kondo T, Alber J, Galldiks N, Kustermann E, Arndt S, Jacobs AH, Krone W, Kahn CR, Bruning JC, 2004 Role for neuronal insulin resistance in neurodegenerative diseases. *Proc Natl Acad Sci U S A* 101, 3100–3105. [PubMed: 14981233]

Schwartz MW, Figlewicz DF, Kahn SE, Baskin DG, Greenwood MR, Porte D, Jr., 1990 Insulin binding to brain capillaries is reduced in genetically obese, hyperinsulinemic Zucker rats. *Peptides* 11, 467–472. [PubMed: 2199946]

Stanley M, Macauley SL, Caesar EE, Koscal LJ, Moritz W, Robinson GO, Roh J, Keyser J, Jiang H, Holtzman DM, 2016 The Effects of Peripheral and Central High Insulin on Brain Insulin Signaling and Amyloid-beta in Young and Old APP/PS1 Mice. *J Neurosci* 36, 11704–11715. [PubMed: 27852778]

Steen E, Terry BM, Rivera EJ, Cannon JL, Neely TR, Tavares R, Xu XJ, Wands JR, de la Monte SM, 2005 Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease--is this type 3 diabetes? *J Alzheimers Dis* 7, 63–80. [PubMed: 15750215]

Stolk RP, Breteler MM, Ott A, Pols HA, Lamberts SW, Grobbee DE, Hofman A, 1997 Insulin and cognitive function in an elderly population. The Rotterdam Study. *Diabetes Care* 20, 792–795. [PubMed: 9135944]

Swerdlow RH, 2007 PET sheds light on Alzheimer's disease genetic risk. *Proc Natl Acad Sci U S A* 104, 18881–18882. [PubMed: 18025455]

Talbot K, Wang HY, Kazi H, Han LY, Bakshi KP, Stucky A, Fuino RL, Kawaguchi KR, Samoyedny AJ, Wilson RS, Arvanitakis Z, Schneider JA, Wolf BA, Bennett DA, Trojanowski JQ, Arnold SE, 2012 Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *J Clin Invest* 122, 1316–1338. [PubMed: 22476197]

Tchilian EZ, Zhelezarov IE, Petkov VV, Hadjiivanova CI, 1990 125I-insulin binding is decreased in olfactory bulbs of aged rats. *Neuropeptides* 17, 193–196. [PubMed: 2093152]

Thienel M, Wilhelm I, Benedict C, Born J, Hallschmid M, 2017 Intranasal insulin decreases circulating cortisol concentrations during early sleep in elderly humans. *Neurobiol Aging* 54, 170–174. [PubMed: 28385552]

Tramutola A, Arena A, Cini C, Butterfield DA, Barone E, 2017 Modulation of GLP-1 signaling as a novel therapeutic approach in the treatment of Alzheimer's disease pathology. *Expert Rev Neurother* 17, 59–75. [PubMed: 27715341]

Tschritter O, Preissl H, Hennige AM, Stumvoll M, Porubsky K, Frost R, Marx H, Klose B, Lutzenberger W, Birbaumer N, Haring HU, Fritsche A, 2006 The cerebrocortical response to hyperinsulinemia is reduced in overweight humans: a magnetoencephalographic study. *Proc Natl Acad Sci U S A* 103, 12103–12108. [PubMed: 16877540]

- Uemura E, Greenlee HW, 2006 Insulin regulates neuronal glucose uptake by promoting translocation of glucose transporter GLUT3. *Exp Neurol* 198, 48–53. [PubMed: 16337941]
- van der Heide LP, Ramakers GM, Smidt MP, 2006 Insulin signaling in the central nervous system: learning to survive. *Prog Neurobiol* 79, 205–221. [PubMed: 16916571]
- Vannucci SJ, Maher F, Simpson IA, 1997 Glucose transporter proteins in brain: delivery of glucose to neurons and glia. *Glia* 21, 2–21. [PubMed: 9298843]
- Velazquez R, Tran A, Ishimwe E, Denner L, Dave N, Oddo S, Dineley KT, 2017 Central insulin dysregulation and energy dyshomeostasis in two mouse models of Alzheimer's disease. *Neurobiol Aging* 58, 1–13. [PubMed: 28688899]
- Vogt MC, Bruning JC, 2013 CNS insulin signaling in the control of energy homeostasis and glucose metabolism - from embryo to old age. *Trends Endocrinol Metab* 24, 76–84. [PubMed: 23265947]
- Wickelgren I, 1998 Tracking insulin to the mind. *Science* 280, 517–519. [PubMed: 9575095]
- Willette AA, Xu G, Johnson SC, Birdsill AC, Jonaitis EM, Sager MA, Hermann BP, La Rue A, Asthana S, Bendlin BB, 2013 Insulin resistance, brain atrophy, and cognitive performance in late middle-aged adults. *Diabetes Care* 36, 443–449. [PubMed: 23069842]
- Woods SC, Lotter EC, McKay LD, Porte D, Jr., 1979 Chronic intracerebroventricular infusion of insulin reduces food intake and body weight of baboons. *Nature* 282, 503–505. [PubMed: 116135]
- Zaia A, Piantanelli L, 2000 Insulin receptors in the brain cortex of aging mice. *Mech Ageing Dev* 113, 227–232. [PubMed: 10714941]
- Zhao W, Chen H, Xu H, Moore E, Meiri N, Quon MJ, Alkon DL, 1999 Brain insulin receptors and spatial memory. Correlated changes in gene expression, tyrosine phosphorylation, and signaling molecules in the hippocampus of water maze trained rats. *J Biol Chem* 274, 34893–34902. [PubMed: 10574963]
- Zhao WQ, Chen H, Quon MJ, Alkon DL, 2004 Insulin and the insulin receptor in experimental models of learning and memory. *Eur J Pharmacol* 490, 71–81. [PubMed: 15094074]
- Zhao WQ, De Felice FG, Fernandez S, Chen H, Lambert MP, Quon MJ, Krafft GA, Klein WL, 2008 Amyloid beta oligomers induce impairment of neuronal insulin receptors. *FASEB J* 22, 246–260. [PubMed: 17720802]

Highlights:

- Increase in published articles containing keywords “insulin resistance” and “brain”
- Evidence for preserved brain insulin sensitivity with age and AD is presented
- Glucose transport is an important component of insulin resistance in the brain

Figure 1.

Number of articles containing the keywords “insulin resistance,” “brain,” and either “aged” or “Alzheimer’s disease” between the years 1998 and 2018. Number of articles were determined by querying the PubMed online database via the EndNote search engine (EndNote™; v.8.2, build 11343). **A.** Number of individual articles published per year. **B.** Number of cumulative articles in the literature between 1998 and 2018.

Figure 2.

Insulin's acute effect on the slow afterhyperpolarization (sAHP) in young and mid-aged F344 male rats. **A.** Comparison of insulin-mediated sAHP reduction between hippocampal slices obtained from young (2.5–4 mo) and mid-aged (12.5–13.5 mo) animals following 15 min perfusion of Humalog® (500nM). A significant reduction in the sAHP was observed in both groups of animals ($F(1,10) = 7.1$; $p < 0.05$). Reduction in the sAHP of midaged animals suggests that insulin-sensitivity in the hippocampus is preserved, not diminished, in early aging. * indicates $p < 0.05$. All data represent means $\pm$ SEM. *Reprinted from Neurobiology of Aging, Vol 34(8), Pancani, et. al., "Effect of high fat diet on metabolic indices, cognition and neuronal physiology in aging F344 rats," pg. 1977–1987, Copyright (2013), with permission from Elsevier.*

Figure 3.

Effect of a zinc-free insulin formulation (Apidra®) on the AHP *ex vivo*. **A.** Measures of the medium AHP (mAHP) in hippocampal slices obtained from young (3 mo) and aged (21 mo) F344 male rats following administration of 1nM Apidra®. Apidra® significantly reduced the mAHP in slices from both young ($n = 8$) and aged ($n = 9$) animals, indicated by a main effect of treatment ($F(1,24) = 32$, $p < 0.0001$). **B.** Measures of the sAHP in hippocampal slices obtained from young and aged F344 male rats following administration of 1nM Apidra®. As with the mAHP, Apidra® significantly reduced the sAHP in both young and aged animals ($F(1,24) = 29.1$, $p < 0.0001$). This effect was larger in aged animals compared to young, evidenced by a significant interaction term following 2-way ANOVA ($[F(1,24) = 10.9$, $p < 0.005$ for the mAHP). This suggests a preservation of insulin-sensitivity in aged

animals. # indicates $p < 0.10$; * indicates $p < 0.05$. All data represent means $\pm$ SEM.
Reprinted from: Maimaiti, et. al., "Intranasal Insulin Improves Age-Related Cognitive Deficits and Reverses Electrophysiological Correlates of Brain Aging," Journal of Gerontology, 2016, Vol. 71, No. 1, 30–39, by permission of Oxford University Press.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 4.

2-NBDG imaging of primary hippocampal neurons with or without expression of a constitutively active form of the human insulin receptor (IR β). **A.** Analysis of indirect measures of glucose uptake using initial 2-NBDG fluorescent signal. Images were taken 3 minutes postwash. All data were normalized prior to statistical analysis using background subtraction. Results indicate a significant increase in initial 2-NBDG signal in IR β -expressing neurons (IR β mCherry, n=27) compared to controls (mCherry, n=25). * indicates $p < 0.05$. All data represent means $\pm$ SEM (*for Methods*, see Pancani, et al., 2013).