Published in final edited form as:

Auton Neurosci. 2019 March ; 217: 35–40. doi:10.1016/j.autneu.2018.12.008.

A time to fight: circadian control of aggression and associated autonomic support

William D. Todd^{#*} and Natalia L. Machado[#]

Department of Neurology, Division of Sleep Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts, 02215, USA

[#] These authors contributed equally to this work.

Abstract

The central circadian clock, located in the suprachiasmatic nucleus of the mammalian hypothalamus (SCN), regulates daily behavioral rhythms including the temporal propensity for aggressive behavior. Such aggression propensity rhythms are regulated by a functional circuit from the SCN to neurons that drive attack behavior in the ventromedial hypothalamus (VMH), via a relay in the subparaventricular zone (SPZ). In addition to this pathway, the SCN also regulates sleep-wake and locomotor activity rhythms, via the SPZ, in a circuit to the dorsomedial hypothalamus (DMH), a structure that is also known to play a key role in autonomic function and the sympathetic “fight-or-flight” response (which prepares the body for action in stressful situations such as an agonistic encounter). While the autonomic nervous system is known to be under pronounced circadian control, it is less apparent how such autonomic rhythms and their underlying circuitry may support the temporal propensity for aggressive behavior. Additionally, it is unclear how circadian and autonomic dysfunction may contribute to aberrant social and emotional behavior, such as agitation and aggression. Here we review the literature concerning interactions between the circadian and autonomic systems and aggression, and we discuss the implications of these relationships for human neural and behavioral pathologies.

Keywords

circadian; aggression; autonomic; fight-or-flight; hypothalamus

1. Introduction

Circadian clocks prepare the brain and body to execute particular physiological and behavioral processes during different times of day, conveying an evolutionarily adaptive

* corresponding author Contact: William D. Todd, Beth Israel Deaconess Medical Center, Department of Neurology, 330 Brookline Ave, E/CLS 7, Boston, MA 02215, wtodd@bidmc.harvard.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations of interest
None

advantage for an organism's survival. Indeed, the central circadian clock regulates baseline rhythms of the autonomic nervous system (Buijs et al., 2013), which is the system that primarily controls another adaptive mechanism, the so-called "fight-or-flight" response. This term was coined by Cannon to refer to physiological responses induced by stress that prepare the body to face adversity, such as an agonistic encounter (Cannon, 1929), and thus the autonomic system's role in situations that promote aggressive behavior has long been recognized. We recently demonstrated that the central circadian system directly regulates a rhythm in the propensity for aggressive behavior through a novel neural pathway (Todd et al., 2018). However, it is less apparent how the circadian control of autonomic regulation and its underlying circuitry may support this temporal propensity for aggression, as well as the implications of these connections for human disorders in which circadian and autonomic dysfunction are comorbid factors with agitation and aggressive behavior. Here we review the literature on interactions between the circadian system, the autonomic system, and aggressive behavior, and we present opportunities for new lines of research that could potentially inform human disorders characterized by circadian and autonomic disruptions and aberrant social and emotional behaviors such as aggression.

2. Anatomy of the circadian system and its direct neural control of aggression

The suprachiasmatic nucleus (SCN) of the hypothalamus is the central circadian clock of the mammalian brain and is required for daily rhythms of physiology and behavior (Reppert and Weaver, 2002). The SCN entrains such rhythms to the daily light-dark cycle (Rusak et al., 1990), via photic input from the retina by way of the retinohypothalamic tract (Gooley et al., 2003, Todd et al., 2012). SCN neurons, which are highly coupled *in vivo*, can function as individual oscillators with rhythmic electrical activity on a period of about 24h (Welsh et al., 1995). This rhythmic activity is under the control of clock genes, via a transcriptional-translational-posttranslational negative feedback loop (Jin et al., 1999). While this same genetic machinery is present in cells throughout the brain and body, the SCN is necessary to synchronize peripheral oscillators and maintain rhythmic behavior (Mohawk et al., 2012). Importantly, axonal output appears to be the primary method by which the SCN establishes such synchrony *in vivo* (Gall et al., 2012, Saper, 2013), and the majority of SCN axons target a nearby region known as the subparaventricular zone (SPZ) (Watts et al., 1987, Watts and Swanson, 1987, Vujovic et al., 2015). Indeed, circadian rhythms of sleep-wake, locomotor activity, and feeding behavior are known to be regulated by a pathway from the SCN, through the SPZ, to the dorsomedial nucleus of the hypothalamus (DMH) (Lu et al., 2001, Chou et al., 2003, Saper, 2013). Pathways by which the circadian system directly regulates more complex behaviors, such as particular social and emotional behaviors, have remained more elusive. For example, while evidence from rodents (Sofia and Salama, 1970, Landau, 1975b, Haller et al., 2000a) and humans (Manfredini et al., 2001) previously demonstrated temporal differences in aggressive behavior, it was unclear whether the SCN clock directly regulates a rhythm in aggression (Bronsard and Bartolomei, 2013, Hood and Amir, 2018). However, we recently demonstrated that the propensity for aggressive behavior follows a daily rhythm in male mice, with the highest levels during the early active phase and the lowest during the early resting phase, and we showed that this rhythm is contingent

on proper functioning of GABAergic SPZ (SPZ^{GABA}) neurons (Todd et al., 2018). Genetic deletion of the vesicular GABA transporter (VGAT) in these neurons, which renders them unable to release GABA, results in increased behavioral aggression during the early light period, the resting phase for nocturnal mice. Importantly, while these deletions reduce locomotor activity and plasma corticosterone levels during the early dark period, when both are normally high in intact mice, they do not increase them during the early light period, suggesting that the increase in aggression at that time results from a direct neural effect. Attack behavior in male mice is known to be directly regulated by neurons within the ventromedial nucleus of the hypothalamus (VMH), and specifically those that express estrogen (Esr1) and progesterone receptors in the ventrolateral VMH (VMHvl) (Lin et al., 2011, Yang et al., 2013, Lee et al., 2014, Yang et al., 2017). We showed that SPZ^{GABA} neurons make functional synaptic connections onto Esr1-expressing VMHvl neurons, and that vasoactive intestinal polypeptide (VIP) neurons of the SCN make functional connections onto SPZ^{GABA} neurons that project to the VMH (Todd et al., 2018). We showed that these SPZ neurons are also more active during the early light period compared to the early dark period, and acutely inhibiting these neurons using a novel inhibitory chemogenetic vector reversibly increased aggression in a phase-dependent manner. Altogether, the SCN→SPZ→VMH circuit constitutes a critical pathway through which the central circadian clock gates aggression propensity across the 24h day (see Figure 1). While this circuit represents the basis for the central circadian control of aggression, the circadian system is also known to modulate the autonomic system, and through these pathways likely supports the body's ability to respond accordingly during an aggressive act.

3. Circadian regulation of the autonomic system

The autonomic system exhibits a distinct circadian pattern, as shown by the early rise in sympathetic nerve activity, blood pressure (BP), heart rate (HR) and increases in body temperature (Tb), cortisol release, and glucose levels that prepare the body for the daily active phase (Buijs et al., 2013, Smolensky et al., 2016). In humans, the well-known phenomenon of cardiovascular events, such as ventricular tachycardia, ventricular fibrillation and sudden cardiac death, most commonly occurring in the morning after waking, reinforces a critical role for the circadian system in autonomic function (Takeda and Maemura, 2016). Additionally, professions involving night-shift work, which causes chronic circadian disruption, are associated with increased risks of cardiovascular disease (Thosar et al., 2018). HR regulation appears to transition from the influence of high sympathetic nerve activity in the early active phase to a state of relative vagal dominance during the resting period (Verrier and Tan, 2009, Takeda and Maemura, 2016). Changes in autonomic regulation are also found during sleep. For instance, non-rapid eye movement (NREM) sleep is characterized by vagal dominance, while sympathetic nerve activity is stable, and during this stage individuals show relative hypotension, bradycardia, and reduced cardiac output and systemic vascular resistance. This pattern is disrupted during rapid eye movement (REM) sleep, which is characterized by vagal withdrawal, while sympathetic nerve activity can reach even higher levels than during wake (Verrier and Josephson, 2009).

Circadian control of the autonomic system has been shown to be coordinated by the SCN, as bilateral thermal ablation of the SCN in mice abolishes HR circadian rhythms, an important

indirect indicator of autonomic control (Warren et al., 1994). Additionally, global knockouts of the clock genes *Clock* and *Bmal1* result in a loss of the HR circadian rhythm (Rahman et al., 2018). However, cardiomyocyte-specific knockout of *Clock* or *Bmal1* reduces but does not abolish the HR circadian rhythm, suggesting that the local cardiac clock contributes, but is not essential for this rhythm (Bray et al., 2008, Schroeder et al., 2011). These results support the hypothesis that the SCN works as an autonomous clock regulating HR rhythmicity. Indeed, VIP neurons within the SCN, which represent only about 10% of all SCN neurons (Mazuski et al., 2018), have specifically been shown to modulate circadian rhythmicity in the heart, but do not appear to eliminate the circadian rhythm of Tb (Schroeder et al., 2011).

The pathway by which the SCN regulates circadian rhythms of Tb is unclear (Saper, 2013), but one possibility could be through the SCN's modulation of brown adipose tissue (BAT) activity (Amir et al., 1989). BAT thermogenesis is regulated by the sympathetic system and exhibits an ultradian rhythm (less than 24h) that corresponds to phase-linked changes in Tb (Ootsuka et al., 2009). Indeed, the DMH region, and specifically the glutamatergic neurons within the dorsal hypothalamic area (DHA^{Vglut2}), regulates BAT activity and thermogenesis induced by stressful stimuli, as part of the fight-or-flight response (Morrison, 2004, Machado et al., 2018, Kataoka et al., 2014). However, DMH/DHA lesions do not abolish circadian rhythms of Tb (Chou et al., 2003), and thus a neuro-circuit for the circadian control of Tb remains to be determined.

In addition to its role in modulating sleep-wake centers and Tb, there is substantial evidence that the DMH projects to sympathetic outflow areas to regulate cardiovascular and respiratory function (see below, and Figure 1). It is therefore likely that the SCN→SPZ→DMH pathway also underlies circadian cardiac and respiratory autonomic rhythms, although this remains to be demonstrated definitively. What is clear is that the hypothalamus regulates the fight-or-flight stress response, and in this context, the DMH is thought to be the critical hypothalamic region mediating the sympathetic activation and cardiovascular responses induced by stressful stimuli in humans and animals (DiMicco et al., 2002, Fontes et al., 2011, Fontes et al., 2017, Lisa et al., 1989, James et al., 2013, Dampney, 2015). Injection of muscimol (an agonist of GABA_A receptors) into the DMH of rats prevents increases in BP, HR, and adrenal corticotrophic hormone (ACTH) induced by air-jet stress (Stotz-Potter et al., 1996), and reduces psychogenic fever induced by the resident-intruder protocol (Kataoka et al., 2014). DMH neurons innervate brainstem regions also critically involved in the regulation of the autonomic system. The rostral ventrolateral medulla (RVLM) and raphe pallidus (RPa), which have premotor neurons that regulate sympathetic output, are considered to be key relays of the stress-induced autonomic responses. Several lines of evidence suggest that the DMH/DHA modulates RVLM and RPa activity in order to produce stress induced fight-or-flight responses, wherein the RVLM provides vasoconstrictor drive to muscle, splanchnic, and renal vascular beds while the RPa provides the cardiac drive and the sympathetic drive to BAT (Fontes et al., 2001, Cao and Morrison, 2003, Horiuchi et al., 2004, Kataoka et al., 2014, Machado et al., 2018). Evidence for a similar DMH mechanism in humans has been demonstrated by the strong resting fMRI signal coupling between the DMH and the RVLM, suggesting strong DMH-RVLM connectivity (Macefield and Henderson, 2016). In addition to its direct pathways to the

RVLM and RPa, the DMH may also regulate circadian autonomic rhythms via its projections to orexin neurons within the lateral hypothalamus (LH) (Saper, 2013). These LH orexin neurons regulate wakefulness and been shown to project to cardiovascular and respiratory centers (Geerling et al., 2003), and they also play a key role in hyperarousal during fight-or-flight responses (Kuwaki et al., 2008).

4. Autonomic support during aggressive behavior

Aggression is considered a normal component of a species' behavioral repertoire, whether it be used to ward off predators, catch prey, or compete for resources or mates, and in both animals and humans, aggressive acts are associated with a state of hyperarousal (Falkner and Lin, 2014). However, in most experimental studies that assess aggression in association with autonomic indicators, these two variables are usually temporally segregated (Haller et al., 2014). For example, human studies often examine individuals with violent criminal pasts or children with a history of antisocial behavior, but such subjects are almost always peaceful during the actual autonomic tests. Other studies that do assess real-time aggression and autonomic function usually employ symbolic or mild forms of aggressive responding, due to the obvious ethical and technical constraints of such work. However, Haller et al. devised a highly realistic model of human aggression in real world scenarios with the concomitant measurement of cardiac autonomic variables (Haller et al., 2014), involving police trainees performing operations under extreme stress. Under these conditions, high levels of inappropriate aggression (in the form of unprovoked gun use) were associated with high HR and low heart rate variability (HRV), whereas refraining from unprovoked aggression under such circumstances was associated with low HR and high HRV. HRV is a valuable tool for investigating changes in autonomic function in humans, as it is an indirect and noninvasive measure of parasympathetic and sympathetic influences on the sinoatrial node. In this regard, parasympathetic activity is indexed by the high-frequency component of HRV, while sympathetic activity can be determined through lower frequency components (Verrier and Tan, 2009). Since changes in autonomic function are highly associated with neuropsychiatric conditions, HRV has become an emerging measure that links physiological responses and cognitive features (Cohen and Benjamin, 2006, Cohen et al., 2000, Ulmer et al., 2018).

Although the autonomic aspects underlying pathological aggression (such as in individuals with violent criminal pasts) are not fully elucidated, the majority of human studies have demonstrated physiological patterns that suggest a correlation of low baseline HR with previous aggressive behavior. One meta-analysis identified that aggression is consistently associated with low resting HR and high HR reactivity in adults (18 years or older) and low HR in child aggression (1–11 years old), when data from 16 different studies was analyzed (Lorber, 2004). In this regard, and as described above, cardiac autonomic function is thought to be a valuable external indicator of internal motivational states and raises the possibility that cardiac measurements could be used as predictors of the responsiveness and potential agitation of individuals to stressful stimuli. HR is shown to be lower in aggressive adults and children, but the mechanisms underlying this phenomenon are not well elucidated. Experiments in rats do not support the idea that vagal tone/parasympathetic activity is higher in aggressive animals. A study identified that highly aggressive rats show resistance in response to vagal blockade administration and are more susceptible to stress-induced

tachyarrhythmias and pharmacologically-induced arrhythmias compared to non-aggressive rats (Carnevali et al., 2013). These results suggest that an impairment of the cardiac autonomic modulation occurs in aggression that is likely due to a vagal dysfunction, which may also be the basis for the low HR associated with aggression in humans. The results from this rodent model also suggest that the disruption of vagal regulation may facilitate the sympathetic action on the heart, leading to arrhythmic events. Supporting this argument, a study using post-weaning social isolation as a stressor model demonstrated that stressed rats exhibit autonomic overreaction, in the form of higher increases in HR and corticosterone levels, during aggression compared to non-stressed rats (Toth et al., 2011). In humans, there is evidence that suppressed vagal control and anger can contribute to arrhythmogenesis and predict premature cardiac morbidity and mortality (Tsuiji et al., 1994, Tsuiji et al., 1996, Lampert, 2010). These observations suggest that in aggressive individuals a vagal disruption associated with overreaction of the sympathetic arm of the autonomic system can increase propensity to arrhythmogenic events and even sudden death.

It is important to note that none of the studies mentioned above assessed time of day as a primary factor when measuring aggressive outcomes or autonomic output. Given that, as described above, circadian and autonomic circuits clearly influence neural processes underlying aggression, as well as each other, we argue that assessments which incorporate real-time measures of autonomic output during tests of aggressive responding at different circadian time points will be crucial to better understand such interactions. Additionally, strong evidence suggests a reciprocal influence from neural processes underlying aggression upon the circadian circuit regulating autonomic rhythms, as repeated aggressive arousal has been shown to be able to entrain HR rhythms (Tornatzky et al., 1998). Indeed, the VMHvl, which contains the neurons that promote attack behavior, sends a strong input to the DMH (Canteras et al., 1994), and this is likely the pathway by which aggression itself can influence circadian autonomic function (see Figure 1). Finally, because agitation and aggression are difficult to quantify in humans, experiments in animal models delineating the various interactions between circadian, autonomic, and aggression systems could prove vital in determining the efficacy of autonomic measures (which are readily assessed with wearable devices) as indicators of underlying agitated states in disorders associated with circadian disruption and aggressive behavior. Such work could thus lead to future examinations that reveal the mechanisms underlying these neural and behavioral pathologies.

5. Circadian and autonomic regulation of hormonal modulation and aggression

The circadian and autonomic systems both modulate a number of hormonal rhythms, and there is a large body of evidence to support various hormonal influences on aggression across the animal kingdom (Haller et al., 1998, Nelson and Trainor, 2007, Haller, 2014). The paraventricular nucleus (PV) of the hypothalamus is the primary neural regulator of the hypothalamic-pituitary-adrenal (HPA) axis (Whitnall, 1993), which plays a complementary role in modulating the fight-or-flight stress response via neuroendocrine output (Li and Sawchenko, 1998). Indeed, the circadian, autonomic, and neuroendocrine systems are highly

interconnected and each plays a role in influencing emotional state and likely behavioral outcomes of an agonistic encounter.

PV neurons expressing corticotropin-releasing-hormone (CRH) regulate the release of plasma corticosterone (Vale et al., 1981), which as mentioned above, demonstrates a pronounced circadian pattern (Engeland and Arnhold, 2005). These CRH neurons stimulate the synthesis of ACTH by the pituitary gland, which further stimulates the secretion of glucocorticoids by the adrenal cortex. This rhythm has been shown to be regulated by the SCN (Engeland and Arnhold, 2005), however the SCN sends only sparse direct projections to the PV (Vrang et al., 1995). Instead, the SCN→SPZ→DMH circuit appears to modulate the corticosterone rhythms through DMH projections to the PV (Chou et al., 2003), and perhaps by direct projections from the SPZ (Vujovic et al., 2015, Todd et al., 2018) (see Figure 1). In our assessment of the circadian control of aggression, SPZ VGAT deletions did not increase plasma corticosterone during the early resting period, when they are normally low in intact mice, in conjunction with the increase in aggression seen at this time point. However, these deletions did decrease corticosterone levels during the early active phase, when they are normally high in intact mice. This suggests that the high levels of corticosterone in intact mice during the early active period may contribute to the normally high levels of aggression seen at this time point. Indeed, studies in rats and hamsters have shown that high levels of attack behavior correspond to the rising phase of these corticosterone rhythms (Landau, 1975b, Haller et al., 2000a), and that this pattern is disrupted by adrenalectomy (Landau, 1975a, Haller et al., 2000b). As further evidence of the interconnectedness of these systems, sympathetic preganglionic fibers project to the adrenal medulla and stimulate the release of catecholamines into the bloodstream to support bodily action during the fight-or-flight response (Kvetnansky et al., 1995). Indeed, lower than normal levels of catecholamine activity have been shown to be associated with greater levels of anger and agitation in adult humans (Schwartz and Portnoy, 2017).

6. Human neural disorders characterized by aggressive behavior and circadian and autonomic dysfunction.

The autonomic system is regulated by the central nervous system and is tightly modulated by circadian rhythms, as discussed above. Because aggressive behavior is also under the influence of the circadian system, disruption of circadian rhythms may lead to facilitation of aggressive arousal. For instance, Alzheimer's disease and related dementias are associated by progressive circadian dysfunction, including "sundowning syndrome", a poorly understood clinical phenomenon associated with agitation and aggression during the early evening that has been described in medical literature for over 70 years (Bachman and Rabins, 2006, Khachiyants et al., 2011, Bedrosian and Nelson, 2013, Canevelli et al., 2016). Interestingly, we found that disrupting the SCN→SPZ→VMH pathway leads to increased behavioral aggression during the early resting phase (Todd et al., 2018), which is temporally analogous to when patients that exhibit sundowning display increased agitation and aggression. Additionally, AD and dementia patients have also been found to exhibit autonomic dysfunction (Jensen-Dahm et al., 2015, Santos et al., 2017), although this has never been examined in the context of sundowning. Similarly, other behavioral disorders

such as schizophrenia, autism, bipolar disorder, and post-traumatic stress disorder, are usually associated with autonomic imbalance as well as circadian/sleep-wake disruption, and these disorders often present with agitation and aggression (Boettger et al., 2006, Henry et al., 2010, Heiler et al., 2011, Panju et al., 2015, Tye et al., 2015, Cohen et al., 2018). In this regard, the identification of the autonomic profile of aggressive behavior may generate physiologically sensitive indices that could be used as possible markers for the neuronal conditions underlying aggressive behavior, pathologies associated with aggression, and behavioral states across the antisocial spectrum.

7. Conclusion and future directions

Interactions between the circadian and autonomic systems likely influence the behavioral outcomes of an agonistic encounter, and may shift a response closer to “fight” rather than “flight” depending on the time of day. Additionally, the degree of interconnectivity between these systems (Figure 1) underscores the emotional and behavioral implications of circadian disruption in both healthy individuals and in pathological conditions. More work in both animal models and humans is needed to better understand such interactions. For instance, strategies for determining the directionality or relative weight of these interactions could be examined in shift workers, or in simulated shift work schedules in rodent models. Indeed, while shift work has been previously associated with autonomic dysfunction (Thosar et al., 2018), we are unaware of any studies examining agitation or aggression in shift workers. Additionally, studies in transgenic mice manipulating the synaptic terminals of particular SCN or SPZ cell types in regions involved with fight-or-flight responses, such as the DMH and the PV, will be critical in order to elucidate the central circuit of autonomic system rhythmicity. Such experiments are likely to inform our understanding of the regulation of autonomic rhythms and how they influence agitation and aggression in humans, considering the evidence that SCN projections are evolutionary conserved from rodents to humans (Buijs et al., 2013). Translational work could also more definitively establish circadian autonomic measures, which can be easily assessed in humans using wearable monitors, as reliable indicators of an underlying agitated state. Most importantly, such examinations could thus prove vital to a better understanding of human neural and behavioral pathologies where aggressive outcomes can be more difficult to study and quantify.

Acknowledgements

We thank C.B. Saper for helpful comments during the preparation of this manuscript. W.D.T. was supported by Alzheimer's Association grant AARF-16-443613 and NIH grants NS084582-01A1 and HL00701-15. N.L.M. was supported by grants NS085477 and HL095491

References

- AMIR S, SHIZGAL P & ROMPRE PP 1989 Glutamate injection into the suprachiasmatic nucleus stimulates brown fat thermogenesis in the rat. *Brain Res*, 498, 140–4. [PubMed: 2571394]
- BACHMAN D & RABINS P 2006 “Sundowning” and other temporally associated agitation states in dementia patients. *Annu Rev Med*, 57, 499–511. [PubMed: 16409163]
- BEDROSIAN TA & NELSON RJ 2013 Sundowning syndrome in aging and dementia: research in mouse models. *Exp Neurol*, 243, 67–73. [PubMed: 22627081]

- BOETTGER S, HOYER D, FALKENHAHN K, KAATZ M, YERAGANI VK & BAR KJ 2006 Altered diurnal autonomic variation and reduced vagal information flow in acute schizophrenia. *Clin Neurophysiol*, 117, 2715–22. [PubMed: 17029956]
- BRAY MS, SHAW CA, MOORE MW, GARCIA RA, ZANQUETTA MM, DURGAN DJ, JEONG WJ, TSAI JY, BUGGER H, ZHANG D, ROHRWASSER A, RENNISON JH, DYCK JR, LITWIN SE, HARDIN PE, CHOW CW, CHANDLER MP, ABEL ED & YOUNG ME 2008 Disruption of the circadian clock within the cardiomyocyte influences myocardial contractile function, metabolism, and gene expression. *Am J Physiol Heart Circ Physiol*, 294, H1036–47. [PubMed: 18156197]
- BRONSARD G & BARTOLOMEI F 2013 Rhythms, rhythmicity and aggression. *J Physiol Paris*, 107, 327–34. [PubMed: 23542545]
- BUIJS RM, ESCOBAR C & SWAAB DF 2013 The circadian system and the balance of the autonomic nervous system. *Handb Clin Neurol*, 117, 173–91. [PubMed: 24095125]
- CANEVELLI M, VALLETTA M, TREBBASTONI A, SARLI G, D'ANTONIO F, TARICIOTTI L, DE LENA C & BRUNO G 2016 Sundowning in Dementia: Clinical Relevance, Pathophysiological Determinants, and Therapeutic Approaches. *Front Med (Lausanne)*, 3, 73. [PubMed: 28083535]
- CANNON W 1929 Bodily changes in pain, hunger, fear and rage, Appleton and Co.
- CANTERAS NS, SIMERLY RB & SWANSON LW 1994 Organization of projections from the ventromedial nucleus of the hypothalamus: a Phaseolus vulgaris-leucoagglutinin study in the rat. *J Comp Neurol*, 348, 41–79. [PubMed: 7814684]
- CAO WH & MORRISON SF 2003 Disinhibition of rostral raphe pallidus neurons increases cardiac sympathetic nerve activity and heart rate. *Brain Res*, 980, 1–10. [PubMed: 12865154]
- CARNEVALI L, TROMBINI M, PORTA A, MONTANO N, DE BOER SF & SGOIFO A 2013 Vagal withdrawal and susceptibility to cardiac arrhythmias in rats with high trait aggressiveness. *PLoS One*, 8, e68316. [PubMed: 23861886]
- CHOU TC, SCAMMELL TE, GOOLEY JJ, GAUS SE, SAPER CB & LU J 2003 Critical role of dorsomedial hypothalamic nucleus in a wide range of behavioral circadian rhythms. *J Neurosci*, 23, 10691–702. [PubMed: 14627654]
- COHEN H & BENJAMIN J 2006 Power spectrum analysis and cardiovascular morbidity in anxiety disorders. *Auton Neurosci*, 128, 1–8. [PubMed: 16731048]
- COHEN H, BENJAMIN J, GEVA AB, MATAR MA, KAPLAN Z & KOTLER M 2000 Autonomic dysregulation in panic disorder and in post-traumatic stress disorder: application of power spectrum analysis of heart rate variability at rest and in response to recollection of trauma or panic attacks. *Psychiatry Res*, 96, 1–13. [PubMed: 10980322]
- COHEN S, FULCHER BD, RAJARATNAM SMW, CONDUIT R, SULLIVAN JP, ST HILAIRE MA, PHILLIPS AJK, LODDENKEMPER T, KOTHARE SV, MCCONNELL K, BRAGA-KENYON P, AHEARN W, SHLESINGER A, POTTER J, BIRD F, CORNISH KM & LOCKLEY SW 2018 Sleep patterns predictive of daytime challenging behavior in individuals with low-functioning autism. *Autism Res*, 11, 391–403. [PubMed: 29197172]
- DAMPNEY RA 2015 Central mechanisms regulating coordinated cardiovascular and respiratory function during stress and arousal. *Am J Physiol Regul Integr Comp Physiol*, 309, R429–43. [PubMed: 26041109]
- DIMICCO JA, SAMUELS BC, ZARETSKAIA MV & ZARETSKY DV 2002 The dorsomedial hypothalamus and the response to stress: part renaissance, part revolution. *Pharmacol Biochem Behav*, 71, 469–80. [PubMed: 11830181]
- ENGELAND WC & ARNHOLD MM 2005 Neural circuitry in the regulation of adrenal corticosterone rhythmicity. *Endocrine*, 28, 325–32. [PubMed: 16388123]
- FALKNER AL & LIN D 2014 Recent advances in understanding the role of the hypothalamic circuit during aggression. *Front Syst Neurosci*, 8, 168. [PubMed: 25309351]
- FONTES MA, TAGAWA T, POLSON JW, CAVANAGH SJ & DAMPNEY RA 2001 Descending pathways mediating cardiovascular response from dorsomedial hypothalamic nucleus. *Am J Physiol Heart Circ Physiol*, 280, H2891–901. [PubMed: 11356650]

- FONTES MA, XAVIER CH, DE MENEZES RC & DIMICCO JA 2011 The dorsomedial hypothalamus and the central pathways involved in the cardiovascular response to emotional stress. *Neuroscience*, 184, 64–74. [PubMed: 21435377]
- FONTES MAP, FILHO ML, SANTOS MACHADO NL, DE PAULA CA, SOUZA CORDEIRO LM, XAVIER CH, MARINS FR, HENDERSON L & MACEFIELD VG 2017 Asymmetric sympathetic output: The dorsomedial hypothalamus as a potential link between emotional stress and cardiac arrhythmias. *Auton Neurosci*, 207, 22–27. [PubMed: 28131565]
- GALL AJ, TODD WD & BLUMBERG MS 2012 Development of SCN connectivity and the circadian control of arousal: a diminishing role for humoral factors? *PLoS One*, 7, e45338. [PubMed: 23028945]
- GEERLING JC, METTENLEITER TC & LOEWY AD 2003 Orexin neurons project to diverse sympathetic outflow systems. *Neuroscience*, 122, 541–50. [PubMed: 14614918]
- GOOLEY JJ, LU J, FISCHER D & SAPER CB 2003 A broad role for melanopsin in nonvisual photoreception. *J Neurosci*, 23, 7093–106. [PubMed: 12904470]
- HALLER J 2014 The glucocorticoid/aggression relationship in animals and humans: an analysis sensitive to behavioral characteristics, glucocorticoid secretion patterns, and neural mechanisms. *Curr Top Behav Neurosci*, 17, 73–109. [PubMed: 24515548]
- HALLER J, HALASZ J, MIKICS E, KRUK MR & MAKARA GB 2000a Ultradian corticosterone rhythm and the propensity to behave aggressively in male rats. *J Neuroendocrinol*, 12, 937–40. [PubMed: 11012834]
- HALLER J, MAKARA GB & KRUK MR 1998 Catecholaminergic involvement in the control of aggression: hormones, the peripheral sympathetic, and central noradrenergic systems. *Neurosci Biobehav Rev*, 22, 85–97. [PubMed: 9491941]
- HALLER J, MILLAR S, VAN DE SCHRAAF J, DE KLOET RE & KRUK MR 2000b The active phase-related increase in corticosterone and aggression are linked. *J Neuroendocrinol*, 12, 431–6. [PubMed: 10792582]
- HALLER J, RACZKEVY-DEAK G, GYIMESINE KP, SZAKMARY A, FARKAS I & VEGH J 2014 Cardiac autonomic functions and the emergence of violence in a highly realistic model of social conflict in humans. *Front Behav Neurosci*, 8, 364. [PubMed: 25374519]
- HEILER S, LEGENBAUER T, BOGEN T, JENSCH T & HOLTMANN M 2011 Severe mood dysregulation: in the “light” of circadian functioning. *Med Hypotheses*, 77, 692–5. [PubMed: 21831530]
- HENRY BL, MINASSIAN A, PAULUS MP, GEYER MA & PERRY W 2010 Heart rate variability in bipolar mania and schizophrenia. *J Psychiatr Res*, 44, 168–76. [PubMed: 19700172]
- HOOD S & AMIR S 2018 Biological Clocks and Rhythms of Anger and Aggression. *Front Behav Neurosci*, 12, 4. [PubMed: 29410618]
- HORIUCHI J, MCALLEN RM, ALLEN AM, KILLINGER S, FONTES MA & DAMPNEY RA 2004 Descending vasomotor pathways from the dorsomedial hypothalamic nucleus: role of medullary raphe and RVLM. *Am J Physiol Regul Integr Comp Physiol*, 287, R824–32. [PubMed: 15205184]
- JAMES C, MACEFIELD VG & HENDERSON LA 2013 Real-time imaging of cortical and subcortical control of muscle sympathetic nerve activity in awake human subjects. *Neuroimage*, 70, 59–65. [PubMed: 23287526]
- JENSEN-DAHM C, WALDEMAR G, STAEHELIN JENSENT, MALMQVIST L, MOELLER MM, ANDERSEN BB, HOGH P & BALLEGAARD M 2015 Autonomic Dysfunction in Patients with Mild to Moderate Alzheimer’s Disease. *J Alzheimers Dis*, 47, 681–9. [PubMed: 26401703]
- JIN X, SHEARMAN LP, WEAVER DR, ZYLKA MJ, DE VRIES GJ & REPPERT SM 1999 A molecular mechanism regulating rhythmic output from the suprachiasmatic circadian clock. *Cell*, 96, 57–68. [PubMed: 9989497]
- KATAOKA N, HIOKI H, KANEKO T & NAKAMURA K 2014 Psychological stress activates a dorsomedial hypothalamus-medullary raphe circuit driving brown adipose tissue thermogenesis and hyperthermia. *Cell Metab*, 20, 346–58. [PubMed: 24981837]
- KHACHIYANTS N, TRINKLE D, SON SJ & KIM KY 2011 Sundown syndrome in persons with dementia: an update. *Psychiatry Investig*, 8, 275–87.

- KUWAKI T, ZHANG W, NAKAMURA A & DENG BS 2008 Emotional and state-dependent modification of cardiorespiratory function: role of orexinergic neurons. *Auton Neurosci*, 142, 11–6. [PubMed: 18440283]
- KVETNANSKY R, PACAK K, FUKUHARA K, VISKUPIC E, HIREMAGALUR B, NANKOVA B, GOLDSTEIN DS, SABBAN EL & KOPIN IJ 1995 Sympathoadrenal system in stress. Interaction with the hypothalamic-pituitary-adrenocortical system. *Ann N Y Acad Sci*, 771, 131–58. [PubMed: 8597393]
- LAMPERT R 2010 Anger and ventricular arrhythmias. *Curr Opin Cardiol*, 25, 46–52. [PubMed: 19864944]
- LANDAU IT 1975a Effects of adrenalectomy on rhythmic and non-rhythmic aggressive behavior in the male golden hamster. *Physiol Behav*, 14, 775–80. [PubMed: 1237904]
- LANDAU IT 1975b Light-dark rhythms in aggressive behavior of the male golden hamster. *Physiol Behav*, 14, 767–74. [PubMed: 1237903]
- LEE H, KIM DW, REMEDIOS R, ANTHONY TE, CHANG A, MADISEN L, ZENG H & ANDERSON DJ 2014 Scalable control of mounting and attack by Esr1+ neurons in the ventromedial hypothalamus. *Nature*, 509, 627–32. [PubMed: 24739975]
- LI HY & SAWCHENKO PE 1998 Hypothalamic effector neurons and extended circuitries activated in “neurogenic” stress: a comparison of footshock effects exerted acutely, chronically, and in animals with controlled glucocorticoid levels. *J Comp Neurol*, 393, 244–66. [PubMed: 9548700]
- LIN D, BOYLE MP, DOLLAR P, LEE H, LEIN ES, PERONA P & ANDERSON DJ 2011 Functional identification of an aggression locus in the mouse hypothalamus. *Nature*, 470, 221–6. [PubMed: 21307935]
- LISA M, MARMO E, WIBLE JH, JR. & DIMICCO JA 1989 Injection of muscimol into posterior hypothalamus blocks stress-induced tachycardia. *Am J Physiol*, 257, R246–51. [PubMed: 2750964]
- LORBER MF 2004 Psychophysiology of aggression, psychopathy, and conduct problems: a meta-analysis. *Psychol Bull*, 130, 531–52. [PubMed: 15250812]
- LU J, ZHANG YH, CHOU TC, GAUS SE, ELMQUIST JK, SHIROMANI P & SAPER CB 2001 Contrasting effects of ibotenate lesions of the paraventricular nucleus and subparaventricular zone on sleep-wake cycle and temperature regulation. *J Neurosci*, 21, 4864–74. [PubMed: 11425913]
- MACEFIELD VG & HENDERSON LA 2016 “Real-time” imaging of cortical and subcortical sites of cardiovascular control: concurrent recordings of sympathetic nerve activity and fMRI in awake subjects. *J Neurophysiol*, 116, 1199–207. [PubMed: 27334958]
- MACHADO NLS, ABBOTT SGB, RESCH JM, ZHU L, ARRIGONI E, LOWELL BB, FULLER PM, FONTES MAP & SAPER CB 2018 A Glutamatergic Hypothalamomedullary Circuit Mediates Thermogenesis, but Not Heat Conservation, during Stress-Induced Hyperthermia. *Curr Biol*, 28, 2291–2301 e5. [PubMed: 30017482]
- MANFREDINI R, VANNI A, PERON L, LA CECILIA O, SMOLENSKY MH & GRASSI L 2001 Day-night variation in aggressive behavior among psychiatric inpatients. *Chronobiol Int*, 18, 503–11. [PubMed: 11475419]
- MAZUSKI C, ABEL JH, CHEN SP, HERMANSTYNE TO, JONES JR, SIMON T, DOYLE FJ, 3RD & HERZOG ED 2018 Entrainment of Circadian Rhythms Depends on Firing Rates and Neuropeptide Release of VIP SCN Neurons. *Neuron*, 99, 555–563 e5. [PubMed: 30017392]
- MOHAWK JA, GREEN CB & TAKAHASHI JS 2012 Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci*, 35, 445–62. [PubMed: 22483041]
- MORRISON SF 2004 Central pathways controlling brown adipose tissue thermogenesis. *News Physiol Sci*, 19, 67–74. [PubMed: 15016906]
- NELSON RJ & TRAINOR BC 2007 Neural mechanisms of aggression. *Nat Rev Neurosci*, 8, 536–46. [PubMed: 17585306]
- PANJU S, BRIAN J, DUPUIS A, ANAGNOSTOU E & KUSHKI A 2015 Atypical sympathetic arousal in children with autism spectrum disorder and its association with anxiety symptomatology. *Mol Autism*, 6, 64. [PubMed: 26693000]

- RAHMAN A, HASAN AU, NISHIYAMA A & KOBORI H 2018 Altered Circadian Timing System-Mediated Non-Dipping Pattern of Blood Pressure and Associated Cardiovascular Disorders in Metabolic and Kidney Diseases. *Int J Mol Sci*, 19.
- REPPERT SM & WEAVER DR 2002 Coordination of circadian timing in mammals. *Nature*, 418, 935–41. [PubMed: 12198538]
- RUSAK B, ROBERTSON HA, WISDEN W & HUNT SP 1990 Light pulses that shift rhythms induce gene expression in the suprachiasmatic nucleus. *Science*, 248, 1237–40. [PubMed: 2112267]
- SANTOS CY, MACHAN JT, WU WC & SNYDER PJ 2017 Autonomic Cardiac Function in Preclinical Alzheimer's Disease. *J Alzheimers Dis*, 59, 1057–1065. [PubMed: 28697563]
- SAPER CB 2013 The central circadian timing system. *Curr Opin Neurobiol*, 23, 747–51. [PubMed: 23706187]
- SCHROEDER A, LOH DH, JORDAN MC, ROOS KP & COLWELL CS 2011 Circadian regulation of cardiovascular function: a role for vasoactive intestinal peptide. *Am J Physiol Heart Circ Physiol*, 300, H241–50. [PubMed: 20952671]
- SCHWARTZ JA & PORTNOY J 2017 Lower catecholamine activity is associated with greater levels of anger in adults. *Int J Psychophysiol*, 120, 33–41. [PubMed: 28711699]
- SMOLENSKY MH, HERMIDA RC, REINBERG A, SACKETT-LUNDEEN L & PORTALUPPI F 2016 Circadian disruption: New clinical perspective of disease pathology and basis for chronotherapeutic intervention. *Chronobiol Int*, 33, 1101–19. [PubMed: 27308960]
- SOFIA RD & SALAMA AI 1970 Circadian rhythm for experimentally-induced aggressive behavior in mice. *Life Sci*, 9, 331–8. [PubMed: 5462761]
- STOTZ-POTTER EH, WILLIS LR & DIMICCO JA 1996 Muscimol acts in dorsomedial but not paraventricular hypothalamic nucleus to suppress cardiovascular effects of stress. *J Neurosci*, 16, 1173–9. [PubMed: 8558246]
- TAKEDA N & MAEMURA K 2016 Circadian clock and the onset of cardiovascular events. *Hypertens Res*, 39, 383–90. [PubMed: 26888119]
- THOSAR SS, BUTLER MP & SHEA SA 2018 Role of the circadian system in cardiovascular disease. *J Clin Invest*, 128, 2157–2167. [PubMed: 29856365]
- TODD WD, FENSELAU H, WANG JL, ZHANG R, MACHADO NL, VENNER A, BROADHURST RY, KAUR S, LYNAGH T, OLSON DP, LOWELL BB, FULLER PM & SAPER CB 2018 A hypothalamic circuit for the circadian control of aggression. *Nat Neurosci*, 21, 717–724. [PubMed: 29632359]
- TODD WD, GALL AJ, WEINER JA & BLUMBERG MS 2012 Distinct retinohypothalamic innervation patterns predict the developmental emergence of species-typical circadian phase preference in nocturnal Norway rats and diurnal Nile grass rats. *J Comp Neurol*, 520, 3277–92. [PubMed: 22431036]
- TORNATZKY W, COLE JC & MICZEK KA 1998 Recurrent aggressive episodes entrain ultradian heart rate and core temperature rhythms. *Physiol Behav*, 63, 845–53. [PubMed: 9618008]
- TOTH M, MIKICS E, TULOGDI A, ALICZKI M & HALLER J 2011 Post-weaning social isolation induces abnormal forms of aggression in conjunction with increased glucocorticoid and autonomic stress responses. *Horm Behav*, 60, 28–36. [PubMed: 21316368]
- TSUJI H, LARSON MG, VENDITTI FJ, JR., MANDERS ES, EVANS JC, FELDMAN CL & LEVY D 1996 Impact of reduced heart rate variability on risk for cardiac events. The Framingham Heart Study. *Circulation*, 94, 2850–5. [PubMed: 8941112]
- TSUJI H, VENDITTI FJ, JR., MANDERS ES, EVANS JC, LARSON MG, FELDMAN CL & LEVY D 1994 Reduced heart rate variability and mortality risk in an elderly cohort. The Framingham Heart Study. *Circulation*, 90, 878–83. [PubMed: 8044959]
- TYE S, VAN VOORHEES E, HU C & LINEBERRY T 2015 Preclinical perspectives on posttraumatic stress disorder criteria in DSM-5. *Harv Rev Psychiatry*, 23, 51–8. [PubMed: 25563569]
- ULMER CS, HALL MH, DENNIS PA, BECKHAM JC & GERMAIN A 2018 Posttraumatic stress disorder diagnosis is associated with reduced parasympathetic activity during sleep in US veterans and military service members of the Iraq and Afghanistan wars. *Sleep*, 41.

- VALE W, SPIESS J, RIVIER C & RIVIER J 1981 Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin. *Science*, 213, 1394–7. [PubMed: 6267699]
- VERRIER RL & JOSEPHSON ME 2009 Impact of sleep on arrhythmogenesis. *Circ Arrhythm Electrophysiol*, 2, 450–9. [PubMed: 19808502]
- VERRIER RL & TAN A 2009 Heart rate, autonomic markers, and cardiac mortality. *Heart Rhythm*, 6, S68–75. [PubMed: 19880076]
- VRANG N, LARSEN PJ & MIKKELSEN JD 1995 Direct projection from the suprachiasmatic nucleus to hypophysiotrophic corticotropin-releasing factor immunoreactive cells in the paraventricular nucleus of the hypothalamus demonstrated by means of Phaseolus vulgaris-leucoagglutinin tract tracing. *Brain Res*, 684, 61–9. [PubMed: 7583205]
- VUJOVIC N, GOOLEY JJ, JHOU TC & SAPER CB 2015 Projections from the subparaventricular zone define four channels of output from the circadian timing system. *J Comp Neurol*, 523, 2714–37. [PubMed: 26010698]
- WARREN WS, CHAMPNEY TH & CASSONE VM 1994 The suprachiasmatic nucleus controls the circadian rhythm of heart rate via the sympathetic nervous system. *Physiol Behav*, 55, 1091–9. [PubMed: 8047576]
- WATTS AG & SWANSON LW 1987 Efferent projections of the suprachiasmatic nucleus: II. Studies using retrograde transport of fluorescent dyes and simultaneous peptide immunohistochemistry in the rat. *J Comp Neurol*, 258, 230–52. [PubMed: 2438309]
- WATTS AG, SWANSON LW & SANCHEZ-WATTS G 1987 Efferent projections of the suprachiasmatic nucleus: I. Studies using anterograde transport of Phaseolus vulgaris leucoagglutinin in the rat. *J Comp Neurol*, 258, 204–29. [PubMed: 3294923]
- WELSH DK, LOGOTHETIS DE, MEISTER M & REPPERT SM 1995 Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron*, 14, 697–706. [PubMed: 7718233]
- WHITNALL MH 1993 Regulation of the hypothalamic corticotropin-releasing hormone neurosecretory system. *Prog Neurobiol*, 40, 573–629. [PubMed: 8484004]
- YANG CF, CHIANG MC, GRAY DC, PRABHAKARAN M, ALVARADO M, JUNTITI SA, UNGER EK, WELLS JA & SHAH NM 2013 Sexually dimorphic neurons in the ventromedial hypothalamus govern mating in both sexes and aggression in males. *Cell*, 153, 896–909. [PubMed: 23663785]
- YANG T, YANG CF, CHIZARI MD, MAHESWARANATHAN N, BURKE KJ, JR., BORRUS M, INOUE S, CHIANG MC, BENDER KJ, GANGULI S & SHAH NM 2017 Social Control of Hypothalamus-Mediated Male Aggression. *Neuron*, 95, 955–970 e4. [PubMed: 28757304]

Figure 1. A schematic representation of a central circuit for the circadian control of aggression and its associated autonomic support during aggressive encounters.

The suprachiasmatic nucleus (SCN) controls circadian rhythms through its projections to hypothalamic regions. Rhythms in aggression propensity are regulated by the SCN via the subparaventricular zone (SPZ), which receives the majority of SCN axonal output and projects to neurons in the ventromedial hypothalamus (VMH) that promote attack behavior, including those that express estrogen receptor 1 (Esr1). SPZ neurons also innervate the dorsomedial hypothalamus (DMH), forming a circuit by which the SCN regulates circadian rhythms of sleep-wake, locomotor activity, and feeding behavior. DMH neurons also innervate two key centers for sympathetic responses during “fight-or-flight” stress, the rostral ventrolateral medulla (RVLM) and raphe pallidus (RPa), forming a pathway that likely underlies the rhythmic autonomic support during temporal changes in aggression propensity. Similarly, DMH projections to the paraventricular nucleus (PV) are thought to modulate the corticotropin-releasing-hormone (CRH) response to stressful stimuli and also modulate circadian rhythms of corticosterone. The DMH also projects to orexin neurons in the lateral hypothalamus (LH), which have been suggested to regulate the hyperarousal during fight-or-flight responses. Finally, DMH neurons also receive innervation from the region of the VMH associated with promoting aggression, representing the possible pathway by which aggressive behavior reciprocally influences circadian autonomic regions. Hatched

lines represent relatively sparser projections from the SCN to the DMH and PV, compared to SCN projections to the SPZ.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript