Published in final edited form as:

J Int Neuropsychol Soc. 2019 February ; 25(2): 156–164. doi:10.1017/S1355617718001054.

Action control deficits in patients with Essential Tremor

Shelby Hughes¹, Daniel O. Claassen¹, Wery P.M. van den Wildenberg^{3,4}, Fenna T. Phibbs¹, Elise B. Bradley¹, Scott A. Wylie², and Nelleke C. van Wouwe^{1,2}

¹Department of Neurology, Vanderbilt University Medical Center, Nashville, TN, USA ²Department of Neurosurgery, University of Louisville, Louisville, KY, USA ³Department of Psychology, University of Amsterdam, Amsterdam, the Netherlands ⁴Amsterdam Brain and Cognition (ABC), University of Amsterdam, Amsterdam, The Netherlands

Abstract

Objective: Essential Tremor (ET) is a movement disorder characterized by action tremor which impacts motor execution. Given the disrupted cerebellar-thalamo-cortical networks in ET, we hypothesized that ET could interfere with the control mechanisms involved in regulating motor performance. The ability to inhibit or stop actions is critical for navigating many daily life situations like driving or social interactions. The current study investigated the speed of action initiation and two forms of action control, response stopping and proactive slowing in ET.

Method: Thirty-three ET patients and 25 healthy controls (HCs) completed a choice reaction task and a stop signal task, and measures of going speed, proactive slowing and stop latencies were assessed.

Results: Going speed was significantly slower in ET patients (649 ms) compared to HCs (526 ms, $F(1, 56) = 42.37$, $p < .001$, $\eta^2 = .43$), whereas proactive slowing did not differ between groups. ET patients exhibited slower stop signal reaction times (320 ms) compared to HCs (258 ms, $F(1, 56) = 15.3$, $p < .001$, $\eta^2 = .22$) and more severe motor symptoms of ET were associated with longer stopping latencies in a subset of patients ($Spearman\ rho = .48$, $p < .05$).

Conclusions: In line with previous studies, ET patients showed slower action initiation. Additionally, inhibitory control was impaired whereas proactive slowing remained intact relatively to HCs. More severe motor symptoms of ET were associated with slower stopping speed, and may reflect more progressive changes to the cerebellar-thalamo-cortical network. Future imaging studies should specify which structural and functional changes in ET can explain changes in inhibitory action control.

Keywords

Tremor; Cerebellum; Choice Behavior; Reaction Time; Movement Disorders; Inhibitory Control

---

Corresponding Author: Nelleke C. van Wouwe, Assistant Professor, Human Cognition and Neurophysiology Group, Department of Neurological Surgery, University of Louisville, 220 Abraham Flexner Way, 15th Floor, Louisville, KY 40202.

Disclosure statements

Dr. Phibbs has done consulting for Boston Scientific, Medtronic and Teva. There are no other conflicts of interest to report.

INTRODUCTION

Essential Tremor (ET) is a neurological movement disorder characterized by uncontrollable shaking or tremor that is commonly expressed during the execution of voluntary upper extremity movements (Heldman et al., 2011; Louis, 2009; Pauletti et al., 2015). Efforts to control the tremor generally exacerbate the tremor, suggesting that ET may interfere with cognitive control mechanisms involved in regulating motor performance in addition to the clear impact on motor execution processes.

We investigated the hypothesis that ET disrupts the proficiency of inhibiting motor actions independent of any effects on response execution. The ability to inhibit or stop actions is critical for navigating many dynamic, action-oriented daily life environments, and deficits in inhibitory motor control can be just as detrimental to quality of life and everyday functioning as deficits initiating and executing actions (Leroi, McDonald, Pantula, & Harbishettar, 2012).

Key brain networks involved in inhibitory action control include circuitries linking the inferior frontal cortex (IFC, often right lateralized), pre-supplementary motor area (preSMA), anterior insula, striatum, and subthalamic nucleus (Aron, Herz, Brown, Forstmann & Zaghoul, 2016; Swick et al., 2011; Cai et al., 2014). However, what aspect of stop-signal performance is reflected by each of those nodes in the circuitry in inhibitory action control (actual inhibitory process versus processing salient cues or updating of action plans), their temporal activation profile related to action cancelation, and whether the inhibition network is right-lateralized is still under debate (Aron et al., 2006; Bartoli, Aron & Tandon, 2018; Hampshire et al., 2010; Hung, Yamak, Gaillard & Arsalidou, 2018).

Inhibitory action control deficits in diseases that affect those cortico-striatal circuitries (like Parkinson's Disease, Tourette's and ADHD) have received a lot of attention (Manza, Amandola, , Tatineni, Li & Leung, 2017; Alderson, Rapport & Kofler, 2007; Wylie, Claassen, Kanoff, van Wouwe & van den Wildenberg, 2016). ET, in contrast, is more commonly associated with disruption to cerebellar-thalamo-cortical circuitries (Bagepally et al., 2012; Cerasa & Quattrone, 2016). Since the thalamus has wide spread cortical connections including areas crucial for inhibition like IFC, insula and pre SMA (O'Muircheartaigh, Keller, Barker & Richardson, 2015; Sun et al., 2018), disruptions along cerebellar-thalamo-cortical circuits as found in ET could hamper inhibitory control. Behavioral investigations of inhibitory action control in ET are relatively sparse though; most cognitive studies have focused on performance on neuropsychological tasks of working memory, language, attention, and visuospatial function (Bhalsing et al., 2014; Troster et al., 2002). Interestingly, recent fMRI and patient lesion studies (cerebellar lesions) have linked cerebellar-thalamic-cortical circuitries to the proficiency of response inhibition (measured by the stop task and Go/No-Go task; Brunamonti et al., 2014; Hirose et al., 2014). These putative linkages between cerebellar-thalamic-cortical circuitries and inhibitory action control further bolster the hypothesis that ET may interfere with these circuitries and the proficiency of inhibitory control directly.

The current study investigated the effects of ET on response going and inhibition by combining two well-established experimental cognitive paradigms. A basic two-choice reaction task quantified latencies (i.e., reaction times) of go reactions in contexts free of the need to inhibit or control these reactions. Participants also completed a stop-signal task, which required similar go reactions, but included occasional and unpredictable stop signals that required the inhibition or stopping of go reactions. The stop-signal task provides an estimate of an individual's stopping latency (i.e., stop-signal reaction time) as well as strategic slowing of go reactions (i.e., proactive slowing) induced by the expectation of having to stop on occasion. Slower stopping latencies indicate difficulties inhibiting actions, and pronounced proactive slowing is considered an adaptive consequence of difficulties inhibiting actions (i.e., go reactions are slowed to compensate for difficulties inhibiting them) (Bisset & Logan, 2011). Notably, both forms of action control have been linked to frontal-basal ganglia and cerebellar-thalamo-cortical circuitries (Aron et al., 2016; Kunimatsu, Suzuki & Tanaka 2016; Hubner et al., 2007). Given the presumed cerebellar-thalamo-cortical dysfunction caused by ET, we predicted that ET patients would show prolonged stopping latencies and exaggerated proactive slowing independent of any effects of ET on response going latencies.

METHOD

Participants

We recruited two groups of participants: Healthy Controls (HC, n=25) and ET patients (n=33). All ET participants for this study were recruited from the Movement Disorders clinic in the Neurology department at Vanderbilt University Medical Center. An age-matched HC group was recruited by community advertisement (see Table 1 for demographic information). All patients were diagnosed with ET by a board-certified neurologist. Motor severity was evaluated in 29 ET patients using the Washington Heights-Inwood Genetic Study of ET (Louis et al., 1997, WHIGET, n= 10) or the Fahn-Tolosa-Marin (FTM, Fahn, Tolosa & Marín, 1988) rating scale (n=19). The DBS movement disorders clinic switched from using the WHIGET to FTM for ET motor symptom assessment because it provides a more comprehensive evaluation of the tremor (covers midline tremor, voice, face). Thus depending on the time tested (before or after the FTM implementation), patients either have a WHIGHET or an FTM score.

As part of the routine pre- and post-DBS neuropsychological evaluations, patients completed a battery of measures spanning several cognitive domains, including: Mini Mental-State Examination (MMSE, Folstein, Folstein, & McHugh, 1975); Repeatable Battery for the Assessment of Neuropsychological Status (RBANS, Randolph, Tierney, Mohr & Chase, 1998): Line Orientation, Picture Naming, Word List; Delis-Kaplan Executive Function System (DKEFS, Delis, Kaplan, & Kramer, 2001): Trail Switching, Color Word, Verbal Fluency; Beck Depression Inventory (BDI; Beck, Steer, & Brown, 1996) and Digit Span Forward. See Table 2 for an overview of neuropsychological information.

Both ET and HC subjects were excluded if they had a medical history of (i) a neurological condition besides ET; (ii) bipolar affective disorder, schizophrenia, or other psychiatric condition known to compromise executive cognitive functions; or (iii) a medical condition

known to interfere with cognition (e.g., diabetes, pulmonary disease). They scored 25 or higher on the Mini-Mental Status Examination to rule out severe gross cognitive deficits. ET patients completed the study on their regular medication (beta blockers, anti-epileptic drugs, and occasionally benzodiazepines, see Table 3 for medication information in the ET group).

All participants had corrected-to-normal vision. The participants provided informed consent prior to participation in any study procedures in full compliance with the standards of ethical conduct in human investigation as regulated by the Vanderbilt University Institutional Review Board (IRB).

Experimental Tasks and Procedures

Participants completed two tasks: a choice reaction task (Figure 1a) and a stop-signal task (Figure 1b). All subjects completed the choice reaction task first, and subsequently performed the stop-signal task. Figure 1 presents trial examples of each task.

Choice Reaction Task: Participants first saw a fixation square in the center of the computer screen. They were instructed to focus their attention on this fixation point. Next, a series of dark gray-colored arrows appeared, one at a time, in the same central location. An arrow pointed in the rightward or leftward direction. Participants were instructed to respond to the direction of the arrow using handheld grips in their left and right hands, with left pointing arrows calling for a left hand button press and right pointing arrows calling for a right hand button press. An arrow appeared and remained on the screen until a participant pressed a button. Next, an intertrial interval randomly selected between 750–1250 milliseconds (ms) in increments of 50 ms transpired before the next arrow appeared. The fixation point was visible at all times except when an arrow was being presented on the screen. Participants were instructed to respond as quickly and accurately as possible to each arrow. The task was counterbalanced so that participants were exposed to an equal number of left and right pointing arrows (randomly presented). A practice session of 16 arrows was completed before participants completed 168 experimental trials across two blocks of 84 arrows.

Stop-Signal Task: The stop-signal task was identical to the choice reaction task with one key difference. On 30% of trials, a gray-colored arrow first appeared, but after a brief delay, the arrow changed color from gray to purple. This color change served as a stop signal that instructed participants to try to stop or inhibit their button response to the arrow's direction. Participants were instructed to respond fast to the direction of gray arrows and not to delay responding in anticipation of the presentation of stop signals. On these stop trials, the delay between the onset of the gray arrow and the color change (i.e., stop-signal delay, or SSD) adjusted dynamically using a 50 ms staircase-tracking procedure that was based on the participant's success or failure to stop on the previous stop trial (Levitt, 1971). If the participant successfully stopped on a stop trial, the SSD on the next stop trial would increase by 50 ms to increase the difficulty of stopping, while a failure to stop a reaction on a stop trial led to a shortening of the SSD on the next stop trial in order to make stopping easier. The dynamic tracking procedure converged on an overall 50% response stopping rate on stop trials, which is optimal for obtaining a reliable estimate of SSRT (Band, van der Molen,

& Logan, 2003). For the stop-signal task, participants completed a practice session of 40 trials followed by 3 blocks of 90 experimental trials comprised of 189 go trials and 81 stop trials.

Data Analyses

Our first set of analyses compared *go* RT and accuracy rates from the choice reaction task with *go* RT times and accuracy rates from the stop-signal task. RT slowing in the latter compared to the former provided a measure of *proactive slowing* in situations when the need to inhibit action may occur. We performed repeated-measures ANOVAs with *Group* (ET and HC) as a between-subject factor and *Task* (Choice reaction time task, Stop-signal task) as a within-subjects factor.

The second set of analyses used ANOVAs to compare response inhibition latencies (SSRT) as a function of *Group* (ET and HC). SSRT was estimated using the well-established horse-race model and integration method (Logan, 1994). The horse-race model assumes an independent race between the go and stop processes. The onset of the stop process is experimentally controlled by the length of the SSD. The finishing time of the stop process is inferred from the time point at which the internal response to the stop-signal occurs and subtracting the SSD from this point. The time point in the go reaction time (GoRT) distribution that corresponds to P (failure-to-stop) is assumed to equal (SSD + SSRT). Individual GoRTs were rank ordered and the *n*th GoRT was selected, where *n* is the point at which P (failure-to-stop). The mean SSD was then subtracted from this finishing time to obtain an estimate of SSRT.

We first verified a few critical assumptions of the method requiring that the probability of stopping responses on stop-signal trials approximated 50% and that failed inhibition trials were associated with shorter RT than average go trials (Band, van der Molen & Logan, 2003; Logan, 1994).

Additionally, in the patient group, we correlated ratings of clinical motor impairments, as measured by the FTM, with measures of going and action control (SSRT and proactive slowing), corrected for age (by using partial correlation analyses). Patients with WHIGET scores were not included in the correlation due to the small sample size (*n*=10) and the absence of a valid transformation between the motor scales to allow combining the scores.

RESULTS

Go Trials and Proactive Slowing

Mean go RT in ET patients (649 ms) was significantly slower than in HC (526 ms), (*Group*: $F(1,56) = 42.37, p < .001, \eta^2 = .43$)¹ and patients made more omission errors (5%) on Go trials compared to HC (0.8%, $F(1,56) = 7.17, p = .01, \eta^2 = .11$); see Figure 2a for go RTs. Proactive slowing was revealed by reactions to go stimuli that slowed significantly in a context that included the unpredictable need to stop reactions abruptly (stop-signal task, 657 ms)

¹Note that go RTs were not significantly different from a normal distribution in either group, HC: Shapiro-Wilk = .95, *p* = .15, ET: Shapiro-Wilk = .98, *p* = .9. Variances were not significantly different between groups either, Levene's statistic = .10, *p* = .92.

compared to a context that did not require stopping (choice reaction task, 518 ms, *Task*, $F(1, 56) = 152.30$, $p < .001$, $\eta^2 = .73$). The magnitude of proactive slowing did not differ between ET patients and controls (*Group x Task*, $F(1, 56) = .74$, $p = .39$, $\eta^2 = .01$). No effects of *Group*, *Task* or an interaction between *Group* and *Task* were found for mean accuracy rates (*all* $F_s < 1.04$, $p_s > .31$).

Stop' Signal Dynamics

The tracking algorithm controlling stop-signal onset successfully converged to overall stop percentages near 50% (min 43% and max 56% for both groups) and was similar across groups (ET: 48%, HC: 48%; $F(1, 56) = .02$, $p = .89$, $\eta^2 < .001$). The SSD was shorter for HC (304 ms) compared to ET (383 ms, $F(1, 56) = 6.0$, $p = .02$, $\eta^2 = .10$).

Also consistent with a key assumption of the horse-race model (Logan, 1994), mean RT for failed stop trials (i.e., those that escaped inhibition) was shorter than the overall mean go RT in all subjects (570 vs. 657 ms; $F(1, 56) = 458.45$, $p < .001$, $\eta^2 = .89$). The difference between GoRT and Signal respond RT was significant in both groups: in ET patients it was 94 ms ($t(32) = 18.9$, $p < .001$, *Cohen's d* = 1.03, min 49 ms-max 162 ms,) and in HC = 81 ms ($t(24) = 12.05$, $p < .001$, *Cohen's d* = .97, min 17 ms-max 150 ms). These patterns satisfy key assumptions needed to reliably estimate SSRT.

Mean SSRT2, shown in Figure 2b, differed between the groups ($F(1, 56) = 15.3$, $p < .001$, $\eta^2 = .22$). Specifically, SSRT was significantly longer among ET patients (320 ms) compared to HC (258 ms). In line with the horse-race model, predictions that going and stopping processes are independent, go RT did not correlate with SSRT across all subjects (*Spearman's rho* = 0.08, $p = 0.56$), nor did go RT correlate with SSRT within each group separately (*Spearman's rho's* = <.3, $p_s > .13$)

Stop task performance and relation with clinical variables

For a subset of ET patients (patients with a motor symptom severity score, the FTM score, $n = 19$) we correlated go RT, proactive slowing, and SSRT with motor severity, corrected for age. FTM scores were not related to go RT (*Spearman's rho* = -.04, $p = .87$) or proactive slowing (*Spearman's rho* = -.38, $p = .11$). Increased severity of motor impairment (higher FTM scores) was associated with longer SSRT (*Spearman's rho* = .48, $p < .05$), see Figure 3a.

Additionally, across all 58 participants (both patients and HC), less proactive slowing was associated with impaired stopping, i.e. prolonged SSRT (corrected for age, *Spearman's rho* = -.33, $p < .05$), see Figure 3b. When looking at the groups separately, we found similar correlations between proactive slowing and SSRT (HC *Spearman's rho* = -.50, $p < .05$; ET *Spearman's rho* = -.42, $p < .05$)

²When separating go RT and SSRT by handedness in the ET group, no difference was found between left and right hand performance ($t_{Go}(32) = 1.12$, $p = .27$; $t_{SSRT}(32) = 1.04$, $p = .31$), nor were the left and right FTM motor scores different ($t_{FTM}(18) = .37$, $p = .72$), confirming the literature suggesting that ET symptoms are generally bilateral (Benito-Leon & Louis, 2006). All analyses were thus based on go RTs and SSRTs collapsed across hands.

DISCUSSION

ET is a neurological movement disorder characterized by action tremor that interferes with the execution of voluntary movements. We hypothesized that cerebellar-thalamo-cortical circuitry dysfunction produced by ET would also impair the ability to inhibit or stop voluntary movements. The current study demonstrated that ET independently slows going and stopping latencies, revealing that ET compromises both the execution and the inhibition of movement.

ET slowed responses to go stimuli across both reaction contexts. In the choice reaction task, reactions to go stimuli were on average 123 milliseconds slower among ET patients compared to controls, revealing a clear impact of ET on response speed. When occasional stop signals were introduced in the stop-signal task, both ET and controls slowed their reactions to go stimuli, consistent with a strategic, proactive slowing of go reactions in order to accommodate the expected but unpredictable need to stop reactions. While this form of proactive slowing has been linked to cerebellar dysfunction, we did not observe differential magnitudes of proactive slowing in ET compared to controls. Both groups slowed go reactions by about 140 milliseconds on average. This suggests that ET patients were able to make normal strategic adjustments in response speed to accommodate demands on inhibitory action control.

Independent of the slowing of go reactions, ET selectively impaired stopping latency. Moreover, ET patients with more severe tremor ratings (i.e., higher FTM motor scores) showed the most pronounced slowing of stopping latencies. Notably, tremor severity was unrelated to the magnitude of slowing of go reactions and of proactive slowing, suggesting a more direct linkage between tremor severity and inhibitory action control. As an interesting side note, given that ET patients show intact proactive slowing, but within the ET group larger proactive slowing was associated with faster stopping latencies, one practical strategy in ET would be to promote greater proactive slowing in functional situations that place greater demands on motor control. This certainly represents an intriguing area for future investigation. Additionally, future studies that systematically manipulate going speeds or inhibitory demands would inform our understanding of how strategic adjustments in control impact motor performance in ET (Bissett & Logan, 2011; van Wouwe et al., 2014; Wylie et al., 2009).

Potential neural mechanism of inhibitory control changes in ET

Inhibitory control impairments and exacerbated tradeoffs between going and stopping mechanisms have been well reported in diseases characterized by cortico-striatal network dysfunction (e.g. Parkinson's Disease, Tourette's and ADHD; Manza et al., 2017; Alderson, Rapport & Kofler, 2007; Wylie et al., 2016). The underlying neuropathology in ET is less clearly understood; it includes degradation of cerebellar gray and white matter along with abnormal signaling in cerebellar-thalamic-cortical circuitries (Cerasa & Quattrone, 2016). Although inhibitory control has repeatedly been linked to activity in frontal-striatal networks including the preSMA, IFC, striatum, premotor cortex, anterior insula and STN (Aron et al., 2016; Mirabella, Pani, Ferraina, 2011, ref), more recent imaging work with healthy adults has suggested that the cerebellum might be critically involved in inhibitory control as well

(Hirose et al., 2014). ET patients in the current study showed prolonged inhibition latencies, which is in line with the presumed cerebellar-thalamo-cortical dysfunction caused by ET. Future imaging studies in ET should investigate whether structural or functional changes in the frontal-striatal circuitries or the cerebellar-thalamo-cortical circuitries are linked to impaired inhibitory control in ET.

How might we understand the link between tremor severity in ET and dysfunctional inhibitory control? Bhalsing et al. (2014) found that hampered performance on Wisconsin Card Sorting Test and Stroop interference correlated with gray matter volume loss of anterior cingulate, right inferior frontal gyrus and right inferior parietal lobe in ET patients. Bagepally et al. (2012) also found a positive relation between the amount of frontal cortical and striatal gray matter atrophy and disease severity in ET patients (FTM tremor severity). Cerebellar degeneration might additionally affect functional connectivity in ET patients. For example, Schnitzler and colleagues (Schnitzler, Munks, Butz, Timmerman & Gross, 2009) identified a network of brain areas (including M1, premotor cortex, thalamus, cerebellum and brainstem) that show coupled oscillatory activity at the frequency of the tremor in ET patients, suggesting a pathological alteration of communication between areas that are also important to inhibitory control (Mirabella, Pani, Ferraina, 2011).

Benito-Leon et al. (2016) showed that resting state activity changes across a range of functional networks (both increased and decreased activity), suggesting dysfunctional networks in ET patients compared to HC. More specifically, enhanced activity in the resting state networks (the default mode network and frontoparietal network) correlated with increased disease severity, disease duration, and reduced cognitive ability; i.e. reduced performance on tasks measuring executive (Stroop interference and frontal assessment battery), attention (WAIS digit span), verbal memory (word list learning and recall) and visuo-spatial ability (line orientation). Consistent with the above described imaging studies, the relationship between cognitive impairment and disease severity in ET in our study could be mediated by gray matter volume reductions in cerebellum and fronto-cortical areas, and enhanced activity in resting state networks (Bhalsing et al., 2014; Bagepally et al., 2012; Benito-Leon et al., 2016).

In contrast to inhibitory control deficits, proactive slowing was similar for ET patients and HC. Previous work suggests that the cerebellum, an area crucially impacted by ET, might play a role in proactive control processes (Brunamonti et al., 2014). Olivito et al. (2017) found that patients with focal cerebellar atrophy showed increased post error slowing (indicative of proactive trial-by-trial adjustments) and increased stop latencies when compared to healthy controls thus implicating the cerebellum in both proactive control and reactive motor inhibition. Future fMRI studies could provide insight in which cerebellar-thalamo-cortical loops are critically involved in proactive control and how these networks are recruited in ET.

Limitations and future studies

In the current study, data collection in ET was performed in participants who remained on medications for tremor. We did not test cognitive performance OFF medication, thus, we cannot account for possible medication effects on action control. However, a post-hoc

comparison between ET patients using anti-epileptics ($n=23$), which may alter cognitive performance (Kay, Schwartz, Wingertzahn, Jayawardena & Rosenberg, 2016), versus patients that did not use this medication ($n=10$), did not reveal a difference in SSRT ($F(1, 32) = .90, p=.35$) or proactive slowing ($(F(1,32)=.73, p=.40)$).

Additionally, we acknowledge that the use of different motor scales in our sample is a limitation and reduced the sample size for the correlational analysis between motor symptoms and cognitive measures. Future studies should replicate the current findings in a larger sample to improve generalizability.

Another limitation with respect to generalizability of the current study was the age range of the patients. Based on a meta-analysis of 51 studies on the role of the cerebellum in ET neuropathology, Cerasa et al. (2016) concluded that the age of ET patients in most studies ranged from 53–70 years with the exception of a few studies including younger patients. Future studies should replicate ET effects on inhibitory control in a wider age range (including more younger patients) to improve generalizability.

Moreover, the patient group tended to have a slightly different sex distribution compared to the control group (more males in patient group), although there is no evidence that this should strongly impact stop signal performance (for a meta-analysis on sex effects on the SSRT see Lipszyc & Schachar, 2010).

Finally, it is unlikely that physical manipulation of the response device impacted performance. Responding to the computer task required a simple button press while the button grips were resting comfortably in the participant's lap; no overt limb movement was needed. Additionally, tremor severity did not correlate with going speed in either left or right hand.

Given the results of the current study, and previous studies that examined functional and structural changes in ET, there are multiple directions for further study. Functional imaging studies investigating the involvement of the cerebellar-thalamo-cortical loops in ET during performance of an inhibitory control paradigm like a stop-signal task could provide insight into the underlying neural mechanism that explains changes in inhibitory control in ET. Longitudinal studies could also help advance our understanding of ET's disease pathology (disease severity) in relation to changes in going and stopping latencies. Future studies might compare the effects of medical and surgical interventions (i.e., deep-brain stimulation of the ventral intermediate nucleus of the thalamus) on inhibitory action control in ET (see also van den Wildenberg et al., 2006).

Conclusion

The current study demonstrates that ET patients show a selective impairment in their ability to inhibit, or stop, movements. The ability to stop was also linked to tremor severity and independent of generalized response slowing. Clinically, an improved understanding of how ET impacts action control processes might enable optimization of therapeutic strategies and interventions.

Acknowledgements

We thank Bert van Beek for programming the computer task. The authors would also like to acknowledge support with patient testing from Maxim Turchan.

Funding

This work was supported by the National Institutes of Health RO1NS097783-01 (to D.O.C)

References

- Alderson RM, Rapport MD, & Kofler MJ (2007). Attention-deficit/hyperactivity disorder and behavioral inhibition: a meta-analytic review of the stop-signal paradigm. *J Abnorm Child Psychol*, 35(5), 745–758. doi:10.1007/s10802-007-9131-6 [PubMed: 17668315]
- Aron A, Herz D, Brown P, Forstmann B, & Zaghoul K (2016). Frontosubthalamic Circuits for Control of Action and Cognition. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 36(45), 11489–11495. [PubMed: 27911752]
- Bagepally BS, Bhatt MD, Chandran V, Saini J, Bharath RD, Vasudev MK, ... Pal PK (2012). Decrease in cerebral and cerebellar gray matter in essential tremor: a voxel-based morphometric analysis under 3T MRI. *J Neuroimaging*, 22(3), 275–278. doi:10.1111/j.1552-6569.2011.00598.x [PubMed: 21447032]
- Band GPH, van der Molen MW, & Logan GD (2003). Horse-race model simulations of the stop-signal procedure. *Acta Psychologica*, 112(2), 105–142. [PubMed: 12521663]
- Bartoli E, Aron AR, & Tandon N (2018). Topography and timing of activity in right inferior frontal cortex and anterior insula for stopping movement. *Hum Brain Mapp*, 39(1), 189–203. [PubMed: 29024235]
- Beck AT, Steer RA, & Brown GK (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Benito-León J, Louis E, Romero J, Hernández-Tamames J, Manzanedo E, Álvarez-Linera J, ... Onofrij M (2015). Altered Functional Connectivity in Essential Tremor. *Medicine*, 94(49), Medicine, 2015, Vol.94(49).
- Benito-León J, & Louis ED (2006). Essential tremor: emerging views of a common disorder. *Nat Clin Pract Neurol*, 2(12), 666–678 [PubMed: 17117170]
- Bhalsing K, Upadhyay N, Kumar K, Saini J, Yadav R, Gupta A, & Pal P (2014). Association between cortical volume loss and cognitive impairments in essential tremor. *European Journal of Neurology*, 21(6), 874–883. [PubMed: 24612409]
- Bissett PG, & Logan GD (2011). Balancing cognitive demands: control adjustments in the stop-signal paradigm. *J Exp Psychol Learn Mem Cogn*, 37(2), 392–404. doi:10.1037/a0021800 [PubMed: 21171806]
- Brunamonti E, Chiricozzi FR, Clausi S, Olivito G, Giusti MA, Molinari M, ... Leggio M (2014). Cerebellar damage impairs executive control and monitoring of movement generation. *PLoS One*, 9(1), e85997. doi:10.1371/journal.pone.0085997 [PubMed: 24465830]
- Cai W, Ryali S, Chen T, Li CS, & Menon V (2014). Dissociable roles of right inferior frontal cortex and anterior insula in inhibitory control: evidence from intrinsic and task-related functional parcellation, connectivity, and response profile analyses across multiple datasets. *J Neurosci*, 34(44), 14652–14667. [PubMed: 25355218]
- Cerasa A, & Quattrone A (2016). Linking Essential Tremor to the Cerebellum-Neuroimaging Evidence. *Cerebellum*, 15(3), 263–275. doi:10.1007/s12311-015-0739-8 [PubMed: 26626626]
- Delis DC, Kaplan E and Kramer J 2001 *Delis Kaplan Executive Function System.*, San Antonio, TX: The Psychological Corporation.
- Fahn S, Tolosa E, Marín C. Clinical rating scale for tremor In: Jankovic J, Tolosa E, eds. *Parkinson's disease and movement disorders*. Baltimore: Urban and Schwarzenberg, 1988:225–34.
- Folstein MF, Folstein SE, & McHugh PR (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189–198. [PubMed: 1202204]

- Hampshire A, Chamberlain SR, Monti MM, Duncan J, & Owen AM (2010). The role of the right inferior frontal gyrus: inhibition and attentional control. *Neuroimage*, 50(3), 1313–1319. [PubMed: 20056157]
- Heldman D, Jankovic J, Vaillancourt D, Prodoehl J, Elble R, & Giuffrida J (2011). Essential tremor quantification during activities of daily living. *Parkinsonism & Related Disorders*, 17(7), 537–542. doi:10.1016/j.parkreldis.2011.04.017 [PubMed: 21570891]
- Hirose S, Jimura K, Kunimatsu A, Abe O, Ohtomo K, Miyashita Y, & Konishi S (2014). Changes in cerebro-cerebellar interaction during response inhibition after performance improvement. *Neuroimage*, 99, 142–148. doi:10.1016/j.neuroimage.2014.05.007 [PubMed: 24830836]
- Hübner J, Sprenger A, Klein C, Hagenah J, Rambold H, Zühlke C, ... Helmchen C (2007). Eye movement abnormalities in spinocerebellar ataxia type 17 (SCA17). *Neurology*, 69(11), 1160–1168. doi:10.1212/01.wnl.0000276958.91986.89 [PubMed: 17846415]
- Hung Y, Gaillard SL, Yarmak P, & Arsalidou M (2018). Dissociations of cognitive inhibition, response inhibition, and emotional interference: Voxelwise ALE meta-analyses of fMRI studies. *Hum Brain Mapp*.
- Kay GG, Schwartz HI, Wingertzahn MA, Jayawardena S, & Rosenberg RP (2016). Next-day residual effects of gabapentin, diphenhydramine, and triazolam on simulated driving performance in healthy volunteers: a phase 3, randomized, double-blind, placebo-controlled, crossover trial. *Hum Psychopharmacol*, 31(3), 217–226. doi:10.1002/hup.2530 [PubMed: 27018419]
- Kunimatsu J, Suzuki TW, & Tanaka M (2016). Implications of Lateral Cerebellum in Proactive Control of Saccades. *J Neurosci*, 36(26), 7066–7074. doi:10.1523/JNEUROSCI.0733-16.2016 [PubMed: 27358462]
- Leroi I, McDonald K, Pantula H, & Harbisetar V (2012). Cognitive impairment in Parkinson disease: impact on quality of life, disability, and caregiver burden. *J Geriatr Psychiatry Neurol*, 25(4), 208–214. doi:10.1177/0891988712464823 [PubMed: 23172765]
- Levitt H (1971). Transformed up-down methods in psychoacoustics. *The Journal of the Acoustical Society of America*, 49(2), Suppl 2:467–477.
- Lipszyc J, & Schachar R (2010). Inhibitory control and psychopathology: A meta-analysis of studies using the Stop-Signal task. *Journal of the International Neuropsychological Society*, 16, 1064–1076. [PubMed: 20719043]
- Logan GD (1994). Spatial attention and the apprehension of spatial relations. *Journal of Experimental Psychology: Human Perception and Performance*, 20, 1015–1036. [PubMed: 7964527]
- Louis ED (2009). Essential tremors: a family of neurodegenerative disorders? *Arch Neurol*, 66(10), 1202–1208. doi:10.1001/archneurol.2009.217 [PubMed: 19822775]
- Louis ED, Ottman R, Ford B, Pullman S, Martinez M, Fahn S, & Hauser WA (1997). The Washington Heights-Inwood Genetic Study of Essential Tremor: methodologic issues in essential-tremor research. *Neuroepidemiology*, 16(3), 124–133. [PubMed: 9159767]
- Manza P, Amandola M, Tatineni V, Li CR, & Leung HC (2017). Response inhibition in Parkinson's disease: a meta-analysis of dopaminergic medication and disease duration effects. *NPJ Parkinsons Dis*, 3, 23. doi:10.1038/s41531-017-0024-2 [PubMed: 28702504]
- Mirabella G, Pani P, & Ferraina S (2011). Neural correlates of cognitive control of reaching movements in the dorsal premotor cortex of rhesus monkeys. *J Neurophysiol*, 106(3), 1454–1466. doi:10.1152/jn.00995.2010 [PubMed: 21697448]
- Olivito G, Brunamonti E, Clausi S, Pani P, Chiricozzi FR, Giamundo M, ... Ferraina S (2017). Atrophic degeneration of cerebellum impairs both the reactive and the proactive control of movement in the stop signal paradigm. *Exp Brain Res*, 235(10), 2971–2981. doi:10.1007/s00221-017-5027-z [PubMed: 28717819]
- O'Muirheartaigh J, Keller SS, Barker GJ, & Richardson MP (2015). White Matter Connectivity of the Thalamus Delineates the Functional Architecture of Competing Thalamocortical Systems. *Cereb Cortex*, 25(11), 4477–4489. [PubMed: 25899706]
- Pauletti C, Mannarelli D, De Lucia MC, Locuratolo N, Currà A, Missori P, ... Fattapposta F (2015). Selective attentional deficit in essential tremor: Evidence from the attention network test. *Parkinsonism Relat Disord*, 21(11), 1306–1311. doi:10.1016/j.parkreldis.2015.08.035 [PubMed: 26365776]

- Randolph C, Tierney MC, Mohr E, & Chase TN (1998). The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): preliminary clinical validity. *J Clin Exp Neuropsychol*, 20(3), 310–319. [PubMed: 9845158]
- Schnitzler A, Münks C, Butz M, Timmermann L, & Gross J (2009). Synchronized brain network associated with essential tremor as revealed by magnetoencephalography. *Mov Disord*, 24(11), 1629–1635. [PubMed: 19514010]
- Sun C, Wang Y, Cui R, Wu C, Li X, & Bao Y (2018). Human Thalamic-Prefrontal Peduncle Connectivity Revealed by Diffusion Spectrum Imaging Fiber Tracking. *Front Neuroanat*, 12, 24. [PubMed: 29719502]
- Swick D, Ashley V, & Turken U (2011). Are the neural correlates of stopping and not going identical? Quantitative meta-analysis of two response inhibition tasks. *Neuroimage*, 56(3), 1655–1665. [PubMed: 21376819]
- Tröster A, Woods S, Fields J, Lyons K, Pahwa R, Higginson C, & Koller W (2002). Neuropsychological deficits in essential tremor: An expression of cerebello-thalamo- cortical pathophysiology? *European Journal of Neurology*, 9(2), 143–151. [PubMed: 11882055]
- van den Wildenberg WPM, van Boxtel GJM, van der Molen MW, Bosch DA, Speelman JD, & Brunia CHM (2006). Stimulation of the subthalamic region facilitates the selection and inhibition of motor responses in Parkinson's disease. *Journal of Cognitive Neuroscience*, 18, 626–636. [PubMed: 16768365]
- van Wouwe NC, van den Wildenberg WP, Claassen DO, Kanoff K, Bashore TR, & Wylie SA (2014). Speed pressure in conflict situations impedes inhibitory action control in Parkinson's disease. *Biol Psychol*, 101, 44–60. doi:10.1016/j.biopsycho.2014.07.002 [PubMed: 25017503]
- Wylie SA, Claassen DO, Kanoff KE, van Wouwe NC, & van den Wildenberg WP (2016). Stopping Manual and Vocal Actions in Tourette's Syndrome. *J Neuropsychiatry Clin Neurosci*, appineuropsych15110387. doi:10.1176/appi.neuropsych.15110387
- Wylie SA, van den Wildenberg WP, Ridderinkhof KR, Bashore TR, Powell VD, Manning CA, & Wooten GF (2009). The effect of speed-accuracy strategy on response interference control in Parkinson's disease. *Neuropsychologia*, 47(8–9), 1844–1853. doi:10.1016/j.neuropsychologia.2009.02.025 [PubMed: 19428416]

Figure 1.

Trial examples of **a)** the choice reaction task where participants manually respond to the direction of dark-gray arrows presented on the screen with a left or right button press, and **b)** the stop-signal task in which subjects were instructed to respond to the direction of the dark-gray arrow with a left or right button press and to stop their response when the arrow turned purple (30% of the trials).

Figure 2.

a) Mean go RT for the choice reaction task and for the stop-signal task for ET patients and HC. b) Mean SSRT for ET patients and HC. Error bars represent standard errors.

Figure 3.

a) Correlation of SSRT in ET patients and motor severity as measured by the Fahn-Tolosa-Marín (FTM) rating scale **b).** Correlation of SSRT and proactive slowing in both ET patients and HC. Note that for clarity and interpretation of the scales, the values represented in the graph are the original scales (not corrected for age).

Table 1.

Characteristics of healthy control (HC) and Essential Tremor (ET) subjects (age matched), mean and standard deviation.

	HC (N=25)	ET (N=33)	p-value	Test statistic
Age (years)	64.1 (8.2)	66.57(5.3)	.21	t(56) = 1.27
Education (years)	15.8(3.9)	13.1 (2.5)	.003	t(51) =3.1
Sex (M:F)	10 (40%)	19 (57.6%)	.08	$\chi^2 = 3.16$
MMSE	29.6(.87)	28.6 (1.5)	.002	t(56) = 3.2
Disease Duration (years)		26.5(15.6)		
BDI (n= 29)		9.8 (7.8)		
WHIGET left (n = 10/33)		14.9 (5.0)		
WHIGET right		11.6 (4.8)		
FTM left (n= 19/33)		16.5 (12.6)		
FTM right		18.0 (12.1)		

Table 2.

Neuropsychological characteristics of Essential Tremor subjects (scaled scores, mean and SD). Across neuropsychological functions, subject's performance ranged from low average (6–8) to average (8 and up).

	Scaled score mean (SD)
RBANS naming	10.8 (1.3)
RBANS word learning (n=28/33)	8.1 (3.1)
RBANS word learning recall (n=29/33)	8.5 (2.3)
RBANS judgement of line	9.1 (3.1)
RBANS visual discrimination	8.2 (1.7)
DKEFS trails switching	8.9 (2.8)
DKEFS color-word naming	8.4 (3.1)
DKEFS color-word reading	7.0 (3.8)
DKEFS color-word inhibition	8.1 (3.8)
DKEFS color-word switching	8.6 (4.0)
DKEFS fluency phonemic (letters)	7.9 (3.2)
DKEFS fluency semantic (categories)	9.2 (3.1)
WAIS-IV Digit span forward	9.5 (3.3)

RBANS=Repeatable Battery for the Assessment of Neuropsychological Status; DKEFS=Delis-Kaplan Executive Function; WAIS=Wechsler Adult Intelligence Scale

Table 3.

Frequency of medication use in the Essential Tremor subjects.

	Proportion of patients on medication
Anti-epileptica	23/33
Beta-blocker	15/33
Benzodiazepine	8/33
Antipsychotics	4/33
Antidepressants	11/33
Levodopa	5/33
Muscle relaxant	3/33
Opioid	6/33