Comparative transcriptomics reveals potential genes involved in the vegetative growth of *Morchella importuna*

Wei Liu^{1,2} · Yingli Cai³ · Peixin He⁴ · Lianfu Chen^{1,2} · Yinbing Bian^{1,2}

Received: 21 September 2018 / Accepted: 2 February 2019 / Published online: 13 February 2019
© King Abdulaziz City for Science and Technology 2019

Abstract

True morels (*Morchella* spp.) are edible, medicinal mushrooms which have recently been artificially cultivated in China but stable production remains a problem. Here, we describe complete and comprehensive transcriptome of *Morchella importuna* at the stages of vegetative mycelium (VM), initial sclerotium (IS) and mature sclerotium (MS) by deep transcriptional sequencing and *de novo* assembly for the first time and which will potentially provide useful information for improving its cultivation. A total of 26,496 genes were identified with a contig N50 length of 1763 bp and an average length of over 1064 bp. Additionally, 11,957 open reading frames (ORFs) were predicted and 9676 of them (80.9%) were annotated. The 2605 differentially expressed genes (DEGs) identified by gene expression clustering were mainly involved with energy metabolism and could be divided into three broad clusters, of which genes in cluster_1 and cluster_2 were involved in the metabolic process of carbohydrate, polysaccharide, hydrolase, caprolactam, beta-galactosidase, and disaccharide, respectively. Genes in cluster_3 were the largest category, mainly identified with the catalytic activity and transporter activity. Overall, the enzymes involved in the carbohydrate metabolism were highly expressed, and the CAZyme (carbohydrate-active enzyme) genes were significantly expressed within cluster_3. For expression verification, 16 CAZyme genes were selected for qRT-PCR, and the results suggested that the catabolism of carbohydrates occurs mainly in the vegetative mycelium stage, and the anabolism of the energy-rich substances is the main event of mycelial growth and sclerotial morphogenesis of *M. importuna*.

Keyword Morel · Carbohydrate metabolism · Differential expression analysis · CAZymes, qRT-PCR

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13205-019-1614-y>) contains supplementary material, which is available to authorized users.

✉ Peixin He
peixinhe191@163.com

✉ Yinbing Bian
bianyb.123@163.com

¹ Institute of Applied Mycology, Huazhong Agricultural University, 430070 Wuhan, China

² Key Laboratory of Agro-Microbial Resource Comprehensive Utilization, Ministry of Agriculture, Huazhong Agricultural University, 430070 Wuhan, Hubei, China

³ Institute of Vegetable, Wuhan Academy of Agricultural Sciences, 430070 Wuhan, China

⁴ School of Food and Biological Engineering, Zhengzhou University of Light Industry, 450001 Zhengzhou, China

Introduction

True morels (*Morchella* spp., Pezizales, Ascomycota) are edible medicinal mushrooms. They are appreciated for their savory flavor and multiple bioactivities, including anti-oxidative, anti-inflammatory, anti-microbial, immunostimulatory and anti-tumor properties (He et al. 2012; Heleno et al. 2013; Rosbach et al. 2017; Su et al. 2013; Tietel and Masaphy 2018; Vieira et al. 2016). Efforts to domesticate and cultivate morel by enthusiasts and scientists all over the world failed until 2012 when field cultivation of *Morchella* was achieved in China (He et al. 2017; Liu et al. 2017) based on the pioneering work of Ower (1982). The successful cultivation of morel in China was achieved due to two factors: the acquisition of suitable varieties (*M. importuna*, *M. sextelata* and *M. septimelata*) that are prone to fructify (He et al. 2015a, b) and the implementation of exogenous nutrient “feeding technology”, which provided sufficient nutrients for the mycelial network or sclerotial cells in the soil to support sexual reproduction in the later period (Liu et al. 2017;

Tan 2016). The potential mechanism of the “feeding technology” may be the initial accumulation of energy material in the foreign nutrition bag and its subsequent transfer to the mycelial network and sclerotium-like cells in the soil via a high-energy nutritional potential “flow” to a low-energy nutritional potential. However, this hypothesis has not been experimentally proven.

Like many other fungi, sclerotium formation is an essential phase during the life cycle of *Morchella* spp. (Alvarado-Castillo et al. 2008, 2011; Volk and Leonard 1989). Sclerotium is a highly resistant and asexual dormant structure produced by fungi, which can avoid adverse physiological and environmental conditions. Unlike the sclerotium of *Sclerotinia sclerotiorum* and other plant pathogenic fungi, the sclerotium of *Morchella* spp. has no obvious differentiation of tissue structure, so it is called a “pseudosclerotium” (Volk and Leonard 1990). The sclerotia of *Morchella* are assumed to serve as nutrient storage organs that can tolerate adverse conditions such as low temperature and desiccation, thus enhancing survival over winter (Alvarado-Castillo et al. 2011; Liu et al. 2017; Pilz 2007; Volk and Leonard 1990). A large number of sclerotia or sclerotium-like cells can be produced in both agar medium and soils under field cultivation (Liu et al. 2017). As indicated above, the exogenous nutrient “feeding technology” in morel field cultivation is actually a speculation based on extensive cultivation experience and the principle of sclerotia formation (Liu et al. 2017; Tan 2016) and little is known about the nutrition and metabolism in morel artificial cultivation.

In this report, the next generation sequencing (NGS) technology was used to analyze the whole transcriptome and expression pattern of sclerotium development of *M. importuna* at three vegetative growth stages (vegetative mycelium [VM], initial sclerotium [IS] and mature sclerotium [MS]). The genome-wide data mining will improve our understanding of morel sclerotial development and nutrient metabolism and further promote the improvement of *Morchella* cultivation techniques.

Materials and methods

Fungal strain, growth conditions, and development stages

The cultivated strain of *M. importuna* No.1 was used in this study (He et al. 2015a, b). A single mycelial disk was inoculated in the center of autoclaved cellophane membranes which had been placed on the surface of petri plates containing complete yeast extract medium (CYM). CYM contained g l⁻¹ glucose 20, yeast extracts 2 g, peptone 2 g, K₂HPO₄ 1 g, MgSO₄ 0.5 g, KH₂PO₄ 0.46 g and agar 20 g and incubated at 23 °C. The mycelia initially covered the petri dish

with vegetative mycelia (VM) after 3 days of incubation, followed by the occurrence of initial sclerotia (IS) on the surface of the medium after 7 days, and the development into mature sclerotia (MS) after 14 days. Cultures were scraped from the membranes using a clean nipper on days 3, 7 and 14 which were represented, respectively, as the VM, IS and MS stages, followed by rinsing with clean pre-cooled distilled water, drying with filter paper, freezing immediately in liquid nitrogen and storing at − 80 °C until RNA extraction. Two biological repetitions for each stage and a total of six samples were used for transcriptomic analysis.

RNA extraction

Total RNA was extracted by TRIzol® Reagent (Invitrogen) and purified with Plant RNA Purification Reagent (Invitrogen) as described by the manufacturer. The integrity of the RNA was detected by agarose gel electrophoresis and the concentration was determined via a NanoDrop 8000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA). The RNA quality for RNA-seq was further verified using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA).

cDNA library construction and deep sequencing

RNA-seq libraries were prepared using the TruSeq Stranded mRNA Library Prep Kit (Illumina, USA) essentially as per the manufacturer’s instructions with the exception that the oligo (dT)-primed magnetic bead purified poly-A mRNA was fragmented into small pieces using an ultrasonicator (Covaris S220, USA) and following cDNA synthesis, end repair and adapter ligation, a fragment size of 400-bp was selected for PCR amplification to obtain the final library. The libraries were sequenced with paired-end 101 bp. All reads have been deposited in the Sequence Read Archive (SRA) at NCBI (<https://www.ncbi.nlm.nih.gov/sra>) under the accession number SRR8082042-SRR8082047.

De novo assembly and completeness evaluation of *M. importuna* transcriptome

Before assembly, the 101 bp raw paired-end reads were filtered to obtain the high-quality clean reads by removing adapters, low-quality sequences and reads by Trimmomatic (Bolger et al. 2014) and fastuniq (Xu et al. 2012). The clean reads of six data files were mixed together and subjected to *de novo* assembly via short reads assembling program Trinity-v2.4.0 (Grabherr et al. 2011). The TransDecoder package integrating pfam domain by the hmmscan search was used for the prediction of open reading frames (ORFs) with 150 bp as the minimum ORF length (Sanders and Cartwright 2015). The quality and completeness of the *de novo*

assembly were evaluated by BUSCO v3.0 based on evolutionarily informed expectations of gene content from near-universal single-copy orthologs (Simão et al. 2015).

RNAseq analysis and differential expression

Transcript quantification was carried out using BOWTIE (Langmead and Salzberg 2012) and RSEM (Li and Dewey 2011). Briefly, RSEM was used for calculating maximum likelihood abundance estimates as well as posterior mean estimates and 95% credibility intervals for genes/isoforms. Abundance estimates were generated in the form of two measures: the number of fragments that can be derived from an isoform or gene (the expected counts) and the estimated fraction of transcripts within the sample that was represented by the given isoform or gene. To adjust for library sizes and skewed expression of transcripts, the EC values were normalized using the transcripts Per million (TPM) normalization method by edgeR script from Bioconductor (Grabherr et al. 2011; Li and Dewey 2011; Robinson et al. 2010).

All transcriptome contigs were used as the reference for the evaluation of the expression values. The transcriptome contigs with an equal expression among samples were calculated and the false discovery rate (FDR) method was used to determine the threshold p value in multiple tests. An $FDR \leq 0.001$ and an absolute value of the $\log_2$ -ratio ≥ 2 were used as thresholds to determine the significance of gene expression differences in the three developmental stages (Grabherr et al. 2011; Wang et al. 2010).

Annotation and classification of transcripts

The sequence orientations of the transcripts were determined by BLASTx alignment (E value, 10^{-5}) against the NCBI non-redundant protein (nr) database, the Protein sequence analysis and classification (InterPro) database, the Swiss-Prot protein database, the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway database, Gene Ontology (GO) function databases and the Cluster of Orthologous Groups database (COG). The GO annotation was performed using *interpro2go* according to the molecular function, biological process and cellular component ontologies (<http://www.ebi.ac.uk/interpro/>) (Burge et al. 2012). The transcriptome sequences were also aligned against the COG/KOG database to predict and classify the possible functions (Tatusov et al. 2003). KEGG annotation was performed using the single directional best-hit (SBH) method in KAAS web server (<http://www.genome.jp/tools/kaas/>) (Moriya et al. 2007). CAZyme-encoding genes were identified by BLAST analysis of the whole transcriptome against the amino acid sequences derived from the CAZyme database (<http://www.cazy.org>) using a cutoff E value of 10^{-5} (Cantarel et al. 2009). For functional enrichment analysis, the

p values < 0.05 for a significant over representation of a particular GO category were calculated using the *blast2go* software. Pathway enrichment analysis was performed with p values < 0.05 using KEGG annotation results.

qRT-PCR analysis

qRT-PCR was used to validate the assembled transcriptome from the RNA-seq experiment. RNA was extracted as described above and treated with DNase I (Invitrogen, 18068-015) and cDNA was prepared using the SuperScript III First-Strand Synthesis System for RT-PCR (Invitrogen, 18080400) according to the manufacturer's instructions. The qRT-PCR was carried out using SYBR green (Super-Real Premix Plus; TIANGEN, China) methodology and the Bio-RAD CFX real-time PCR system (Bio-RAD, USA). The cyclophilin-type peptidyl-prolyl cis-trans isomerase gene (TRINITY_DN9811_c0_g1, with NR *nblast* $E = 7.87243e-89$) was used as an internal reference for all the qRT-PCR analyses, which has been evaluated as a stable housekeeping reference gene for more growth and development (Zhang et al. 2018). Each gene was analyzed in two biological replicates and three technical repeats. The specificity of qRT-PCR primers was confirmed by melting curve analyses. A relative gene expression was calculated according to the $2^{-(\Delta\Delta CT)}$ method with the mycelial stages as reference for every specific gene (Livak and Schmittgen 2001). Pearson's correlation between transcriptome and qRT-PCR was performed by SPSS 19.0 for Windows (Field 2009).

Results

Sequencing and *de novo* assembly and ORF prediction

To generate an overview of the *M. importuna* gene expression profiles of different sclerotial development stages, cultures of VM, IS and MS that represented sclerotial undifferentiated, differentiated and matured samples were used for *de novo* transcriptome sequencing and assembling (Figure S1). A total of 217.1 M raw data were generated for 101 bp paired-end reads from the 400-bp insert library. After quality filtering, more than 174 M clean paired-end sequence reads were obtained, with the average Q20 and Q30 values being over 98% and 93%, respectively (Table S1). A total of 79,186,911 bp *de novo* assembled bases were obtained using the Trinity software package. The assembly included 56,438 "trinity transcripts" and 26,496 "trinity genes", with a contig N50 length of 1763 bp, an average contig length over 1064.45 bp, and a GC content 47.56%. The contig size distribution revealed that 51.8% transcripts were more than 1000 bp in length (Table 1). The transcriptome completeness

Table 1 Statistics of the *de novo* transcriptome assembly of *M. importuna*

Trinity transcriptome assembly	
Total trinity transcripts (<i>n</i>)	56,438
Total trinity ‘genes’ (<i>n</i>)	26,496
Percent GC (%)	47.56
Stats based on ONLY LONGEST ISOFORM per ‘GENE’	
Contig N50 (bp)	1763
Median contig length (bp)	671
Average contig (bp)	1064.45
Total assembled bases (bp)	28,203,755
Stats based on ALL transcript contigs	
Contig N50 (bp)	2149
Median contig length (bp)	1050
Average contig (bp)	1403.08
Total assembled bases (bp)	79,186,911

was further quantitatively assessed by BUSCO software (version 3.0.2: Creation date: 13 Feb 2016, number of species: 85, number of BUSCOs: 290) with default settings, which is based on evolutionarily informed expectations of gene content (Simão et al. 2015). The results showed that 275 (93.8%) nucleic acid sequences and 271 (92.8%) proteins were identified, with only four BUSCOs missing (Table S2), suggesting that the *de novo* transcriptome assembly of *M. importuna* is highly accurate. The final assembly data submitted to the Transcriptome Shotgun Assembly (TSA) Database at NCBI can be accessed under accession number GGGK00000000 and the version described in this paper is the version 1 (GGGK01000000).

11,957 ORFs were predicted, with most of them in the length between 150 and 300 bp ($n=2978$, 24.9%), and the longest ORF was 13,113 bp (TRINITY_DN13036_c0_g1_orf1 type: complete len: 4371). The complete ORF was about 52.4%, followed by 5' prime partial and internal ORFs with 28.7% and 10.27%, respectively (Table S3).

Functional annotation and characterization

BLASTX alignments (e value cutoff of 10^{-5}) of the predicted protein sequences were performed in Nr, Interpro, GO, KOG, Swiss-Prot and KAAS databases, and 9676 (80.92%) predicted proteins could be annotated with known function genes (Table S4). Nr annotation suggested that about 5250 genes of *Tuber melanosporum* were well matched to *M. importuna* transcripts (Martin et al. 2010). Truffle and morel belong to the same order while in different families (Tuberaceae and Morchellaceae, respectively) (Murat et al. 2004; Bunyard et al. 1995), followed by *Pyronema omphalodes* that belongs to Pezizales and Pyronemataceae, with 4132 homogeneous genes (Traeger et al. 2013), then *Arthrobotrys oligospora* and *Xylona heveae*

with 2230 and 2225 homogeneous genes to *M. importuna*, respectively (Gazis et al. 2016; Yang et al. 2011) (Table S5).

Based on InterPro annotation, GO assignments were used to classify the functions of *M. importuna* transcripts. The 5834 transcripts were assigned at least one GO term and categorized into 62 functional groups (Table S4, Fig. 1), and seven categories mainly related to molecular function and cellular component were missing, including mitochondrion-associated adherens complex, viral occlusion body, D-alanyl carrier, chemoattractant, chemorepellent, morphogen and toxin. Under the main category of cellular component, cell and cell part occupied the dominant position (4635 transcripts), followed by organelle (4131 transcripts) and organelle part (2949 transcripts). Binding (4115 transcripts) and catalytic (3072 transcripts) were the dominant categories of molecular function. Among the various biological processes, cellular process (4515 transcripts) and metabolic process (4204 transcripts) were most highly represented (Fig. 1).

Furthermore, 5506 (46.05%) transcripts were assigned to one or more KOG functional categories. Dominated in 25 KOG categories were the two groups of “General function prediction only” (1422 transcripts, 13.81%) and “Post-translational modification, protein turnover, chaperones” (886 transcripts, 8.61%). Meanwhile, only a few transcripts were assigned to “Cell motility” (12 transcripts, 0.12%) and “Extracellular structures” (20 transcripts, 0.19%). No category was missing in the KOG classifications (Figure S2).

To identify the biological pathways of *M. importuna*, the 3247 annotated sequences were mapped and assigned to 336 reference canonical pathways in the KEGG database (Table S4). The pathways containing the most transcripts were involved in “Ribosome” (108 transcripts), followed by “Spliceosome” (85 transcripts), “Purine metabolism” (81 transcripts), “RNA transport” (81 transcripts) and “Oxidative phosphorylation” (78 transcripts). All the pathways were related to the mycelial growth and sclerotial development of *M. importuna* (Table S4).

Differential expression genes (DEGs) during the three stages

The RNAseq methodology allowed a transcript-wide analysis of the expression patterns to distinguish the three phenotypes from mycelial growth to sclerotial development of *M. importuna*. The final set of DEGs between phenotypes was represented by hierarchical clustering of the expression patterns. The DEGs of six samples could be divided into three main clusters as illustrated by the hierarchical distance tree, representing the genes involved in the phenotypic plasticity of different development statuses (Fig. 2). The clustering analysis exhibited similar transcript profiles between IS stage and MS stage, while VM showed the greatest difference from the other two stages in the transcript profile.

Fig. 1 GO function of the whole transcriptome in *M. importuna*. (The results are summarized in three main categories: biological process, cellular component and molecular function. A total of 5834 transcripts matched to the known proteins by BLAST were assigned into 62 functional groups)

26,496 trinity genes were obtained by *de novo* transcriptional deep-sequencing and assembly in the vegetative growth of *M. importuna*. DEGs were analyzed by integrating DEG and edgeR software package, and the resultant common DEGs were used for the subsequent differential expression analysis. The most differential expression was observed between the stages of mycelial growth and

sclerotial development. Between the VM and IS stages, 1107 DEGs were identified, with 445 of them up-regulated and 652 down-regulated at the VM phase. More obvious changes occurred between the VM and MS stages, of which 2364 DEGs were identified with 994 of them down-regulated and 1370 of them up-regulated at the MS stages. In contrast, only 302 DEGs were identified between the two sclerotial

Fig. 2 Heatmap of differentially expressed genes (DEGs) during the three stages

development stages (IS and MS stages), with 82 of them down-regulated and 220 of them up-regulated at the MS stage. The results indicated that there was no obvious eco-physiological performance at the later two stages of inward development.

Clustering analysis of the expression genes in the six samples was performed by extracting the genes with similar expression patterns that could be regulated by the same

transcription factor or participate in the same regulatory pathways. 2605 DEGs were detected and divided into three clusters. The cluster_1 consisted of 1206 genes with the expression pattern of up-regulation from VM to IS and then to MS stage. The cluster_2 contained 360 genes, with the all genes up-regulated from VM to IS, and then down-regulated from IS to MS, with the final expression levels of these genes still higher than the initial value at the MS stage

(Fig. 2). In cluster_3, 1039 genes were overall continuously down-regulated from VM to IS and then to MS stage.

Under GO function classification, the three categories (biological process, cell component and molecular function) could be divided into 58 subcategories. In the category of biological process, the three largest subcategories consisted of the metabolic, single-organism and cellular process, including 497, 487 and 486 DEGs, respectively. In the category of cell component, the three largest subcategories were cell, cell part and organelle with 487, 487 and 398 DEGs, respectively. The molecular function was the smallest category including 14 subcategories, with binding ($n=454$) and catalytic ($n=453$) as the largest subcategories. Under all subcategories, the GO subcategory of cluster_3 (except for the subcategory of antioxidant and structural molecule) had a larger number of DEGs than cluster_1 or cluster_2, and even their totality (Fig. 3). The GO functional classification of DEGs indicated that the genes in cluster_3 dominated the biological process.

GO and KEGG annotations and enrichment of DEGs

Functional annotation of DEGs by Blast2Go indicated that the three expression patterns of the 2,605 DEGs can be enriched in 54 functional groups with $p < E^{-5}$ (Fig. 4). Cluster_1 mainly included carbohydrate, polysaccharide, hydrolase, caprolactam, beta-galactosidase, and disaccharide metabolic process. In this cluster, 192 DEGs were

enriched in the three GO function categories, with carbohydrate binding and pattern binding dominated in the category of molecular function ($118/192$, p value $8.73E^{-10}$ and $1.37E^{-05}$, respectively), followed by the carbohydrate metabolic process ($105/192$, p value $8.84E^{-07}$). In cluster_2, 106 of the 360 DEGs were mainly enriched in the carbohydrate metabolic process, such as pattern binding ($71/106$, p value $= 6.83E^{-07}$), polysaccharide metabolic process ($44/106$, p value $= 7.30E^{-06}$), polysaccharide catabolic process ($18/106$, p value $= 1.10E^{-05}$), hydrolase activity, acting on glycosyl bonds ($29/106$, p value $= 1.94E^{-05}$), carbohydrate metabolic process ($61/106$, p value $= 4.08E^{-05}$) and carbohydrate catabolic process ($28/106$, p value $= 4.80E^{-05}$). In cluster_3, 423 genes were enriched in 36 categories, with the most significant categories being catalytic activity ($397/423$, p value $= 3.86E^{-08}$) and transporter activity ($397/423$, p value $= 6.34E^{-06}$). In this cluster, 163 genes (p value $= 4.96E^{-05}$) involved in lipid biosynthesis were also enriched.

Enrichment was detected in the 32 metabolic pathways of KAAS belonging to the four functional groups of metabolism, cellular processes, organismal systems and human diseases (Table S6). Both cluster_1 and cluster_2 belong to the metabolism group, indicating that the up-regulated genes from mycelium to sclerotium mainly function in metabolism, such as biosynthesis of secondary metabolites, metabolism of amino acids, carbohydrate metabolism and energy metabolism. Cluster_3 included genes with a

Fig. 3 GO function classification of DEGs. (Green blue and red bars indicate the DEGs of cluster_1, cluster_2 and cluster_3, respectively)

Fig. 4 GO and KAAS functional enrichment analysis of DEGs. (Left: GO functional enrichment analysis of the three clusters of DEGs. Right: KAAS functional enrichment analysis of the three clusters of

DEGs. The digital numbers of the circle correspond to the three clusters. The abscissa is a p value of enrichment)

continuous down-regulated expression from mycelial growth to sclerotial development, with the enrichment of 23 subgroups under four functional groups. The biggest group was metabolism, including eight subgroups such as amino acid, carbohydrate, energy, glycan, lipid, xenobiotics and other secondary metabolites. Enrichment was also detected in some DEGs involved in cell growth and death, secondary metabolites related to disease, and melanogenesis during vegetative growth of *M. importuna* (Fig. 3).

Annotations and enrichment of CAZyme

Since a large number of DEGs were enriched in basic metabolic pathways, the CAZY enzyme system was used for a further analysis of the vegetative growth of *M. importuna*. CAZymes are families of enzymes that degrade, modify, or generate glycosidic bonds. In the whole transcriptome, 295 transcripts belonging to 111 CAZymes families were annotated as CAZymes, and 194 (65%) of them showed a high expression level (total TPM at three stages ≥ 30). The

111 CAZyme families could be divided into six categories. Among them, the largest one was glycoside hydrolases (GHs), containing 51 transcripts, about 38.35% of the known CAZyme GHs family. followed by the category of glycosyl transferases (GTs), which consisted of 26 transcripts, about 26.80% of the known CAZyme family. In the six categories, the category of polysaccharide lyases (PLs) had the least family number ($n = 3$), about 13.04% of the known family. The categories of carbohydrate esterases (CEs) and auxiliary activities (AAs) had the same family number ($n = 10$), about 62.50% and 76.92% of the known families, respectively. The last category of carbohydrate-binding modules (CBMs) included 11 families, covering about 15.49% of the known families (Table S7).

Among the 295 CAZyme genes, the TRINITY_DN13121_c0_g1 genes showed the highest expression (a total TPM of 8247.964 at the three stages) and accounted for 15.48% of the total, which was further conformed by qRT-PCR analysis (refer to the following section). While the annotation of this transcript showed no definite results,

three conserved domains were identified in conserved domain annotation (Blastp/NCBI Nr protein sequences). The first one was the N-terminal early set domain, a glycogen-binding domain, associated with the catalytic domain of AMP-activated protein kinase beta subunit (the accession cd02859, 5-78aa, 2.06E^{-21}), the second one was the glycogen recognition site of AMP-activated protein kinase (the accession pfam16561, 6-62aa, 2.33E^{-17}), which was contained within the first domain, and the last one was the PRK10811 ribonuclease E/G family (85-277aa, 4.58E^{-03}), which showed 53% (e value 1.73117E^{-61}) similarity to the *Colletotrichum tofieldiae* CMB48 protein (Accession Number: KZL76669.1) (Hacquard et al. 2016). The CMB48 was established containing the putative starch-binding domains of the pullulanase subfamily, the β -subunit of AMP-activated protein kinase and some other GH13 enzymes with pullulanase and/or α -amylase-pullulanase specificity. This domain may thus be related to glycogen or starch degradation in *M. importuna*. The second highest expressed transcript was TRINITY_DN12035_c0_g1, with a total TPM of 5646.325, accounting for 10.60% of the total CAZymes gene expression. The domain was annotated as CH31 family and functioned as alpha-1,4-glucan lyase. The two aforementioned transcripts had similar expression patterns, with both of them up-regulated at the VM stage and down-regulated at the SI and SM stages. The relationship of these two genes with the starches of *M. importuna* needs to be further confirmed. The third highest expressed transcript was TRINITY_DN6176_c0_g1, which belongs to AA3 CAZyme family as the function of alcohol oxidase accounted for 3.16% of the total CAZymes gene expression. This transcript showed a different expression pattern from that of the two aforementioned transcripts, with a low-level expression at the VM stage, an up-regulated expression at the SI stage and a slightly down-regulated expression at the SM stage (Table 2).

Using all the *M. importuna* transcripts as background and all the DEGs as reference, the enrichment analysis revealed 74 CAZyme DEGs in cluster_3 ($p = 2.72\text{E}^{-24}$), which showed higher correlation from mycelia to sclerotia with a similar expression pattern. In cluster_2 ($p = 9.18\text{E}^{-05}$), 18 CAZyme DEGs showed an up-regulated expression pattern first and then a down-regulated expression pattern during morel mycelial growth. However, no differentially expressed CAZyme genes were assigned to cluster_1 category (Table S8).

Quantitative real-time PCR

Despite cultivation on a simple complete yeast extracts medium, a large number of genes were identified to be involved in carbohydrate metabolism. These genes varied in their expression at different stages of vegetative

growth development. To confirm the RNA-seq results, qRT-PCR was performed of the top 14 genes belonging to five categories and the other two amylases (Amy_1 TRINITY_DN22515_c0_g1_orf1, glucoamylase, e value = 5.80433E^{-18} ; Amy_2 TRINITY_DN13204_c1_g1_orf1, alpha-glucosidase, e value = 0, BLAST to Swissprot database, respectively) of the CAZymes in *M. importuna* (Fig. 5; Table 2, Table S9). The RNA-seq and qRT-PCR results were compared with the log2 ratio at each test time point. The two results showed strong correlation with each other and were statistically significant (Pearson correlation, $r = 0.8604$, $p = 4.6031\text{E}^{-15}$), indicating that the sequencing results were reliable.

Discussion

To our best knowledge, this is probably the first report about the transcriptomic analysis of the vegetative growth of *M. importuna*. Six samples collected at the three stages from mycelium to mature sclerotium were analyzed by transcriptional deep-sequencing and *de novo* assembly. Given that the genome size of *M. importuna* is about 54 Mb (Liu et al. 2018), the ~17 Gb of transcriptomic sequence clean data represent more than 300-fold coverage (Table 1). The *de novo* assembly of lower eukaryotic transcriptome is straight forward and more than 30-fold coverage is required for the full length of most yeast transcripts (Gygi et al. 1999). However, the *de novo* assembly of higher eukaryotic transcriptome is much more challenging because of the larger sizes of the datasets and the difficulties involved in identifying the alternatively spliced variants (Maier et al. 2009; Treangen and Salzberg 2011). Here, the over 300-fold genome coverage would ensure the comprehensiveness of the transcriptional data for the morel vegetative growth.

To obtain a complete gene expression pattern, the transcriptome of *M. importuna* was conducted by Trinity *de novo* assembly software package. Based on paired-end reads, the transcripts for all potential splice isoforms in each group can be constructed, leading to the assembly of some low-abundance genes, which may also increase the risks of assembling error or pseudogenes (Grabherr et al. 2011). So, the assembly results were filtered by comparing at least 10 reads to every 1000 bp potential assembly gene sequence. Finally, a total of 11,957 effective proteins were obtained, which was consistent with the recently reported genomic data (Liu et al. 2018). BUSCO quantitative assessment identified 93.8% nucleotide sequences and 92.8% protein conserved genes in the fungi BUSCO database, indicating that the transcriptome for the vegetative growth of *M. importuna* was nearly complete (Table S2). Using six public databases as references, 9676 transcripts were annotated to known function at BLAST e value cutoff of 10^{-5} . This study has

Table 2 Top 50 highest expressed CAZyme transcripts in the vegetative growth of *M. importuna*

Transcript number	VM_1	VM_2	IS_1	IS_2	MS_1	MS_2	CAZyme family	Nr potential annotation results
TRINITY_DN13121_c0_g1	2040.124	1110.507	1182.937	1903.176	956.381	1054.839	CBM48	Hypothetical protein
TRINITY_DN12035_c0_g1	2008.44	1067.147	931.827	697.731	428.772	512.318	CH31	Alpha-glucosidase
TRINITY_DN6176_c0_g1	11.609	6.619	820.445	503.334	184.784	160.263	AA3	Alcohol oxidase
TRINITY_DN6152_c0_g1	226.41	715.31	177.22	111.127	35.904	34.955	CE7	Uncharacterized
TRINITY_DN13197_c4_g3	119.14	165.943	218.357	82.657	239.402	289.031	GT8	Glycogenin-2
TRINITY_DN11299_c0_g1	314.729	255.86	126.606	50.321	162.651	170.264	GH13	1,4-Alpha-glucan-branching enzyme
TRINITY_DN14171_c0_g1	411.061	336.355	94.453	23.983	51.557	51.399	CE4	Bifunctional xylanase/deacetylase
TRINITY_DN1195_c0_g1	409.328	290.779	84.804	49.441	22.86	26.014	AA3	Alcohol dehydrogenase
TRINITY_DN10163_c1_g1	594.858	258.167	4.932	4.114	0.286	0.972	AA3	Pyranose 2-oxidase
TRINITY_DN5481_c0_g1	169.375	185.765	176.212	125.682	91.463	101.515	AA2	Cytochrome c peroxidase
TRINITY_DN5712_c0_g1	98.073	57.387	372.077	132.187	92.571	93.979	CE1	S-formylglutathione hydrolase
TRINITY_DN11326_c0_g1	120.969	146.296	196.904	142.699	86.984	113.583	GT20	Alpha, alpha-trehalose-phosphate synthase
TRINITY_DN13373_c1_g10	99.406	51.884	84.097	154.97	194.04	183.515	CE4	Peptidoglycan deacetylase
TRINITY_DN13422_c3_g4	427.112	192.11	61.357	28.328	24.527	27.241	AA6	NAD(P)H dehydrogenase
TRINITY_DN9575_c0_g1	138.945	132.15	96.211	86.256	95.847	139.023	GH72	1,3-Beta-glucanosyltransferase
TRINITY_DN11744_c0_g1	122.119	105.545	34.565	152.473	92.035	164.109	GT2	Chitin synthase
TRINITY_DN2550_c0_g1	0.139	0	137.874	54.746	228.729	212.877	CBM52	Endo-1,3(4)-beta-glucanase
TRINITY_DN12947_c2_g1	203.715	322.347	27.392	10.743	3.621	4.697	CE4	Bifunctional xylanase/deacetylase
TRINITY_DN12622_c0_g1	267.953	129.935	51.365	23.805	26.171	29.551	CE4	Peptidoglycan-N-acetylglucosamine deacetylase
TRINITY_DN8657_c0_g2	360.896	86.437	3.388	2.408	2.514	2.199	CBM18	Endochitinase 1
TRINITY_DN6820_c0_g1	70.317	61.104	91.419	62.246	76.775	84.298	GT28	UDP-N-acetylglucosamine transferase
TRINITY_DN27788_c0_g1	66.738	52.058	113.955	89.952	52.95	55.178	CE5	Cutinase
TRINITY_DN13341_c1_g4	118.139	108.31	57.358	64.645	34.558	35.861	GH133	Glycogen debranching enzyme
TRINITY_DN22948_c0_g1	0.157	0.366	88.632	163.083	88.104	78.01	GH7	1,4-Beta-D-glucan cellobiohydrolase
TRINITY_DN7980_c0_g1	86.838	88.68	69.334	46.998	59.49	60.505	GT2	Dolichol-phosphate mannosyltransferase
TRINITY_DN10700_c0_g1	84.722	64.912	77.567	81.724	46.458	56.206	GT3	Glycogen [starch] synthase
TRINITY_DN12510_c0_g1	252.86	98.293	14.141	7.295	10.257	14.212	GH31	Probable alpha/beta-glucosidase
TRINITY_DN11808_c0_g2	90.122	151.295	52.073	46.34	25.993	29.584	GT20	Trehalose-phosphatase
TRINITY_DN13416_c0_g1	27.678	15.784	105.121	109.874	68.043	64.45	GH95	Alpha-L-fucosidase
TRINITY_DN8487_c0_g1	35.455	25.324	68.197	72.989	88.52	86.143	GH37	Neutral trehalase
TRINITY_DN7885_c0_g1	121.527	96.407	43.12	48.535	25.862	26.567	GH76	Mannan endo-1,6-alpha-mannosidase
TRINITY_DN12072_c0_g1	52.943	36.485	30.051	125.54	47.399	51.343	CBM21	Protein phosphatase 1
TRINITY_DN11002_c0_g2	47.526	161.1	23.072	54.702	13.008	14.775	GH3	Beta-glucosidase
TRINITY_DN5863_c0_g1	7.098	7.535	31.81	51.84	121.101	89.028	GH3	Probable beta-glucosidase
TRINITY_DN11817_c0_g1	2.665	1.95	21.303	39.898	101.148	135.089	CBM1	Mannan endo-1,4-beta-mannosidase
TRINITY_DN13252_c0_g1	134.303	77.236	18.987	14.697	18.547	24.39	GH71	Thiamine biosynthetic bifunctional enzyme
TRINITY_DN9739_c0_g1	56.566	40.513	45.844	45.851	42.908	40.768	GH38	Alpha-mannosidase

Table 2 (continued)

Transcript number	VM_1	VM_2	IS_1	IS_2	MS_1	MS_2	CAZyme family	Nr potential annotation results
TRINITY_DN9260_c0_g1	64.813	54.759	26.46	39	35.916	47.829	GT33	Chitobiosyldiphosphodolichol beta-mannosyltransferase
TRINITY_DN2753_c0_g1	66.59	50.026	32.871	46.207	32.402	39.01	GH20	Beta-hexosaminidase 2
TRINITY_DN7873_c0_g1	89.948	66.213	22.257	45.283	17.428	20.179	GH17	Probable glucan endo-1,3-beta-glucosidase
TRINITY_DN10702_c0_g1	87.047	134.76	11.214	7.073	6.73	6.907	CE4	Peptidoglycan-N-acetylglucosamine deacetylase
TRINITY_DN9742_c0_g1	91.916	47.957	26.524	37.943	21.728	21.461	GT62	Mannan polymerase
TRINITY_DN8381_c0_g1	4.59	4.047	55.9	98.154	43.885	38.414	GH6	Probable 1,4-beta-D-glucan cellobiohydrolase
TRINITY_DN11588_c0_g1	10.991	10.923	70.03	50.774	48.459	52.283	AA7	6-Hydroxy-D-nicotine oxidase
TRINITY_DN25184_c0_g1	65.031	152.045	3.645	4.63	5.218	7.448	GH17	Glucan 1,3-beta-glucosidase
TRINITY_DN10413_c0_g1	21.337	11.847	33.579	27.226	73.606	66.66	GH2	Beta-galactosidase
TRINITY_DN11951_c0_g1	95.077	32.31	38.703	49.397	8.112	7.802	AA3	Choline dehydrogenase
TRINITY_DN9380_c0_g2	37.763	39.14	36.173	35.055	36.904	44.414	GT15	Glycolipid 2-alpha-mannosyltransferase
TRINITY_DN27871_c0_g1	0	0	12.201	58.594	81.075	77.258	AA9	Endoglucanase-4
TRINITY_DN13054_c4_g4	30.012	18.668	9.392	36.477	49.567	77.302	GT32	Mannosyl phosphorylinositol ceramide synthase

provided quite complete and comprehensive transcriptomic information of *M. importuna* (Table S4).

The sclerotial development from vegetative mycelium to mature sclerotium underwent significant morphological changes (Figure S1). However, the whole transcriptomic analysis and functional enrichment showed that the DEGs in the transcriptional level were mainly involved in the primary metabolism process, including carbohydrates, fatty acids, proteins and energy metabolism (Fig. 3). During this process, 295 CAZyme transcripts were obtained and assigned to 111 families (Table S7). More importantly, 65% of the CAZyme transcripts were highly expressed (TPM $\geq$ 30), with 67% of them showing significant difference at the three vegetative growth stages of sclerotial development. The overall results indicated that decomposition and utilization of exogenous substances might mainly occur at the VM stage, and internal energy metabolisms could be the main events of sclerotial development of *M. importuna*.

Although several studies revealed the ectomycorrhizal status of morel with the species of Pinaceae (Buscot and Kottke 1990; Buscot 1993; Dahlstrom et al. 2000) or with the terrestrial orchid *Gymnadenia conopsea* (Stark et al. 2009), *M. importuna* and many other *Morchella* spp. are generally considered to be saprotrophic (Fujimura et al. 2005; Kellner et al. 2007; Pion et al. 2013; Hobbie et al. 2016). The pseudosclerotium of *Morchella* fungi was speculated as the nutrient storage organs (Volk and Leonard 1989, 1990; Liu et al. 2017; Hobbie et al. 2001). However, the accumulation and utilization of the stored nutrients still remain unclear. Lipids have been recorded as the main

cytoplasmic reserves in sclerotia of *Morchella* fungi (Stott and Mohammed 2004; Volk and Leonard 1990). Recently, we have confirmed that more lipids were accumulated as energy-rich substance in sclerotia than in hyphal cells in *M. importuna* (He et al. 2018). In morel sclerotial development, a preliminary analysis was performed concerning the content changes of internal monosaccharides and disaccharides, such as glucose, galactose, mannose and trehalose (Amir et al. 1993; Palma and Kiraly 2011; Kiraly and Czovek 2007). Further research efforts should be made on the nutrition and metabolism of *Morchella* mushrooms.

Morel sclerotial formation is a global regulation of the complex network involving genetic and physiological regulation, such as light, humidity, oxygen, nutrient and salt concentration (osmotic potential), (Liu et al. 2017; He et al. 2018; Amir et al. 1992, 1993, 1994, 1995a, b). Amir et al. (1994) identified the two essential conditions for the sclerotial formation: the high expansion potential required for the growth of mycelial tips and the expansion potential generated by energy metabolism such as hexose and hexanol. In *Rhizoglyphus solani* and *S. rolfii*, the nutrients directly flow into the developing sclerotium. However, in *S. sclerotiorum*, the sclerotia began to form upon the migration of the hyphae from a rich nutrient medium to a poor one (Punja 1985; Little and Rahe 1991). As in *S. sclerotiorum*, exogenous and endogenous cAMP inhibited sclerotia formation (Rollins and Dickman 1998), and the intracellular cAMP concentration increased with the prolongation of the culture time of *M. crassipes* (He et al. 2015a, b). cAMP plays a role in sclerotial formation in fungal as a key modulator of

Fig. 5 qRT-PCR analysis of sixteen CAZyme genes in *M. importuna*

cAMP-dependent protein kinase A (PKA) activity (Jurick and Rollins 2007). The relative PKA activity increased during the white sclerotium stage in the wild-type strain, but was at a low-level in the nonsclerotium-producing mutants of *S. sclerotiorum*, suggesting the involvement of PKA activity in sclerotial development (Harel et al. 2005). TRINITY_DN13405_c4_g1 was identified as PKA pathway protein, with higher expression in the sclerotial development stages, but the other five putative PKA genes had a different expression pattern (Table S10), which could be related to the

complex regulatory functions of PKA (Dürrenberger et al. 1998; D'Souza et al. 2001). Light can affect the fungal sclerotial development (Miller and Liberta 1997). Yager (1992) pointed out that *veA* mediates a developmental light response in *Aspergillus nidulans*. The *A. parasiticus veA* gene and its homologue in *A. flavus* have been shown to be involved in sclerotial formation (Calvo et al. 2004; Duran et al. 2006; Calvo 2008). A certain amount of light stimulation can induce sclerotia formation in the spawn production of morel cultivation (Liu et al. 2017). Two homologous genes

TRINITY_DN3052_c0_g1 and TRINITY_DN13008_c2_g1 have been identified as *veA*, with the same gene expression pattern, decreased gradually from the VM to MS stages, but no significant difference ($p < 0.05$) (Table S10). Unlike sclerotia formation under an acidic ambient pH of *S. sclerotiorum* (Rollins and Dickman 2001), neutral or slightly alkaline pH conditions facilitate the mycelial growth and sclerotial formation favor of morels and, in the course of growth, the initial acidic or alkaline pH tends to be neutral (Brock 1951; Liu et al. 2017). A common zinc finger transcription factor *PacC* in *A. nidulans* has been reported to regulate many gene products with pH-sensitive activities (Caddick et al. 1986), and the homologue gene *pac1* mutation affects the sclerotial development in *S. sclerotiorum*. The homologous gene TRINITY_DN13300_c14_g1 has been annotated as transcription factor *pacC/RIM101* with e value $7.44E^{-83}$, which was highly expressed at the SI stage, but with no significant difference from the other two stages ($p < 0.05$) (Table S10). Chen et al. (2004) demonstrated that the homologue of an ERK-type mitogen-activated protein kinase (MAPK) in *S. sclerotiorum* (*Smk1*) regulates sclerotial development via a pH-dependent signalling pathway involving the accumulation of oxalic acid. The transcriptional regulator *LaeA*, a member of the conserved transcriptional regulatory velvet complex and also a global regulator of fungal secondary metabolite, regulated sclerotial development via a cellular fusion cascade (Zhao et al. 2017). Similar transcription factors were also detected in the present study (Table S10), but whether these genes (e.g., *VeA*, *LaeA*, *pacC*, cAMP kinase, MAPK) regulate the sclerotial development in the same manner needs to be further studied.

Sclerotium can be formed from the primary hyphae and the secondary hyphae (Liu et al. 2017; He et al. 2018). Volk and Leonard (1990) pointed out that only the sclerotia formed from the secondary hyphae may develop into the ascocarps (Volk and Leonard 1990). In 1982, Ower achieved the cultivation of *M. rufobrunnea* under controlled condition firstly (Ower 1982; Pilz 2007; Liu et al. 2017). Then, three American patents were applied successively by Ower and his partners in the cultivation schemes with the culture and treatment of morel sclerotia as the pivotal events (Ower 1982, 1986, 1988, 1989). Meanwhile, they also paid attention to exogenous nutrient supplement (Ower 1986, 1988, 1989), but morel artificial cultivation was not explored (Liu et al. 2017). Fortunately, The technology of exogenous nutrient supplement has been gradually improved in recent years in China, which ultimately facilitated the successful cultivation of *Morchella* mushrooms. According to this technology, after 7–20 days of inoculation, mycelia will cover the soil surface; the sterilized nutrient bags containing substrates which can be easily assimilated by morel hypha were punched with holes and placed on the soil surface. Then, the mycelia in the soil can enter the nutrient bags through the punched holes to utilize the nutritive materials. Importantly,

the nutrients accumulated in mycelial cells of the bags can be transferred into and stored in the mycelial and sclerotial cells in the soil (Liu et al. 2017). The detailed metabolic process of this energy substance flowing from exogenous nutrient bags to mycelia network, sclerotia or sclerotium-like cells in soil and facilitating the later sexual reproduction needs to be further studied.

Conclusions

The vegetative mycelia (VM), initial sclerotia (IS) and mature sclerotia (MS) of *M. importuna* were subjected to deep transcriptome sequencing analysis and *de novo* assembly, resulting in the first complete and comprehensive transcriptome of *M. importuna*. 56,438 transcripts and 26,496 trinity genes, with an average contig of 1064.45 bp and a contig N50 length of 1763 bp, were obtained. Additionally, 11,957 ORFs were predicted, and more than 80.9% of them (9676 ORFs) were annotated. The gene expressions of the six samples at three vegetative growth stages were calculated, and the edgeR and DESeq results were integrated for DEGs analysis, leading to the identification of 2605 DEGs in three categories. Enrichment analysis showed that the DEGs at the three developmental stages were mainly involved in the primary metabolism. All the CAZyme genes were highly expressed as validated by qRT-PCR of 16 CAZyme genes and the correlation index between the qRT-PCR and the transcriptome data was 0.8604, indicating the reliability of transcriptional data in our study. The overall results showed that the anabolism of the energy-rich substances was the main event of the three biological processes.

Acknowledgements This study was supported by the Key Technologies R & D Program of Henan Province (No. 172102310553), the Major Technology Innovation Projects of Hubei Province (Grant No. 2016ABA100) and the Industry (Agriculture) Science and Technology Program (Grant No. 201503107).

Author contributions WL and PH conceived and designed the experiments. YC and WL prepared the experiment materials. WL and YC carried out the experiments and collected the data. Data analysis was carried out by LC and WL. WL and PH wrote the manuscript. PH and YB provided intellectual input and revised the manuscript. All authors have read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Alvarado-Castillo G, Mata G, Tablada MEN, Martinez-Carrera D, Rosado DEP (2008) Obtaining sclerotia of morel mushroom

- (*Morchella esculenta*) in different culture media. *Interciencia* 33(7):528–531
- Alvarado-Castillo G, Mata G, Perez-Vazquez A, Martinez-Carrera D, Tablada MEN, Gallardo-Lopez F, Osorio-Acosta F (2011) *Morchella sclerotia* production through grain supplementation. *Interciencia* 36:768–773
- Amir R, Levanon D, Hadar Y, Chet I (1992) Formation of sclerotia by *Morchella esculenta*: relationship between media composition and turgor potential in the mycelium. *Mycol Res* 96(11):943–948. [https://doi.org/10.1016/S0953-7562\(09\)80595-9](https://doi.org/10.1016/S0953-7562(09)80595-9)
- Amir R, Levanon D, Hadar Y, Chet I (1993) Morphology and physiology of *Morchella esculenta* during sclerotial formation. *Mycol Res* 97(6):683–689. [https://doi.org/10.1016/S0953-7562\(09\)80148-2](https://doi.org/10.1016/S0953-7562(09)80148-2)
- Amir R, Levanon D, Hadar Y, Chet I (1994) The role of source-sink relationships in translocation during sclerotial formation by *Morchella esculenta*. *Mycol Res* 98(12):1409–1414. [https://doi.org/10.1016/S0953-7562\(09\)81071-X](https://doi.org/10.1016/S0953-7562(09)81071-X)
- Amir R, Levanon D, Hadar Y, Chet I (1995a) Factors affecting translocation and sclerotial formation in *Morchella esculenta*. *Exp Mycol* 19(1):61–70. <https://doi.org/10.1006/emyc.1995.1007>
- Amir R, Steudle E, Levanon D, Hadar Y, Chet I (1995b) Turgor changes in *Morchella esculenta* during translocation and sclerotial formation. *Exp Mycol* 19(2):129–136. <https://doi.org/10.1006/emyc.1995.1015>
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30(15):2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
- Brock TD (1951) Studies on the nutrition of *Morchella esculenta* Fries. *Mycologia* 43(4):402–422. <https://doi.org/10.2307/3755649>
- Bunyard BA, Nicholson MS, Royse DJ (1995) Phylogenetic resolution of *Morchella*, *Verpa*, and *Disciotis* [Pezizales: Morchellaceae] based on restriction enzyme analysis of the 28S ribosomal RNA gene. *Exp Mycol* 19(3):223–233. <https://doi.org/10.1006/emyc.1995.1027>
- Burge S, Kelly E, Lonsdale D, Mutowo-Mueller P, McAnulla C, Mitchell A, Sangrador-Vegas A, Yong S-Y, Mulder N, Hunter S (2012) Manual GO annotation of predictive protein signatures: the InterPro approach to GO curation. *Database* <https://doi.org/10.1093/database/bar068>
- Buscot F (1993) Synthesis of two types of association between *Morchella esculenta* and *Picea abies* under controlled culture conditions. *J Plant Physiol* 141(1):12–17. [https://doi.org/10.1016/S0176-1617\(11\)80845-X](https://doi.org/10.1016/S0176-1617(11)80845-X)
- Buscotand F, Kottke I (1990) The association of *Morchella rotunda* (Pers.) boudier with roots of *Picea abies* (L.) Karst. *New Phytol* 116(3):425–430. <https://doi.org/10.1111/j.1469-8137.1990.tb00528.x>
- Caddick MX, Brownlee AG, Arst HN Jr (1986) Regulation of gene expression by pH of the growth medium in *Aspergillus nidulans*. *Mol Gen Genet* 203:346–353. <https://doi.org/10.1007/BF00333978>
- Calvo AM (2008) The VeA regulatory system and its role in morphological and chemical development in fungi. *Fungal Genet Biol* 45(7):1053–1061. <https://doi.org/10.1016/j.fgb.2008.03.014>
- Calvo AM, Bok J, Brooks W, Keller NP, Hinze LL, Gardner HW (2004) veA is required for toxin and sclerotial production in *Aspergillus parasiticus*. *Appl Environ Microbiol* 70:4733–4739. <https://doi.org/10.1128/AEM.70.8.4733-4739.2004>
- Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B (2009) The Carbohydrate-active enzymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res* 37(suppl_1):D233–D238. <https://doi.org/10.1093/nar/gkn663>
- Chen CB, Harel A, Gorovits R, Yarden O, Dickman MB (2004) MAPK regulation of sclerotial development in *Sclerotinia sclerotiorum* is linked with pH and cAMP sensing. *Mol Plant Microbe Interact* 17:404–413. <https://doi.org/10.1094/MPMI.2004.17.4.404>
- D'Souza CA, Alspaugh JA, Yue C, Harashima T, Cox GM, Perfect JR, Heitman J (2001) Cyclic AMP-dependent protein kinase controls virulence of the fungal pathogen *Cryptococcus neoformans*. *Mol Cell Biol* 21(9):3179–3191. <https://doi.org/10.1128/MCB.21.9.3179-3191.2001>
- Dahlstrom J, Smith J, Weber S N (2000) Mycorrhiza-like interaction by *Morchella* with species of the Pinaceae in pure culture synthesis. <https://doi.org/10.1007/PL00009992>
- Duran RM, Cary JW, Calvo AM (2006) Production of cyclopiazonic acid, aflatoxin, and aflatoxin by *Aspergillus flavus* is regulated by veA, a gene necessary for sclerotial formation. *Appl Genet Mol Biotechnol* 74:1158–1168. <https://doi.org/10.1007/s00253-006-0581-5>
- Dürrenberger F, Wong K, Kronstad JW (1998) Identification of a cAMP-dependent protein kinase catalytic subunit required for virulence and morphogenesis in *Ustilago maydis*. *Proc Natl Acad Sci* 95(10), 5684–5689. <https://doi.org/10.1073/pnas.95.10.5684>
- Field A (2009) Discovering statistics using SPSS. Sage publications, London
- Fujimura KF, Smith JE, Horton TR, Weber NS, Spatafora JW (2005) Pezizalean mycorrhizas and sporocarps in ponderosa pine (*Pinus ponderosa*) after prescribed fires in eastern Oregon, USA. *Mycorrhiza* 15(2):79–86. <https://doi.org/10.1007/s00572-004-0303-8>
- Gazis R, Kuo A, Riley R, LaButti K, Lipzen A, Lin J, Amirebrahimi M, Hesse CN, Spatafora JW, Henrissat B, Hainaut M, Grigoriev IV, Hibbett DS (2016) The genome of *Xylona heveae* provides a window into fungal endophytism. *Fungal Biol* 120(1):26–42. <https://doi.org/10.1016/j.funbio.2015.10.002>
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng QD, Chen ZH, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* 29(7):644–652. <https://doi.org/10.1038/nbt.1883>
- Gygi SP, Rochon Y, Franz A, Aebersold R (1999) Correlation between Protein and mRNA Abundance in Yeast. *Mol Cell Biol* 19(3):1720–1730. <https://doi.org/10.1128/mcb.19.3.1720>
- Hacquard S, Kracher B, Hiruma K, Munch PC, Garrido-Oter R, Thon MR, Weimann A, Damm U, Dallery JF, Hainaut M, Henrissat B, Lespinet O, Sacristan S, van Themaat E, Kemen E, McHardy AC, Schulze-Lefert P, O'Connell RJ (2016) Survival trade-offs in plant roots during colonization by closely related beneficial and pathogenic fungi. *Nat Commun* 7:11362. <https://doi.org/10.1038/ncomms11362>
- Harel A, Gorovits R, Yarden O (2005) Changes in protein kinase A activity accompany sclerotial development in *Sclerotinia sclerotiorum*. *Phytopathology* 95(4):397–404. <https://doi.org/10.1094/PHYTO-95-0397>
- He PX, Geng LJ, Mao DB, Xu CP (2012) Production, characterization and antioxidant activity of exopolysaccharides from submerged culture of *Morchella crassipes*. *Bioproc Biosyst Eng* 35(8):1325–1332. <https://doi.org/10.1007/s00449-012-0720-6>
- He PX, Huang LN, Liu W, Liu SM, Cheng CK, Chen LT (2015a) Effects of cAMP on mycelial growth and sclerotial morphogenesis in *Morchella crassipes*. *J Fungal Res* 13(3):155–159
- He PX, Liu W, Cai YL, He XSH (2015b) Strain identification and phylogenetic analysis of cultivated and wild strains of *Morchella* belonging to Elata Clade in China. *J Zhengzhou Univ Light Ind (Nat Sci)* 30(3):26–29
- He PX, Wang K, Cai YL, Liu W (2017) Live cell confocal laser imaging studies on the nuclear behavior during meiosis and ascosporeogenesis in *Morchella importuna* under artificial

- cultivation. *Micron* 101:108–113. <https://doi.org/10.1016/j.micron.2017.06.012>
- He PX, Wang K, Cai YL, Hu XL, Zheng Y, Zhang JJ, Liu W (2018) Involvement of autophagy and apoptosis and lipid accumulation in sclerotial morphogenesis of *Morchella importuna*. *Micron* 109:34–40. <https://doi.org/10.1016/j.micron.2018.03.005>
- Heleno SA, Stojkovic D, Barros L, Glamoclija J, Sokovic M, Martins A, Queiroz MJRP, Ferreira ICFR (2013) A comparative study of chemical composition, antioxidant and antimicrobial properties of *Morchella esculenta* (L.) Pers. from Portugal and Serbia. *Food Res Int* 51(1):236–243. <https://doi.org/10.1016/j.foodres.2012.12.020>
- Hobbie EA, Weber NS, Trappe JM (2001) Mycorrhizal vs saprotrophic status of fungi: the isotopic evidence. *New Phytol* 150(3):601–610. <https://doi.org/10.1046/j.1469-8137.2001.00134.x>
- Hobbie EA, Rice SF, Weber NS, Smith JE (2016) Isotopic evidence indicates saprotrophy in post-fire *Morchella* in Oregon and Alaska. *Mycologia* 108(4):638–645. <https://doi.org/10.3852/15-281>
- Jurick IJWM, Rollins JA (2007) Deletion of the adenylate cyclase (*sac1*) gene affects multiple developmental pathways and pathogenicity in *Sclerotinia sclerotiorum*. *Fungal Genet Biol* 44(6):521–530. <https://doi.org/10.1016/j.fgb.2006.11.005>
- Kellner H, Luis P, Buscot F (2007) Diversity of laccase-like multicopper oxidase genes in *Morchellaceae*: identification of genes potentially involved in extracellular activities related to plant litter decay. *Fems Microbiol Ecol* 61(1):153–163. <https://doi.org/10.1111/j.1574-6941.2007.00322.x>
- Kiraly I, Czovek P (2007) Oxidative burst induced pseudosclerotium formation of *Morchella steppicola* Zerova on different malt agar media. *Can J Microbiol* 53(8):975–982. <https://doi.org/10.1139/W07-055>
- Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9(4):357–359. <https://doi.org/10.1038/nmeth.1923>
- Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform* 12(1):323. <https://doi.org/10.1186/1471-2105-12-323>
- Little ER, Rahe JE (1991) In vitro initiation of sclerotia of *Sclerotium cepivorum*. *Can J Plant Pathol* 13(1):45–49. <https://doi.org/10.1080/07060669109500964>
- Liu W, Zhang Y, He PX (2017) Morel biology and cultivation. Jilin science and Technology Press, Changchun, pp 1–340
- Liu W, Chen LF, Cai YL, Zhang QQ, Bian YB (2018) Opposite polarity monospore genome *de novo* sequencing and comparative analysis reveal the possible heterothallic life cycle of *Morchella importuna*. *Int J Mol Sci* 19(9):2525. <https://doi.org/10.3390/ijms19092525>
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25(4):402–408. <https://doi.org/10.1006/meth.2001.1262>
- Maier T, Güell M, Serrano L (2009) Correlation of mRNA and protein in complex biological samples. *FEBS Lett* 583(24):3966–3973. <https://doi.org/10.1016/j.febslet.2009.10.036>
- Martin F, Kohler A, Murat C, Balestrini R, Coutinho PM, Jaillon O, Montanini B, Morin E, Noel B, Percudani R, Porcel B, Rubini A, Amicucci A, Amselem J, Anthouard V, Arcioni S, Artiguenave F, Aury J-M, Ballario P, Bolchi A, Brenna A, Brun A, Buée M, Cantarel B, Chevalier G, Couloux A, Da Silva C, Denoeud F, Duplessis S, Ghignone S, Hilselberger B, Iotti M, Marçais B, Mello A, Miranda M, Pacioni G, Quesneville H, Riccioni C, Ruotolo R, Splivallo R, Stocchi V, Tisserant E, Viscomi AR, Zambonelli A, Zampieri E, Henrissat B, Lebrun M-H, Paolocci F, Bonfante P, Ottonello S, Wincker P (2010) Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:1033. <https://doi.org/10.1038/nature08867>
- Miller RM, Liberta AE (1977) The effects of light and tyrosinase during sclerotium development in *Sclerotium rolfii* Sacc. *Can J Microbiol* 23(3):278–287. <https://doi.org/10.1139/m77-041>
- Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M (2007) KAAAS: an automatic genome annotation and pathway reconstruction server. *Nucleic Acids Res* 35(suppl_2):W182–W185. <https://doi.org/10.1093/nar/gkm321>
- Murat C, Díez J, Luis P, Delaruelle C, Dupré C, Chevalier G, Bonfante P, Martin F (2004) Polymorphism at the ribosomal DNA ITS and its relation to postglacial re-colonization routes of the Périgord truffle *Tuber melanosporum*. *New Phytol* 164(2):401–411. <https://doi.org/10.1111/j.1469-8137.2004.01189.x>
- Ower R (1982) Notes on the Development of the Morel Ascocarp: *Morchella esculenta*. *Mycologia* 74(1):142–144. <https://doi.org/10.2307/3792639>
- Ower RD, Mills GL, Malachowski JA (1986) Cultivation of *Morchella*. U.S. Patent No. 4,594,809. U.S. Patent and Trademark Office, Washington, DC
- Ower RD, Mills GL, Malachowski JA (1988) Cultivation of *Morchella*. U.S. Patent No. 4,757,640. U.S. Patent and Trademark Office, Washington, DC
- Ower RD, Mills GL, Malachowski JA (1989) Cultivation of *Morchella*. U.S. Patent No. 4,866,878. Patent and Trademark Office, Washington, DC
- Palma C, Kiraly I (2011) Inducible trehalase enzyme activity of *Morchella conica* Persoon mycelium. *Acta Microbiol Imm H* 58(1):1–11. <https://doi.org/10.1556/AMicr.58.2011.1.1>
- Pilz D, McLain R, Alexander S, Villarreal-Ruiz L, Berch S, Wurtz TL, Smith JE (2007) Ecology and management of morels harvested from the forests of western North America. Gen. Tech. Rep. PNW-GTR-710. US Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland
- Pion M, Spangenberg JE, Simon A, Bindschedler S, Flury C, Chatelain A, Bshary R, Job D, Junier P (2013) Bacterial farming by the fungus *Morchella crassipes*. *P Roy Soc B Biol Sci* <https://doi.org/10.1098/rspb.2013.2242>
- Punja ZK (1985) The biology, ecology, and control of *Sclerotium rolfii*. *Ann Rev Phytopathol* 23(1):97–127. <https://doi.org/10.1146/annurev.py.23.090185.000525>
- Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1):139–140. <https://doi.org/10.1093/bioinformatics/btp616>
- Rollins JA, Dickman MB (1998) Increase in endogenous and exogenous cyclic AMP levels inhibits sclerotial development in *Sclerotinia sclerotiorum*. *Appl Environ Microbiol* 64(7):2539–2544
- Rollins JA, Dickman MB (2001) pH signaling in *Sclerotinia sclerotiorum*: identification of a *pacC/RIM1* homolog. *Appl Environ Microbiol* 67:75–81. <https://doi.org/10.1128/AEM.67.1.75-81.2001>
- Rosbach M, Kummerle E, Schmidt S, Gohmert M, Stieghorst C, Revay Z, Wiehl N (2017) Elemental analysis of *Morchella esculenta* from Germany. *J Radioanal Nucl Ch* 313(1):273–278. <https://doi.org/10.1007/s10967-017-5298-7>
- Sanders SM, Cartwright P (2015) Interspecific differential expression analysis of RNA-Seq data yields insight into life cycle variation in *Hydractinia hydrozoans*. *Genome Biol Evol* 7(8):2417–2431. <https://doi.org/10.1093/gbe/evv153>
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31(19):3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>
- Stark C, Babik W, Durka W (2009) Fungi from the roots of the common terrestrial orchid *Gymnadenia conopsea*. *Mycol Res* 113(Pt 9):952–959. <https://doi.org/10.1016/j.mycres.2009.05.002>

- Stott K, Mohammed C (2004) Specialty mushroom production systems: maitake and morels. RIRDC, Australia
- Su CA, Xu XY, Liu DY, Wu M, Zeng FQ, Zeng MY, Wei W, Jiang N, Luo X (2013) Isolation and characterization of exopolysaccharide with immunomodulatory activity from fermentation broth of *Morchella conica*. DARU <https://doi.org/10.1186/2008-2231-21-5>
- Tan F (2016) The history, current situation and prospect of artificial cultivation of morels. *Edible Med Mushrooms* 24(3):140–144
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA (2003) The COG database: an updated version includes eukaryotes. *BMC Bioinform* 4(1):41. <https://doi.org/10.1186/1471-2105-4-41>
- Tietel Z, Masaphy S (2018) Aroma-volatile profile of black morel (*Morchella importuna*) grown in Israel. *J Sci Food Agr* 98(1):346–353. <https://doi.org/10.1002/jsfa.8477>
- Traeger S, Altegoer F, Freitag M, Gabaldon T, Kempken F, Kumar A, Marcet-Houben M, Poggeler S, Stajich JE, Nowrousian M (2013) The genome and development-dependent transcriptomes of *Pyronema confluens*: a window into fungal evolution. *PLoS Genet* 9(9):e1003820. <https://doi.org/10.1371/journal.pgen.1003820>
- Treangen TJ, Salzberg SL (2011) Repetitive DNA and next-generation sequencing: computational challenges and solutions. *Nat Rev Genet* 13(1):36–46. <https://doi.org/10.1038/nrg3117>
- Vieira V, Fernandes A, Barros L, Glamoclija J, Ciric A, Stojkovic D, Martins A, Sokovic M, Ferreira ICFR (2016) Wild *Morchella conica* Pers. from different origins: a comparative study of nutritional and bioactive properties. *J Sci Food Agr* 96(1):90–98. <https://doi.org/10.1002/jsfa.7063>
- Volk TJ, Leonard TJ (1989) Physiological and Environmental Studies of sclerotium formation and maturation in isolates of *Morchella crassipes*. *Appl Environ Microbiol* 55(12):3095–3100
- Volk TJ, Leonard TJ (1990) Cytology of the life-cycle of *Morchella*. *Mycol Res* 94(3):399–406. [https://doi.org/10.1016/S0953-7562\(09\)80365-1](https://doi.org/10.1016/S0953-7562(09)80365-1)
- Wang L, Feng Z, Wang X, Wang X, Zhang X (2010) DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26(1):136–138. <https://doi.org/10.1093/bioinformatics/btp612>
- Xu HB, Luo X, Qian J, Pang XH, Song JY, Qian GR, Chen JH, Chen S (2012) FastUniq: a fast de novo duplicates removal tool for paired short reads. *PloS one* 7(12):e52249. <https://doi.org/10.1371/journal.pone.0052249>
- Yager LN (1992) Early developmental events during asexual and sexual sporulation in *Aspergillus nidulans*. *Biotechnology* 23:19–41
- Yang J, Wang L, Ji X, Feng Y, Li X, Zou C, Xu J, Ren Y, Mi Q, Wu J, Liu S, Liu Y, Huang X, Wang H, Niu X, Li J, Liang L, Luo Y, Ji K, Zhou W, Yu Z, Li G, Liu Y, Li L, Qiao M, Feng L, Zhang KQ (2011) Genomic and proteomic analyses of the fungus *Arthrobotrys oligospora* provide insights into nematode-trap formation. *PLoS Pathog* 7(9):e1002179. <https://doi.org/10.1371/journal.ppat.1002179>
- Zhang QQ, Liu W, Cai YL, Lan AF, Bian YB (2018) Validation of internal control genes for quantitative real-time PCR gene expression analysis in *Morchella*. *Molecules* 23(9):2331. <https://doi.org/10.3390/molecules23092331>
- Zhao X, Spraker JE, Bok JW, Velk T, He ZM, Keller NP (2017) A cellular fusion cascade regulated by laea is required for sclerotial development in *Aspergillus flavus*. *Front Microbiol*, 8, 1925. <https://doi.org/10.3389/fmicb.2017.01925>