Published in final edited form as:

Trends Parasitol. 2019 February ; 35(2): 129–139. doi:10.1016/j.pt.2018.11.008.

Important Extracellular Interactions between *Plasmodium* Sporozoites and Host Cells Required for Infection.

Kirsten Dundas¹, Melanie J. Shears², Photini Sinnis², and Gavin J. Wright^{1,*}

¹Cell Surface Signalling Laboratory and Parasites and Microbes Programme, Wellcome Trust Sanger Institute, Cambridge CB10 1SA, United Kingdom.

²Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe Street, Baltimore, MD 21205.

Abstract

Malaria is an infectious disease caused by *Plasmodium* parasites that remains a major global health problem. Infection begins when salivary gland sporozoites are transmitted through the bite of an infected mosquito. Once within the host, sporozoites navigate through the dermis, into the bloodstream and eventually invade hepatocytes. While we have an increasingly sophisticated cellular description of this journey, our molecular understanding of the extracellular interactions between the sporozoite and mammalian host that regulate migration and invasion remain comparatively poor. Here, we review the current state of our understanding, highlight the technical limitations that have frustrated progress, and outline how new approaches will help address this knowledge gap with the ultimate aim of improving malaria treatments.

Keywords

Plasmodium; malaria; sporozoite; host-parasite interactions

Sporozoites and malaria infection

Malaria is an infectious disease caused by *Plasmodium* parasites, of which *P. falciparum* causes most morbidity and mortality. Around 40% of the global population is considered at risk of infection and almost half a million deaths were attributed to malaria in 2016 [1]. *Plasmodium* parasites are unicellular eukaryotes in the phylum Apicomplexa. They have motile extracellular forms called ‘-zoites’ that can actively invade host cells where they multiply. Infection is initiated when an infected female anopheline mosquito searches for blood and transmits *Plasmodium* sporozoites into the dermis of the host. Sporozoites move within the skin in apparently random directions to locate and then enter blood vessels, and from there, are passively transported in the bloodstream and rapidly reach the liver.

*Correspondence: gw2@sanger.ac.uk (G. J. Wright).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Sporozoites leave the circulation by crossing the liver sinusoidal capillary barrier and enter the liver parenchyma where they traverse through several hepatocytes before productively invading a cell and developing into a liver stage form. Here, they multiply into thousands of merozoites which are released into the bloodstream to invade erythrocytes and initiate the blood stage of infection. The asexual blood stage involves the rapid and iterative invasion of erythrocytes and is the symptomatic part of the life cycle.

Sporozoites are considered an attractive target for therapeutic intervention, particularly for vaccines. This is firstly because sporozoites constitute a bottleneck in the parasite life cycle, with only around ten to a hundred sporozoites transmitted by an infected mosquito bite [2]. Secondly, the extracellular sporozoites are exposed to host antibodies for minutes to hours compared with just minutes for the blood stages [3]. Finally, the sporozoite stage is asymptomatic, and so targeting sporozoites could prevent clinical disease. Molecular interactions between sporozoites and the host are likely to be important for the parasite's journey to the liver, and these interactions would be ideal vaccine targets. Indeed, subunit vaccines based on sporozoite surface proteins and attenuated whole sporozoites show some protection in animal models and humans [4].

The development of *in vitro* sporozoite culture methods, advances in microscopy techniques, and the use of tractable animal models of malaria have resulted in a detailed cellular description of how sporozoites interact with their host; however, we know little at the molecular level, leaving many questions unanswered. How do sporozoites recognise the different cell types that act as signposts along its journey to the hepatocyte? Are there host cues that instruct sporozoites to switch behaviour between cell traversal and invasion? What are the host-parasite receptor-ligand interactions involved in guiding sporozoites? With the advent of new techniques and technologies, we are now starting to uncover some of these important sporozoite-host interactions. This review focuses on sporozoite behaviors and their cellular and extracellular molecular interactions with host cells.

Cellular architecture of sporozoites

Sporozoites are slender, crescent-shaped cells that measure approximately 10-12 μm by 1-2 μm . They develop in oocysts on the mosquito midgut wall and are released into the hemocoel where they are carried throughout the open circulatory system of the mosquito, with ~20% ultimately invading salivary glands. Oocyst and salivary gland sporozoites differ, with the latter competent for infection of the mammalian host, a phenotype accompanied by changes in the sporozoite cytoskeleton [5], motility [6, 7] and surface proteins [8]. Transcriptomic and proteomic studies of oocyst and salivary gland sporozoites have identified transcriptional changes associated with maturation [9], and described a repertoire of ~2000 proteins expressed by mature sporozoites [10]. Similar to other motile zoites, sporozoites have distinct cell polarity, specialized motility apparatus, and apical secretory organelles called micronemes, rhoptries and dense granules that have separate yet complementary roles in motility and invasion [11] (Figure 1A). The sporozoite surface is densely covered by the circumsporozoite protein (CSP), an essential and multifunctional protein that is likely linked to the plasma membrane via a GPI anchor [12]. The thrombospondin-related anonymous protein (TRAP), a micronemal protein secreted to the

sporozoite surface during motility, likewise plays a critical role in motility [13]. Targeted proteomic analysis of the surface of mature sporozoites has further identified several dozen surface proteins in addition to CSP and TRAP that might interact with host factors [14]. Alongside this, several other important secreted or sporozoite surface proteins such as CelTOS, P36 and P52 have also been characterized. As outlined below, ongoing genetic, biochemical and molecular analyses are helping link this knowledge of the sporozoite's cellular architecture to specific sporozoite behaviors and their underlying molecular effectors.

Cellular behaviours of sporozoites

Sporozoites display several specialised behaviours on their journey to the liver. Sporozoites move by **gliding motility** (see Glossary), a type of substrate-based motility characterised by a lack of cilia or flagella, and the absence of any overt change in cell shape [15]. After inoculation into the skin, sporozoites must breach several cell barriers to reach the hepatocyte. This, in part, relies on a process known as **cell traversal**, which enables sporozoites to cross host cell membranes and cell barriers. Using motility and cell traversal, sporozoites ultimately reach and **invade** a hepatocyte (Figure 1A, B). This is characterised by the formation of a parasitophorous vacuole: an intracellular membrane within which the parasite resides and develops until the completion of the liver stage.

Sporozoite gliding motility

Sporozoites are among the fastest eukaryotic cells, moving at 1-3 μm per second *in vitro* and *in vivo*. In the mammalian host, motility is essential for sporozoite exit from the dermal inoculation site and subsequent hepatocyte invasion [13, 16, 17]. Critical to gliding motility are the TRAP family of transmembrane proteins. These proteins share several features in their domain architecture, including extracellular adhesion domains and a cytoplasmic domain that couples to an actinomyosin molecular motor [13]. To propel itself, the parasite must adhere to a substrate or cell to provide traction for displacement [15]. A study using reflection interference contrast microscopy (RICM) to examine the contact points between sporozoites and the substrate revealed that “gliding motility” is in fact saltatory, involving the stepwise formation and release of discrete adhesion sites [18]. TRAP is the most critical TRAP family member with targeted mutations of *trap* significantly affecting gliding motility and infectivity [17, 19], revealing the importance of gliding motility for these events.

Advances in microscopy and the accessibility of the skin enabled sporozoite gliding within the dermis to be visualised *in vivo*. Quantitative imaging of sporozoites from the rodent malaria model *P. berghei* in the mouse ear demonstrated that within one hour, approximately half of the sporozoites enter the blood (20%) or lymphatic (30%) vessels [16], with the rest remaining in the dermis for up to seven hours after transmission. Two distinct patterns of sporozoite motility were identified: within the dermis, they moved quickly and were displaced over relatively large distances, while within the vicinity of a blood vessel, their movement was slower and the curvature of their path more constrained, a motility pattern that may optimise blood vessel contact [17]. This suggests that a specific, but unidentified

host-parasite recognition event occurs at the blood vessels, initiating a change in sporozoite behavior, although further work is required to fully understand this critical process.

Sporozoite cell traversal

The ability of sporozoites to traverse host cells is important at several points along its journey from the site of deposition to the hepatocyte. Cell traversal was first described when *P. berghei* sporozoites were observed to transit through host macrophages and later hepatocytes and epithelial cells [20]. There is debate about whether cell traversal involves the formation of a transient parasitophorous vacuole or if it ruptures the host cell membrane [21]. Several parasite proteins involved in cell traversal have been identified, with knockout of genes encoding either SPECT1 (Sporozoite Protein Essential for Cell Traversal) [22] and SPECT2 [23, 24] found to affect sporozoite traversal. The sequence similarities of SPECT2 with membrane pore-forming proteins and the ability of purified recombinant protein to disrupt host membranes suggest that SPECT2 functions by interacting with host phospholipids to perturb membranes [25]. The role of SPECT1, however, is less clear. In the liver, sporozoites cross the sinusoidal barrier mainly by traversing Kupffer cells: specialised macrophages that line the endothelial vessels of the liver sinusoids. SPECT1-deficient sporozoites are weakly infective *in vivo*, but regain normal infectivity if Kupffer cells are depleted [22, 24]. A more recent study has shown that when endothelial cells are loaded with dye that is dispersed upon cell wounding, these cells can also be used to enter the liver parenchyma and that cell traversal enables the parasites to avoid phagocytic destruction by Kupffer cells (Figure 1B) [26]. Given the large size of Kupffer cells compared to endothelial cells, sporozoites are more likely to contact these cells and use their ability to traverse cells to avoid phagocytosis and successfully navigate entry into the liver [26]. In addition, the same mechanisms that are used to traverse cells in the mammalian host are likely to operate in the mosquito and a small secreted sporozoite protein called CeLTOS has been shown to be necessary for crossing the cytoplasm of both hepatocytes and the insect midgut epithelium [27]. The CeLTOS crystal structure and ability of the recombinant protein to bind to lipids suggests that it may also act by helping to perturb the host cell membrane during traversal [28]. Together these studies highlight important roles for cell traversal in transmission and infection that are independent of invasion.

Sporozoite invasion

After crossing the sinusoidal barrier, sporozoites switch to an invasive state and invade hepatocytes with the concurrent formation of a parasitophorous vacuole membrane [29-31]. Invasion involves the regulated secretion of parasite ligands from apical organelles to form a membranous connection called a tight junction between the sporozoite apical tip and the host cell, which opens up in a ring-like structure through which the sporozoite invades using the same molecular machinery that powers gliding motility [32]. The parasite surface proteins AMA1 and RON2 are thought to be essential for tight junction formation [33] but knockdown of AMA1 to undetectable levels had no effect on sporozoite invasion of hepatocytes [34]. This finding was later confirmed by complete knockout of AMA1 in *P. berghei* sporozoites [35]. In addition, two proteins which contain 6-cysteine domains, P36 and P52, have been implicated in hepatocyte invasion although their precise roles are still unclear. *P. berghei* and *P. yoelli* sporozoites with targeted mutations in both *p36* and *p52*

were able to traverse cells but did not commit to productive invasion [36]; however, *P. falciparum* sporozoites lacking both genes appeared to invade cells but were instead arrested during liver stage development [37]. It is likely that P36 and P52 are required after invasion for proper formation of the parasitophorous vacuole [37], which protects it from destruction once in the host hepatocyte. Thus, the molecules involved in sporozoite invasion of hepatocytes and their precise functions are still incompletely defined. As we will discuss, a small number of host cell surface proteins such as CD81, SR-BI, and CD68 are now known to be important for sporozoite traversal and invasion, but exactly how these interact with specific parasite ligands is also still undetermined.

Switching between behaviors

Major unanswered questions are how the sporozoite switches between different cellular behaviors during its migration from the dermis to the liver, and how the sporozoite distinguishes the liver from other host tissues. The most abundant protein displayed on the surface of the sporozoite, CSP, is thought to sense the relatively higher level of sulfation of heparan sulfate proteoglycans (HSPGs) displayed by the stellate cells that protrude into the vascular lumen through fenestrations in the hepatic vascular endothelium [20, 38] (Figure 1B). How sporozoites switch from a traversal to an invasive phenotype is also the subject of debate. Early work suggested that it was the traversal of hepatocytes that allowed the sporozoites to switch, either by increasing secretion of organelles required for invasion [39], or by increased exposure to intracellular potassium [40]. It was also suggested that wounded hepatocytes secrete hepatocyte growth factor, which, after interacting with its receptor on neighboring cells, made them more susceptible to infection [41]; however, this was found to apply only in *P. berghei* and not *P. yoelii* or *P. falciparum* [42]. These theories are contradicted by the fact that mutant sporozoites deficient in cell traversal can still infect hepatocytes when Kupffer cells are depleted [22, 24]. One model that does not require cell traversal to trigger the switch to invasion again invokes the presence of highly sulfated HSPGs within the liver, but to trigger the pro-invasive proteolytic cleavage of CSP [30]; however, this too remains to be unequivocally demonstrated. Overall, the host molecular cues and how these are perceived by the sporozoite in the extracellular environment remain poorly defined.

Identifying extracellular host-parasite interactions

Extracellular recognition events are critical for host-pathogen interactions and for defining sporozoite cell and host tropisms; however, we know little about the host factors that act as molecular guide-posts for sporozoites. Given their importance for infection biology and their potential for therapeutic intervention, why is this so? Likely reasons include the limited amounts of sporozoite material which, by contrast to blood stages, cannot be expanded *in vitro*, and must be dissected from mosquito salivary glands. Furthermore, proteins that bridge the plasma membrane are typically biochemically intractable: their amphipathic nature makes them difficult to solubilize in their native conformation, and extracellular receptor-ligand interactions are often highly transient such that they cannot withstand wash steps common to biochemical purification protocols [43]. These challenges could be tackled by using soluble recombinant proteins corresponding to ectodomains which often retain their

receptor binding activity; however, a suitable protein expression system must be selected because these regions occupy an oxidizing environment and structurally-critical post-translational modifications such as disulfide bonds must be formed correctly.

Since many of these challenges are not specific to parasites, recent advances in eukaryotic recombinant protein expression systems have enabled soluble versions of the ectodomains of *Plasmodium* ligands to be produced in biochemically active forms [44]. These can be used as binding probes to detect interactions with host tissues and molecules; however, so far, no widely accepted host-sporozoite interactions have been identified using standard biochemical approaches. Some success has been achieved using a bespoke protein interaction assay designed to detect transient extracellular protein interactions called AVEXIS (for AVidity-based EXtracellular protein Interaction Screening) [45]. This method uses direct binding within libraries of recombinant proteins constituting the extracellular regions of host and parasite proteins in an ELISA-style assay. This has successfully identified host receptors for blood stage *Plasmodium* ligands [46] and more recently identified alpha-v-containing host integrins as a ligand for TRAP in sporozoites [47] (Figure 1A). One challenge with this approach is that the sporozoite interacts with many host cell types, making the library of recombinant proteins impractically large for most laboratories. Other approaches include screening cellular microarrays that have been transfected with cDNA constructs representing a large number host receptors [48]. Again, such large collections of reagents are usually beyond the reach of most laboratories, but the recent development of cell-based genome-wide genetic loss of function screens provide the possibility of identifying host receptors without needing large reagent collections [49].

Important sporozoite surface proteins and known host interactions

The two most abundant proteins displayed at the surface of the sporozoite are CSP and TRAP. The functional role of both proteins has been investigated in detail.

Circumsporozoite protein (CSP)

The overall structure of CSP is conserved among *Plasmodium* species, with a central repeat region whose sequence varies between species, a C-terminal type I thrombospondin repeat (TSR) adhesion domain, and a conserved five amino acid sequence N-terminal to the repeats that serves as a proteolytic cleavage site [29]. Deletion of the *csp* gene results in a severe defect in sporozoite development [50], and studies indicate the TSR domain is likely involved in this process [51]. The efficiency of sporozoite salivary gland invasion and infection of the mammalian host critically depends on interactions between CSP and both mosquito and mammalian host tissues. During sporozoite migration, CSP is folded such that the TSR is masked, a conformation that enables sporozoites to efficiently navigate through tissues [17, 29]. Once in the liver, a parasite protease cleaves CSP, leading to the removal of the N-terminal third of the protein and exposure of the TSR [29, 52]. The signal for this cleavage event is, in part, reliant upon sporozoite contact with HSPGs, and the resulting exposure of the TSR is associated with a switch from a migratory to an invasive phenotype [30].

The controlled exposure of CSP domains suggests that different portions of CSP may function in the recognition of host tissues; however, there remains much to be learnt about these events. Several studies have found that in the mosquito, the CSP N-terminus binds to the salivary glands [53, 54] and inhibition of this leads to lower salivary gland infections [55, 56]. Previous studies describing CSP binding to HSPGs in the mammalian host (described below) spurred the search for similar molecules in the mosquito. HSPGs are indeed found on mosquito salivary glands [57] and these can mediate salivary gland invasion [58]; however, in their absence, other surface molecules compensate. This suggests that HSPGs may mediate initial adhesion to mosquito salivary glands via the N-terminus of CSP, but that other interactions are involved; for example, a yet to be fully characterized CSP-binding protein on mosquito salivary glands has been described [54]. Other secreted or sporozoite surface proteins such as TREP [59], a member of the TRAP family, and MAEBL [8, 60], are also essential for salivary gland invasion although their binding partners have not been identified.

In the mammalian host, a considerable body of work has demonstrated that CSP binds to HSPGs. Initial studies demonstrated that recombinant CSP bound to hepatocyte HSPGs [61] and that this was more important under shear stress conditions [38], suggesting it may mediate localization of sporozoites to the liver but not hepatocyte invasion *per se*. Indeed, *in vitro* studies demonstrated that heparin had a negligible effect on sporozoite invasion into hepatocytes under static conditions [38, 62]. The identity of the heparin-binding domain of CSP remains controversial. Initial studies using peptides representing portions of the TSR in binding competition assays suggested that the TSR had heparin-binding activity [61]; however, a recent crystal structure of the CSP TSR was described and this protein had no heparin-binding activity [63], suggesting that previous data with peptides did not reflect the activity of the native protein. In subsequent studies, peptides and recombinant protein representing the N-terminus of CSP were shown to have high affinity heparin-binding activity [64], raising the possibility that the CSP N-terminus binds to hepatic HSPGs, a scenario consistent with what we currently know of CSP domain conformation. Since CSP densely coats the sporozoite, one can imagine that CSP cleavage and subsequent exposure of an array of TSRs would make for a very adhesive sporozoite that could, upon binding to a yet-to-be identified receptor, provide the signal for it to switch to an invasive phenotype. How can hepatic HSPGs provide a signal to sporozoites that they are in the liver when HSPGs are found in many tissues and cell types? Previous studies found that it was only the highly sulfated HSPGs that bound to CSP, and the degree of sulfation but not a specific pattern of sulfation or specific saccharide that was critical for sporozoite binding and induction of CSP cleavage [30, 38]. In the mammalian host, the only highly sulfated HSPGs exposed to the circulation are in the liver sinusoids, explaining how they can account for the specific arrest of sporozoites in this organ.

Thrombospondin-related anonymous protein (TRAP)

TRAP is a type I transmembrane protein containing a von Willebrand factor A (VWA) domain and a TSR domain in its extracellular region. It is the founding member of a family of TRAP-like proteins that are expressed in sporozoites (such as TLP, TREP), and other invasive *Plasmodium* stages (MTRAP and CTRP) [32]. TRAP is mainly localized to

micronemes [65, 66] but is trafficked to the surface and shed from the posterior end of gliding sporozoites [19]. TRAP-deficient *P. berghei* parasites exhibit multiple defects: they are unable to glide *in vitro*, and are significantly impaired in their ability to invade mosquito salivary glands and infect rodent hosts. These and other data provide evidence that TRAP is an essential component of the sporozoite gliding motility and invasion machinery [13]. Consistent with this, parasites with mutations in the cytoplasmic region of TRAP exhibit abnormal gliding and are unable to infect mice or invade hepatocytes *in vitro* [67].

Early work suggested that TRAP interacted directly with HSPGs, identifying a conserved motif within the TSR domain that is important for TRAP binding to glycans in plate-based assays and on hepatocytes [68]. Genetic studies of TRAP in *P. berghei* have further dissected the roles of the VWA and TSR domains, with both domains being found to be important in host cell invasion, but not gliding motility [69]. The structure of the TRAP ectodomain was recently solved and broadened our understanding of TRAP function since it found TRAP adopted at least two conformations: a compact “closed” form and an extended “open” conformation [70]. One possibility is that binding of TRAP to a host receptor triggers this conformational change, generating additional force that is transmitted along the extended structure. Such a phenomenon might provide insight into the mechanism for the proposed saltatory ‘stick and slip’ model of gliding motility [18].

Our recent study identified α_v -containing integrins as host receptors for *P. falciparum* TRAP [47]. The interaction between TRAP and integrin was shown to be high affinity, and binding was enhanced by manganese ions, a known integrin activator, demonstrating that TRAP may bind preferentially to integrins in their activated ligand binding state (Figure 1A). This preference may restrict TRAP-mediated interactions to clusters of activated integrins such as at focal adhesion sites. There was no evidence that this interaction was involved in hepatocyte invasion, but *P. falciparum* sporozoites did glide more rapidly and with a greater displacement in integrin-deficient mice. This suggests that the role for this interaction might be to act as ‘stop’ signal in the dermis in response to a localized signal from the host [47]. However, further work is this required to more fully understand the role of this interaction in sporozoite migration and infection.

Other host proteins important for sporozoite infections

Erythrocyte invasion by *Plasmodium* merozoites provide an important paradigm for understanding molecular interactions required for host cell invasion. Here, we know ~50 different cell surface or secreted parasite proteins are involved [11] and have begun to delineate the order in which they are deployed to interact with host receptors during the multi-step invasion process [71]. There is every reason to believe that traversal and invasion of hepatocytes by sporozoites is equally complex, but so far, only a handful of host receptor proteins have been identified including CD81, SR-B1, CD68 and potentially EPHA2.

In a remarkable finding, *Cd81*-deficient mice were shown to be refractory to *P. yoelii* liver, but not blood stage infection, leading the authors to conclude that host CD81 was required for both *P. yoelii* and *P. falciparum* sporozoites to invade hepatocytes [72]. CD81 belongs to the class of tetraspanin receptors, so-named because they traverse the membrane four times

[73]. CD81 and other tetraspanins such as CD9 have documented roles in manipulation of membranes and are involved in membrane fusion events including fertilization [74] and viral infections [75]. Host CD81 was later shown to be necessary for the discharge of *P. yoelii* sporozoite rhoptry contents and may be the source of the host signal that switches sporozoites from cell traversal to invasion, since parasites preferentially traverse cells lacking CD81, but require it for productive invasion [76, 77]. Because of these cell-type specific functions but broad expression pattern, CD81 is thought to be required for the organisation of other tissue-restricted cell surface proteins, and/or to stabilise intermediates in membrane curvature needed for rhoptry discharge and parasitophorous vacuole formation. However, while undoubtedly critical for both *P. yoelii* and *P. falciparum* sporozoite invasion [72] the precise role of CD81 in invasion is currently unknown.

Using clues that the host lipoprotein clearance pathway was important for liver infection [78], the scavenger receptor BI (SR-BI) was found to have dual roles in *P. berghei* and *P. falciparum* sporozoite invasion and development within hepatocytes [79, 80]. The amount of CD81 detectable at the cell surface was significantly decreased in cells lacking SR-BI, possibly due to a reduction in cellular cholesterol levels. In light of the recent crystal structure of CD81 that showed a cholesterol molecule bound within the intramembranous pocket formed by the four transmembrane-spanning helices, these data could suggest this is an important feature of CD81 and hence SB-RI function [73].

Searching for host surface proteins important for sporozoite infection, Cha and colleagues [81] selected peptide-displaying phages from a large library that bound intact Kupffer cells and prevented *P. berghei* sporozoite entry. They identified the target of these entry-blocking peptides as CD68: a large heavily O-glycosylated type I membrane protein mainly localised to the endosomes and lysosomes of macrophages and monocytes. *Cd68*-deficient mice exhibited a strong but incomplete reduction in *P. berghei* liver infection suggesting an important but redundant role for CD68 and Kupffer cells in the exit of sporozoites from circulation, consistent with other sporozoite routes to the liver [26]. Host CD68, therefore, appears to play at least some role in sporozoite traversal prior to hepatocyte invasion.

Exploiting differences in susceptibility of different mouse strains and hepatocytes to *P. yoelii* infection, the Ephrin receptor A2 (EPHA2) was implicated in sporozoite invasion of hepatocytes, with higher levels of EPHA2 expression correlated with increased invasion by *P. yoelii* and *P. falciparum* sporozoites [82]. The same study also found *Epha2*-deficient mice were less susceptible to infection with *P. yoelii* sporozoites, and that antibodies to EPHA2 could markedly inhibit sporozoite invasion *in vitro*. Together with other data, these findings supported the conclusion that EPHA2 was important for sporozoite invasion [82]. Recent studies using siRNA to knockdown EPHA2 expression in hepatocytes and similar antibody inhibition studies, however, did not support a role for EPHA2 in hepatocyte invasion for either *P. yoelii* or *P. berghei* [83]. EPHA2 knockdown appeared to have no clear effect on sporozoite invasion regardless of the hepatocyte source, and antibodies to EPHA2 showed no greater ability to inhibit sporozoite invasion compared to the control [83]. In light of these conflicting results, the precise role of EPHA2 in sporozoite invasion remains an open question.

For all of these host receptors, their role in sporozoite infection would be greatly clarified by identifying the parasite molecules they interact with. While there are data to suggest that EPHA2 might interact with P36 [82], so far there has been no conclusive evidence for any of these receptors interacting directly with a parasite protein. Clarifying the molecular events involved in host-sporozoite interactions is further complicated because experiments suggest, as might be expected, that different species of *Plasmodium* use different host factors. For example, CD81 is important for the invasion of hepatocytes by *P. yoelii* and *P. falciparum*, but is not required by *P. vivax*, and is redundant with SR-BI for *P. berghei* [36]. Similarly, SR-BI is required for *P. vivax* but not *P. falciparum* or *P. yoelii* [36]. These differences make the defining of paradigms that can be extrapolated from experimentally more tractable rodent models to the parasites that cause disease in humans difficult.

Concluding remarks

We are far from having a good molecular understanding of how sporozoites recognise and interact with mammalian host cells (see Outstanding Questions). Progress has been hard-won because of experimental challenges associated with manipulation of sporozoites and membrane-embedded cell surface proteins. Sporozoites can only be produced in mosquitoes, and the relatively modest numbers that can be obtained together with the inevitable mosquito contamination have significantly hindered our ability to cleanly define host-pathogen interactions of this stage of the *Plasmodium* life cycle. Future advances will rely on the development and application of new techniques and models including *in vivo* imaging technologies, humanised mice, and methods to identify extracellular protein interactions. There are many reasons to be optimistic, particularly in light of recent genome-scale genetic techniques to manipulate the parasite [84, 85] and host [86]. Future discoveries will not only provide a molecular explanation of how sporozoites interact with host cells, but also aid the development of pre-erythrocytic vaccines that might ultimately lead to reducing and eliminating the burden of malaria.

Acknowledgements

This work was supported by the Wellcome Trust grant 206194, Medical Research Council Doctoral Training Grant (MR/J004111/1), EU Horizon 2020 (Agreement 733273), and the National Institutes of Health (RO1 AI056840 and RO1 AI132359 to PS).

Glossary

Gliding motility:

A type of substrate-dependent motility that does not involve changes to the cell shape, but instead relies on the parasite cytoskeleton and secreted cell surface proteins. It is required for sporozoite migration, cell traversal, and invasion.

Cell traversal:

A sporozoite behaviour that allows for transient passage through a host cell in order to migrate through cells or tissues and escape phagocytic destruction. As sporozoites move through and exit host cells they wound the host cell plasma membrane.

Cell invasion:

A sporozoite behaviour that allows for entry into a host cell to establish an infection. It involves the formation of a parasitophorous vacuole membrane around the parasite that separates it from the host cell cytoplasm.

Thrombospondin-related anonymous protein (TRAP):

An abundant transmembrane protein that is secreted onto the sporozoite surface during gliding motility. TRAP is a critical component of the sporozoite's motility machinery, serving to link intracellular cytoskeletal elements with extracellular substrates or ligands. It is essential for sporozoite motility and hence normal traversal and invasion.

Circumsporozoite protein (CSP):

A highly abundant, multifunctional protein that densely covers the surface of sporozoites. CSP is essential for sporozoite formation in the mosquito host and guiding the migration of sporozoites from mosquito to mammalian host. CSP is highly immunogenic, and is the target of the current leading anti-sporozoite vaccine.

Avidity-based Extracellular Interaction Screen (AVEXIS):

A method designed to detect low affinity extracellular protein interactions that can be scaled to systematically screen many thousands of potential interactions. It has been used to identify novel endogenous interactions between receptor proteins displayed on cells of the same species and to identify host-pathogen interactions.

Heparan sulfate proteoglycans (HSPGs)

Cell surface proteins on animal cells that contain linear sulfated glycosaminoglycan polysaccharides. HSPGs are a major constituent of the glycocalyx and are thought to be more highly sulfated on the surface of stellate cells in the liver and in the loose basement membrane of the liver called the Space of Disse.

References

1. WHO, World malaria report. 2017.
2. Medica DL and Sinnis P (2005) Quantitative dynamics of *Plasmodium yoelii* sporozoite transmission by infected anopheline mosquitoes. *Infect Immun.* 73, 4363–9. [PubMed: 15972531]
3. Sinnis P and Zavala F (2012) The skin: where malaria infection and the host immune response begin. *Semin Immunopathol.* 34, 787–92. [PubMed: 23053392]
4. Draper SJ et al. (2018) Malaria Vaccines: Recent Advances and New Horizons. *Cell Host Microbe.* 24, 43–56. [PubMed: 30001524]
5. Kudryashev M et al. (2012) Structural basis for chirality and directional motility of *Plasmodium* sporozoites. *Cell Microbiol.* 14, 1757–68. [PubMed: 22776715]
6. Hegge S et al. (2009) Automated classification of *Plasmodium* sporozoite movement patterns reveals a shift towards productive motility during salivary gland infection. *Biotechnol J.* 4, 903–13. [PubMed: 19455538]
7. Sato Y et al. (2014) *Plasmodium berghei* sporozoites acquire virulence and immunogenicity during mosquito hemocoel transit. *Infect Immun.* 82, 1164–72. [PubMed: 24379288]
8. Kariu T et al. (2002) MAEBL is essential for malarial sporozoite infection of the mosquito salivary gland. *J Exp Med.* 195, 1317–23. [PubMed: 12021311]
9. Aly AS et al. (2009) Malaria parasite development in the mosquito and infection of the mammalian host. *Annu Rev Microbiol.* 63, 195–221. [PubMed: 19575563]

10. Lindner SE et al. (2013) Total and putative surface proteomics of malaria parasite salivary gland sporozoites. *Mol Cell Proteomics*. 12, 1127–43. [PubMed: 23325771]
11. Cowman AF et al. (2017) The Molecular Basis of Erythrocyte Invasion by Malaria Parasites. *Cell Host Microbe*. 22, 232–245. [PubMed: 28799908]
12. Wang Q et al. (2005) Mutational analysis of the GPI-anchor addition sequence from the circumsporozoite protein of Plasmodium. *Cell Microbiol*. 7, 1616–26. [PubMed: 16207248]
13. Sultan AA et al. (1997) TRAP is necessary for gliding motility and infectivity of plasmodium sporozoites. *Cell*. 90, 511–22. [PubMed: 9267031]
14. Swearingen KE et al. (2016) Interrogating the Plasmodium Sporozoite Surface: Identification of Surface-Exposed Proteins and Demonstration of Glycosylation on CSP and TRAP by Mass Spectrometry-Based Proteomics. *PLoS Pathog*. 12, e1005606. [PubMed: 27128092]
15. Heintzelman MB (2015) Gliding motility in apicomplexan parasites. *Semin Cell Dev Biol*. 46, 135–42. [PubMed: 26428297]
16. Amino R et al. (2006) Quantitative imaging of Plasmodium transmission from mosquito to mammal. *Nat Med*. 12, 220–4. [PubMed: 16429144]
17. Hopp CS et al. (2015) Longitudinal analysis of Plasmodium sporozoite motility in the dermis reveals component of blood vessel recognition. *Elife*. 4, e07789.
18. Munter S et al. (2009) Plasmodium sporozoite motility is modulated by the turnover of discrete adhesion sites. *Cell Host Microbe*. 6, 551–62. [PubMed: 20006843]
19. Ejigiri I et al. (2012) Shedding of TRAP by a rhomboid protease from the malaria sporozoite surface is essential for gliding motility and sporozoite infectivity. *PLoS Pathog*. 8, e1002725. [PubMed: 22911675]
20. Menard R et al. (2013) Looking under the skin: the first steps in malarial infection and immunity. *Nat Rev Microbiol*. 11, 701–12. [PubMed: 24037451]
21. Yang AS and Boddey JA (2017) Molecular mechanisms of host cell traversal by malaria sporozoites. *Int J Parasitol*. 47, 129–136. [PubMed: 27825827]
22. Ishino T et al. (2004) Cell-passage activity is required for the malarial parasite to cross the liver sinusoidal cell layer. *PLoS Biol*. 2, e4. [PubMed: 14737184]
23. Kaiser K et al. (2004) A member of a conserved Plasmodium protein family with membrane-attack complex/perforin (MACPF)-like domains localizes to the micronemes of sporozoites. *Mol Biochem Parasitol*. 133, 15–26. [PubMed: 14668008]
24. Ishino T et al. (2005) A Plasmodium sporozoite protein with a membrane attack complex domain is required for breaching the liver sinusoidal cell layer prior to hepatocyte infection. *Cell Microbiol*. 7, 199–208. [PubMed: 15659064]
25. Garg S et al. (2013) Calcium-dependent permeabilization of erythrocytes by a perforinlike protein during egress of malaria parasites. *Nat Commun*. 4, 1736. [PubMed: 23591903]
26. Tavares J et al. (2013) Role of host cell traversal by the malaria sporozoite during liver infection. *J Exp Med*. 210, 905–15. [PubMed: 23610126]
27. Kariu T et al. (2006) CelTOS, a novel malarial protein that mediates transmission to mosquito and vertebrate hosts. *Mol Microbiol*. 59, 1369–79. [PubMed: 16468982]
28. Jimah JR et al. (2016) Malaria parasite CelTOS targets the inner leaflet of cell membranes for pore-dependent disruption. *Elife*. 5, e20621. [PubMed: 27906127]
29. Coppi A et al. (2011) The malaria circumsporozoite protein has two functional domains, each with distinct roles as sporozoites journey from mosquito to mammalian host. *J Exp Med*. 208, 341–56. [PubMed: 21262960]
30. Coppi A et al. (2007) Heparan sulfate proteoglycans provide a signal to Plasmodium sporozoites to stop migrating and productively invade host cells. *Cell Host Microbe*. 2, 316–27. [PubMed: 18005753]
31. Mota MM et al. (2001) Migration of Plasmodium sporozoites through cells before infection. *Science*. 291, 141–4. [PubMed: 11141568]
32. Morahan BJ et al. (2009) No TRAP, no invasion. *Trends Parasitol*. 25, 77–84. [PubMed: 19101208]
33. Besteiro S et al. (2011) The moving junction of apicomplexan parasites: a key structure for invasion. *Cell Microbiol*. 13, 797–805. [PubMed: 21535344]

34. Giovannini D et al. (2011) Independent roles of apical membrane antigen 1 and rhoptry neck proteins during host cell invasion by apicomplexa. *Cell Host Microbe*. 10, 591–602. [PubMed: 22177563]
35. Bargieri DY et al. (2013) Apical membrane antigen 1 mediates apicomplexan parasite attachment but is dispensable for host cell invasion. *Nat Commun*. 4, 2552. [PubMed: 24108241]
36. Manzoni G et al. (2017) Plasmodium P36 determines host cell receptor usage during sporozoite invasion. *Elife*. 6, e25903. [PubMed: 28506360]
37. VanBuskirk KM et al. (2009) Preerythrocytic, live-attenuated Plasmodium falciparum vaccine candidates by design. *Proc Natl Acad Sci U S A*. 106, 13004–9. [PubMed: 19625622]
38. Pinzon-Ortiz C et al. (2001) The binding of the circumsporozoite protein to cell surface heparan sulfate proteoglycans is required for plasmodium sporozoite attachment to target cells. *J Biol Chem*. 276, 26784–91. [PubMed: 11352923]
39. Mota MM et al. (2002) Migration through host cells activates Plasmodium sporozoites for infection. *Nat Med*. 8, 1318–22. [PubMed: 12379848]
40. Kumar KA et al. (2007) Exposure of Plasmodium sporozoites to the intracellular concentration of potassium enhances infectivity and reduces cell passage activity. *Mol Biochem Parasitol*. 156, 32–40. [PubMed: 17714805]
41. Carrolo M et al. (2003) Hepatocyte growth factor and its receptor are required for malaria infection. *Nat Med*. 9, 1363–9. [PubMed: 14556002]
42. Kaushansky A and Kappe SH (2011) The crucial role of hepatocyte growth factor receptor during liver-stage infection is not conserved among Plasmodium species. *Nat Med*. 17, 1180–1. [PubMed: 21988987]
43. Wright GJ (2009) Signal initiation in biological systems: the properties and detection of transient extracellular protein interactions. *Mol Biosyst*. 5, 1405–12. [PubMed: 19593473]
44. Crosnier C et al. (2013) A library of functional recombinant cell-surface and secreted P. falciparum merozoite proteins. *Mol Cell Proteomics*. 12, 3976–86. [PubMed: 24043421]
45. Bushell KM et al. (2008) Large-scale screening for novel low-affinity extracellular protein interactions. *Genome Res*. 18, 622–30. [PubMed: 18296487]
46. Bartholdson SJ et al. (2013) Identifying novel Plasmodium falciparum erythrocyte invasion receptors using systematic extracellular protein interaction screens. *Cell Microbiol*. 15, 1304–12. [PubMed: 23617720]
47. Dundas K et al. (2018) Alpha-v-containing integrins are host receptors for the Plasmodium falciparum sporozoite surface protein, TRAP. *Proc Natl Acad Sci U S A*. 115, 4477–4482. [PubMed: 29632205]
48. Turner L et al. (2013) Severe malaria is associated with parasite binding to endothelial protein C receptor. *Nature*. 498, 502–5. [PubMed: 23739325]
49. Sharma S et al. (2018) Genome-scale identification of cellular pathways required for cell surface recognition. *Genome Res*. 28, 1372–1382. [PubMed: 29914970]
50. Menard R et al. (1997) Circumsporozoite protein is required for development of malaria sporozoites in mosquitoes. *Nature*. 385, 336–40. [PubMed: 9002517]
51. Ferguson DJ et al. (2014) The repeat region of the circumsporozoite protein is critical for sporozoite formation and maturation in Plasmodium. *PLoS One*. 9, e113923. [PubMed: 25438048]
52. Coppi A et al. (2005) The Plasmodium circumsporozoite protein is proteolytically processed during cell invasion. *J Exp Med*. 201, 27–33. [PubMed: 15630135]
53. Sidjanski SP et al. (1997) Anopheles stephensi salivary glands bear receptors for region I of the circumsporozoite protein of Plasmodium falciparum. *Mol Biochem Parasitol*. 90, 33–41. [PubMed: 9497030]
54. Wang J et al. (2013) Anopheles gambiae circumsporozoite protein-binding protein facilitates plasmodium infection of mosquito salivary glands. *J Infect Dis*. 208, 1161–9. [PubMed: 23801601]
55. Myung JM et al. (2004) The Plasmodium circumsporozoite protein is involved in mosquito salivary gland invasion by sporozoites. *Mol Biochem Parasitol*. 133, 53–9. [PubMed: 14668012]

56. Kojin BB et al. (2016) Endogenously-expressed NH₂-terminus of circumsporozoite protein interferes with sporozoite invasion of mosquito salivary glands. *Malar J.* 15, 153. [PubMed: 26964736]
57. Sinnis P et al. (2007) Mosquito heparan sulfate and its potential role in malaria infection and transmission. *J Biol Chem.* 282, 25376–84. [PubMed: 17597060]
58. Armistead JS et al. (2011) A role for heparan sulfate proteoglycans in *Plasmodium falciparum* sporozoite invasion of anopheline mosquito salivary glands. *Biochem J.* 438, 475–83. [PubMed: 21663594]
59. Combe A et al. (2009) TREP, a novel protein necessary for gliding motility of the malaria sporozoite. *Int J Parasitol.* 39, 489–96. [PubMed: 19000911]
60. Saenz FE et al. (2008) The transmembrane isoform of *Plasmodium falciparum* MAEBL is essential for the invasion of *Anopheles* salivary glands. *PLoS One.* 3, e2287. [PubMed: 18509478]
61. Frevert U et al. (1993) Malaria circumsporozoite protein binds to heparan sulfate proteoglycans associated with the surface membrane of hepatocytes. *J Exp Med.* 177, 1287–98. [PubMed: 8478608]
62. Frevert U et al. (1996) Cell surface glycosaminoglycans are not obligatory for *Plasmodium berghei* sporozoite invasion in vitro. *Mol Biochem Parasitol.* 76, 257–66. [PubMed: 8920011]
63. Doud MB et al. (2012) Unexpected fold in the circumsporozoite protein target of malaria vaccines. *Proc Natl Acad Sci U S A.* 109, 7817–22. [PubMed: 22547819]
64. Ancsin JB and Kisilevsky R (2004) A binding site for highly sulfated heparan sulfate is identified in the N terminus of the circumsporozoite protein: significance for malarial sporozoite attachment to hepatocytes. *J Biol Chem.* 279, 21824–32. [PubMed: 15007056]
65. Cowan G et al. (1992) Expression of thrombospondin-related anonymous protein in *Plasmodium falciparum* sporozoites. *Lancet.* 339, 1412–3. [PubMed: 1350818]
66. Rogers WO et al. (1992) Characterization of *Plasmodium falciparum* sporozoite surface protein 2. *Proc Natl Acad Sci U S A.* 89, 9176–80. [PubMed: 1409621]
67. Kappe S et al. (1999) Conservation of a gliding motility and cell invasion machinery in Apicomplexan parasites. *J Cell Biol.* 147, 937–44. [PubMed: 10579715]
68. Muller HM et al. (1993) Thrombospondin related anonymous protein (TRAP) of *Plasmodium falciparum* binds specifically to sulfated glycoconjugates and to HepG2 hepatoma cells suggesting a role for this molecule in sporozoite invasion of hepatocytes. *EMBO J.* 12, 2881–9. [PubMed: 8392935]
69. Matuschewski K et al. (2002) *Plasmodium* sporozoite invasion into insect and mammalian cells is directed by the same dual binding system. *EMBO J.* 21, 1597–606. [PubMed: 11927544]
70. Song G et al. (2012) Shape change in the receptor for gliding motility in *Plasmodium* sporozoites. *Proc Natl Acad Sci U S A.* 109, 21420–5. [PubMed: 23236185]
71. Weiss GE et al. (2015) Revealing the sequence and resulting cellular morphology of receptor-ligand interactions during *Plasmodium falciparum* invasion of erythrocytes. *PLoS Pathog.* 11, e1004670. [PubMed: 25723550]
72. Silvie O et al. (2003) Hepatocyte CD81 is required for *Plasmodium falciparum* and *Plasmodium yoelii* sporozoite infectivity. *Nat Med.* 9, 93–6. [PubMed: 12483205]
73. Zimmerman B et al. (2016) Crystal Structure of a Full-Length Human Tetraspanin Reveals a Cholesterol-Binding Pocket. *Cell.* 167, 1041–1051 e11. [PubMed: 27881302]
74. Rubinstein E et al. (2006) Reduced fertility of female mice lacking CD81. *Dev Biol.* 290, 351–8. [PubMed: 16380109]
75. Sharma NR et al. (2011) Hepatitis C virus is primed by CD81 protein for low pH-dependent fusion. *J Biol Chem.* 286, 30361–76. [PubMed: 21737455]
76. Risco-Castillo V et al. (2014) CD81 is required for rhoptry discharge during host cell invasion by *Plasmodium yoelii* sporozoites. *Cell Microbiol.* 16, 1533–48. [PubMed: 24798694]
77. Risco-Castillo V et al. (2015) Malaria Sporozoites Traverse Host Cells within Transient Vacuoles. *Cell Host Microbe.* 18, 593–603. [PubMed: 26607162]
78. Sinnis P et al. (1996) Remnant lipoproteins inhibit malaria sporozoite invasion of hepatocytes. *J Exp Med.* 184, 945–54. [PubMed: 9064354]

79. Rodrigues CD et al. (2008) Host scavenger receptor SR-BI plays a dual role in the establishment of malaria parasite liver infection. *Cell Host Microbe*. 4, 271–82. [PubMed: 18779053]
80. Yalaoui S et al. (2008) Scavenger receptor BI boosts hepatocyte permissiveness to Plasmodium infection. *Cell Host Microbe*. 4, 283–92. [PubMed: 18779054]
81. Cha SJ et al. (2015) CD68 acts as a major gateway for malaria sporozoite liver infection. *J Exp Med*. 212, 1391–403. [PubMed: 26216124]
82. Kaushansky A et al. (2015) Malaria parasites target the hepatocyte receptor EphA2 for successful host infection. *Science*. 350, 1089–92. [PubMed: 26612952]
83. Langlois AC et al. (2018) Plasmodium sporozoites can invade hepatocytic cells independently of the Ephrin receptor A2. *PLoS One*. 13, e0200032. [PubMed: 29975762]
84. Bushell E et al. (2017) Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. *Cell*. 170, 260–272 e8. [PubMed: 28708996]
85. Zhang M et al. (2018) Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis. *Science*. 360, eaap7847. [PubMed: 29724925]
86. Puschnik AS et al. (2017) A CRISPR toolbox to study virus-host interactions. *Nat Rev Microbiol*. 15, 351–364. [PubMed: 28420884]

Highlights

- Cellular descriptions of sporozoite behaviors and host cell interactions are increasingly well defined; however, signals triggering sporozoite behavioral switches remain unknown.
- The molecular basis of sporozoite-host recognition is poorly characterised due to technical challenges that have limited progress.
- New genetic and biochemical techniques are addressing some of these challenges and have identified extracellular sporozoite-host interactions involved in host infection.
- Using these new tools to understand the molecular basis of host-sporozoite interactions will help improve anti-malarial treatments and vaccines.

Outstanding questions

- What are the molecular identities of the host-sporozoite receptor-ligand interactions responsible for cellular recognition events within the host?
- What are the interactions that signal to the sporozoite to become motile and to switch to an invasive state?
- Specifically, what is the identity of the interactions that are responsible for:
 - Providing a substrate for sporozoite gliding;
 - Specific recognition of vascular endothelial cells;
 - Recognition and arrest within the liver sinusoid;
 - Cell traversal of Kupffer cells and hepatic endothelium;
 - Productive invasion of hepatocytes;
 - Switching from traversal to invasion.
- What are the identities of the proteins and complexes formed on the sporozoite surface that are required for interacting with host receptors?
- What is the order of interactions required for host cell invasion by sporozoites? Are they released from intracellular organelles in a specific schedule? How are these interactions related and different from erythrocyte invasion by *Plasmodium* merozoites, and are they conserved across other Apicomplexan parasites?

Figure 1. Schematic illustrating the extracellular interactions involved in *Plasmodium* sporozoite migration and invasion.

Sporozoites are deposited in the skin of the host when an infected mosquito takes a blood meal, and move within the dermis by an active process known as gliding motility. Sporozoites recognize and traverse the vascular endothelium to enter the bloodstream where they are transported to the liver. (A) The abundant sporozoite surface protein, TRAP, which is essential for gliding motility, has been recently found to interact with αv -containing integrins in their extended active conformation displayed on the surface of host cells, and may be involved in cell host recognition events in the dermis. (B) Sporozoites are transported to the liver where they are arrested within the liver sinusoids: specialized capillary beds bordered by a highly porous epithelium which contain fenestrations that are clustered into structures known as sieve plates. Hepatocytes are not separated from the vascular endothelial cells by the usual basement membrane, but rather by a conspicuous gap known as the space of Disse that are occupied by stellate cells; the lumen of the sinusoid is patrolled by hepatic macrophages called Kupffer cells. Parasites are thought to be arrested in the liver by interactions between the abundant sporozoite surface protein CSP and unusually high levels of sulfated HSPGs displayed by stellate cells that protrude as tufts through the fenestrations in the sinusoid endothelium. The arrested sporozoite enters the liver parenchyma either by cell traversal of Kupffer cells, which appears to involve CD68, or by traversal of endothelial cells. Sporozoites then traverse through several hepatocytes, a process which requires the action of the sporozoite proteins SPECT1, SPECT2 and CelTOS. Eventually, sporozoites productively invade a hepatocyte requiring the parasite proteins P36 and P52, and the host receptors SR-BI, and/or CD81, and potentially EPHA2.