Published in final edited form as:

Cell Calcium. 2019 March ; 78: 48–55. doi:10.1016/j.ceca.2018.12.010.

CRAC Channels in Secretory Epithelial Cell Function and Disease

Haiping Liu, Ahmed Kabrah, Malini Ahuja, and Shmuel Muallem

Epithelial Signaling and Transport Section, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD 20892,

Abstract

The receptor-evoked Ca^{2+} signal in secretory epithelia mediate many cellular functions essential for cell survival and their most fundamental functions of secretory granules exocytosis and fluid and electrolyte secretion. Ca^{2+} influx is a key component of the receptor-evoked Ca^{2+} signal in secretory cell and is mediated by both TRPC and the STIM1-activated Orai1 channels that mediates the Ca^{2+} release-activated current (CRAC) I_{Crac} . The core components of the receptor-evoked Ca^{2+} signal are assembled at the ER/PM junctions where exchange of materials between the plasma membrane and internal organelles take place, including transfer of lipids and Ca^{2+} . The Ca^{2+} signal generated at the confined space of the ER/PM junctions is necessary for activation of the Ca^{2+} -regulated proteins and ion channels that mediate exocytosis with high fidelity and tight control. In this review we discuss the general properties of Ca^{2+} signaling, $\text{PI}(4,5)\text{P}_2$ and other lipids at the ER/PM junctions with regard to secretory cells function and disease caused by uncontrolled Ca^{2+} influx.

Graphical abstract

Introduction

The change in the free cytoplasmic Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) encodes information and controls virtually all functions of secretory epithelial cells, including the cardinal functions of secretory granules exocytosis, fluid and electrolyte secretion [1, 2]. The tight regulation of $[\text{Ca}^{2+}]_i$ occurs in all cellular compartments, from across the plasma membrane (PM) to the

Address for correspondence: Shmuel Muallem, Ph.D., Epithelial Signaling and Transport Section, National Institute of Health, National Institute of Dental and Craniofacial Research, Building 10, Room 1N-112, Bethesda MD 20892., Phone: 301-402-0262, shmuel.muallem@nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

cytoplasmic organelles, including endoplasmic reticulum (ER), mitochondria, lysosomes, endosomes, nucleus and secretory granules. The regulation of $[Ca^{2+}]_i$ is done by Ca^{2+} pumps and Ca^{2+} channels that form and dissipate Ca^{2+} gradients on demands.

The Ca^{2+} gradients across the PM and the ER are generated by the PM Ca^{2+} ATPase (PMCA) pump [3] and the sarco-endoplasmic reticulum ATPase (SERCA) pump [4], respectively. The secretory pathway Ca^{2+} ATPase pumps (SPCA) mainly pump Ca^{2+} into the Golgi [5] while mitochondrial Ca^{2+} uptake is mediated by the mitochondrial Ca^{2+} uniporter complex [6–8]. How Ca^{2+} accumulates in lysosomes, endosomes and secretory granules is not well understood, let alone the identity of the proteins mediating the Ca^{2+} accumulation. Known Ca^{2+} channels that regulate $[Ca^{2+}]_i$ include IP_3 receptors (IP_3Rs) and Ryanodine receptors ($RyRs$), which mediate the release of Ca^{2+} from the ER [9, 10], and store-operated Orai channels [11], the receptor-stimulated TRPC [12, 13], and other TRP channels [13] which control Ca^{2+} influx across the PM. In mitochondria, Ca^{2+} is cleared by the mitochondrial Na^+/Ca^{2+} exchanger NCLX (NCX-9) [14] and Ca^{2+} release from lysosomes and endosomes is mediated by TRPML1 [15] and TRPML3 [16], respectively. Two additional channels involved in lysosomal and endosomal Ca^{2+} release are the TPC channels, TPC1 (mainly endosomes) and TPC2 (mainly lysosomes) [17, 18]. The TPC channels appear to be Na^+ selective and activated by $PI(3,5)P_2$ [19] and NAADP [18, 20] to mediate the NAADP-triggered Ca^{2+} release. IP_3Rs , Ryanodine receptors ($RyRs$), and the TPC channels have all been implicated in Ca^{2+} release from secretory granules [21].

The receptor-evoked Ca^{2+} signal makes use of all Ca^{2+} transporting elements to generate specific Ca^{2+} signals for various cellular functions. The signal starts by interaction of ligand with receptors to activate phospholipase C that hydrolyses $PI(4,5)P_2$ at the PM to release IP_3 and diacylglycerol (DAG). IP_3 binds to the IP_3 receptors, which are concentrated at the apical pole of secretory cells [22, 23], to trigger Ca^{2+} release at the apical region that initiates a propagated Ca^{2+} wave to the basal pole along the lateral membrane [24, 25]. Several receptors, like the cholecystokinin (CCK) receptor in pancreatic acinar cells, also generate NAADP to cause additional Ca^{2+} release from endosome, lysosomes and likely secretory granules at the apical pole [26–28]. Ca^{2+} release from the ER causes clustering of the ER Ca^{2+} sensor STIM1 [29, 30], which binds to $PI(4,5)P_2$ at the ER/PM junctions [29, 31] where it activates the Orai1 [11, 32] and TRPC channels [33] leading to Ca^{2+} influx.

The increase in cytoplasmic Ca^{2+} rapidly activates Ca^{2+} clearance mechanisms to prevent Ca^{2+} toxicity. The Ca^{2+} -mediated inhibition of the IP_3 receptors [34] and of the Orai [11, 35] and TRPC channels [36] act to restrict Ca^{2+} flow into the cytosol. In addition, cytoplasmic Ca^{2+} is taken up by mitochondria to activate energy production and regulate other mitochondrial functions [37–39]. However, the bulk of the Ca^{2+} is incorporated back into the ER by the SERCA pumps and part is extruded out of the cells by the PMCA pumps, including the Ca^{2+} that leaves the mitochondria during the Ca^{2+} spikes. The process of Ca^{2+} release and clearance generate a Ca^{2+} transient that is repeated during the period of cell stimulation, resulting in Ca^{2+} oscillations. The amplitude, frequency and duration of the Ca^{2+} oscillations determine the information of the Ca^{2+} signal, and combinations of which can activate specific Ca^{2+} -dependent cellular processes.

A nodal point in sustaining the receptor-evoked Ca^{2+} signal is Ca^{2+} influx across the PM through the Orai channels. Orai1-mediated Ca^{2+} influx provides the source of Ca^{2+} to sustain Ca^{2+} loads into the various stores and sustain the Ca^{2+} signal during receptor stimulation. Moreover, aberrant Ca^{2+} signal and Ca^{2+} toxicity is almost invariably caused by excessive Ca^{2+} influx by TRPC [40–43] and Orai1 channels [44–46], but mostly the latter. Therefore, in this short review we discuss Ca^{2+} influx by Orai1 in secretory cells, and the role of Orai1-mediated Ca^{2+} influx in several key cellular functions and toxicity in secretory cells.

Orai1 and Ca^{2+} signaling at the ER/PM junctions

Secretory cell receptor-evoked Ca^{2+} signal initiates at the apical pole and propagate to the basal pole. Propagation to the basal pole depends on polarized expression of Ca^{2+} signaling proteins and organelles that store Ca^{2+} , such as secretory granules, lysosomes [47, 48] and mitochondria [37, 38]. The Ca^{2+} signaling proteins at the apical pole include GPCRs [49, 50], IP_3Rs [22], SERCA [51] and PMCA4 [52] pumps, and the sensitivity of the apical pole to receptor stimulation [53] has been demonstrated. The polarized localization requires targeting and arrangement of the Ca^{2+} signaling proteins into complexes at discreet cellular domain, the ER/PM junctions. How the junctions are formed and organized in exocrine cells or any cell *in vivo* is not known at this time. However, recent findings in model mammalian cells [12, 54, 55], that followed previous findings in yeast [56, 57], identified several key components required for assembly of the ER/PM junctions.

Briefly, the studies of lipid transport between the ER and plasma membrane (PM) in yeast revealed the role of the tethers (the three tricalbins [57], Ist2 [58], non-vesicular lipid transporters [59] and VAPA/B [60]) as essential components of the ER/PM junctions [56]. The mammalian homologues of the tricalbins are the three Extended synaptotagmins (E-Syt1–3) [61] and the lipid transport proteins expressed at the ER/PM junctions are the Nir proteins [62, 63], ORP5 and ORP8 [64] and the Aster proteins [65]. It is likely that other lipid transfer proteins bind to VAPA/B are also targeted to the ER/PM junctions which have yet to be defined. The tether proteins (E-Syt1–3) are required to maintain the ER/PM junctions [56] and control their size and likely location [55]. The role of the lipid transfer proteins in assembly of the ER/PM junctions has not been fully examined, but they may not be essential components of the junction since they appear to be transferred to the junction in response to changes in cellular lipids [59].

What is known about the localization and function of the lipid transfer proteins is illustrated in Figure 1. The Nir proteins maintain and replenish the level of PM $\text{PI}(4,5)\text{P}_2$ [62, 63]. They localize at the ER/PM junctions by interacting with the ER localized VAPA through two phenylalanines (FF) in an Acidic Tract (FFAT) motif [60]. At the junctions, the Nir proteins interact with lipids through their Phosphatidylinositol-Transfer Domain (PITD) that binds PI and their Lipin/Ndel/Smp2 (LNS2) domain that binds phosphatidic acid (PA) to mediate an exchange between ER PI and PM PA, thus supplying the ER with PA while maintaining PM PI for the synthesis of $\text{PI}(4,5)\text{P}_2$ [62]. The ORP5 and ORP8 proteins have an ER spanning single C-terminal transmembrane domain and N-terminal PH and polybasic domain that interacts with PM phosphatidylinositol 4-phosphate (PI4P) [64, 66] and

PI(4,5)P₂ [67]. Their ORD lipid transfer domain binds PM PI4P and PI(4,5)P₂ or ER phosphatidylserine (PS) to mediate PI4P/PS exchange, which is supported by the PI4P gradient that is maintained by its conversion to PI(4,5)P₂ in the PM and to PI in the ER. The Aster proteins have an ER localized N-terminal single transmembrane domain, a lipid transfer ASTER domain that binds cholesterol and a GRAM domain that interacts with PM PS and PA. The Aster proteins are recruited to the ER/PM junctions in response to increased PM cholesterol and transfer it to the ER. In addition, the Aster proteins interact with the E-Syts and ORP5/8 proteins at the junction [65].

The generation of PI(4,5)P₂-rich domain (and perhaps PS-rich domain) and of cholesterol at the ER/PM junctions by lipid transfer proteins has important implications for Ca²⁺ influx and Ca²⁺ signaling [54]. Cholesterol affects many cellular processes by altering membrane fluidity and has a direct effect on many proteins, including ion channels, where Orai1 and TRP channels are no exception [68, 69]. Depletion of PM cholesterol impacts TRPC channel localization and thus activation by various stimuli [70]. Opposite effects have been reported for Orai1. Depletion of PM cholesterol resulted in internalization of Orai1 and inhibition of channel activity [71]. In addition, the SOAR domain of STIM1 [72] was reported to possess a cholesterol binding site and cholesterol binding to STIM1 restricts binding of the STIM1 SOAR domain to Orai1 [73]. On the other hand, a cholesterol binding domain was detected in Orai1 and depletion of PM cholesterol and mutation of the cholesterol sensing site in Orai1 increased Orai1 current and Ca²⁺ influx [74]. Although some of the conflicting results may be explained by differential effects of cholesterol on the plasma membrane physical parameters and direct effects on Orai1 and STIM1, this topic needs significant further mechanistic clarification.

The importance of the plasma membrane PI(4,5)P₂ is demonstrated by the additional roles it plays in receptor-evoked Ca²⁺ signaling besides providing IP₃ to initiate the Ca²⁺ signal. PI(4,5)P₂ is a substrate for lipid transfer proteins like ORP5 and ORP8 [67] which control PM lipid composition. PI(4,5)P₂ interacts with and directly regulates the function of many channels, including several K⁺, Cl⁻ and voltage-gated Ca²⁺ channels, thereby affecting cytosolic ionic composition and the plasma membrane potential [54, 75]. PI(4,5)P₂ is a prominent PM site for interaction of proteins with Pleckstrin Homology (PH) domains, polybasic domains and other PI(4,5)P₂ interacting sites. One of the C2 domains of the E-Syts tethers [61], the PH domain of ORP5 and ORP8 [64, 66] and the polybasic domains of Ist2 [58] and the ORP5 and ORP8 [66] anchor the proteins to the PM by binding to PI(4,5)P₂.

The polybasic domains are essential for targeting of STIM1 [29, 76] and STIM2 [77] to the ER/PM junctions and for stabilization of the puncta. Moreover, presence of the STIM1-Orai1 complex at PI(4,5)P₂-poor and PI(4,5)P₂-rich domain determines channel function [31]. Orai1 undergoes two forms of Ca²⁺-dependent inactivation, a fast inactivation with a time constant of 10 msec and slow inactivation that is completed in 1–2 min [11, 35]. The two forms of inactivation are mediated by a STIM1 sequence [78–80] within the STIM1 CTID domain [20] that guides the SARAF protein [81] to the STIM1 SOAR domain [20]. Interaction of SARAF with STIM1 is dependent on the STIM1-Orai1 complex being in a PI(4,5)P₂-rich domain [31]. Depletion of ER Ca²⁺ results in the clustered STIM1 and its

interaction with Orai1. In a PI(4,5)P₂-poor domain where SARAF does not interact with STIM1, the Ca²⁺ influx through the STIM1-Orai1 complex is maximally active. Subsequently, the STIM1-Orai1 complex translocates to a PI(4,5)P₂-rich domain (or PI(4,5)P₂ translocates to the STIM1-Orai1 complex) where SARAF interacts with STIM1 to initiate Ca²⁺-dependent inactivation that limits Ca²⁺ influx and protect against Ca²⁺-dependent toxicity [31]. The role of the PI(4,5)P₂ domains in Ca²⁺ influx is illustrated in Figure 2.

Another protein that regulates STIM1 function is STIMATE [82] that is encoded by TMEM110 [83]. STIMATE is an ER protein that markedly facilitates STIM1 clustering and puncta formation at the ER/PM junctions, by stabilizing the open conformation of STIM1. Whether STIMATE interacts with STIM1 at PI(4,5)P₂-poor or PI(4,5)P₂-rich domain is not known at present. However, STIMATE has a polybasic domain [82], which may interact with acidic lipids, including PI(4,5)P₂ and thus aid in translocation of the STIM1-Orai1 complex to the PI(4,5)P₂-rich domain. Facilitation of STIM1 clustering by STIMATE to increase Ca²⁺ influx adds another level of regulation of the influx and thus excessive Ca²⁺ influx may also result from excessive activation of STIM1 by STIMATE. Additional information is needed on the mechanism by which STIMATE activates STIM1 to better understand a potential role of STIMATE in Ca²⁺ influx pathology.

Orai1 and Ca²⁺ influx in secretory gland functions

Ca²⁺ influx plays a critical role in secretory cell functions. The cardinal functions of secretory gland cells are exocytotic secretion of macromolecules, such as digestive enzymes and mucins, and fluid secretion with defined ionic composition [1]. Both functions are primarily activated by changes in cytoplasmic Ca²⁺ ([Ca²⁺]_i), but [Ca²⁺]_i synergizes with cAMP to mediate the physiological response ([2, 84] and see below). As shown in Figure 3, both TRPC channels and Orai1 contribute to receptor- and store-dependent Ca²⁺ influx and physiological Ca²⁺ oscillations [33, 85–88]. However, Orai1-mediated influx is essential for all forms of Ca²⁺ influx, likely because it regulates the TRPC channels dependent function and Ca²⁺ influx [89, 90].

Exocytosis by secretory gland cells occurs in two phases, a fast phase that is completed within one minute of receptor stimulation and a slow phase that continues for the length of the stimulation period and lasts for many tens of minutes [1, 91]. The first phase depends on Ca²⁺ release from internal stores next to the apical pole, consisting of the ready-release granules, while the second phase strictly depends on Ca²⁺ influx [92]. Ca²⁺ influx by Orai1 is essential for the exocytotic secretion function. Exocytosis by lacrimal glands of Orai1 knockout mice is markedly reduced [93] and targeted deletion of Orai1 in pancreatic acini reduced exocytosis of secretory granules, particularly the sustained phase of exocytosis [87]. Several Ca²⁺-regulated and Ca²⁺ binding proteins translate the Ca²⁺ signal to an exocytotic response [94]. The best understood Ca²⁺ binding proteins are the synaptotagmins (Syts), which are found in neuronal and nonneuronal tissues [95]. They sense Ca²⁺ by C2A and C2B domains located in the cytoplasmic C-terminus [96] and several Syts have been identified in various acinar cells [52]. Other Ca²⁺ binding proteins are the exocytosis tethers Munc13/CAPS, which have C2 and PH domains that bind Ca²⁺ and PI(4,5)P₂ [97]. In

addition there are the Synaptotagmin-like proteins (SLPs) which contain two C2 domains [98] and interact with Syntaxin 2 to control exocytosis in exocrine cells [99]. The specific role of Ca^{2+} influx by Orai1 in relation to the function of these proteins remains to be established.

Fluid and electrolyte secretion is a fundamental function of all epithelia and in secretory glands it occurs in two stages. The serous cells secrete plasma-like fluid that flows to a ductal or ductallike system that determines the final volume and ionic composition of the secreted fluid [1]. Fluid and electrolyte secretion are regulated by synergy between the Ca^{2+} and cAMP/PKA signaling pathways [2]. Many of the ion transporters mediating epithelial fluid and electrolyte secretion are activated by the scaffolding protein IRBIT (IP_3 receptors binding protein released with IP_3) [100, 101]. In the resting state IRBIT is associated with the IP_3 receptors. Upon cell stimulation with Ca^{2+} and cAMP mobilizing agonists, the IP_3 Rs are phosphorylated by PKA, which increases their affinity for IP_3 and reduces their affinity for IRBIT [84]. A small physiological increase in IP_3 triggers Ca^{2+} oscillations and at the same time releases IRBIT from the IP_3 Rs. IRBIT will then interact with and activate ion transporters to promote fluid and electrolyte secretion [84].

In addition to activation by IRBIT, ion channels mediating fluid and electrolyte secretion are directly activated by $[\text{Ca}^{2+}]_i$. The key transporters involved in acinar cell fluid and electrolyte secretion are the volume-sensitive basolateral membrane $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporters NKCC1 and the Ca^{2+} -activated K^+ channel Kcnma1 and the luminal membrane Ca^{2+} -activated Cl^- channel ANO1 and the Ca^{2+} -activated K^+ channel Kcnn4 [1, 102, 103] (illustrated in Fig. 4). The receptor stimulated increase in $[\text{Ca}^{2+}]_i$ activates the Ca^{2+} -activated K^+ and Cl^- channels to evoke K^+ efflux across the basolateral membrane and Cl^- efflux across the luminal membrane and the obligatory water flow through the luminal aquaporin water channels [1, 102]. This results in cell shrinkage that participates in termination of the Ca^{2+} spiking due to inhibition of Ca^{2+} influx channels [104] and perhaps Ca^{2+} release [105]. Cell shrinkage, in turn, activates NKCC1 to replenish the released KCl and restore cell volume [106]. Electroneutrality is maintained by Na^+ flow through the tight junctions (Fig. 4).

The fundamental importance of Ca^{2+} influx by Orai1 in epithelial fluid and electrolyte secretion is evident from the anhidrosis observed in patients with non-functional mutations in Orai1 and STIM1 [107, 108]. In addition, deletion of Orai1 in mice impaired lacrimal glands tear secretion [93] and lactation [109] and targeted deletion of Orai1 and STIM1 and STIM2 in sweat glands resulted in anhidrosis in mice [107]. The critical role of Orai1 mediated Ca^{2+} influx in many fluid and electrolyte secretory epithelia is due to the direct and specific activation of ANO1 and likely K^+ channels by the Orai1 mediated Ca^{2+} influx. The importance of Ca^{2+} influx for sustained activation of the Ca^{2+} -activated Cl^- current in epithelia was established long before the identification of Orai1 and ANO1 [110]. However, the relationship between them was well documented in sweat glands, in which activation of Orai1 by receptor-stimulated or passive store depletion resulted in sustained activation of ANO1. By contrast, in the absence of Orai1, activation of ANO1 was transient and lasted only until the Ca^{2+} stored in the ER was exhausted [107]. It is likely that Orai1 and ANO1 are present at the same ER/PM junctions at the luminal membrane, because activation of

ANO1 is mediated by Ca^{2+} in the same microdomain [111]. In addition, SOC-mediated Ca^{2+} influx activates the native and expressed KCa3.1 in HEK cells [112]. Moreover, in mast cells the non-functional Orai1 mutants that inhibited Ca^{2+} influx also inhibited activation of KCa3.1 [113].

It is important to note that in cells where TRPC channels participate or influence store-dependent Ca^{2+} influx, Ca^{2+} influx through TRPC channels is essential for fluid secretion and activation of the channels. For example, deletion of TRPC1 in mice reduced store-dependent Ca^{2+} influx and salivation in response to muscarinic stimulation, which was attributed to inhibition of the Ca^{2+} -activated K^+ channel in salivary glands acinar cells [88]. A recent study examined the role of Ca^{2+} influx in the activation of endogenous ANO1 in salivary gland cells and reported that Ca^{2+} influx by TRPC1 increased plasma membrane expression of ANO1 and was required for activation of ANO1 [114]. In smooth muscle cells, ANO1 colocalized and immunoprecipitated with TRPC6 and Ca^{2+} influx by TRPC6 specifically activated ANO1 [115]. A potential explanation for activation of ANO1 by Ca^{2+} influx from both Orai1 and TRPC channels can be due to cell specific coupling of ANO1 to the two types of Ca^{2+} influx channels. Another alternative is an interdependence of Ca^{2+} influx by the two channels. TRPC1 was suggested to regulate the level of plasma membrane Orai1 [116], and the channel function of Orai1 and TRPC channels was required for physiological-receptor-mediated Ca^{2+} influx [89]. Further studies on the relationships between Orai1, STIM1, TRPC channels and ANO1 and K^+ channel, in particular within specialized microdomain of the ER/PM junctions, are required to better understand these important regulatory functions that are critical for epithelial fluid and electrolyte secretion.

Ca^{2+} influx in secretory glands pathology

It is well established that both, reduced and excessive Ca^{2+} influx are associated with cell dysfunction and pathology in various cell types, including secretory glands [12, 45, 108, 117, 118]. As mentioned above, non-functional mutations in Orai1 and STIM1 in humans and their deletion in mice resulted in anhidrosis [108]. In addition, targeted deletion of Orai1 in pancreatic acinar cells of adult mice revealed a critical role in the pancreas with the control of the gut microbiome [87]. Deletion of Orai1 in pancreatic acinar cells inhibited receptor-stimulated Ca^{2+} influx and secretion of antimicrobials by acinar cells to the jejunum. This resulted in dysbiosis, damaged intestinal barrier, sepsis and death, despite the full activation of the intestinal innate response [87]. In this respect, germline deletion of Orai1 resulted in lacrimal glands inflammation [93]. These findings argue for an important function of acinar cells in epithelial innate response that depends on Ca^{2+} influx mediated by Orai1. These findings may have important clinical implications because acute and chronic pancreatitis influence the human gut microbiome [119, 120].

Excessive activity of Orai1 and TRPC channels is the initiator of Ca^{2+} overload and cell toxicity [40, 46, 85, 86, 121, 122]. Excessive Ca^{2+} influx results in accumulation of Ca^{2+} in the mitochondria of salivary gland cells [123], enterocytes [124], hepatocytes [125] and pancreatic acinar [46] and stellate cells [126]. The Ca^{2+} overload causes excessive mitochondrial ROS generation, cell damage and cell death [127–129]. Chronic excessive ROS has several deleterious effects. ROS activates TRPC [130, 131] and TRPM2 channels

[132], modulates STIM1-Orai1 interaction and function [133] and is a potent inhibitor of PMCA pumps in epithelial cells [134, 135]. This results in a feedforward effect that further increases Ca^{2+} influx and cell toxicity. This scenario has been worked out in great details in the pancreas in relation to acute pancreatitis [136]. The two types of Ca^{2+} influx channels shown to mediate the excessive pathological Ca^{2+} were the TRPC [40, 85] and Orai1 channels [86, 87, 121].

There is no effective treatment to either chronic or acute pancreatitis. The importance of Ca^{2+} influx channels in the pathology of pancreatitis suggested Ca^{2+} influx channels as potential targets for treatment. Several studies attempted to evaluate this option. The first was genetic [40] and pharmacological inhibition of TRPC3 [85], which reduced receptor stimulated and store-operated Ca^{2+} influx by about 50%. Inhibition of TRPC channel Ca^{2+} influx was quite effective in reducing the pathology of acute pancreatitis, but only when the channels were inhibited prior to induction of pancreatitis, while inhibition of the channels after induction of acute pancreatitis was not effective in reducing disease symptoms [85]. Greater success was observed by inhibition of Orai1 [121], in which inhibition of Orai1 prior to induction of acute pancreatitis prevented the disease, and ameliorated the symptoms of the disease when applied after induction of the pancreatitis in several mouse models [121].

As promising is the use of Orai1 inhibitors to treat Ca^{2+} overload in epithelia and inflammatory diseases in other tissues [45, 118], it is important to note the plethora of crucial roles of Orai1 in cellular and tissue physiology and their intricate regulation by Ca^{2+} . A significant change in the Ca^{2+} signal due to inhibition of Orai1 can be as toxic as excessive Ca^{2+} influx. Good examples are the many aberrant functions of the immune system caused by deletion of Orai1 and the STIM proteins [118] and the dysbiosis caused by deletion of pancreatic acinar cells Orai1 [87]. Perhaps a better approach will be to use a combination of partial inhibitors of TRPC and Orai1 channels that may prevent undesired effects due to prominent inhibition of the channels. Another option might be modulation of proteins that regulate the Ca^{2+} influx channels. One such protein is SARAF [31, 81], an inhibitor of Ca^{2+} influx channel and the other is STIMATE [82], which controls STIM1 clustering. Increasing SARAF activity and reducing STIMATE activity can be an alternative, perhaps safer approach to reducing pathological Ca^{2+} influx and its associated toxicity.

Acknowledgments

We thank Dr. Beth McNally for careful reading of the manuscript and suggestions. The authors are supported by intramural NIH grant DE000735–7.

References

- [1]. Lee MG, Ohana E, Park HW, Yang D, Muallem S, Molecular mechanism of pancreatic and salivary gland fluid and HCO_3 secretion, *Physiol Rev*, 92 (2012) 39–74. [PubMed: 22298651]
- [2]. Ahuja M, Jha A, Maleth J, Park S, Muallem S, cAMP and Ca^{2+} signaling in secretory epithelia: crosstalk and synergism, *Cell Calcium*, 55 (2014) 385–393. [PubMed: 24613710]
- [3]. Cali T, Brini M, Carafoli E, Regulation of Cell Calcium and Role of Plasma Membrane Calcium ATPases, *Int Rev Cell Mol Biol*, 332 (2017) 259–296. [PubMed: 28526135]

- [4]. Chemaly ER, Troncone L, Lebeche D, SERCA control of cell death and survival, *Cell Calcium*, 69 (2018) 46–61. [PubMed: 28747251]
- [5]. Dang D, Rao R, Calcium-ATPases: Gene disorders and dysregulation in cancer, *Biochim Biophys Acta*, 1863 (2016) 1344–1350. [PubMed: 26608610]
- [6]. Nemani N, Shanmughapriya S, Madesh M, Molecular regulation of MCU: Implications in physiology and disease, *Cell Calcium*, 74 (2018) 86–93. [PubMed: 29980025]
- [7]. Baughman JM, Perocchi F, Girgis HS, Plovanich M, Belcher-Timme CA, Sancak Y, Bao XR, Strittmatter L, Goldberger O, Bogorad RL, Kotliansky V, Mootha VK, Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter, *Nature*, 476 (2011) 341–345. [PubMed: 21685886]
- [8]. De Stefani D, Raffaello A, Teardo E, Szabo I, Rizzuto R, A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter, *Nature*, 476 (2011) 336–340. [PubMed: 21685888]
- [9]. Mikoshiba K, Role of IP₃ receptor signaling in cell functions and diseases, *Adv Biol Regul*, 57 (2015) 217–227. [PubMed: 25497594]
- [10]. Seo MD, Enomoto M, Ishiyama N, Stathopulos PB, Ikura M, Structural insights into endoplasmic reticulum stored calcium regulation by inositol 1,4,5-trisphosphate and ryanodine receptors, *Biochim Biophys Acta*, 1853 (2015) 1980–1991. [PubMed: 25461839]
- [11]. Prakriya M, Lewis RS, Store-Operated Calcium Channels, *Physiol Rev*, 95 (2015) 1383–1436.
- [12]. Chung WY, Jha A, Ahuja M, Muallem S, Ca(2+) influx at the ER/PM junctions, *Cell Calcium*, 63 (2017) 29–32. [PubMed: 28262262]
- [13]. Nishida M, Kuwahara K, Kozai D, Sakaguchi R, Mori Y, TRP Channels: Their Function and Potentiality as Drug Targets, in: Nakao K, Minato N, Uemoto S (Eds.) *Innovative Medicine: Basic Research and Development*, Tokyo, 2015, pp. 195–218.
- [14]. Palty R, Sekler I, The mitochondrial Na(+)/Ca(2+) exchanger, *Cell Calcium*, 52 (2012) 9–15. [PubMed: 22430014]
- [15]. Di Paola S, Scotto-Rosato A, Medina DL, TRPML1: The Ca(2+) retractor of the lysosome, *Cell Calcium*, 69 (2018) 112–121. [PubMed: 28689729]
- [16]. Grimm C, Barthmes M, Wahl-Schott C, Trpml3, *Handb Exp Pharmacol*, 222 (2014) 659–674. [PubMed: 24756725]
- [17]. Grimm C, Butz E, Chen CC, Wahl-Schott C, Biel M, From mucopolidosis type IV to Ebola: TRPML and two-pore channels at the crossroads of endo-lysosomal trafficking and disease, *Cell Calcium*, 67 (2017) 148–155. [PubMed: 28457591]
- [18]. Grimm C, Chen CC, Wahl-Schott C, Biel M, Two-Pore Channels: Catalyzers of Endolysosomal Transport and Function, *Front Pharmacol*, 8 (2017) 45. [PubMed: 28223936]
- [19]. Xu H, Ren D, Lysosomal physiology, *Annu Rev Physiol*, 77 (2015) 57–80. [PubMed: 25668017]
- [20]. Jha A, Ahuja M, Patel S, Brailoiu E, Muallem S, Convergent regulation of the lysosomal two-pore channel-2 by Mg(2+), NAADP, PI(3,5)P(2) and multiple protein kinases, *EMBO J*, 33 (2014) 501–511. [PubMed: 24502975]
- [21]. Petersen OH, Ca(2+) signalling in the endoplasmic reticulum/secretory granule microdomain, *Cell Calcium*, 58 (2015) 397–404. [PubMed: 25662795]
- [22]. Lee MG, Xu X, Zeng W, Diaz J, Wojcikiewicz RJ, Kuo TH, Wuytack F, Racymaekers L, Muallem S, Polarized expression of Ca²⁺ channels in pancreatic and salivary gland cells. Correlation with initiation and propagation of [Ca²⁺]_i waves, *J Biol Chem*, 272 (1997) 15765–15770. [PubMed: 9188472]
- [23]. Yule DI, Ernst SA, Ohnishi H, Wojcikiewicz RJ, Evidence that zymogen granules are not a physiologically relevant calcium pool. Defining the distribution of inositol 1,4,5-trisphosphate receptors in pancreatic acinar cells, *J Biol Chem*, 272 (1997) 9093–9098. [PubMed: 9083036]
- [24]. Hong JH, Li Q, Kim MS, Shin DM, Feske S, Birnbaumer L, Cheng KT, Ambudkar IS, Muallem S, Polarized but differential localization and recruitment of STIM1, Orai1 and TRPC channels in secretory cells, *Traffic*, 12 (2011) 232–245. [PubMed: 21054717]
- [25]. Sneyd J, Means S, Zhu D, Rugis J, Won JH, Yule DI, Modeling calcium waves in an anatomically accurate three-dimensional parotid acinar cell, *J Theor Biol*, 419 (2017) 383–393. [PubMed: 27155044]

- [26]. Cancela JM, Van Coppenolle F, Galione A, Tepikin AV, Petersen OH, Transformation of local Ca^{2+} spikes to global Ca^{2+} transients: the combinatorial roles of multiple Ca^{2+} releasing messengers, *EMBO J*, 21 (2002) 909–919. [PubMed: 11867519]
- [27]. Gerasimenko JV, Sherwood M, Tepikin AV, Petersen OH, Gerasimenko OV, NAADP, cADPR and IP_3 all release Ca^{2+} from the endoplasmic reticulum and an acidic store in the secretory granule area, *J Cell Sci*, 119 (2006) 226–238. [PubMed: 16410548]
- [28]. Gerasimenko JV, Charlesworth RM, Sherwood MW, Ferdek PE, Mikoshiba K, Parrington J, Petersen OH, Gerasimenko OV, Both RyRs and TPCs are required for NAADP-induced intracellular Ca^{2+} release, *Cell Calcium*, 58 (2015) 237–245. [PubMed: 26100948]
- [29]. Liou J, Kim ML, Heo WD, Jones JT, Myers JW, Ferrell JE, Jr., Meyer T, STIM is a Ca^{2+} sensor essential for Ca^{2+} -store-depletion-triggered Ca^{2+} influx, *Curr Biol*, 15 (2005) 12351241.
- [30]. Roos J, DiGregorio PJ, Yeromin AV, Ohlsen K, Lioudyno M, Zhang S, Safrina O, Kozak JA, Wagner SL, Cahalan MD, Velicelebi G, Stauderman KA, STIM1, an essential and conserved component of store-operated Ca^{2+} channel function, *J Cell Biol*, 169 (2005) 435–445. [PubMed: 15866891]
- [31]. Maleth J, Choi S, Muallem S, Ahuja M, Translocation between $\text{PI}(4,5)\text{P}_2$ -poor and $\text{PI}(4,5)\text{P}_2$ -rich microdomains during store depletion determines STIM1 conformation and Orai1 gating, *Nat Commun*, 5 (2014) 5843. [PubMed: 25517631]
- [32]. Zhang SL, Yeromin AV, Zhang XH, Yu Y, Safrina O, Penna A, Roos J, Stauderman KA, Cahalan MD, Genome-wide RNAi screen of Ca^{2+} influx identifies genes that regulate Ca^{2+} release-activated Ca^{2+} channel activity, *Proc Natl Acad Sci U S A*, 103 (2006) 9357–9362. [PubMed: 16751269]
- [33]. Bodnar D, Chung WY, Yang D, Hong JH, Jha A, Muallem S, STIM-TRP Pathways and Microdomain Organization: Ca^{2+} Influx Channels: The Orai-STIM1-TRPC Complexes, *Adv Exp Med Biol*, 993 (2017) 139–157. [PubMed: 28900913]
- [34]. Mak DO, Foskett JK, Inositol 1,4,5-trisphosphate receptors in the endoplasmic reticulum: A single-channel point of view, *Cell Calcium*, 58 (2015) 67–78. [PubMed: 25555684]
- [35]. Parekh AB, Regulation of CRAC channels by Ca^{2+} -dependent inactivation, *Cell Calcium*, 63 (2017) 20–23. [PubMed: 28043696]
- [36]. Beech DJ, Characteristics of transient receptor potential canonical calcium-permeable channels and their relevance to vascular physiology and disease, *Circ J*, 77 (2013) 570–579. [PubMed: 23412755]
- [37]. Gunter TE, Yule DI, Gunter KK, Eliseev RA, Salter JD, Calcium and mitochondria, *FEBS Lett*, 567 (2004) 96–102. [PubMed: 15165900]
- [38]. Petersen OH, Specific mitochondrial functions in separate sub-cellular domains of pancreatic acinar cells, *Pflugers Arch*, 464 (2012) 77–87. [PubMed: 22491894]
- [39]. Giorgi C, Marchi S, Pinton P, The machineries, regulation and cellular functions of mitochondrial calcium, *Nat Rev Mol Cell Biol*, 19 (2018) 713–730. [PubMed: 30143745]
- [40]. Kim MS, Hong JH, Li Q, Shin DM, Abramowitz J, Birnbaumer L, Muallem S, Deletion of TRPC3 in mice reduces store-operated Ca^{2+} influx and the severity of acute pancreatitis, *Gastroenterology*, 137 (2009) 1509–1517. [PubMed: 19622358]
- [41]. Sachdeva R, Schlotterer A, Schumacher D, Matka C, Mathar I, Dietrich N, Medert R, Kriebs U, Lin J, Nawroth P, Birnbaumer L, Fleming T, Hammes HP, Freichel M, TRPC proteins contribute to development of diabetic retinopathy and regulate glyoxalase 1 activity and methylglyoxal accumulation, *Mol Metab*, 9 (2018) 156–167. [PubMed: 29373286]
- [42]. He X, Li S, Liu B, Susperreguy S, Formoso K, Yao J, Kang J, Shi A, Birnbaumer L, Liao Y, Major contribution of the 3/6/7 class of TRPC channels to myocardial ischemia/reperfusion and cellular hypoxia/reoxygenation injuries, *Proc Natl Acad Sci U S A*, 114 (2017) E4582–E4591. [PubMed: 28526717]
- [43]. Zhou J, Jia Y, TRPC Channels and Programmed Cell Death, *Adv Exp Med Biol*, 976 (2017) 47–60. [PubMed: 28508312]
- [44]. Maher P, van Leyen K, Dey PN, Honrath B, Dolga A, Methner A, The role of Ca^{2+} in cell death caused by oxidative glutamate toxicity and ferroptosis, *Cell Calcium*, 70 (2018) 47–55. [PubMed: 28545724]

- [45]. Vaeth M, Feske S, Ion channelopathies of the immune system, *Curr Opin Immunol*, 52 (2018) 39–50. [PubMed: 29635109]
- [46]. Criddle DN, Reactive oxygen species, Ca(2+) stores and acute pancreatitis; a step closer to therapy?, *Cell Calcium*, 60 (2016) 180–189. [PubMed: 27229361]
- [47]. Motta PM, Macchiarelli G, Nottola SA, Correr S, Histology of the exocrine pancreas, *Microsc Res Tech*, 37 (1997) 384–398. [PubMed: 9220418]
- [48]. Murakami M, Yoshimura K, Sugiya H, Seo Y, Loffredo F, Riva A, Relationship of fluid and mucin secretion to morphological changes in the perfused rat submandibular gland, *Eur J Morphol*, 40 (2002) 203–207. [PubMed: 14566596]
- [49]. Shin DM, Luo X, Wilkie TM, Miller LJ, Peck AB, Humphreys-Beher MG, Muallem S, Polarized expression of G protein-coupled receptors and an all-or-none discharge of Ca2+ pools at initiation sites of [Ca2+]i waves in polarized exocrine cells, *J Biol Chem*, 276 (2001) 4414644156.
- [50]. Fischer A, Kummer W, Couraud JY, Adler D, Branscheid D, Heym C, Immunohistochemical localization of receptors for vasoactive intestinal peptide and substance P in human trachea, *Lab Invest*, 67 (1992) 387–393. [PubMed: 1328764]
- [51]. Lee MG, Xu X, Zeng W, Diaz J, Kuo TH, Wuytack F, Racymaekers L, Muallem S, Polarized expression of Ca2+ pumps in pancreatic and salivary gland cells. Role in initiation and propagation of [Ca2+]i waves, *J Biol Chem*, 272 (1997) 15771–15776. [PubMed: 9188473]
- [52]. Zhao XS, Shin DM, Liu LH, Shull GE, Muallem S, Plasticity and adaptation of Ca2+ signaling and Ca2+-dependent exocytosis in SERCA2(+/-) mice, *EMBO J*, 20 (2001) 2680–2689. [PubMed: 11387203]
- [53]. Li Q, Luo X, Muallem S, Functional mapping of Ca2+ signaling complexes in plasma membrane microdomains of polarized cells, *J Biol Chem*, 279 (2004) 27837–27840. [PubMed: 15123684]
- [54]. Muallem S, Chung WY, Jha A, Ahuja M, Lipids at membrane contact sites: cell signaling and ion transport, *EMBO Rep*, 18 (2017) 1893–1904. [PubMed: 29030479]
- [55]. Saheki Y, De Camilli P, Endoplasmic Reticulum-Plasma Membrane Contact Sites, *Annu Rev Biochem*, 86 (2017) 659–684. [PubMed: 28301744]
- [56]. Henne WM, Liou J, Emr SD, Molecular mechanisms of inter-organelle ER-PM contact sites, *Curr Opin Cell Biol*, 35 (2015) 123–130. [PubMed: 26025028]
- [57]. Lahiri S, Toulmay A, Prinz WA, Membrane contact sites, gateways for lipid homeostasis, *Curr Opin Cell Biol*, 33 (2015) 82–87. [PubMed: 25569848]
- [58]. Fischer MA, Temmerman K, Ercan E, Nickel W, Seedorf M, Binding of plasma membrane lipids recruits the yeast integral membrane protein Ist2 to the cortical ER, *Traffic*, 10 (2009) 1084–1097. [PubMed: 19453974]
- [59]. Kentala H, Weber-Boyvat M, Olkkonen VM, OSBP-Related Protein Family: Mediators of Lipid Transport and Signaling at Membrane Contact Sites, *Int Rev Cell Mol Biol*, 321 (2016) 299–340. [PubMed: 26811291]
- [60]. Selitrennik M, Lev S, The role of phosphatidylinositol-transfer proteins at membrane contact sites, *Biochem Soc Trans*, 44 (2016) 419–424. [PubMed: 27068949]
- [61]. Giordano F, Saheki Y, Idevall-Hagren O, Colombo SF, Pirruccello M, Milosevic I, Gracheva EO, Bagriantsev SN, Borgese N, De Camilli P, PI(4,5)P(2)-dependent and Ca(2+)-regulated ER-PM interactions mediated by the extended synaptotagmins, *Cell*, 153 (2013) 1494–1509. [PubMed: 23791178]
- [62]. Kim YJ, Guzman-Hernandez ML, Wisniewski E, Balla T, PhosphatidylinositolPhosphatidic Acid Exchange by Nir2 at ER-PM Contact Sites Maintains Phosphoinositide Signaling Competence, *Dev Cell*, 33 (2015) 549–561. [PubMed: 26028218]
- [63]. Chang CL, Liou J, Phosphatidylinositol 4,5-Bisphosphate Homeostasis Regulated by Nir2 and Nir3 Proteins at Endoplasmic Reticulum-Plasma Membrane Junctions, *J Biol Chem*, 290 (2015) 14289–14301. [PubMed: 25887399]
- [64]. Chung J, Torta F, Masai K, Lucast L, Czapla H, Tanner LB, Narayanaswamy P, Wenk MR, Nakatsu F, De Camilli P, INTRACELLULAR TRANSPORT. PI4P/phosphatidylserine countertransport at ORP5- and ORP8-mediated ER-plasma membrane contacts, *Science*, 349 (2015) 428–432. [PubMed: 26206935]

- [65]. Sandhu J, Li S, Fairall L, Pfisterer SG, Gurnett JE, Xiao X, Weston TA, Vashi D, Ferrari A, Orozco JL, Hartman CL, Strugatsky D, Lee SD, He C, Hong C, Jiang H, Bentolila LA, Gatta AT, Levine TP, Ferng A, Lee R, Ford DA, Young SG, Ikonen E, Schwabe JWR, Tontonoz P, Aster Proteins Facilitate Nonvesicular Plasma Membrane to ER Cholesterol Transport in Mammalian Cells, *Cell*, 175 (2018) 514–529 e520. [PubMed: 30220461]
- [66]. Sohn M, Korzeniowski M, Zewe JP, Wills RC, Hammond GRV, Humpolickova J, Vrzal L, Chalupska D, Veverka V, Fairn GD, Boura E, Balla T, PI(4,5)P2 controls plasma membrane PI4P and PS levels via ORP5/8 recruitment to ER-PM contact sites, *J Cell Biol*, 217 (2018) 1797–1813. [PubMed: 29472386]
- [67]. Ghai R, Du X, Wang H, Dong J, Ferguson C, Brown AJ, Parton RG, Wu JW, Yang H, ORP5 and ORP8 bind phosphatidylinositol-4, 5-bisphosphate (PtdIns(4,5)P₂) and regulate its level at the plasma membrane, *Nat Commun*, 8 (2017) 757. [PubMed: 28970484]
- [68]. Rosenhouse-Dantsker A, Insights Into the Molecular Requirements for Cholesterol Binding to Ion Channels, *Curr Top Membr*, 80 (2017) 187–208. [PubMed: 28863816]
- [69]. Levitan I, Singh DK, Rosenhouse-Dantsker A, Cholesterol binding to ion channels, *Front Physiol*, 5 (2014) 65. [PubMed: 24616704]
- [70]. Pacheco J, Ramirez-Jarquín JO, Vaca L, Microdomains Associated to Lipid Rafts, *Adv Exp Med Biol*, 898 (2016) 353–378. [PubMed: 27161236]
- [71]. Bohorquez-Hernandez A, Gratton E, Pacheco J, Asanov A, Vaca L, Cholesterol modulates the cellular localization of Orai1 channels and its disposition among membrane domains, *Biochim Biophys Acta Mol Cell Biol Lipids*, 1862 (2017) 1481–1490. [PubMed: 28919480]
- [72]. Yuan JP, Zeng W, Dorwart MR, Choi YJ, Worley PF, Muallem S, SOAR and the polybasic STIM1 domains gate and regulate Orai channels, *Nat Cell Biol*, 11 (2009) 337–343. [PubMed: 19182790]
- [73]. Pacheco J, Dominguez L, Bohorquez-Hernandez A, Asanov A, Vaca L, A cholesterol-binding domain in STIM1 modulates STIM1-Orai1 physical and functional interactions, *Sci Rep*, 6 (2016) 29634.
- [74]. Derler I, Jardin I, Stathopoulos PB, Muik M, Fahrner M, Zayats V, Pandey SK, Poteser M, Lackner B, Absolonova M, Schindl R, Groschner K, Ettrich R, Ikura M, Romanin C, Cholesterol modulates Orai1 channel function, *Sci Signal*, 9 (2016) ra10.
- [75]. Hille B, Dickson EJ, Kruse M, Vivas O, Suh BC, Phosphoinositides regulate ion channels, *Biochim Biophys Acta*, 1851 (2015) 844–856. [PubMed: 25241941]
- [76]. Liou J, Fivaz M, Inoue T, Meyer T, Live-cell imaging reveals sequential oligomerization and local plasma membrane targeting of stromal interaction molecule 1 after Ca²⁺ store depletion, *Proc Natl Acad Sci U S A*, 104 (2007) 9301–9306. [PubMed: 17517596]
- [77]. Ong HL, de Souza LB, Zheng C, Cheng KT, Liu X, Goldsmith CM, Feske S, Ambudkar IS, STIM2 enhances receptor-stimulated Ca²⁺(+) signaling by promoting recruitment of STIM1 to the endoplasmic reticulum-plasma membrane junctions, *Sci Signal*, 8 (2015) ra3.
- [78]. Lee KP, Yuan JP, Zeng W, So I, Worley PF, Muallem S, Molecular determinants of fast Ca²⁺-dependent inactivation and gating of the Orai channels, *Proc Natl Acad Sci U S A*, 106 (2009) 14687–14692. [PubMed: 19706554]
- [79]. Derler I, Fahrner M, Muik M, Lackner B, Schindl R, Groschner K, Romanin C, A Ca²⁺(+)release-activated Ca²⁺(+) (CRAC) modulatory domain (CMD) within STIM1 mediates fast Ca²⁺(+)dependent inactivation of ORAI1 channels, *J Biol Chem*, 284 (2009) 24933–24938. [PubMed: 19622747]
- [80]. Mullins FM, Park CY, Dolmetsch RE, Lewis RS, STIM1 and calmodulin interact with Orai1 to induce Ca²⁺-dependent inactivation of CRAC channels, *Proc Natl Acad Sci U S A*, 106 (2009) 15495–15500. [PubMed: 19706428]
- [81]. Palty R, Raveh A, Kaminsky I, Meller R, Reuveny E, SARAF inactivates the store operated calcium entry machinery to prevent excess calcium refilling, *Cell*, 149 (2012) 425–438. [PubMed: 22464749]
- [82]. Jing J, He L, Sun A, Quintana A, Ding Y, Ma G, Tan P, Liang X, Zheng X, Chen L, Shi X, Zhang SL, Zhong L, Huang Y, Dong MQ, Walker CL, Hogan PG, Wang Y, Zhou Y, Proteomic mapping

of ER-PM junctions identifies STIMATE as a regulator of Ca(2)(+) influx, *Nat Cell Biol*, 17 (2015) 1339–1347. [PubMed: 26322679]

- [83]. Quintana A, Rajanikanth V, Farber-Katz S, Gudlur A, Zhang C, Jing J, Zhou Y, Rao A, Hogan PG, TMEM110 regulates the maintenance and remodeling of mammalian ER-plasma membrane junctions competent for STIM-ORAI signaling, *Proc Natl Acad Sci U S A*, 112 (2015) E7083–7092. [PubMed: 26644574]
- [84]. Park S, Shcheynikov N, Hong JH, Zheng C, Suh SH, Kawaai K, Ando H, Mizutani A, Abe T, Kiyonari H, Seki G, Yule D, Mikoshiba K, Muallem S, Irbit mediates synergy between ca(2+) and cAMP signaling pathways during epithelial transport in mice, *Gastroenterology*, 145 (2013) 232–241. [PubMed: 23542070]
- [85]. Kim MS, Lee KP, Yang D, Shin DM, Abramowitz J, Kiyonaka S, Birnbaumer L, Mori Y, Muallem S, Genetic and pharmacologic inhibition of the Ca2+ influx channel TRPC3 protects secretory epithelia from Ca2+-dependent toxicity, *Gastroenterology*, 140 (2011) 2107–2115, 2115 e2101–2104. [PubMed: 21354153]
- [86]. Gerasimenko JV, Gryshchenko O, Ferdek PE, Stapleton E, Hebert TO, Bychkova S, Peng S, Begg M, Gerasimenko OV, Petersen OH, Ca2+ release-activated Ca2+ channel blockade as a potential tool in antipancreatitis therapy, *Proc Natl Acad Sci U S A*, 110 (2013) 13186–13191. [PubMed: 23878235]
- [87]. Ahuja M, Schwartz DM, Tandon M, Son A, Zeng M, Swaim W, Eckhaus M, Hoffman V, Cui Y, Xiao B, Worley PF, Muallem S, Orai1-Mediated Antimicrobial Secretion from Pancreatic Acini Shapes the Gut Microbiome and Regulates Gut Innate Immunity, *Cell Metab*, 25 (2017) 635–646. [PubMed: 28273482]
- [88]. Liu X, Cheng KT, Bandyopadhyay BC, Pani B, Dietrich A, Paria BC, Swaim WD, Beech D, Yildirim E, Singh BB, Birnbaumer L, Ambudkar IS, Attenuation of store-operated Ca2+ current impairs salivary gland fluid secretion in TRPC1(–/–) mice, *Proc Natl Acad Sci U S A*, 104 (2007) 17542–17547. [PubMed: 17956991]
- [89]. Kim MS, Zeng W, Yuan JP, Shin DM, Worley PF, Muallem S, Native Store-operated Ca2+ Influx Requires the Channel Function of Orai1 and TRPC1, *J Biol Chem*, 284 (2009) 97339741.
- [90]. Cheng KT, Liu X, Ong HL, Ambudkar IS, Functional requirement for Orai1 in storeoperated TRPC1-STIM1 channels, *J Biol Chem*, 283 (2008) 12935–12940. [PubMed: 18326500]
- [91]. Williams JA, Receptor-mediated signal transduction pathways and the regulation of pancreatic acinar cell function, *Curr Opin Gastroenterol*, 24 (2008) 573–579. [PubMed: 19122497]
- [92]. Pandol SJ, Schoeffield MS, Sachs G, Muallem S, Role of free cytosolic calcium in secretagogue-stimulated amylase release from dispersed acini from guinea pig pancreas, *J Biol Chem*, 260 (1985) 10081–10086. [PubMed: 2410419]
- [93]. Xing J, Petranks JG, Davis FM, Desai PN, Putney JW, Bird GS, Role of Orai1 and store-operated calcium entry in mouse lacrimal gland signalling and function, *J Physiol*, 592 (2014) 927–939. [PubMed: 24297846]
- [94]. Messenger SW, Falkowski MA, Groblewski GE, Ca(2)(+)-regulated secretory granule exocytosis in pancreatic and parotid acinar cells, *Cell Calcium*, 55 (2014) 369–375. [PubMed: 24742357]
- [95]. Pang ZP, Sudhof TC, Cell biology of Ca2+-triggered exocytosis, *Curr Opin Cell Biol*, 22 (2010) 496–505. [PubMed: 20561775]
- [96]. Wang Z, Liu H, Gu Y, Chapman ER, Reconstituted synaptotagmin I mediates vesicle docking, priming, and fusion, *J Cell Biol*, 195 (2011) 1159–1170. [PubMed: 22184197]
- [97]. James DJ, Martin TF, CAPS and Munc13: CATCHRs that SNARE Vesicles, *Front Endocrinol (Lausanne)*, 4 (2013) 187. [PubMed: 24363652]
- [98]. Fukuda M, Rab27 effectors, pleiotropic regulators in secretory pathways, *Traffic*, 14 (2013) 949–963. [PubMed: 23678941]
- [99]. Fukuda M, Imai A, Nashida T, Shimomura H, Slp4-a/granuphilin-a interacts with syntaxin2/3 in a Munc18–2-dependent manner, *J Biol Chem*, 280 (2005) 39175–39184. [PubMed: 16186111]
- [100]. Ando H, Mizutani A, Matsu-ura T, Mikoshiba K, IRBIT, a novel inositol 1,4,5-trisphosphate (IP3) receptor-binding protein, is released from the IP3 receptor upon IP3 binding to the receptor, *J Biol Chem*, 278 (2003) 10602–10612. [PubMed: 12525476]

- [101]. Park S, Hong JH, Ohana E, Muallem S, The WNK/SPAK and IRBIT/PP1 pathways in epithelial fluid and electrolyte transport, *Physiology* (Bethesda), 27 (2012) 291–299. [PubMed: 23026752]
- [102]. Catalan MA, Pena-Munzenmayer G, Melvin JE, Ca(2+)-dependent K(+) channels in exocrine salivary glands, *Cell Calcium*, 55 (2014) 362–368. [PubMed: 24559652]
- [103]. Jang Y, Oh U, Anoctamin 1 in secretory epithelia, *Cell Calcium*, 55 (2014) 355–361. [PubMed: 24636668]
- [104]. Reetz G, Reiser G, [Ca²⁺]_i oscillations induced by bradykinin in rat glioma cells associated with Ca²⁺ store-dependent Ca²⁺ influx are controlled by cell volume and by membrane potential, *Cell Calcium*, 19 (1996) 143–156. [PubMed: 8689672]
- [105]. Park K, Lee S, Elliott AC, Kim JS, Lee JH, Swelling-induced Ca²⁺ release from intracellular calcium stores in rat submandibular gland acinar cells, *J Membr Biol*, 186 (2002) 165–176. [PubMed: 12148843]
- [106]. Shekarabi M, Zhang J, Khanna AR, Ellison DH, Delpire E, Kahle KT, WNK Kinase Signaling in Ion Homeostasis and Human Disease, *Cell Metab*, 25 (2017) 285–299. [PubMed: 28178566]
- [107]. Concepcion AR, Vaeth M, Wagner LE, 2nd, Eckstein M, Hecht L, Yang J, Crottes D, Seidl M, Shin HP, Weidinger C, Cameron S, Turvey SE, Issekutz T, Meyts I, Lacruz RS, Cuk M, Yule DI, Feske S, Store-operated Ca²⁺ entry regulates Ca²⁺-activated chloride channels and eccrine sweat gland function, *J Clin Invest*, 126 (2016) 4303–4318. [PubMed: 27721237]
- [108]. Concepcion AR, Feske S, Regulation of epithelial ion transport in exocrine glands by store-operated Ca(2+) entry, *Cell Calcium*, 63 (2017) 53–59. [PubMed: 28027799]
- [109]. Davis FM, Janoshazi A, Janardhan KS, Steinckwich N, D'Agostin DM, Petranksa JG, Desai PN, Roberts-Thomson SJ, Bird GS, Tucker DK, Fenton SE, Feske S, Monteith GR, Putney JW, Jr., Essential role of Orai1 store-operated calcium channels in lactation, *Proc Natl Acad Sci U S A*, 112 (2015) 5827–5832. [PubMed: 25902527]
- [110]. Hartzell C, Putzier I, Arreola J, Calcium-activated chloride channels, *Annu Rev Physiol*, 67 (2005) 719–758. [PubMed: 15709976]
- [111]. Jin X, Shah S, Liu Y, Zhang H, Lees M, Fu Z, Lippiat JD, Beech DJ, Sivaprasadarao A, Baldwin SA, Zhang H, Gamper N, Activation of the Cl-channel ANO1 by localized calcium signals in nociceptive sensory neurons requires coupling with the IP3 receptor, *Sci Signal*, 6 (2013) ra73.
- [112]. Sforna L, Megaro A, Pessia M, Franciolini F, Catacuzzeno L, Structure, Gating and Basic Functions of the Ca²⁺-activated K Channel of Intermediate Conductance, *Curr Neuropharmacol*, 16 (2018) 608–617. [PubMed: 28875832]
- [113]. Duffy SM, Ashmole I, Smallwood DT, Leyland ML, Bradding P, Orai/CRACM1 and KCa3.1 ion channels interact in the human lung mast cell plasma membrane, *Cell Commun Signal*, 13 (2015) 32. [PubMed: 26177720]
- [114]. Sun Y, Birnbaumer L, Singh BB, TRPC1 regulates calcium-activated chloride channels in salivary gland cells, *J Cell Physiol*, 230 (2015) 2848–2856. [PubMed: 25899321]
- [115]. Wang Q, Leo MD, Narayanan D, Kuruvilla KP, Jaggar JH, Local coupling of TRPC6 to ANO1/TMEM16A channels in smooth muscle cells amplifies vasoconstriction in cerebral arteries, *Am J Physiol Cell Physiol*, 310 (2016) C1001–1009. [PubMed: 27147559]
- [116]. Ambudkar IS, de Souza LB, Ong HL, TRPC1, Orai1, and STIM1 in SOCE: Friends in tight spaces, *Cell Calcium*, 63 (2017) 33–39. [PubMed: 28089266]
- [117]. Avila-Medina J, Mayoral-Gonzalez I, Dominguez-Rodriguez A, Gallardo-Castillo I, Ribas J, Ordonez A, Rosado JA, Smani T, The Complex Role of Store Operated Calcium Entry Pathways and Related Proteins in the Function of Cardiac, Skeletal and Vascular Smooth Muscle Cells, *Front Physiol*, 9 (2018) 257. [PubMed: 29618985]
- [118]. Lacruz RS, Feske S, Diseases caused by mutations in ORAI1 and STIM1, *Ann N Y Acad Sci*, 1356 (2015) 45–79. [PubMed: 26469693]
- [119]. Zhang XM, Zhang ZY, Zhang CH, Wu J, Wang YX, Zhang GX, Intestinal Microbial Community Differs between Acute Pancreatitis Patients and Healthy Volunteers, *Biomed Environ Sci*, 31 (2018) 81–86. [PubMed: 29409589]

- [120]. Hamada S, Masamune A, Nabeshima T, Shimosegawa T, Differences in Gut Microbiota Profiles between Autoimmune Pancreatitis and Chronic Pancreatitis, *Tohoku J Exp Med*, 244 (2018) 113–117. [PubMed: 29434076]
- [121]. Wen L, Voronina S, Javed MA, Awais M, Szatmary P, Latawiec D, Chvanov M, Collier D, Huang W, Barrett J, Begg M, Stauderman K, Roos J, Grigoryev S, Ramos S, Rogers E, Whitten J, Velicelebi G, Dunn M, Tepikin AV, Criddle DN, Sutton R, Inhibitors of ORAI1 Prevent Cytosolic Calcium-Associated Injury of Human Pancreatic Acinar Cells and Acute Pancreatitis in 3 Mouse Models, *Gastroenterology*, 149 (2015) 481–492 e487. [PubMed: 25917787]
- [122]. Correll RN, Goonasekera SA, van Berlo JH, Burr AR, Accornero F, Zhang H, Makarewich CA, York AJ, Sargent MA, Chen X, Houser SR, Molkentin JD, STIM1 elevation in the heart results in aberrant Ca(2+)(+) handling and cardiomyopathy, *J Mol Cell Cardiol*, 87 (2015) 38–47. [PubMed: 26241845]
- [123]. Liu X, Gong B, de Souza LB, Ong HL, Subedi KP, Cheng KT, Swaim W, Zheng C, Mori Y, Ambudkar IS, Radiation inhibits salivary gland function by promoting STIM1 cleavage by caspase-3 and loss of SOCE through a TRPM2-dependent pathway, *Sci Signal*, 10 (2017).
- [124]. Kolar SS, Barhoumi R, Lupton JR, Chapkin RS, Docosahexaenoic acid and butyrate synergistically induce colonocyte apoptosis by enhancing mitochondrial Ca2+ accumulation, *Cancer Res*, 67 (2007) 5561–5568. [PubMed: 17545640]
- [125]. Bernardi P, Rasola A, Calcium and cell death: the mitochondrial connection, *Subcell Biochem*, 45 (2007) 481–506. [PubMed: 18193649]
- [126]. Gerasimenko JV, Peng S, Tsugorka T, Gerasimenko OV, Ca(2+) signalling underlying pancreatitis, *Cell Calcium*, 70 (2018) 95–101. [PubMed: 28552244]
- [127]. Ambudkar IS, Muallem S, ROS and Ca(2+)-Partners in sickness and in health, *Cell Calcium*, 60 (2016) 51–54. [PubMed: 27474448]
- [128]. Ambudkar IS, Muallem S, ROS in Ca(2+) signaling and disease-part 2, *Cell Calcium*, 60 (2016) 153–154. [PubMed: 27553171]
- [129]. Criddle DN, Murphy J, Fistetto G, Barrow S, Tepikin AV, Neoptolemos JP, Sutton R, Petersen OH, Fatty acid ethyl esters cause pancreatic calcium toxicity via inositol trisphosphate receptors and loss of ATP synthesis, *Gastroenterology*, 130 (2006) 781–793. [PubMed: 16530519]
- [130]. Poteser M, Graziani A, Rosker C, Eder P, Derler I, Kahr H, Zhu MX, Romanin C, Groschner K, TRPC3 and TRPC4 associate to form a redox-sensitive cation channel. Evidence for expression of native TRPC3-TRPC4 heteromeric channels in endothelial cells, *J Biol Chem*, 281 (2006) 13588–13595. [PubMed: 16537542]
- [131]. Malczyk M, Veith C, Schermuly RT, Gudermann T, Dietrich A, Sommer N, Weissmann N, Pak O, NADPH oxidases-do they play a role in TRPC regulation under hypoxia?, *Pflugers Arch*, 468 (2016) 23–41. [PubMed: 26424109]
- [132]. Liu X, Ong HL, Ambudkar I, TRP Channel Involvement in Salivary Glands-Some Good, Some Bad, *Cells*, 7 (2018).
- [133]. Bhardwaj R, Hediger MA, Demareux N, Redox modulation of STIM-ORAI signaling, *Cell Calcium*, 60 (2016) 142–152. [PubMed: 27041216]
- [134]. Baggaley EM, Elliott AC, Bruce JJ, Oxidant-induced inhibition of the plasma membrane Ca2+-ATPase in pancreatic acinar cells: role of the mitochondria, *Am J Physiol Cell Physiol*, 295 (2008) C1247–1260. [PubMed: 18787078]
- [135]. Kim MJ, Choi KJ, Yoon MN, Oh SH, Kim DK, Kim SH, Park HS, Hydrogen peroxide inhibits Ca(2+) efflux through plasma membrane Ca(2+)-ATPase in mouse parotid acinar cells, *Korean J Physiol Pharmacol*, 22 (2018) 215–223. [PubMed: 29520174]
- [136]. Gryshchenko O, Gerasimenko JV, Peng S, Gerasimenko OV, Petersen OH, Calcium signalling in the acinar environment of the exocrine pancreas: physiology and pathophysiology, *J Physiol*, 596 (2018) 2663–2678. [PubMed: 29424931]

Highlights

- Ca^{2+} signaling in secretory epithelia
- Ca^{2+} influx and cell functions in secretory epithelia
- Ca^{2+} influx and disease in secretory epithelia

Fig. 2: Ca^{2+} signaling and Ca^{2+} influx at the ER/PM junctions

At the resting state the plasma membrane (PM) and endoplasmic reticulum (ER) components of the receptor-evoked Ca^{2+} signal are not in close contacts and several of them, in particular Orai1, STIM1, and IP_3 receptors and the Ca^{2+} pumps are outside the ER/PM junctions. Upon cell stimulation and ER Ca^{2+} release from the ER STIM1 clusters and interact with Orai1 in a $\text{PI}(4,5)\text{P}_2$ poor domain at which SARAF cannot interact with STIM1 and Ca^{2+} influx is maximally active. Shortly thereafter, within few seconds, the STIM1-Orai1 complex translocates to a $\text{PI}(4,5)\text{P}_2$ -rich ER/PM junctions to where all components of the Ca^{2+} signal clusters, including PMCA in the PM and SERCA and IP_3 Rs in the ER. At the junctions STIM1 polybasic domain interacts with $\text{PI}(4,5)\text{P}_2$ to stabilize the junctions and SARAF interacts with STIM1 to initiate the fast and initiate the slow Ca^{2+} -dependent Orai1 inactivation and reduce Ca^{2+} influx to a minimum necessary to sustain the physiological Ca^{2+} oscillations. Retention of the STIM1-Orai1 complex at the $\text{PI}(4,5)\text{P}_2$ -poor domain or excess hydrolysis of $\text{PI}(4,5)\text{P}_2$ by intense receptor stimulation, results in sustained and maximal activation of Ca^{2+} influx and cell pathology.

Fig. 3: TRPC channels and Orai1 participates in store-operated Ca^{2+} influx in pancreatic acinar cells

Ca^{2+} was measured in freshly isolated pancreatic acini prepared from wild-type mice (black in a, c-e, blue in b) or TRPC1^{-/-} mice (green in a-c), TRPC3^{-/-} mice (red in a-c) or Orai1^{-/-} acini (red in d, e). The cells were stimulated with 100 μM carbachol (a, d) or treated with 25 μM CPA (b, e) to deplete ER stored Ca^{2+} and then exposed to Ca^{2+} -free solution and solution containing 2 or 5 mM Ca^{2+} to assay store-dependent Ca^{2+} influx. Note that deletion of either TRPC1 or TRPC3, which interacts to mediate the same Ca^{2+} influx pathway, reduced influx by about 50%. Deletion of Orai1 in acinar cells almost completely inhibited the influx. The effect of knockout of TRPC1, TRPC3 and Orai1 on the physiological Ca^{2+} oscillations are shown in (c) and (f), respectively. The results with TRPC channels (a-c) were taken from reference [40] and with Orai1 (d-f) were taken from reference [87].

Fig. 4: Mechanism of fluid and electrolyte secretion by secretory glands acinar cells

The key transporters mediating acinar cells fluid and electrolyte secretion are the basolateral $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporters NKCC1 and the Ca^{2+} -activated K^+ Kcnma1 and the luminal Ca^{2+} -activated Cl^- channel ANO1 and the water channel Aqp5. The basolateral membrane Na^+/H^+ exchanger NHE1 and $\text{Cl}^-/\text{HCO}_3^-$ exchanger AE2 control pH_{in} and cell volume and the $\text{Cl}^-/\text{HCO}_3^-$ exchange AE4 participates in Cl^- secretion. Cell stimulation that increases $[\text{Ca}^{2+}]_i$ results in activation of Kcnma1 and ANO1 and K^+/Cl^- efflux, the obligatory water efflux through Aqp5 and cell shrinkage to halt the Ca^{2+} oscillations. Cell shrinkage phosphorylates and activates NKCC1 that restores cell K^+ and Cl^- and cell volume to allow the next Ca^{2+} spike a fluid and electrolyte secretory event.