Published in final edited form as:

J Surg Educ. 2019 ; 76(2): 370–377. doi:10.1016/j.jsurg.2018.08.022.

Social Belonging as a Predictor of Well-being and Surgical Resident Attrition

Arghavan Salles, MD, PhD^a, Robert C. Wright, MA^b, Laurel Milam, MA^a, Roheena Z. Panni, MD^a, Cara A. Liebert, MD^c, James N. Lau, MD, MHPE, FACS^c, Dana T. Lin, MD, FACS^c, and Claudia M. Mueller, PhD, MD, FACS^d

^aDepartment of Surgery, Washington University in St. Louis, Saint Louis, MO

^bPsychology Department, University of California, Riverside, Riverside, CA

^cDepartment of Surgery, Goodman Surgical Education Center, Stanford School of Medicine, Stanford, CA

^dDivision of Pediatric Surgery, Lucile Packard Children's Hospital, Palo Alto, CA

Abstract

Objective: In light of the predicted shortage of surgeons, attrition from surgical residency is a significant problem. Prior data has shown that those who are happier are more productive, and those who are less well have higher rates of absenteeism. This study sought to identify the role of social belonging and its relationship to well-being and risk of attrition.

Design: Surgical residents were invited to participate in an online survey containing measures of social belonging (a 10-item scale adapted from previous studies), well-being (the Dupuy Psychological General Well-Being Scale, Beck Depression Inventory Short Form, and Maslach Burnout Inventory), and risk of attrition (indicated by frequency of thoughts of leaving the program).

Setting: We surveyed residents at two tertiary care centers, Stanford Health Care (2010, 2011, and 2015) and Washington University in St. Louis (2017).

Participants: Categorical general surgery residents, designated preliminary residents going into 7 surgical sub-specialties, and non-designated preliminary residents were included.

Results: 169 residents responded to the survey for a response rate of 66%. Belonging was positively correlated with general psychological well-being ($r=0.56$, $p<0.0001$) and negatively

Corresponding author: Arghavan Salles, MD, PhD, Washington University in St. Louis, Department of Surgery, Saint Louis, Missouri 63110, sallesa@wustl.edu, 314-747-5544.

The authors have no conflicts of interest to report.

Financial Support: The Balance in Life program and the Goodman Surgical Center at Stanford University contributed funding. RZP was supported by NIH5T32CA00962128.

Disclosure of Financial Interests: The authors report no financial conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

correlated with depression ($r=-0.57$, $p<0.0001$), emotional exhaustion ($r=-0.58$, $p<0.0001$), and depersonalization ($r=-0.36$, $p<0.0001$). Further, belonging was negatively correlated with frequency of thoughts of leaving residency ($r=-0.45$, $p<0.0001$). In regression analysis controlling for demographic variables, belonging was a significant positive predictor of psychological well-being ($B=0.95$, $t=8.18$, $p<0.0001$) and a significant negative predictor of thoughts of leaving ($B=-1.04$, $t=-5.44$, $p<0.0001$).

Conclusions: Social belonging has a significant positive correlation with well-being and negative correlation with thoughts of leaving surgical training. Lack of social belonging appears to be a significant predictor of risk of attrition in surgical residency. Efforts to enhance social belonging may protect against resident attrition.

Keywords

Social belonging; burnout; well-being; attrition; residency

Introduction

Research has shown a steady decline in the number of general surgeons in the US.¹ Based on current training rates and population growth models there is an even greater projected shortage of general surgeons in the future.²⁻³ A contributing factor to the shortage is the high attrition rate in training programs. It has been estimated that as many as 20% of those who start surgical residency do not complete it.⁴⁻⁵ One reason for this high attrition rate could be related to poor resident well-being as previous research has shown that fatigue and depression may contribute to attrition.⁶ Thus, improving well-being and reducing burnout could subsequently decrease attrition rates.⁷

While work-hour regulations mandated by the Accreditation Council for Graduate Medical Education (ACGME) may decrease fatigue and depression, data has not yet shown lower rates of burnout with the 80-hour workweek.⁸⁻⁹ A longitudinal investigation found that one in seven surgical interns had thoughts about leaving their program weekly.¹⁰ Additionally, a large multicenter study found that 58% of surgical residents seriously considered leaving during their residency.¹¹ This study found the three most common reasons for remaining in residency were: support from family and significant others, support from other residents, and the perception of getting more rest. While the latter response seems to be addressed by the work-hour regulations, the former responses are social in nature and are likely unrelated to the work-hour regulations. As of yet, there is not a clear understanding of what specific factors lead to attrition in surgical residency. In this study, we aim to understand the relationship between attrition and a social-psychological factor, belonging, which we hypothesize may protect against attrition.

Belonging is a basic human need which is conceptualized as the feeling of social connectedness or a sense of positive association with others.¹²⁻¹³ In other fields, researchers have found strong connections between belonging and attrition. This study seeks to assess the relationship between social belonging and psychological well-being as well as thoughts of attrition among surgical residents. We hypothesize that social belonging is associated with better psychological well-being and is a predictor of attrition.

Material and methods

Following Institutional Review Board approval, surgical residents from two academic medical centers were invited to participate in an online survey. Data collection was part of an ongoing, yearly survey and took place at the first institution in 2011, 2014, and 2015.^a Data from the second institution were collected in 2017. Categorical general surgery residents as well as designated preliminary residents going into cardiothoracic, otolaryngology - head and neck surgery, vascular surgery, orthopedics, plastic surgery, urology, and neurosurgery and non-designated preliminary residents were included ($n=169$).

Belonging was assessed using a ten-item scale adapted from prior rigorous work on social belonging.^{13–16} These items were customized to suit the context of surgical training while maintaining the scale's internal reliability and specifically seeking to measure perceived belonging within the surgical specialty. The text of each item in the belonging scale is presented in Table 1. The responses were graded on a five-point scale (1=always to 5=never) and scored such that a larger value represented a greater sense of belonging ($\alpha=0.84$).

Overall well-being was assessed using a complement of three measures: the 22-item Dupuy General Psychological Well-Being Scale (DPGWB),¹⁷ the 13-item Beck Depression Inventory (BDI) Short Form,¹⁸ and the 22-item Maslach Burnout Inventory (MBI).^{19–20} The emotional exhaustion (EE) and depersonalization (DE) subscales were used in analyses as these are the most clinically relevant.

Attrition was measured in two ways. Risk of attrition was measured with a self-report item which asked, "In the past month, how often have you thought about leaving your current residency program?" Response options ranged from 1 ("Never") to 5 ("Very often"). Actual attrition was recorded for any residents at the first institution who left their program during the time of the study. There was no actual attrition from the second institution, but little time has elapsed since data collection at this site.

Three different types of residents--non-designated preliminary, designated preliminary, and categorical general surgery--were surveyed. Non-designated preliminary residents ($n=23$) were invited to participate in the study along with the other residents in order to minimize perceptions of exclusion or inferior social status. However, their data were excluded from further analyses because the temporary nature of their contracts may affect their sense of belonging and risk of attrition. In other words, since they did not have permanent contracts we would expect them to have lower levels of belonging. Designated preliminary residents were included because the survey items asked about their experience as relates to their primary specialty, not general surgery. To evaluate for non-responder bias, we performed a wave analysis comparing the responses of late responders (those who responded after the last email reminder) to early responders (those who responded to the first email).²¹ We found no statistical differences in any of the examined variables between these two groups.

^a 21 residents took the survey in both 2014 and 2015. In order to eliminate redundancy, their scores from the first year were used in analyses.

Data were first analyzed using Pearson correlations among all the key variables. Then, these relationships were investigated further using linear regression analyses. We tested for differences in the well-being outcomes by gender using t-tests. ANOVAs were performed to test for differences by type of resident (non-designated preliminary, designated preliminary, and categorical general surgery). Statistical significance was defined at the level of $p=0.05$, and all tests of significance were two-tailed. The analyses were conducted in SPSS Version 22 (IBM Corp, Armonk, NY) and SAS Version 9.4 (SAS Institute, Cary, NC).

Results

The overall response rate was 66% (169 residents). Twenty-three of these residents were non-designated preliminary residents, thus leaving 146 residents in our final sample. Participants' post-graduate years at recruitment ranged from one to five ($M=2.0$, $SD=1.6$). The residents' average age was 30.4 ($SD=3.2$). Full demographics are reported in Table 2.

Preliminary Analyses

A sub-group analysis was performed to compare data across different types of residents to assess whether the categories could be collapsed into one. Figure 1 shows the level of belonging, general psychological well-being, depression, emotional exhaustion, depersonalization, and frequency of thoughts of leaving by type of resident. Of all the outcome measures reported here, belonging was the only one that varied significantly by type of resident. There were statistically significant differences in level of belonging among the three groups ($F_{2, 146}=7.00$, $p=0.0013$) such that non-designated preliminary residents had lower levels of belonging ($M=3.3$, $SD=0.1$) than did either the categorical general surgery residents ($M=3.8$, $SD=0.1$, $t=3.63$, $p=0.0011$) or the designated preliminary residents ($M=3.7$, $SD=0.1$, $t=3.32$, $p=0.0032$). There was no significant difference in belonging between categorical general surgery residents and designated preliminary residents ($t=0.26$, $p=0.9646$). There were no significant differences in psychological well-being, depression, burnout, or thoughts about leaving residency among the three groups of residents (general surgery categorical, designated preliminary, and non-designated preliminary; $ps>0.10$).

Additionally, we assessed whether any of the well-being outcomes differed by gender. We found a statistically significant difference in emotional exhaustion, with women having higher emotional exhaustion compared to men ($t=-2.51$, $p=0.0131$). We found no significant differences by gender in belonging, psychological well-being, depression, depersonalization, or thoughts of leaving residency ($ps>0.1354$).

Because non-designated preliminary residents' contracts are by definition short-term, it is not meaningful to assess their responses regarding risk of attrition. Also, as expected, their level of belonging differed significantly from those in general surgery categorical or designated preliminary positions. Thus, all further analyses were performed only on the subset of general surgery categorical and designated preliminary residents.

Correlations Among Variables

The correlations among the main variables are shown in Table 3. As expected, belonging was significantly positively associated with well-being as measured by general

psychological well-being ($r=0.56$, $p<0.0001$). It was also significantly negatively associated with outcomes signifying resident distress such as depression ($r=-0.57$, $p<0.0001$), emotional exhaustion ($r=-0.58$, $p<0.0001$), and depersonalization ($r=-0.36$, $p<0.0001$). Belonging was also negatively correlated with risk of attrition as measured by frequency of thoughts of leaving residency ($r=-0.45$, $p<0.0001$). Lastly, there were seven residents at the first institution who left residency during the study period (7.53%). Five were general surgery categorical residents, and 2 were designated preliminary residents. The point-biserial correlation between belonging and actual attrition was non-significant ($r=0.15$, $p=0.1846$).

Regression Analyses

Next, controlling for gender, age, ethnicity, and post-graduate year (PGY), regression analyses found that belonging was a significant positive predictor of general psychological wellbeing ($B=0.95$, $t=8.18$, $p<0.0001$). Furthermore, while controlling for the same variables, belonging was a significant negative predictor of depression ($B=-0.27$, $t=-3.49$, $p=0.0014$). In similar analyses, belonging was a significant negative predictor of emotional exhaustion ($B=-1.34$, $t=-6.90$, $p<0.0001$) and depersonalization ($B=-0.86$, $t=-3.76$, $p=0.0003$). Thus, belonging was positively predictive of a desirable outcome (general psychological well-being) and negatively predictive of undesirable outcomes (depression, emotional exhaustion, and depersonalization). In the multivariable analyses, gender was not a significant predictor of any of the well-being outcomes ($p=0.0612$).

We also analyzed whether belonging predicted risk of attrition in multivariable regression. Table 4 shows that, while controlling for the same demographic variables, belonging was negatively associated with frequency of thoughts of leaving residency ($B=-1.04$, $t=-5.44$, $p<0.0001$). The covariates gender, age, and post-graduate year were not significantly associated with belonging in the regression model.

Discussion

In this study examining social belonging in the context of surgical training, we demonstrate two main findings: 1) that social belonging is associated with well-being and 2) that a surgical resident's lack of social belonging is associated with more thoughts of leaving surgical training. The correlation between social belonging and well-being suggests that increasing belonging may improve the well-being of residents. Our second major finding suggests that efforts to cultivate and enhance social belonging may protect against residency attrition. As the research on physician well-being shifts from focusing on individual characteristics (such as resilience) and starts to examine systemic and institutional interventions²², our data suggest that efforts should at least in part be aimed at increasing connectedness and belonging.

Several studies have examined the association between belonging and factors that may influence risk of attrition and group dynamics. Notably, in a series of elegant studies, Walton and Cohen have demonstrated how a threat to social belonging in an academic setting can lead to negative outcomes for participants.¹⁴ Specifically, minority students who felt low social connectedness or sense of belonging also felt low perceived fit and lower potential. The researchers subsequently performed interventions to create a sense of belonging in these

students which resulted in better subjective well-being (e.g., happiness) as well as health benefits (e.g., improved perceived health and fewer doctor visits).¹³ In a large multisite investigation of undergraduates in STEM fields researchers found a strong link between the sense of belonging and engagement.²³ Taken together, these results add to a growing body of literature that shows that social belonging has positive influence on members of a group.²⁴

Cultivating a sense of belonging among trainees may be more important now than it has been in the past. Clinical care in the modern era is more isolating than ever before.²⁵ As physicians spend more time in front of computers, there are fewer opportunities to interact with colleagues. Current efforts to improve physician well-being have focused mainly on individual strategies such as increasing resilience and mindfulness.^{26–28} While important, these strategies fail to recognize the fundamental human need to feel connected to others.¹⁴ Opportunities, such as Balint groups, that bring physicians together may be protective against burnout²⁹ by giving physicians a venue in which to engage with each other in a meaningful way. Our data are consistent with these findings and suggest that interventions aimed at increasing engagement and connectedness may be associated with better well-being and lower risk of attrition.

Although the reasons for which surgical residents leave their programs may be varied and complex, belonging appears to be a factor that is predictive of how often these thoughts occur. This, in turn, may be predictive of true attrition. Certainly, multiple other factors, such as lifestyle and job satisfaction, likely contribute to trainees' decisions to leave their residencies. We were unable to demonstrate a significant link between belonging and actual attrition, and the study is underpowered to examine this endpoint in such a small sample. Nonetheless, while elements such as the work environment and job requirements are not likely to change in the short term, interventions aimed at improving residents' well-being should be further investigated.³⁰ We have shown here that belonging is a factor that is highly correlated with well-being and thus presents itself as a promising target for effective intervention.

Of note, some degree of attrition may be appropriate. Not everyone who enters surgical training, or any other field, will or should stay in it. However, the attrition rate in general surgery is higher than that of other specialties, and it seems appropriate to attempt to bring this in line with other specialties. For example, the annual rate of attrition in ophthalmology, orthopaedic surgery, otolaryngology and head and neck surgery, and urology is approximately 1.5%.³¹ General surgery, then, with an annual attrition rate of 5% (overall 20% over the five clinical years of training), is clearly an outlier.

An interesting null finding in this study was the lack of gender differences in most of the raw well-being measures. Population data have suggested that women are more likely to be depressed than men.³² Furthermore, women may be more at risk for leaving the surgical specialties than men.^{33–37} The data here show, however, that there is no difference in any of the well-being measures by gender besides emotional exhaustion. In regression analyses, gender was not predictive of any of the well-being outcomes. The relationship between gender and psychological well-being is complex and is not accounted for simply by gender itself. For example, previous research has shown less stereotyping and more perceived

equality in specialties such as general surgery when there are roughly equal numbers of male and female residents.^{30,38} In this sample, there was a relatively high percentage of women (42%), which may explain the lack of differential well-being outcomes by gender.

There were several limitations to the present study. First, this is a study conducted at only two academic institutions, limiting the generalizability of our conclusions. Additional work, ideally multi-institutional in nature, is needed to confirm the veracity of our findings. Second, PGY 1 and 2 residents made up a significant portion of the sample, reflecting the higher proportion of junior to senior residents in the training programs. However, residents who quit typically do so in the initial or professional development years.³⁹ Therefore, the fact that the study sample contains a higher representation of these residents may actually be beneficial. Lastly, the survey item measuring attrition risk has yet to be validated. However, given the yearly attrition rate, this would only be feasible in a large longitudinal multi-institution study.

To conclude, our findings indicate that social belonging and connectivity are important factors that influence not only resident well-being but also the desire to leave residency training. As programs try to ensure their residents' well-being and satisfaction to improve retention, this research suggests that efforts should focus specifically at improving residents' sense of belonging within their program. Future research should focus on examining belonging as it relates to actual attrition, as well as designing and investigating social belonging interventions in order to effectively optimize well-being and retention among surgical trainees.

Acknowledgements

Jennifer S. Tran, Assistant Operations Manager at the Goodman Surgical Center, helped with data collection for this study.

The Balance in Life program contributed funding for this research which was used for data collection. RZP was supported by NIH5T32CA00962128.

The Goodman Surgical Center contributed funding for this research which was used for data collection.

Arghavan Salles had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Institution:

Stanford Health Care, Department of Surgery, Stanford School of Medicine 300 Pasteur Drive H3552A Stanford, CA 94305

Presentation: A portion of this work was presented orally at the American College of Surgeons 101st Annual Clinical Congress, Chicago, IL, October 2015.

References

1. Lynge DC, Larson EH, Thompson MJ, et al. A longitudinal analysis of the general surgery workforce in the United States, 1981–2005. *Arch Surg* 2008;1434:345–350. [PubMed: 18427021]
2. Fischer JE. "The impending disappearance of the general surgeon." *JAMA* 2007;29818:2191–2193. [PubMed: 18000204]
3. Williams TE, and Ellison EC. Population analysis predicts a future critical shortage of general surgeons. *Surgery* 2008;1444:548–556. [PubMed: 18847638]

4. Longo WE, Seashore J, Duffy A, Udelsman R. Attrition of categoric general surgery residents: results of a 20-year audit. *Am J Surg* 2009;1976:774–778. [PubMed: 19178898]
5. Yeo H, Bucholz E, Ann Sosa J, et al. A national study of attrition in general surgery training: which residents leave and where do they go? *Ann Surg* 2010;2523:529–536. [PubMed: 20739854]
6. Zaré SM, Galanko J, Behms KE, et al. Psychological well-being of surgery residents before the 80-hour work week: a multiinstitutional study. *J Am Coll Surg* 2004;1984:633–640. [PubMed: 15051017]
7. Shanafelt TD, Balch CM, GJ Bechamps, et al. Burnout and career satisfaction among American surgeons. *Ann Surg* 2009;2503:463–471. [PubMed: 19730177]
8. Zaré SM, Galanko J, Behms KE, et al. Psychologic well-being of surgery residents after inception of the 80-hour workweek: a multi-institutional study. *Surgery* 2005;1382:150–157. [PubMed: 16153421]
9. Ahmed N, Devitt KS, Keshet I, et al. A systematic review of the effects of resident duty hour restrictions in surgery: impact on resident wellness, training, and patient outcomes. *Ann Surg* 2014;2596:1041. [PubMed: 24662409]
10. Antiel RM, Reed DA, Van Arendonk KJ, et al. Effects of duty hour restrictions on core competencies, education, quality of life, and burnout among general surgery interns.” *JAMA Surg* 2013;1485:448–455. [PubMed: 23325404]
11. Gifford E, Galante J, Kaji AH, et al. Factors Associated With General Surgery Residents’ Desire to Leave Residency Programs: A Multi-institutional Study. *JAMA Surg* 2014;1499:948–953. [PubMed: 25075473]
12. Baumeister RF, Leary MR. The need to belong: desire for interpersonal attachments as a fundamental human motivation. *Psychol Bull* 1995;1173:497. [PubMed: 7777651]
13. Walton GM, Cohen GL. A brief social-belonging intervention improves academic and health outcomes of minority students. *Science* 2011;3316023:1447–1451. [PubMed: 21415354]
14. Walton GM, Cohen GL. A question of belonging: race, social fit, and achievement. *J Pers Soc Psychol* 2007;921:82. [PubMed: 17201544]
15. Walton GM, Cohen GL, Cwir D, & Spencer SJ. Mere belonging: The power of social connections. *J Pers Soc Psychol* 2012;102(3)513–32. [PubMed: 22023711]
16. Walton GM, Logel C, Peach JM, et al. Two brief interventions to mitigate a “chilly climate” transform women’s experience, relationships, and achievement in engineering. *J Ed Psychol* 2014;107(2):468–85.
17. Dupuy HJ. The psychological general well-being (PGWB) index In Wenger NK, Mattson ME, Furburg CD, Elinson J, editors. Assessment of quality of life in clinical trials of cardiovascular therapies. Le Jacq Publishing; 1984:170–183.
18. Beck AT, Beck RW. Screening depressed patients in family practice: A rapid technique. *Postgrad Med* 1982;52:81–85.
19. Maslach C, Jackson SE. The measurement of experienced burnout. *J Occ Behav* 2001;2:99–113.
20. Maslach C, Jackson SE. Maslach Burnout Inventory-Human Services Survey (MBI-HSS) In Maslach C, Jackson SE, Leiter MP, editors. MBI manual 3rd ed Mountain View, CA: CPP, Inc; 1996.
21. Phillips AW, Reddy S, Durning SJ. Improving response rates and evaluating nonresponse bias in surveys: AMEE Guide No. 102. *Med Teach* 2016;38(3): 217–28. [PubMed: 26648511]
22. Shanafelt TD, Noseworthy JH. Executive Leadership and Physician Well-being: Nine Organizational Strategies to Promote Engagement and Reduce Burnout. *Mayo Clin Proc.* 2017;92(1):129–146. doi:10.1016/j.mayocp.2016.10.004. [PubMed: 27871627]
23. Wilson D, Jones D, Bocell F, et al. Belonging and academic engagement among undergraduate STEM students: A multi-institutional study. *Res High Educ* 2015;56:750–776.
24. Sargent J, Williams RA, Hagerty B, et al. Sense of belonging as a buffer against depressive symptoms. *J Am Psychiatr Nurses Assoc* 2002;84:120–129.
25. Ziegelstein RC. Creating Structured Opportunities for Social Engagement to Promote WellBeing and Reduce Burnout in Medical Students and Residents. *Acad Med.* 2017;1. doi: 10.1097/ACM.0000000000002117.

26. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *Lancet*. 2016;388(10057):2272–2281. doi: 10.1016/S0140-6736(16)31279-X. [PubMed: 27692469]
27. Alobuia W, Salles A, Gibson M, Mueller C. Physician wellness in surgical residency. *Current Surgery Reports* 2018; 6(3).
28. Souba WW J Z CM, et al. Resilience—Back to the Future. *JAMA Surg*. 2016;88(3):382–389. doi: 10.1001/jamasurg.2016.1056.
29. Ghetti C, Chang J, Gosman G. Burnout, Psychological Skills, and Empathy: Balint Training in Obstetrics and Gynecology Residents. *J Grad Med Educ*. 2009;1(2):231–235. doi: 10.4300/JGME-D-09-00049.1. [PubMed: 21975984]
30. Salles A, Mueller CM, Cohen GL. Exploring the Relationship Between Stereotype Perception and Residents' Well-Being. *J Am Coll Surg* 2015;2221:52–58. [PubMed: 26616033]
31. Pugno PA. Residency attrition: trends and themes. 2011 Available at: https://www.aamc.org/download/185478/data/2011_pwc_pugno.pdf. Accessed March 10, 2015.
32. Kessler RC. Epidemiology of women and depression. *J Affect Disord* 2003;741:5–13. [PubMed: 12646294]
33. Bergen PC, Turnage RH, & Carrico CJ. Gender-related attrition in a general surgery training program. *J Surg Res* 1998;77:59–62. [PubMed: 9698534]
34. Yeo H, Viola K, Berg D, et al. Attitudes, training experiences, and professional expectations of US general surgery residents: A national survey. *JAMA* 2009;302:1301–1308. [PubMed: 19773565]
35. Lynch G, Nieto K, Puthenveetil S, et al. Attrition rates in neurosurgery residency: Analysis of 1361 consecutive residents matched from 1990–1999. *JNeurosurg* 2015;122:240–249. [PubMed: 25415065]
36. Walker JL, Janssen H, Hubbard D. Gender differences in attrition from orthopaedic surgery residency. *J Am Med Womens Assoc* 1993;48:182–184.
37. Yeo HL, Abelson JS, Symer MM, et al. Association of Time to Attrition in Surgical Residency With Individual Resident and Programmatic Factors. *JAMA Surg*. 2018. doi: 10.1001/jamasurg.2017.6202.
38. Dahlerup D From a Small to a Large Minority: Women in Scandinavian Politics. *Scan Polit Stud*. 1988;11(4):275–298. doi:10.1111/j.1467-9477.1988.tb00372.x.
39. Yeo H, Bucholz E, Ann Sosa J, et al. A national study of attrition in general surgery training: which residents leave and where do they go? *Ann Surg* 2010;252(3):529–34. [PubMed: 20739854]

Highlights

- Social belonging was negatively correlated with depression
- Belonging was inversely correlated with emotional exhaustion and depersonalization
- In regression, belonging was a positive predictor of psychological well-being
- Belonging was a significant negative predictor of thoughts of leaving residency

Figure 1.

Well-being and risk of attrition by resident type. *Level of belonging was significantly lower for non-designated preliminary residents than categorical or designated preliminary residents ($p < 0.01$).

Table 1.

Belonging scale items.

Number	Question text
1	How often do you feel like you belong in your specialty?
2	How often do you feel like you belong at [institution]?
3	How often do you feel like people in your specialty accept you?
4	How often do you feel like an outsider in your specialty?
5	How often do you feel like people in your specialty are similar to you?
6	How often do you feel like people at [institution] are similar to you?
7	How often do you feel like you know what you need to do to succeed in your specialty?
8	How often do you feel like there is someone in your specialty you can count on?
9	How often do you feel different from the other residents in your specialty?
10	How often do you feel different from other residents at [institution]?

Table 2.

Demographics.

Variables	<i>n</i> (%)
Gender	
Female	71 (42.0)
Male	98 (58.0)
Age	
25–26	10(5.9)
27–29	55 (32.5)
30–32	37(21.9)
33+	31 (18.3)
Unknown	36(21.3)
Ethnicity	
Caucasian	93 (55.0)
African-American	9(5.3)
Hispanic/Latino	11 (6.5)
Asian/Pacific Islander	44 (26.0)
Middle Eastern	6 (3.6)
Mixed	5 (3.0)
Unknown	1 (0.6)
Post-graduate year	
1	96 (56.8)
2	34 (20.1)
3	10(5.9)
4	8 (4.7)
5	10(5.9)
Research/professional development	11 (6.5)
Type of resident	
General surgery categorical	80 (47.3)
Designated preliminary resident	66 (39.1)
Non-designated preliminary resident	23 (13.6)

Table 3.Correlations among key variables.^a

	Belonging	Well-being	Depression	Emotional exhaustion	Depersonalization	Thoughts of leaving
Belonging	1					
Well-being	0.56 ^{**}	1				
Depression	-0.57 ^{**}	-0.80 ^{**}	1			
Emotional exhaustion	-0.58 ^{**}	-0.77 ^{**}	0.68 ^{**}	1		
Depersonalization	-0.36 ^{**}	-0.51 ^{**}	0.55 ^{**}	0.71 ^{**}	1	
Thoughts of leaving residency	-0.45 ^{**}	-0.56 ^{**}	0.46 [*]	0.63 ^{**}	0.48 ^{**}	1

*
 $p < 0.001$,**
 $p < 0.0001$ ^aThe analytical sample includes only categorical general surgical and designated preliminary residents.

Table 4.

Regression analysis of thoughts of leaving residency on belonging controlling for age, gender, ethnicity, and PGY ($n=105$).^a

Variable	Coefficient	<i>t</i>	<i>P</i>
Belonging	-1.04	-5.44	<0.0001
Age	-0.00	-0.02	0.9833
Gender (female)	0.08	0.39	0.6943
Ethnicity			
African-American	-0.42	-0.71	0.4776
Hispanic or Latino	0.34	0.73	0.4649
Asian/Pacific Islander	0.03	0.13	0.9001
Middle Eastern	1.07	1.04	0.2993
Mixed race	1.24	2.40	0.0186
PGY			
PGY 2	0.15	0.46	0.6437
PGY 3	0.27	0.68	0.4966
PGY 4	-0.83	-1.71	0.0912
PGY 5	-0.61	-1.48	0.1422
Research	-0.14	-0.40	0.6909
Constant	5.88	4.09	<0.0001

^aThe analytical sample includes only categorical general surgical and designated preliminary residents