Cochrane
Library

Cochrane Database of Systematic Reviews

Interventions for primary vesicoureteric reflux (Review)

Williams G, Hodson EM, Craig JC

Williams G, Hodson EM, Craig JC.
Interventions for primary vesicoureteric reflux.
Cochrane Database of Systematic Reviews 2019, Issue 2. Art. No.: CD001532.
DOI: [10.1002/14651858.CD001532.pub5](https://doi.org/10.1002/14651858.CD001532.pub5).

www.cochranelibrary.com

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	12
OBJECTIVES	12
METHODS	12
RESULTS	14
Figure 1.	15
Figure 2.	17
Figure 3.	18
DISCUSSION	25
AUTHORS' CONCLUSIONS	26
ACKNOWLEDGEMENTS	27
REFERENCES	28
CHARACTERISTICS OF STUDIES	35
DATA AND ANALYSES	75
Analysis 1.1. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 1 Urinary tract infection.	78
Analysis 1.2. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 2 All symptomatic UTI by 1-2 years: allocation concealment.	79
Analysis 1.3. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 3 All symptomatic UTI by 1-2 years: blinding.	80
Analysis 1.4. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 4 Renal parenchymal abnormality on DMSA scan: unit of analysis (children).	80
Analysis 1.5. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 5 Resolution of VUR.	81
Analysis 1.6. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 6 Adverse events.	82
Analysis 1.7. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 7 Microbial resistance to prophylactic drug.	82
Analysis 1.8. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 8 GFR at 3 years.	82
Analysis 1.9. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 9 Renal growth at 3 years.	82
Analysis 1.10. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 10 All symptomatic UTI by 1-2 years: addressing of incomplete outcome data.	83
Analysis 1.11. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 11 Febrile UTI by 1-2 years: allocation concealment.	83
Analysis 1.12. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 12 Febrile UTI by 1-2 years: blinding.	84
Analysis 1.13. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 13 Febrile UTI by 1-2 years: addressing of incomplete outcome data.	85
Analysis 1.14. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 14 Febrile UTI by 1-2 years according to gender.	85
Analysis 2.1. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 1 Urinary tract infection.	89
Analysis 2.2. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 2 Renal parenchymal defects (scars and thinning) on IVP: unit of analysis (children).	90
Analysis 2.3. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 3 Renal scars on IVP: unit of analysis (children).	91
Analysis 2.4. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 4 Repeat positive urine culture.	92
Analysis 2.5. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 5 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children).	92
Analysis 2.6. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 6 Renal parenchymal defects on IVP: unit of analysis (kidneys).	93
Analysis 2.7. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 7 Hypertension and ESKD.	94

Analysis 2.8. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 8 GFR measured by Schwartz formula.	94
Analysis 2.9. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 9 Height Standard Deviation Score (SDS).	95
Analysis 2.10. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 10 Correction of VUR: unit of analysis (children).	95
Analysis 2.11. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 11 Correction of VUR: unit of analysis (ureters).	95
Analysis 3.1. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 1 Repeat positive urine culture.	96
Analysis 3.2. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 2 Persistent VUR (ureters).	96
Analysis 3.3. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 3 Peri-operative complications.	96
Analysis 4.1. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 1 Urinary tract infection.	98
Analysis 4.2. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 2 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children).	99
Analysis 4.3. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 3 Renal damage on ultrasound.	99
Analysis 4.4. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 4 VUR status after treatment: unit of analysis (children).	100
Analysis 4.5. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 5 Microbial resistance to the prophylactic drug.	100
Analysis 5.1. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 1 Symptomatic urinary tract infection.	101
Analysis 5.2. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 2 Febrile urinary tract infection. ...	101
Analysis 5.3. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 3 Renal parenchymal abnormality.	101
Analysis 6.1. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 1 Macroplastique versus dextranomer/hyaluronic acid polymer (Deflux).	102
Analysis 6.2. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 2 GAX 65 versus GAX 35.	103
Analysis 6.3. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 3 Dexcel versus Vantris.	104
Analysis 7.1. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 1 Urinary tract infection.	105
Analysis 7.2. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 2 E coli resistance to prophylactic drug.	105
Analysis 7.3. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 3 Renal parenchymal defects on DMSA scan.	106
Analysis 7.4. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 4 Probiotics plus antibiotics versus antibiotics alone.	106
Analysis 8.1. Comparison 8 Other interventions, Outcome 1 Reimplantation surgery versus endoscopic treatment.	108
Analysis 8.2. Comparison 8 Other interventions, Outcome 2 Cranberry product versus no treatment.	108
Analysis 8.3. Comparison 8 Other interventions, Outcome 3 Circumcision plus antibiotics versus antibiotics alone.	108
Analysis 8.4. Comparison 8 Other interventions, Outcome 4 Oxybutynin versus placebo.	109
APPENDICES	109
FEEDBACK	111
WHAT'S NEW	111
HISTORY	112
CONTRIBUTIONS OF AUTHORS	112
DECLARATIONS OF INTEREST	112
SOURCES OF SUPPORT	112
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	112
INDEX TERMS	112

[Intervention Review]

Interventions for primary vesicoureteric reflux

Gabrielle Williams¹, Elisabeth M Hodson^{2,3}, Jonathan C Craig^{3,4}

¹Analytics Assist, NSW Ministry of Health, North Sydney, Australia. ²Sydney School of Public Health, The University of Sydney, Sydney, Australia. ³Cochrane Kidney and Transplant, Centre for Kidney Research, The Children's Hospital at Westmead, Westmead, Australia.

⁴College of Medicine and Public Health, Flinders University, Adelaide, Australia

Contact: Elisabeth M Hodson, Cochrane Kidney and Transplant, Centre for Kidney Research, The Children's Hospital at Westmead, Locked Bag 4001, Westmead, NSW, 2145, Australia. elisabeth.hodson@health.nsw.gov.au.

Editorial group: Cochrane Kidney and Transplant Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 2, 2019.

Citation: Williams G, Hodson EM, Craig JC. Interventions for primary vesicoureteric reflux. *Cochrane Database of Systematic Reviews* 2019, Issue 2. Art. No.: CD001532. DOI: [10.1002/14651858.CD001532.pub5](https://doi.org/10.1002/14651858.CD001532.pub5).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Vesicoureteric reflux (VUR) results in urine passing retrograde up the ureter. Urinary tract infections (UTI) associated with VUR have been considered a cause of permanent renal parenchymal damage in children with VUR. Management has been directed at preventing UTI by antibiotic prophylaxis and/or surgical correction of VUR. This is an update of a review first published in 2004 and updated in 2007 and 2011.

Objectives

The aim of this review was to evaluate the available evidence for both benefits and harms of the currently available treatment options for primary VUR: operative, non-operative or no intervention.

Search methods

We searched the Cochrane Kidney and Transplant Specialised Register to 3 May 2018 through contact with the Information Specialist using search terms relevant to this review. Studies contained in the Specialised Register are identified through search strategies specifically designed for CENTRAL, MEDLINE, and EMBASE; handsearching conference proceedings, and searching the International Clinical Trials Register (ICTRP) Search Portal and ClinicalTrials.gov.

Selection criteria

RCTs in any language comparing any treatment of VUR and any combination of therapies.

Data collection and analysis

Two authors independently determined study eligibility, assessed quality and extracted data. Dichotomous outcomes were expressed as risk ratios (RR) with 95% confidence intervals (CI) and continuous data as mean differences (MD) with 95% CI. Data were pooled using the random effects model.

Main results

Thirty four studies involving 4001 children were included. Interventions included; long-term low-dose antibiotics, surgical reimplantation of ureters, endoscopic injection treatment, probiotics, cranberry products, circumcision, and oxybutynin. Interventions were used alone and in combinations. The quality of conduct and reporting of these studies was variable, with many studies omitting crucial methodological information used to assess the risk of bias. Only four of the 34 studies were considered at low risk of bias across all fields of study quality. The majority of studies had many areas of uncertainty in the risk of bias fields, reflecting missing detail rather than stated poor design.

Low-dose antibiotic prophylaxis compared to no treatment/placebo may make little or no difference to the risk of repeat symptomatic UTI (9 studies, 1667 children: RR 0.77, 95% CI 0.54 to 1.09; low certainty evidence) and febrile UTI (RR 0.83, 95% CI 0.56 to 1.21; low certainty evidence) at one to two years. At one to three years, antibiotic prophylaxis made little or no difference to the risk of new or progressive renal damage on DMSA scan (8 studies, 1503 children: RR 0.73, 95% CI 0.33 to 1.61; low certainty evidence). Adverse events were reported in four studies with little or no difference between treatment groups (1056 children: RR 0.94, 95% CI 0.81 to 1.08;), but antibiotics increased the likelihood of bacterial drug resistance threefold (187 UTIs: RR 2.97, 95% CI 1.54 to 5.74; moderate certainty evidence).

Seven studies compared long-term antibiotic prophylaxis alone with surgical reimplantation of ureters plus antibiotics, but only two reported the outcome febrile UTI (429 children). Surgery plus antibiotic treatment may reduce the risk of repeat febrile UTI by 57% (RR 0.43, 95% CI 0.27 to 0.70; moderate certainty evidence). There was little or no difference in the risk of new kidney defects detected using intravenous pyelogram at 4 to 5 years (4 studies, 572 children, RR 1.09, 95% CI 0.79 to 1.49; moderate certainty evidence)

Four studies compared endoscopic injection with antibiotics alone and three reported the outcome febrile UTI. This analysis showed little or no difference in the risk of febrile UTI with endoscopic injection compared to antibiotics (RR 0.74, 95% CI 0.31 to 1.78; low certainty evidence). Four studies involving 425 children compared two different materials for endoscopic injection under the ureters (polydimethylsiloxane (Macroplastique) versus dextranomer/hyaluronic acid polymer (Deflux), glutaraldehyde cross-linked (GAX) collagen (GAX) 35 versus GAX 65 and Deflux versus polyacrylate polyalcohol copolymer (VANTRIS)) but only one study (255 children, low certainty evidence) had the outcome of febrile UTI and it reported no difference between the materials. All four studies reported rates of resolution of VUR, and the two studies comparing Macroplastique with Deflux showed that Macroplastique was probably superior to dextranomer/hyaluronic acid polymer (3 months: RR 0.50, 95% CI 0.33 to 0.78; 12 months: RR 0.54 95% CI 0.35 to 0.83; low certainty evidence)

Two studies compared probiotic treatment with antibiotics and showed little or no difference in risk of repeat symptomatic UTI (RR 0.82 95% CI 0.56 to 1.21; low certainty evidence)

Single studies compared circumcision with antibiotics, cranberry products with no treatment, oxybutynin with placebo, two different surgical techniques and endoscopic injection with no treatment.

Authors' conclusions

Compared with no treatment, the use of long-term, low-dose antibiotics may make little or no difference to the number of repeat symptomatic and febrile UTIs in children with VUR (low certainty evidence). Considerable variation in the study designs and subsequent findings prevented drawing firm conclusions on efficacy of antibiotic treatment.

The added benefit of surgical or endoscopic correction of VUR over antibiotic treatment alone remains unclear since few studies comparing the same treatment and with relevant clinical outcomes were available for analysis.

PLAIN LANGUAGE SUMMARY

Interventions for primary vesicoureteric reflux

What is the issue?

Vesicoureteric reflux (VUR) is the backflow of urine from the bladder up the ureters to the kidney. People with VUR are thought to be more likely to get urinary tract infections (UTIs) involving the kidney tissue, which may cause permanent kidney damage. Current treatment options include surgical reimplantation of the ureters, long-term antibiotics, endoscopic correction by subureteric injection of a substance (injection into the bladder under the ureter), complementary medicines, or a combination of interventions

What did we do?

A literature search identified all randomised studies comparing different treatment options in children with vesicoureteric reflux. Results along with study design details were extracted and compiled. Studies comparing similar treatments and outcomes had data pooled to obtain an estimate of effect on outcomes such as repeat urinary tract infection with illness, urinary tract infection with fever and kidney damage.

What did we find?

A total of 34 randomised studies, involving 4001 children were identified and underwent data extraction and analysis. The most frequent comparisons were for long-term, low-dose antibiotics with no treatment (8 studies) or placebo (4 studies) and antibiotics versus surgical reimplantation of ureters plus antibiotics (7 studies). Other treatments looked at endoscopic correction by injection compared with antibiotics (3 studies), different materials for endoscopic correction (2 studies) circumcision (1 study), probiotics (1 study), cranberry product (1 study), and oxybutynin (2 studies).

Meta-analysis of similar studies found that long-term low-dose antibiotic treatment compared with no treatment may lead to little or no difference in the risk for repeat UTIs in children with VUR. Associated side effects were infrequent and minor, but prophylaxis was associated with a threefold increased risk of bacterial resistance to the treatment drug in later infections. Surgery decreased the number of repeat

UTIs with fever, but did not change the number of children developing UTI with illness or kidney damage. Many studies did not contribute to the meta-analysis as they failed to report relevant outcomes or were single studies examining a treatment option not used by other studies or combinations of treatments.

Conclusions

Long-term low-dose antibiotic treatment in children with VUR makes little or no difference to the risk of repeat UTI causing a person to be unwell. Surgery may reduce the risk of repeat UTI with fever however this is based on two studies of 429 children who may not represent the majority and may not bear true in a more general group of children with VUR. Complementary therapies such as probiotics and cranberry were trialled in single or two studies and do not provide evidence of sufficient certainty to support or deny their use.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Antibiotic prophylaxis versus surveillance or no treatment for primary vesicoureteric reflux

Antibiotic prophylaxis versus surveillance or no treatment for primary VUR

Patient or population: primary VUR
Setting: paediatric clinics
Intervention: antibiotic prophylaxis
Comparison: surveillance or no treatment

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with surveillance or no treatment	Risk with Antibiotic prophylaxis				
Symptomatic UTI by 1 to 2 years: antibiotics versus no prophylaxis	222 per 1,000	171 per 1,000 (120 to 242)	RR 0.77 (0.54 to 1.09)	1667 (9)	⊕⊕⊕⊖ LOW ^{1 2}	Poor clarity of methods in many studies may lead to bias and this could explain the inconsistent findings across these studies. This reduces our confidence in the evidence
Febrile UTI by 1 to 2 years: antibiotics versus no prophylaxis	143 per 1,000	118 per 1,000 (80 to 172)	RR 0.83 (0.56 to 1.21)	1667 (9)	⊕⊕⊕⊖ LOW ^{1 2}	Poor clarity of methods in many studies may lead to bias and this could explain the inconsistent findings across these studies. This reduces our confidence in the evidence
Symptomatic UTI; antibiotics versus placebo in trials with low risk of bias across all domains	203 per 1,000	173 per 1,000 (81 to 364)	RR 0.85 (0.40 to 1.79)	943 (3)	⊕⊕⊕⊖ MODERATE ²	Two relatively large studies showed consistent results but the third small study with small numbers of participants and few events was not consistent
Resolution of VUR after 1 to 2 years: unit of analysis (children)	153 per 1,000	223 per 1,000 (108 to 456)	RR 1.46 (0.71 to 2.99)	262 (3)	⊕⊕⊕⊖ LOW ^{1 3}	Poor clarity of methods in some studies and imprecision due to small numbers of included participants may lead to bias. This reduces our confidence in the evidence
Adverse events	342 per 1,000	321 per 1,000 (277 to 369)	RR 0.94 (0.81 to 1.08)	1056 (4)	⊕⊕⊕⊖ MODERATE ⁴	Design faults in one study reduces our confidence in the evidence
Microbial resistance to prophylactic drug: unit of analysis (UTI)	237 per 1,000	645 per 1,000 (439 to 949)	RR 2.72 (1.85 to 4.00)	259 (6)	⊕⊕⊕⊖ MODERATE ⁵	Small sample sizes of participants with UTI makes the results imprecise and reduces our confidence in the evidence

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio; **UTI:** Urinary tract infection; **VUR:** vesicoureteric reflux

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

- 1 Most studies had serious design faults
- 2 Inconsistency between study results with one study showing results opposite to the other studies
- 3 Small number of events in most studies
- 4 One study had a serious design fault
- 5 Small numbers of urinary tract infections results in imprecision

Summary of findings 2. Surgical reimplantation of ureters plus antibiotics versus antibiotics alone for primary vesicoureteric reflux

Surgical reimplantation of ureters plus antibiotics versus antibiotics alone for primary VUR

Patient or population: primary VUR

Setting: paediatric clinics

Intervention: surgical reimplantation of ureters plus antibiotics

Comparison: antibiotics alone

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with antibiotics alone	Risk with Surgical reimplantation of ureters plus antibiotics				
All symptomatic UTI by 4 to 5 years: surgical correction + antibiotics versus antibiotics alone	307 per 1,000	291 per 1,000 (205 to 414)	RR 0.95 (0.67 to 1.35)	297 (1)	⊕⊕⊕⊙ MODERATE ¹	Only one study evaluated this outcome so our confidence in the evidence is reduced
Febrile UTI by 5 years: surgical correction + antibiotics versus antibiotics alone	220 per 1,000	95 per 1,000 (59 to 154)	RR 0.43 (0.27 to 0.70)	429 (2)	⊕⊕⊕⊙ MODERATE ²	Two studies evaluated this outcome but one of these had risk of bias issues

						so our confidence in the evidence is reduced
Repeat positive urine cultures by 4 to 5 years: surgical correction + antibiotics versus antibiotics alone	369 per 1,000	365 per 1,000 (291 to 465)	RR 0.99 (0.79 to 1.26)	479 (3)	⊕⊕⊕⊕ MODERATE ³	Three studies evaluated this outcome but two of these had risk of bias issues reducing our confidence in the evidence
New renal parenchymal defects (scars and thinning) on IVP: unit of analysis (children) at 4 to 5 years: surgical correction + antibiotics versus antibiotics alone	195 per 1,000	213 per 1,000 (154 to 291)	RR 1.09 (0.79 to 1.49)	572 (4)	⊕⊕⊕⊕ MODERATE ³	Four studies evaluated this outcome but three of these had risk of bias issues reducing our confidence in the issues
ESKD	38 per 1,000	41 per 1,000 (9 to 191)	RR 1.07 (0.23 to 5.04)	154 (2)	⊕⊕⊕⊕ VERY LOW ^{4 5}	Few participants included in the analysis and such imprecise findings are of limited clinical use
Hypertension	76 per 1,000	71 per 1,000 (19 to 260)	RR 0.93 (0.25 to 3.42)	154 (2)	⊕⊕⊕⊕ VERY LOW ^{4 5}	Few participants included in the analysis and such imprecise findings are of limited clinical use
Correction of VUR: unit of analysis (children) after 3 years: surgical correction + antibiotics versus antibiotics alone	250 per 1,000	323 per 1,000 (138 to 763)	RR 1.29 (0.55 to 3.05)	55 (1)	⊕⊕⊕⊕ VERY LOW ^{6 7}	One small study at high risk of bias in all domains evaluated this outcome so our confidence in the evidence is very low
Number of UTI with microbial resistance (resistant organisms)	See comments	See comments	not estimable	-	-	Not reported

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **ESKD:** end-stage kidney disease; **IVP:** intravenous pyelography; **RR:** Risk ratio; **UTI:** Urinary tract infection; **VUR:** vesicoureteric reflux

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹ Single study with relatively small number of included participants

² One included study was at unclear risk of attrition and selection bias

³ Two studies were at unclear risk of selection bias and one had unclear allocation concealment

- ⁴ Both studies had significant risk of bias issues
⁵ Small studies with few events
⁶ Single study at high risk of bias in all domains
⁷ Very small study with few participants

Summary of findings 3. Endoscopic injection treatment versus antibiotics alone for primary vesicoureteric reflux

Endoscopic injection treatment versus antibiotics alone for primary VUR

Patient or population: primary VUR
Setting: paediatric clinics
Intervention: endoscopic injection treatment
Comparison: antibiotics alone

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with antibiotics alone	Risk with endoscopic injection treatment				
All symptomatic UTI by 1 to 2 years: endoscopic correction + antibiotics versus antibiotics alone	225 per 1,000	180 per 1,000 (83 to 393)	RR 0.80 (0.37 to 1.75)	254 (3)	⊕⊕⊕⊕ LOW ^{1 2}	Included studies had risk of bias issues and inconsistency between results limited our confidence in the evidence
Febrile UTI by 1 to 2 years: endoscopic correction + antibiotics versus antibiotics alone	209 per 1,000	155 per 1,000 (65 to 373)	RR 0.74 (0.31 to 1.78)	254 (3)	⊕⊕⊕⊕ LOW ^{1 2}	Included studies had risk of bias issues and inconsistency between results limited our confidence in the evidence
Repeat positive urine culture by 1 year	0 per 1,000	0 per 1,000 (0 to 0)	RR 7.15 (0.42 to 121.04)	60 (1)	⊕⊕⊕⊕ VERY LOW ^{1 3}	Single small study with no events and risk of bias issues reduces our confidence in the results
Total new and progressive Renal parenchymal abnormalities on DMSA scan at 1 to 2 years: endoscopic correction + antibiotics versus antibiotics alone (unit of analysis: children)	85 per 1,000	102 per 1,000 (31 to 336)	RR 1.20 (0.36 to 3.96)	208 (2)	⊕⊕⊕⊕ LOW ^{1 4}	Included studies had risk of bias issues and there was imprecision in the results reducing our confidence in the evidence
Resolution of VUR (grade 0) for endoscopic correction + anti-	156 per 1,000	418 per 1,000 (244 to 720)	RR 2.69 (1.57 to 4.63)	164 (2)	⊕⊕⊕⊕ LOW ^{1 3}	Included studies had risk of bias issues and the small numbers of included par-

otics versus antibiotics alone (unit of analysis: children)						participants resulted in imprecision reducing our confidence in the evidence
Microbial resistance to the prophylactic drug	800 per 1,000	504 per 1,000 (272 to 920)	RR 0.63 (0.34 to 1.15)	24 (1)	⊕⊕⊕⊕ VERY LOW ^{1 5}	Single study reported this outcome with a small of urinary tract infections seriously limiting our confidence in this outcome

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **DMSA:** 99m-technetium dimercaptosuccinic acid; **RR:** Risk ratio; **UTI:** Urinary tract infection; **VUR:** vesicoureteric reflux

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

- 1 Significant risk of bias issues in all studies
- 2 Inconsistent results between studies
- 3 Small numbers of participants resulting in imprecision
- 4 Small numbers of events
- 5 Small number of events and small numbers of UTI resulting in imprecision

Summary of findings 4. Macroplastique versus dextranomer/hyaluronic acid polymer (Deflux) to correct vesicoureteric reflux

Macroplastique versus Deflux to correct VUR

Patient or population: primary VUR

Setting: paediatric clinics

Intervention: Macroplastique

Comparison: Deflux

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with macroplastique	Risk with Deflux				

Number of children with febrile UTI	68 per 1,000	74 per 1,000 (30 to 180)	RR 1.09 (0.45 to 2.66)	255 (1)	⊕⊕⊕⊕ LOW ^{1 2}	Single study with few events and risk of bias issues reduces our confidence in the evidence
Number of children with afebrile UTI	105 per 1,000	177 per 1,000 (55 to 573)	RR 1.68 (0.52 to 5.44)	72 (1)	⊕⊕⊕⊕ LOW ^{1 2}	Single study with few events and risk of bias issues reduces our confidence in the evidence
Number of ureters with persistent VUR > grade 1 at 1 year	210 per 1,000	113 per 1,000 (73 to 174)	RR 0.54 (0.35 to 0.83)	472 (2)	⊕⊕⊕⊖ MODERATE ³	Two studies with risk of bias issues moderately reduces our confidence in the results
Number of kidneys with temporary renal dilatation after injection	214 per 1,000	396 per 1,000 (219 to 718)	RR 1.85 (1.02 to 3.35)	114 (1)	⊕⊕⊕⊕ LOW ^{1 2}	Single study with few events and risk of bias issues reduces our confidence in the evidence
Ureteral obstruction	10 per 1,000	15 per 1,000 (3 to 88)	RR 1.46 (0.25 to 8.66)	399 (1)	⊕⊕⊕⊕ LOW ^{1 2}	Single study with few events and risk of bias issues reduces our confidence in the evidence
Post operative flank pain	23 per 1,000	33 per 1,000 (7 to 143)	RR 1.45 (0.33 to 6.36)	255 (1)	⊕⊕⊕⊕ LOW ^{1 2}	Single study with few events and risk of bias issues reduces our confidence in the evidence
New renal parenchymal defects on DMSA scan (kidney injury) Follow up: range 1 years to 3 years	See comments	See comments	not estimable	-	-	Not reported

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **Deflux:** dextranomer/hyaluronic acid polymer; **DMSA:** 99m-technetium dimercaptosuccinic acid; **RR:** Risk ratio; **UTI:** Urinary tract infection; **VUR:** vesicoureteric reflux

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹ Single study with several risk of bias issues

² Small numbers of events resulting in imprecision

³ Two studies with several risk of bias issues

Summary of findings 5. Probiotics versus antibiotic prophylaxis for primary vesicoureteric reflux

Probiotics compared to antibiotic prophylaxis for primary VUR

Patient or population: primary VUR

Setting: paediatric clinics

Intervention: probiotics

Comparison: antibiotic prophylaxis

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with antibiotic prophylaxis	Risk with probiotics				
All symptomatic UTI by 1 year	315 per 1,000	258 per 1,000 (176 to 381)	RR 0.82 (0.56 to 1.21)	248 (2)	⊕⊕⊕⊕ LOW ^{1 2}	Both studies had significant risk of bias issues and also small numbers of included participants limiting our confidence in the results
Febrile UTI by 1 year	183 per 1,000	150 per 1,000 (68 to 335)	RR 0.82 (0.37 to 1.83)	120 (1)	⊕⊕⊕⊕ LOW ^{2 3}	Both studies had significant risk of bias issues and also small numbers of included participants limiting our confidence in the results
E. coli resistance to prophylactic drug	1,000 per 1,000	380 per 1,000 (210 to 690)	RR 0.38 (0.21 to 0.69)	38 (2)	⊕⊕⊕⊕ LOW ^{1 2}	The single study had significant risk of bias issues and also small numbers of included participants limiting our confidence in the results
Renal parenchymal defects on DMSA scan - Patient data at 1 year	364 per 1,000	251 per 1,000 (156 to 404)	RR 0.69 (0.43 to 1.11)	152 (2)	⊕⊕⊕⊕ LOW ^{1 2}	Both studies had significant risk of bias issues and also small numbers of included participants limiting our confidence in the results
Resolution of VUR	See comments	See comments	not estimable	-	-	Not reported
Adverse events	See comments	See comments	not estimable	-	-	Not reported

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio; **UTI:** Urinary tract infection; **VUR:** vesicoureteric reflux

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

- ¹ Two small studies with significant design issues
- ² Small numbers of included participants
- ³ One small study with significant design issues

BACKGROUND

Description of the condition

Primary vesicoureteric reflux (VUR) is thought to be a maturational abnormality of the vesicoureteric junction, which results in retrograde passing of urine up the ureter during voiding. Although the exact prevalence is unknown, about a third of children investigated after a urinary tract infection (UTI), show evidence of VUR (Smellie 1994). UTI is common, affecting 5% to 10% of all children (Hellstrom 1991), with about 20% of them likely to suffer a recurrence (Smellie 1994, Panaretto 1999). VUR is thought to predispose children to UTI and renal involvement during a UTI can potentially cause permanent renal damage, possibly in up to 15% of patients (Montini 2007b). Retrospective analyses of selected individuals with renal scarring, have reported hypertension and chronic kidney disease (CKD) in approximately 20% and 10% respectively (Martinell 1996; Smellie 1998). However data from a prospective cohort study indicated that these adverse outcomes occur considerably less frequently (Wennerstrom 2000a; Wennerstrom 2000b).

As a result of the hypothesized causal link between VUR and renal scarring, VUR screening and treatment strategies have largely been directed towards avoidance of UTI and therefore any subsequent renal damage (Belman 1995). Preventative treatments have included long-term, low-dose antibiotic, surgical correction, endoscopic injection and more recently alternative strategies such as cranberry products and probiotics. Surgical correction of VUR involves techniques such as Politano-Leadbetter, Lich-Gregoir and Cohen ureteral reimplantation and the less invasive endoscopic periureteric injection of bulking agents such as polydimethylsiloxane (Macroplastique), dextranomer/hyaluronic acid copolymer (Deflux) or glutaraldehyde cross-linked bovine collagen (GAX) (Capozza 2002; Frankenschmidt 1997; Frey 1997; Oswald 2002). Although VUR is a common problem in childhood, there has been no consensus regarding the optimal management strategy and practice varies widely.

Description of the intervention

Interventions have included long-term, low-dose antibiotic treatment, primarily cotrimoxazole, trimethoprim or nitrofurantoin, for an average of 1.5 years, surgical correction using Politano-Leadbetter, Cohen, Lich-Gregoir or Glenn Anderson methods, endoscopic injection of various agents, probiotics, cranberry product, circumcision and oxybutynin,

How the intervention might work

Maintaining a constant low-dose of antibiotic may prevent the overgrowth of pathogenic bacteria that cause a UTI. Surgery to reimplant the ureters should reduce the back flow of urine up the ureters toward the kidney and may reduce the likelihood of infection. Similarly endoscopic injection of an agent to elongate and weld the submucosal tract of the ureter could reduce backward flow of urine toward the kidneys. Cranberry products are thought to contain a substance that reduces the adhesion of bacteria to the bladder wall, reducing the likelihood of overgrowth of bacterial pathogens. Probiotics are thought to increase the number of beneficial bacteria in the body and reduce overgrowth of pathogenic organisms. Circumcision is thought to reduce the carriage of bacteria on the penis and thus reduce the likelihood that bacteria pass up the urethra causing an infection. An

anticholinergic agent, such as oxybutynin, is used to treat unstable bladder to reduce symptoms of frequency, incontinence and urgency which may increase the risk of UTI.

Why it is important to do this review

Many treatment options are available for children with VUR to prevent recurring UTI and there is considerable variability in clinical practice worldwide. Identifying and analysing the available data on prophylactic treatment efficacy will guide the clinical care of children with VUR while also identifying evidence gaps for future focus.

OBJECTIVES

The aim of this review was to evaluate the available evidence for both benefits and harms of the currently available treatment options for primary VUR: operative, non-operative or no intervention.

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials (RCTs) and quasi-RCTs (RCTs in which allocation to treatment was obtained by alternation, use of alternate medical records, date of birth or other predictable methods) which evaluated any treatment for primary VUR were included.

Types of participants

Inclusion criteria

Males and females of any age with primary VUR diagnosed by voiding cystourethrogram (VCUG), with or without UTI.

Exclusion criteria

Patients with VUR associated with posterior urethral valves, spina bifida, other urological abnormalities or kidney transplants.

Types of interventions

Treatments of VUR including surgery (open and endoscopic techniques), antibiotic prophylaxis of any duration, non-invasive techniques such as bladder training, complementary medicines, and any combination of therapies.

Types of outcome measures

Primary outcomes

- Number of patients with symptomatic UTI: defined as symptoms consistent with a UTI which may include local symptoms only e.g. dysuria or systemic symptoms such as fever, together with a positive urine culture. This outcome includes febrile UTI

Secondary outcomes

- Number of patients with UTI accompanied by fever (temperature > 38°C or > 100.4°F) (a subset of the data from the outcome symptomatic UTI)
- Number of patients with at least one repeat positive urine culture during follow-up

- Renal parenchymal abnormality: defined as new, progression from pre-existing damage, resolution, end-stage kidney disease (ESKD) and diagnosed by ultrasound, intravenous pyelography (IVP) or ^{99m}Tc-DMSA (dimercaptosuccinic acid) scintigraphy (DMSA scan)
- Number of previously unaffected subjects who developed hypertension: defined as > 140 mmHg systolic, 90 mmHg diastolic for adults and above the 95th percentiles for systolic and diastolic blood pressures in children
- Kidney function impairment: defined as a glomerular filtration rate (GFR) (measured either directly or calculated from serum creatinine (eGFR)) < 95th percentile for age, or a decrease in kidney function over the duration of the study
- Correction or spontaneous resolution of VUR: defined as the number of children and/or ureters without VUR on follow up VCUG
- Microbial resistance: obstruction following correction of VUR, death or serious injury resulting from the anaesthetic, wound infection, fever, adverse effects of medication including urticaria and gastro-intestinal reaction.
- Any reported adverse events (drug or surgery-related)

Search methods for identification of studies

Electronic searches

We searched the [Cochrane Kidney and Transplant Specialised Register](#) up to 3 May 2018 through contact with the Information Specialist using search terms relevant to this review. The Cochrane Kidney and Transplant Specialised Register contains studies identified from several sources.

1. Monthly searches of the Cochrane Central Register of Controlled Trials (CENTRAL)
2. Weekly searches of MEDLINE OVID SP
3. Handsearching of kidney-related journals and the proceedings of major kidney conferences
4. Searching of the current year of EMBASE OVID SP
5. Weekly current awareness alerts for selected kidney and transplant journals
6. Searches of the International Clinical Trials Register (ICTRP) Search Portal & ClinicalTrials.gov.

Studies contained in the Specialised Register are identified through search strategies for CENTRAL, MEDLINE, and EMBASE based on the scope of Cochrane Kidney and Transplant. Details of these strategies, as well as a list of handsearched journals, conference proceedings and current awareness alerts, are available in the Specialised Register section of information about [Cochrane Kidney and Transplant](#).

See [Appendix 1](#) for search terms used in strategies for this review.

Searching other resources

1. Reference lists of review articles, relevant studies and clinical practice guidelines.
2. Letters seeking information about unpublished or incomplete studies to investigators known to be involved in previous studies.

Data collection and analysis

Selection of studies

For the first three versions of this review ([Hodson 2007](#); [Nagler 2011](#); [Wheeler 2004](#)) titles and abstracts obtained from the above searches were screened for selection independently by at least two authors. In all cases an overly inclusive selection was preferred to avoid losing relevant studies and to ensure additional studies could be identified from the reference lists. Where suitability was uncertain or no abstract available, the full article was obtained and screened by the same authors. Any disagreements were resolved by discussion with a third author. Authors were contacted to obtain raw or missing data where necessary. For the 2018 update a single author reviewed the titles.

Data extraction and management

For the 2004, 2007 and 2011 versions of this review data extraction was conducted independently by at least two authors, using a standardised data extraction form. All studies, reported in a non-English journal, were translated prior to assessment. Any further information required from the original author was requested by written correspondence and any relevant information obtained in this manner was included in the review. Any disagreements were resolved by discussion with a third author. For the 2018 update, two authors extracted five studies and reviewed agreement; 84 of 90 fields were identical across authors so a single author extracted data from the remaining nine studies. Foreign language articles were translated using [Google Translate](#).

Assessment of risk of bias in included studies

The following items were independently assessed by two authors using the risk of bias assessment tool ([Higgins 2011](#)) (see [Appendix 2](#)). In 2018 two authors extracted data from five new studies and compared agreement. A consistent difference between authors was seen and corrected in the remaining studies that were extracted by a single author. Since all three authors are also authors on an included study ([PRIVENT 2009](#)), an independent researcher extracted risk of bias data for that study.

- Was there adequate sequence generation?
- Was allocation adequately concealed?
- Was knowledge of the allocated interventions adequately prevented during the study?
- Were incomplete outcome data adequately addressed?
- Are reports of the study free of suggestion of selective outcome reporting?
- Was the study apparently free of other problems that could put it at a risk of bias?

Measures of treatment effect

Dichotomous outcomes

For dichotomous outcomes, the risk ratio (RR) and corresponding 95% confidence interval (CI) were chosen to describe the treatment effects and the precision of their point estimates. Number needed to treat (NNT) estimates were calculated to compare the benefits and harms of each active treatment.

Continuous outcomes

Where continuous scales of measurement were used to assess effects of treatment such as blood pressure and kidney function measured by eGFR, the mean difference (MD) and 95% CI was used. Where summary statistics were missing, they were derived from accompanying P values.

Planned treatment comparisons

- Antibiotics versus surgery or endoscopic treatment
- Antibiotics versus placebo or no treatment
- One antibiotic treatment versus another
- Surgical or endoscopic correction with no other treatment
- Any combinations of any active treatment

Unit of analysis issues

The unit of analysis for the primary and most secondary outcomes except resolution of reflux, was children only. For resolution of reflux, we included data for ureters as the unit of analysis and for microbial resistance, rates were expressed per UTI.

Dealing with missing data

Email contact was made to authors where there was missing data or methodology details. Responses were infrequent but Professor Jean Smellie provided additional details.

Assessment of heterogeneity

Heterogeneity between studies was analysed using the Cochran's Q statistic with the threshold for statistical significance set at $\alpha = 0.1$ (Lau 1997). It was also tested by means of the I^2 test, reflecting the percentage of total variation across studies that could be ascribed to heterogeneity (Higgins 2003). Due to an insufficient number of studies within each comparison group, formal evaluation of the different sources of heterogeneity was not possible.

Assessment of reporting biases

Publication bias was to be assessed using a funnel plot; there were insufficient studies to do so.

Data synthesis

A random effects model was used, with subsequent testing for robustness of the analysis by applying the fixed effect model.

Subgroup analysis and investigation of heterogeneity

Where there is sufficient data univariate analysis was to be used to explore whether the duration of antibiotic treatment, type of antibiotic, type of surgery and material used for endoscopic treatment have an effect on treatment efficacy. Heterogeneity was analysed using the Q statistic with a threshold for the P value of < 0.1 and the I^2 test (Higgins 2003)

Sensitivity analysis

To determine the effect of study quality on the primary outcome's pooled summary measure, sensitivity and subgroup analysis was to be performed to examine the influence of design features on results but small numbers of studies and participants prevented this study.

'Summary of findings' tables

We presented the main results of the review in 'Summary of findings' tables. These tables present key information concerning the quality of the evidence, the magnitude of the effects of the interventions examined, and the sum of the available data for the main outcomes (Schünemann 2011a). The 'Summary of findings' tables also include an overall grading of the evidence related to each of the main outcomes using the GRADE (Grades of Recommendation, Assessment, Development and Evaluation) approach (GRADE 2008). The GRADE approach defines the quality of a body of evidence as the extent to which one can be confident that an estimate of effect or association is close to the true quantity of specific interest. The quality of a body of evidence involves consideration of within-trial risk of bias (methodological quality), directness of evidence, heterogeneity, precision of effect estimates and risk of publication bias (Schünemann 2011b). We presented the following outcomes in the 'Summary of findings' tables.

- UTI
 - Symptomatic UTI
 - Febrile UTI
 - Asymptomatic (repeat positive urine culture; afebrile UTI)
- Renal parenchymal abnormality
- Spontaneous resolution or surgical correction of VUR; persistent VUR
- Adverse events
 - ESKD
 - Hypertension
 - Drug related (e.g. vomiting, rash, thrush)
 - Surgery related (e.g. ureteral obstruction, post-operative pain)
- Microbial resistance

RESULTS

Description of studies

Results of the search

The first version of this review (Wheeler 2004) included 10 studies (BIRSG 1983; Capozza 2002; Frey 1997; Holland 1982; IRS USA 1992; IRS EUR 1981-2003; Morris 1991; Oswald 2002; Reddy 1997; Smellie 2001). The International Reflux Study was reported in a European (IRS EUR 1981-2003) and an American arm (IRS USA 1992) and these reports were treated as two separate studies. Two studies were published as abstracts in conference proceedings only (Morris 1991; Reddy 1997).

The second version of this review (Hodson 2007), searched literature up to June 2006 and identified one new study (Garin 2006) along with two additional reports of the European arm of the International Reflux Study (IRS EUR 1981-2003).

A third update of this review (Nagler 2011) searched the literature up to August 2010 and included an additional nine studies (Craig 2002; Dite 2007; Lee 2007a; Montini 2008; Pennesi 2008; PRIVENT 2009; Roussey-Kesler 2008; Scott 1968; Swedish Reflux Trial 2010). Craig 2002 was published only as an abstract and Dite 2007 was originally published in Czech and was translated before assessment. We also found two papers that belonged to previously

included studies ([IRS EUR 1981-2003](#); [IRS USA 1992](#)). One ongoing study was identified ([RIVUR 2008](#)).

The current update searched literature up to May 2018 and adds a further 14 studies ([Ahn 2003](#); [Dotis 2014](#); [Espino Hernandez 2012](#);

[Garcia-Aparicio 2013](#); [Gucuk 2013](#); [Habeb Abid 2015](#); [Hari 2015](#); [Kojima 2012](#); [Lee 2015](#); [Moore 2014](#); [Nordenstrom 2016](#); [RIVUR 2008](#); [Takvani 2015](#)). [Figure 1](#) details the study selection process and results.

Figure 1. Study flow diagram.

Included studies

Thirty two of 34 studies included two study arms and two studies ([Swedish Reflux Trial 2010](#); [Reddy 1997](#)) involved three comparisons.

Antibiotic prophylaxis versus no treatment/placebo

In eight studies (796 children) antibiotic treatment was compared with no treatment ([Espino Hernandez 2012](#); [Garin 2006](#); [Montini 2008](#); [Pennesi 2008](#); [Reddy 1997](#); [Roussey-Kesler 2008](#); [Swedish Reflux Trial 2010](#); [Takvani 2015](#)) and with placebo in four studies involving 984 children ([Craig 2002](#); [Hari 2015](#); [PRIVENT 2009](#); [RIVUR 2008](#)). In [Reddy 1997](#) no treatment was compared with two antibiotic prophylaxis regimens (daily or intermittent antibiotic administration). Nine of the 12 studies reported the outcome of repeat symptomatic UTI. In eight of the 12 studies, participants were recruited after at least one symptomatic UTI ([Garin 2006](#); [Hari 2015](#); [Montini 2008](#); [Pennesi 2008](#); [PRIVENT 2009](#); [RIVUR 2008](#); [Roussey-Kesler 2008](#); [Swedish Reflux Trial 2010](#)), in two studies children had not experienced a UTI but had VUR detected after antenatal ultrasonography ([Craig 2002](#); [Takvani 2015](#)), and in two studies, details on prior UTI were not reported ([Espino Hernandez](#)

[2012](#); [Reddy 1997](#)). Two studies included children with the entire spectrum of VUR grade ([Craig 2002](#); [PRIVENT 2009](#)); two studies included children with grades I to IV ([Hari 2015](#); [RIVUR 2008](#)); seven studies included various lower grades of VUR ([Espino Hernandez 2012](#); [Garin 2006](#); [Montini 2008](#); [Pennesi 2008](#); [Roussey-Kesler 2008](#); [Swedish Reflux Trial 2010](#); [Takvani 2015](#)); and one study ([Reddy 1997](#)) did not report VUR grade details. Five studies did not report the gender of participating children ([Craig 2002](#); [Espino Hernandez 2012](#); [Hari 2015](#); [Reddy 1997](#); [Takvani 2015](#)) while in the remaining seven studies girls (990) outnumbered boys (331), with an extreme imbalance in [RIVUR 2008](#) (558 girls to 49 boys). In three studies ([Garin 2006](#); [Montini 2008](#); [PRIVENT 2009](#)), the investigators included both children with and without VUR but for this review we included only the data from children with VUR. Antibiotic treatment duration ranged from one to three years, with the most common practice being one year (5 studies). Two studies did not report antibiotic duration ([Espino Hernandez 2012](#); [Takvani 2015](#))

Adherence was addressed in six studies ([Hari 2015](#); [Montini 2008](#); [Pennesi 2008](#); [PRIVENT 2009](#); [RIVUR 2008](#); [Swedish Reflux Trial 2010](#)).

- [Hari 2015](#) measured adherence by weighing returned medication bottles and compliance was reported as not different between the two treatment groups
- [Pennesi 2008](#) tested the urine samples of children that developed a febrile UTI for presence of the prophylactic drug. He found all patients were adherent.
- [Montini 2008](#) found 71% of participants were adherent when assessed by measuring antimicrobial activity in a sample of screened urine samples. The reported adherence rate was 86% according to the visual analogue questionnaire.
- [PRIVENT 2009](#) tested adherence by weighing the bottles at each clinic visit as well as by direct questioning of the parents. The authors reported no difference in the frequency of measured non-adherence between the groups.
- [Swedish Reflux Trial 2010](#) report stated that adherence to the antibiotic prophylaxis had been assessed by asking questions during every follow-up visit, but no findings were provided.
- [RIVUR 2008](#) measured adherence by collecting used and unused medication bottles, by direct questioning at clinic visits and patient diaries. 467 children were reported as being 75% compliant with medication while 91 children in the antibiotic group and 76 children in the placebo group stopped taking the medication. No significant difference between adherence rates across the two treatment groups was reported.

Surgical reimplantation of the ureters

In seven studies (815 children) the effectiveness of low-dose antibiotic prophylaxis, given for one to five years, was compared with ureteric reimplantation by open surgery, using different surgical methods ([BIRSG 1983](#); [Holland 1982](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Morris 1991](#); [Smellie 2001](#); [Scott 1968](#)). Only one study ([IRS EUR 1981-2003](#)) reported the outcome of repeat symptomatic UTI. Five studies recruited children with at least one UTI while [Morris 1991](#) did not report this detail. Three studies involved children with VUR of grades III and IV ([IRS EUR 1981-2003](#); [IRS USA 1992](#); [Morris 1991](#)). One study ([Smellie 2001](#)) included children with grades III to V VUR, [Holland 1982](#) included children with grades III to IV and [BIRSG 1983](#) included children with grades II (with scarring) and III to V. [Scott 1968](#) did not report VUR grades of participating children. [IRS EUR 1981-2003](#) and [Holland 1982](#) did not report gender. A total of 368 girls and 144 boys were included in the remaining studies.

[Kojima 2012](#) compared two different surgical methods, reimplantation with and without advancement, for the outcomes of positive urine culture and correction of reflux. Children in the study had VUR grades I to V with 13 girls and 17 boys.

Endoscopic injection treatment

Four studies compared endoscopic subureteric injection of Deflux/Dexel ([Capozza 2002](#); [Dite 2007](#); [Nordenstrom 2016](#), [Swedish Reflux Trial 2010](#)) with prophylactic antibiotic. Three of these ([Dite 2007](#); [Nordenstrom 2016](#); [Swedish Reflux Trial 2010](#)) reported the outcome of repeat symptomatic UTI. Four studies ([Frey 1997](#); [Habeeb Abid 2015](#); [Moore 2014](#); [Oswald 2002](#)) compared

different injection products, two of which were Deflux with Macroplastique ([Oswald 2002](#); [Moore 2014](#)), and these studies reported on symptomatic UTI allowing for pooling of data. One study compared GAX 35 with GAX 65 ([Frey 1997](#)) and one study compared dextranomer/hyaluronic acid polymer with polyacrylate polyalcohol copolymer (Vantris) ([Habeeb Abid 2015](#)). One study compared endoscopic injection with surgical reimplantation of ureters ([Garcia-Aparicio 2013](#)). [Swedish Reflux Trial 2010](#) included a third arm of no treatment, within their study allowing a comparison between endoscopic injection and no treatment. Two of the four studies comparing endoscopic injection with antibiotics recruited the majority of their participating children after a UTI ([Nordenstrom 2016](#); [Swedish Reflux Trial 2010](#)), none of the four studies comparing different endoscopic injection polymers detailed prior UTI history of the participating children and the study of endoscopic injection versus surgical reimplantation also did not report prior UTI history ([Garcia-Aparicio 2013](#)). VUR grade was reported by eight of the nine studies, with only [Habeeb Abid 2015](#) not reporting this detail. VUR grade ranged widely from all grades to a focus on high grades; III to V. Gender was reported in eight of the nine studies and showed an over representation of girls (445) compared to boys (225).

Probiotics compared with low dose antibiotic treatment

Two studies involving 248 children compared probiotics (*Lactobacillus acidophilus*) with antibiotic treatment, cotrimoxazole ([Lee 2007a](#); [Lee 2015](#)) and one study of 85 children compared a combination of two strains of probiotic (*Lactobacillus acidophilus* and *Bifidobacterium lactis*) plus antibiotics (nitrofurantoin) with antibiotics alone ([Mohseni 2013](#)). The three studies reported VUR grade and covered the spectrum of grades from I to V while two of the three studies reported a prior history of UTI ([Lee 2015](#); [Mohseni 2013](#)). There were 133 girls and 200 boys who participated.

Other interventions

A study of 76 children (53 girls and 23 boys), with grades I and II VUR and recruited after a previous UTI, compared cranberry capsules with no treatment ([Dotis 2014](#)). One study of 91 boys with one prior UTI and VUR grades I to III, compared antibiotics alone with antibiotics plus circumcision ([Gucuk 2013](#)). A study of 149 children, that did not report on grade of VUR, gender or UTI history, compared oxybutynin with placebo for the single outcome VUR resolution ([Ahn 2003](#)).

Excluded studies

We excluded 10 studies. See [Characteristics of excluded studies](#).

Risk of bias in included studies

Four studies were considered at low risk of bias for all domains within the risk of bias tool ([Craig 2002](#); [Hari 2015](#); [PRIVENT 2009](#); [RIVUR 2008](#)) and all compared low dose antibiotics with placebo. The remaining 30 studies ranged in quality with many study quality fields having a high risk of bias or unclear risk of bias.

An overview is provided in [Figure 2](#) and [Figure 3](#).

Figure 2. Methodological quality graph: review authors' judgements about each methodological quality item presented as percentages across all included studies.

Figure 3. Methodological quality summary: review authors' judgements about each methodological quality item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Ahn 2003	?	?	?	?	?	-	?
BIRSG 1983	?	?	-	+	-	-	?
Capozza 2002	+	+	-	-	-	?	+
Craig 2002	+	+	+	+	+	+	+
Dite 2007	?	?	-	?	+	?	?
Dotis 2014	?	?	?	?	?	-	?
Espino Hernandez 2012	?	?	?	?	?	-	?
Frey 1997	?	+	+	+	?	-	?
Garcia-Aparicio 2013	+	?	?	?	-	-	?
Garin 2006	?	?	-	?	-	-	?
Gucuk 2013	?	?	?	?	?	-	?
Habeeb Abid 2015	?	?	?	?	-	-	?
Hari 2015	+	+	+	+	+	+	+
Holland 1982	?	?	-	?	+	?	-
IRS EUR 1981-2003	+	+	-	+	+	+	+
IRS USA 1992	+	+	-	+	?	?	?
Kojima 2012	?	?	?	?	-	?	?
Lee 2007a	-	-	-	+	+	+	+
Lee 2015	+	?	-	-	+	+	?
Mohseni 2013	+	?	?	?	+	+	-

Figure 3. (Continued)

Mohseni 2013	+	?	?	?	+	+	-
Montini 2008	+	+	-	-	-	+	?
Moore 2014	?	-	?	?	-	-	?
Morris 1991	?	?	-	?	-	-	?
Nordenstrom 2016	+	?	?	?	-	+	?
Oswald 2002	?	?	-	?	?	?	?
Pennesi 2008	+	+	-	+	+	+	?
PRIVENT 2009	+	+	+	+	+	+	+
Reddy 1997	?	?	?	?	+	-	?
RIVUR 2008	+	+	+	+	+	+	+
Roussey-Kesler 2008	+	+	?	?	+	+	?
Scott 1968	-	-	-	-	-	-	-
Smellie 2001	+	?	-	+	?	?	?
Swedish Reflux Trial 2010	+	+	-	+	+	?	-
Takvani 2015	?	?	?	?	?	-	?

Allocation

Sequence generation

The method of sequence generation was at a low risk of bias in 16 studies (Capozza 2002; Craig 2002; Garcia-Aparicio 2013; Hari 2015; IRS EUR 1981-2003; IRS USA 1992; Lee 2015; Mohseni 2013; Montini 2008; Nordenstrom 2016; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Roussey-Kesler 2008; Smellie 2001; Swedish Reflux Trial 2010), at high risk of bias in two studies (Lee 2007a; Scott 1968), and unclear risk in 16 studies (Ahn 2003; BIRSG 1983; Dite 2007; Dotis 2014; Espino Hernandez 2012; Frey 1997; Garin 2006; Gucuk 2013; Habeeb Abid 2015; Holland 1982; Kojima 2012; Moore 2014; Morris 1991; Oswald 2002; Reddy 1997; Takvani 2015).

Allocation concealment

Allocation concealment was at a low risk of bias in 12 studies (Capozza 2002; Craig 2002; Frey 1997; Hari 2015; IRS EUR 1981-2003; IRS USA 1992; Montini 2008; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Roussey-Kesler 2008; Swedish Reflux Trial 2010), high risk of bias in three studies (Lee 2007a; Moore 2014; Scott 1968), and unclear risk in 19 studies (Ahn 2003; BIRSG 1983; Dite 2007; Dotis 2014; Espino Hernandez 2012; Garcia-Aparicio 2013; Garin 2006; Gucuk 2013; Habeeb Abid 2015; Holland 1982; Kojima 2012; Lee 2015; Mohseni 2013; Morris 1991; Nordenstrom 2016; Oswald 2002; Reddy 1997; Smellie 2001; Takvani 2015).

Blinding

While it is understood blinding clinicians and participants/carers to a surgical procedure is not easily feasible, blinding of outcome assessment is feasible.

Performance bias

Risk of performance bias was low in five studies (Craig 2002; Frey 1997; Hari 2015; PRIVENT 2009; RIVUR 2008), high in 16 studies (BIRSG 1983; Capozza 2002; Dite 2007; Garin 2006; Holland 1982; IRS EUR 1981-2003; IRS USA 1992; Lee 2007a; Lee 2015; Montini 2008; Morris 1991; Oswald 2002; Pennesi 2008; Scott 1968; Smellie 2001; Swedish Reflux Trial 2010), and unclear in 13 studies (Ahn 2003; Dotis 2014; Espino Hernandez 2012; Garcia-Aparicio 2013; Gucuk 2013; Habeeb Abid 2015; Kojima 2012; Mohseni 2013; Moore 2014; Nordenstrom 2016; Reddy 1997; Roussey-Kesler 2008; Takvani 2015).

Detection bias

Risk of detection bias was low in 12 studies (BIRSG 1983; Craig 2002; Frey 1997; Hari 2015; IRS EUR 1981-2003; IRS USA 1992; Lee 2007a; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Smellie 2001; Swedish Reflux Trial 2010); high in four studies (Capozza 2002; Lee 2015; Montini 2008; Scott 1968), and unclear in 18 studies (Ahn 2003; Dite 2007; Dotis 2014; Espino Hernandez 2012; Garcia-Aparicio 2013; Garin 2006; Gucuk 2013; Habeeb Abid 2015; Holland 1982; Kojima 2012; Mohseni 2013; Moore 2014; Morris 1991; Nordenstrom 2016; Oswald 2002; Reddy 1997; Roussey-Kesler 2008; Takvani 2015).

Incomplete outcome data

Fourteen studies accounted for all study participants in the analysis and report, giving them a low risk of bias (Craig 2002; Dite 2007; Hari 2015; Holland 1982; IRS EUR 1981-2003; Lee 2007a; Lee 2015; Mohseni 2013; Pennesi 2008; PRIVENT 2009; Reddy 1997; RIVUR 2008; Roussey-Kesler 2008; Swedish Reflux Trial 2010), 11 studies were classified as high risk of bias (BIRSG 1983; Capozza 2002; Garcia-Aparicio 2013; Garin 2006; Habeeb Abid 2015; Kojima 2012; Montini 2008; Moore 2014; Morris 1991; Nordenstrom 2016; Scott 1968). In nine studies it was unclear how many participants were included in the analyses (Ahn 2003; Dotis 2014; Espino Hernandez 2012; Frey 1997; Gucuk 2013; IRS USA 1992; Oswald 2002; Smellie 2001; Takvani 2015), usually because denominators were not reported.

Selective reporting

Twelve studies reported the most clinically appropriate outcomes and were considered at low risk of bias (Craig 2002; Hari 2015; IRS EUR 1981-2003; Lee 2007a; Lee 2015; Mohseni 2013; Montini 2008; Nordenstrom 2016; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Roussey-Kesler 2008). Thirteen studies reported only surrogate or not the most clinically relevant outcomes such as resolution of VUR and positive urine culture, and were considered at high risk of bias (Ahn 2003; BIRSG 1983; Dotis 2014; Espino Hernandez 2012; Frey 1997; Garcia-Aparicio 2013; Garin 2006; Gucuk 2013; Habeeb Abid 2015; Moore 2014; Morris 1991; Scott 1968; Takvani 2015). In nine studies details on outcomes were ambiguous or poorly detailed meaning there was an unclear risk of reporting bias (Capozza 2002; Dite 2007; Holland 1982; IRS USA 1992; Kojima 2012; Oswald 2002; Smellie 2001; Swedish Reflux Trial 2010).

Other potential sources of bias

For many studies it was difficult to discern who the children were and how many were reviewed for possible inclusion in the study protocol, thereby largely limiting the ability to evaluate the extent of selection bias. Seven studies were considered at low risk of bias from other sources because patients and methods were well detailed and comprehensive (Capozza 2002; Craig 2002; Hari 2015; IRS EUR 1981-2003; Lee 2007a; PRIVENT 2009; RIVUR 2008). Four studies were considered at high risk of bias from other sources, primarily from poorly reported details on how participants were selected (Holland 1982; Mohseni 2013; Scott 1968; Swedish Reflux Trial 2010), while in 23 studies it was difficult to determine other potential biases (Ahn 2003; BIRSG 1983; Dite 2007; Dotis 2014; Espino Hernandez 2012; Frey 1997; Garcia-Aparicio 2013; Garin 2006; Gucuk 2013; Habeeb Abid 2015; IRS USA 1992; Kojima 2012; Lee 2015; Montini 2008; Moore 2014; Morris 1991; Nordenstrom 2016; Oswald 2002; Pennesi 2008; Reddy 1997; Roussey-Kesler 2008; Smellie 2001; Takvani 2015).

Effects of interventions

See: [Summary of findings for the main comparison](#) Antibiotic prophylaxis versus surveillance or no treatment for primary vesicoureteric reflux; [Summary of findings 2](#) Surgical reimplantation of ureters plus antibiotics versus antibiotics alone for primary vesicoureteric reflux; [Summary of findings 3](#) Endoscopic injection treatment versus antibiotics alone for primary vesicoureteric reflux; [Summary of findings 4](#) Macroplastique versus dextranomer/hyaluronic acid polymer (Deflux) to correct

vesicoureteric reflux; [Summary of findings 5](#) Probiotics versus antibiotic prophylaxis for primary vesicoureteric reflux

Antibiotic prophylaxis versus surveillance or no treatment

The data were analysed using a random and a fixed effects model, without there being an appreciable difference between the summary estimates. Results are presented using the random effects model. See [Summary of findings for the main comparison](#).

Urinary tract infection

Symptomatic and febrile UTI

Of the 12 studies that compared antibiotic prophylaxis with placebo or no treatment, nine reported repeat symptomatic UTI as an outcome and allowed distinction of febrile UTI as a separate entity. One and two year incidence of symptomatic UTI varied from 11% to 36% in the group treated with antibiotics and 7% to 58% in the placebo/no treatment groups. The point estimate for overall effect suggested antibiotic prophylaxis may slightly reduce the risk of repeat symptomatic UTI but the wide confidence interval, includes the point of no difference between the treatment groups indicating that antibiotic prophylaxis compared with placebo/no treatment may make little or no difference to the risk of repeat symptomatic UTI ([Analysis 1.1.1](#) (9 studies, 1667 children): RR 0.77, 95% CI 0.54 to 1.09; $I^2 = 59\%$; low certainty evidence). Individual studies varied widely in results creating large inconsistencies and study designs were often suboptimal leading to a conclusion of low quality evidence and considerable uncertainty in findings. Similarly for the outcome of febrile UTI by one to two years of follow-up, antibiotic prophylaxis compared with placebo/no treatment may make little or no difference to the outcome of febrile UTI ([Analysis 1.1.2](#) (9 studies, 1667 children): RR 0.83, 95% CI 0.56 to 1.21 $I^2 = 51\%$; low certainty evidence). The individual studies were inconsistent and contained considerable bias leading to the assessment of low certainty evidence.

Two studies (Pennesi 2008; Swedish Reflux Trial 2010) reported outcome data within study arms and separate for girls and boys ([Analysis 1.14](#)) and point estimates suggested that in boys, risk of recurrence of febrile UTI was similar whether on antibiotics or with no treatment (RR 1.26, 95% CI 0.53 to 2.99; $I^2 = 0\%$). In girls the point estimate suggested a lower risk for recurrence for those on antibiotic treatment than with no treatment (RR 0.63, 95% CI 0.17 to 2.33; $I^2 = 86\%$). However both estimates included the point of no difference because sample size was small and events not very common, leading to a lack of precision.

For these outcomes, there was a large amount of unexplained heterogeneity among studies. We could not identify systematic differences in allocation mechanism, blinding or participant characteristics that explained the heterogeneity. Three studies were rated as low risk of bias across all quality assessment fields (Hari 2015; PRIVENT 2009; RIVUR 2008) and while two studies were reasonably similar in their point estimates (PRIVENT 2009: RR 0.66 95% CI 0.35 to 1.24; RIVUR 2008: RR 0.55 95% CI 0.38 to 0.78), Hari 2015 was very different (RR 3.26, 95% CI 0.96 to 11.10). Therefore we concluded that in this subgroup of studies, antibiotic prophylaxis compared with placebo probably leads to little or no difference in the risk of repeat UTI (moderate certainty evidence). An explanation might be that of a chance affect in a small sample size; Hari 2015 included only 93 children while RIVUR 2008 and PRIVENT 2009

included 607 and 243 children with VUR, respectively. (Analysis 1.1.3).

Repeat positive urine culture

In seven studies involving 729 children, the outcome of repeat positive urine culture was reported for urine samples taken in asymptomatic subjects (Craig 2002; Garin 2006; Hari 2015; Montini 2008; Pennesi 2008; Reddy 1997; Roussey-Kesler 2008). The analysis showed a slightly reduced risk of repeat positive urine culture in children taking antibiotics compared to those on placebo/no treatment (Analysis 1.1.4 (7 studies, 729 participants) RR 0.80, 95% CI 0.63 to 1.02; $I^2 = 0\%$). However the confidence interval include the point of no difference so that antibiotic prophylaxis compared with no antibiotics may be little or no difference to the outcome of positive urine culture (low certainty evidence). In addition to comparing a daily antibiotic regimen with surveillance, Reddy 1997 introduced an extra treatment arm to compare intermittent treatment three times/week, with no treatment. In the group treated with intermittent antibiotics 2/14 participants (14%) had a repeat positive urine culture and in the no-treatment group 5/16 patients experienced a repeat positive urine culture (31%) (Analysis 1.1.5 (1 study, 30 children): RR 0.46, 95% CI 0.10 to 2.00).

Renal parenchymal abnormality

In eight studies, the acquisition of new renal abnormalities was evaluated by comparing DMSA scans, taken both at study commencement and completion of follow-up (Craig 2002; Espino Hernandez 2012; Hari 2015; Montini 2008; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Swedish Reflux Trial 2010). Three studies had no events in either group and hence did not contribute data to the analyses (Craig 2002; Espino Hernandez 2012; Pennesi 2008). Overall antibiotic prophylaxis compared with placebo/no treatment may make little or no difference to the risk of new renal abnormalities (Analysis 1.4.1 (8 studies, 1503 children): RR 0.73, 95% CI 0.33 to 1.61; $I^2 = 24\%$), the risk of deterioration of existing renal abnormalities (Analysis 1.4.2 (3 studies, 446 children): RR 0.68, 95% CI 0.27 to 1.73; $I^2 = 0\%$), or total renal abnormalities (Analysis 1.4.3 (7 studies, 1288 participants): RR 0.83, 95% CI 0.42 to 1.62; $I^2 = 42\%$; low certainty evidence).

The study comparing intermittent with continuous prophylaxis (Reddy 1997) showed a very imprecise estimate of the risk of renal parenchymal injury between the two treatments with the effect ranging from a possible 98% reduction to over 8-fold greater risk (Analysis 1.4.5 (1 study, 30 children): RR 0.38, 95% CI 0.02 to 8.59).

Resolution of VUR

Three studies (Pennesi 2008; Reddy 1997; Swedish Reflux Trial 2010) reported the VUR status after a two year follow-up. Overall there was little or no difference in the number of children with persisting VUR (Analysis 1.5 (3 studies, 262 children): RR 1.46, 95% CI 0.71 to 2.99; $I^2 = 0\%$; low certainty evidence).

Adverse events

Five studies reported adverse effects and the findings were very different for each study (Garin 2006; Hari 2015; Montini 2008; PRIVENT 2009; RIVUR 2008) (Analysis 1.6). Antibiotic prophylaxis compared with placebo/no treatment may make little or no difference to the number of adverse effects reported (Analysis 1.6

(4 studies, 1056 children): RR 0.94, 95% CI 0.81 to 1.08; $I^2 = 0\%$; moderate certainty evidence).

- Garin 2006 explicitly stated 'no reported side effects associated with the use of urinary antibiotic prophylaxis'.
- Hari 2015 reported 12 and 11 adverse events in the antibiotic and placebo groups respectively
- PRIVENT 2009 reported two participants developed thrush while on antibiotics and five developed a rash while on placebo.
- Montini 2008 reported 25 minor adverse events (out of 211 participants), mainly vomiting or gastro-intestinal intolerance. These data however included patients without VUR and were not reported per treatment group.
- RIVUR 2008 reported similar frequencies of adverse events across the treatment groups with 153 children in the antibiotic and 165 children in the placebo groups experiencing one or more events.

Microbial resistance

Six studies described bacterial resistance to the prophylactic drug in subsequent symptomatic UTIs (Hari 2015; Pennesi 2008; PRIVENT 2009; RIVUR 2008; Roussey-Kesler 2008; Swedish Reflux Trial 2010). Overall the estimated risk of prophylactic drug resistance in a repeat symptomatic UTI was almost three times higher for children who received antibiotics compared to those on placebo or no treatment (Analysis 1.7.1 (6 studies, 259 urine cultures): RR 2.72, 95% CI 1.85 to 4.00; $I^2 = 24\%$; moderate certainty evidence). Assessment of these studies showed there were some reported design features and led to the conclusion that antibiotic prophylaxis compared with placebo/no treatment probably increases the risk of drug resistance. In Garin 2006, all of the seven pyelonephritis cases in children given antibiotics were caused by a resistant micro-organism but authors did not report whether the single case of pyelonephritis in the placebo group was caused by a bacteria that was sensitive or resistant to commonly used antibiotics.

Other outcomes

One small study (46 children), only published in abstract form, reported on DTPA measured GFR, and renal growth at the end of a three year follow-up period (Craig 2002). Results were only reported as means, but standard deviations could be derived from the accompanying P value. At three year follow-up there was little or no difference in GFR (Analysis 1.8 (1 study, 41 children): MD -11.00 mL/min/1.73 m², 95% CI -31.53 to 9.53) or in renal growth (Analysis 1.9 (1 study, 41 children): MD 0.04 cm, 95% CI -0.04 to 0.12).

The outcomes of ESKD and hypertension were not reported.

Surgical reimplantation of ureters plus antibiotics versus antibiotics alone

There was no appreciable difference between the summary estimates using random and fixed effects models. There were insufficient studies to explore potential effect modification using subgroup analysis or meta-regression. See Summary of findings 2.

Urinary tract infection

Symptomatic and febrile UTI

Symptomatic UTI was reported in only one surgical study with the finding of little or no difference in risk between the two treatment groups, at four to five (moderate certainty evidence), and five to 10 years. Febrile UTI was reported in both arms of the international reflux study ([IRS EUR 1981-2003](#), [IRS USA 1992](#)); at five years the risk of repeat febrile UTI was 57% lower in the surgical group ([Analysis 2.1.4](#) (2 studies, 429 participants): RR 0.43 95% CI 0.27 to 0.70; $I^2 = 0\%$). The consistency in these two studies and precision of the estimates suggested surgical correction with antibiotic prophylaxis compared with antibiotic prophylaxis probably reduces the risk of febrile UTI (moderate certainty evidence). Only the European arm reported febrile UTI at five to 10 years and after 10 years.

Repeat positive urine culture

Repeat positive urine culture was reported in three studies ([IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#)) and the risk estimate showed little or no difference between the treatment groups. ([Analysis 2.1.7](#) (3 studies, 479 participants): RR 0.99 95% CI 0.79 to 1.26; $I^2 = 0\%$; moderate certainty evidence).

Renal parenchymal abnormality

Renal parenchymal abnormalities were reported in five studies ([BIRSG 1983](#); [Holland 1982](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#)) at varying time points. The frequency of renal parenchymal abnormality (scars and renal parenchymal thinning) on IVP at study entry was 56% to 100% with little or no difference between children receiving antibiotic prophylaxis alone and those treated with surgery plus antibiotics. Surgical correction + antibiotic prophylaxis compared with antibiotic prophylaxis probably makes little or no difference to the number of children developing a new renal parenchymal abnormality, either at two years ([Analysis 2.2.1](#) (2 studies, 171 children): RR 1.06, 95% CI 0.33 to 3.42) or at four to five years ([Analysis 2.2.2](#) (4 studies, 572 children): RR 1.09, 95% CI 0.79 to 1.49; moderate certainty evidence). The risk of progression of an existing abnormality was also similar both at two years ([Analysis 2.2.3](#) (1 study, 10 children): RR 7.00, 95% CI 0.45 to 108.26) and at four to five years ([Analysis 2.2.4](#) (3 studies, 468 children): RR 0.99, 95% CI 0.69 to 1.42). When the development of a new or a progressive abnormality were considered as a single outcome at four to five years, there was little or no difference between the two groups ([Analysis 2.2.6](#) (3 studies, 468 children): RR 1.05, 95% CI 0.85 to 1.29).

The European ([IRS EUR 1981-2003](#)) and US arms ([IRS USA 1992](#)) of the International Reflux Study differentiated renal scarring and renal parenchymal thinning on IVP. Surgical correction + antibiotic prophylaxis with antibiotic prophylaxis probably makes little or no difference to the number of patients with renal scars on IVP at zero to five years ([Analysis 2.3.1](#) (2 studies, 418 children): RR 1.28, 95% CI 0.84 to 1.94), at five to 10 years ([Analysis 2.3.2](#) (1 study, 223 children): RR 1.03, 95% CI 0.07 to 16.22) or zero to 10 years in children followed for 10 years in the European arm of the International Reflux Study ([Analysis 2.3.3](#) (1 study, 223 children): RR 1.03, 95% CI 0.53 to 2.00). All risk estimates were fairly imprecise with large confidence intervals that included the point of no effect. In the [IRS EUR 1981-2003](#), renal scarring on IVP was present at entry in 49% of the 306 children originally treated and in 51% of 223 children studied by IVP at 10 years. During the first five years of

follow-up, 40 children (surgery plus antibiotic group (21); antibiotic group (19)) developed new scars. Of these, 28 were among the 223 followed radiologically at 10 years. Only two more children, one from each therapy group, developed new scars between five and 10 years.

When the data were examined according to the total number of kidneys, at two years of follow-up, Surgical correction + antibiotic prophylaxis with antibiotic prophylaxis probably makes little or no difference to the number of new abnormalities at 2 years ([Analysis 2.6.1](#) (2 studies, 235 children): RR 1.03, 95% CI 0.31 to 3.37), progressive abnormalities ([Analysis 2.6.2](#) (2 studies, 235 children): RR 1.56, 95% CI 0.24 to 10.08) or total renal parenchymal abnormalities ([Analysis 2.6.3](#) (2 studies, 235 children): RR 1.54, 95% CI 0.24 to 9.95) or at 5 years for new abnormalities ([Analysis 2.6.4](#) (2 studies, 319 children): RR 0.85, 95% CI 0.24 to 3.09), progression in abnormality ([Analysis 2.6.5](#) (2 studies, 319 children): RR 0.84, 95% CI 0.50 to 1.41) or total abnormality ([Analysis 2.6.6](#) (2 studies, 319 children): RR 0.84, 95% CI 0.53 to 1.34). All confidence intervals were wide showing imprecision in the estimate and included the point of no effect.

One study evaluated renal parenchymal abnormality with DMSA scan ([IRS EUR 1981-2003](#)); 97% of children had a scintigraphy performed at five years and 73% of children at 10 years. Parenchymal abnormalities were present in 83% of children at study entry. Relative to the antibiotics-only group, in the surgical plus antibiotic treatment group there was probably little or no increased risk of new or progressive DMSA scan abnormalities ([Analysis 2.5.2](#) (1 study, 287 children): RR 0.97, 95% CI 0.58 to 1.62) or of deterioration in DMSA scan appearance between five to 10 years ([Analysis 2.5.3](#) (1 study, 216 children): RR 0.71, 95% CI 0.31 to 1.58). All confidence intervals included the point of no effect.

Renal growth was evaluated in four studies ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#)) at two to 10 years by measurements of changes in renal length standard deviation score (SDS) (3 studies, 510 children) or renal area (1 study, 82 children) on IVP. No real differences between groups were found at any time point or in any age group. Combining data in meta-analysis was not possible because of differences in reporting.

ESKD and hypertension

The two outcomes of greatest clinical importance, ESKD and hypertension, were reported in three studies ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [Smellie 2001](#)). Six children developed ESKD and 14 developed hypertension during follow-up. It is uncertain whether surgical correction plus antibiotics compared with antibiotics alone reduced the risk for ESKD at five years ([Analysis 2.7.1](#) (2 studies, 154 children): RR 1.07, 95% CI 0.23 to 5.04; $I^2 = 0\%$; very low certainty evidence) or of hypertension overall ([Analysis 2.7.2](#) (2 studies, 154 children): RR 0.93, 95% CI 0.25 to 3.42; $I^2 = 13\%$; very low certainty evidence), or for hypertension at 10 years ([Analysis 2.7.3](#) (1 study, 252 children): RR 0.15, 95% CI 0.01 to 2.78). Studies were small and imprecise and all confidence intervals were wide and crossed the point of no effect.

GFR

Four studies ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [Morris 1991](#); [Smellie 2001](#)) reported on GFR but these were unable to be combined because of insufficiently reported point estimate and variance

data. Individually, no study reported any differences between groups. Data from [IRS EUR 1981-2003](#) showed no differences in GFR measured by the Schwartz formula at study entry ([Analysis 2.8.1](#)), at five years ([Analysis 2.8.2](#)) and at 10 years ([Analysis 2.8.3](#)).

Height

Growth was investigated in [IRS EUR 1981-2003](#). There was no difference in height SDS at study entry ([Analysis 2.9.1](#)) or at 10 years ([Analysis 2.9.2](#)).

Correction of VUR

Resolution of VUR was an outcome described in five studies ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#); [Scott 1968](#)). We could not combine individual study data for the surgical studies because of differences in reporting practices (patients and ureters), not all patients having had follow-up VCUGs and missing data. In four studies ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#)) the postoperative resolution rate at four to five years for ureters was 93% to 99%. Over a follow-up period of three to five years, 16% to 49% of patients had spontaneous resolution of VUR ([BIRSG 1983](#); [IRS EUR 1981-2003](#); [IRS USA 1992](#); [Smellie 2001](#); [Scott 1968](#)). In [IRS EUR 1981-2003](#), 130/155 children in the antibiotics-only group had persisting VUR at five years though in 50 other children VUR grade had diminished. Among 102 children undergoing voiding VCUGs at 10 years, VUR was still present in 27 children (22 with grade IV and five with grade III). In [Scott 1968](#), 6/31 had persistent VUR three years postoperatively (very low certainty evidence) and 4/31 had successful operations, but developed VUR in the opposite ureter at a later date.

Adverse events

Adverse events for either group were generally not well reported. Postoperative obstruction to the urinary tract occurred in 7% of children (10/151) in the [IRS EUR 1981-2003](#) study. The [BIRSG 1983](#) study stated that no cases of postoperative obstruction were found after five years. The authors did report six other adverse events. One boy had transient ureteral and renal pelvic dilatation on ultrasound at one month, one boy developed urine retention after endoscopic injection, and one boy aspirated during anaesthesia and required overnight observation in the intensive care unit. One girl suffered abdominal pain with pelvic dilatation and decreasing kidney split function. The authors stated this resulted from a crossing vessel at the pelviureteral junction and was not related to the intervention. Finally, in one boy a fibrous narrowing of the bulbar urethra without obstruction was detected during the first endoscopic procedure. A weakening urine stream and obstructive flow curve pattern led to repeat endoscopic investigation, which revealed deterioration of bulbar narrowing. Ultimately internal urethrotomy was done. No other study referred to obstruction. No other adverse outcomes of surgery were reported.

No study reported on whether bacterial resistance developed to the prophylactic drug in subsequent symptomatic UTIs

Surgical reimplantation type 1 versus surgical reimplantation type 2

One study of 30 children ([Kojima 2012](#)) compared surgical reimplantation-advancement technique with surgical reimplantation-no advancement technique and reported the outcomes repeat positive urine culture, persistent VUR and

perioperative complications. Little or no differences were seen in any outcomes ([Analysis 3.1](#); [Analysis 3.2](#); [Analysis 3.3](#)).

Endoscopic injection treatment with/without antibiotics versus antibiotics alone

See [Summary of findings 3](#).

Urinary tract infection

Symptomatic and febrile UTI

Three studies ([Dite 2007](#); [Nordenstrom 2016](#); [Swedish Reflux Trial 2010](#)) compared endoscopic injection with antibiotic prophylaxis and reported the outcomes of repeat symptomatic UTI and febrile UTI. Compared with antibiotics alone, endoscopic correction of VUR may lead to little or no reduction in the risk of symptomatic UTI ([Analysis 4.1.1](#) (3 studies, 254 participants): RR 0.80, 95% CI 0.37 to 1.75; $I^2 = 58\%$; low certainty evidence) and in the risk of febrile UTI ([Analysis 4.1.2](#) (3 studies, 254 participants): RR 0.74 95% CI 0.31 to 1.78; $I^2 = 58\%$; low certainty evidence). The three studies included only 254 children and may reflect chance events. Assessment of these studies showed many study design fields of unclear or high bias and they were small studies with few events and large imprecision.

Repeat positive urine culture

One study ([Capozza 2002](#)), involving 60 children, reported the outcome of repeat positive urine culture with a very imprecise estimate that cannot be used to guide clinical decisions ([Analysis 4.1.3](#) (1 study, 60 participants) RR 7.15 95% CI 0.42 to 121.04; very low certainty evidence).

Renal parenchymal abnormality

Two studies ([Nordenstrom 2016](#); [Swedish Reflux Trial 2010](#)) reported the outcome of renal parenchymal defects on DMSA scan. Compared with antibiotic prophylaxis, endoscopic correction of VUR may make little or no difference to the number of children with new or deteriorating renal damage on DMSA scan ([Analysis 4.2.1](#) (2 studies, 208 participants): RR 1.20, 95% CI 0.36 to 3.96; $I^2 = 45\%$; low certainty evidence). The inconsistency, bias and small sample size with few events in these studies leads to an assessment of low certainty evidence. One study ([Capozza 2002](#)) measured renal damage using ultrasound and reported a very imprecise estimate of treatment effect ([Analysis 4.3.1](#) (1 study, 81 participants: RR 0.34 95% CI 0.04, 3.31) for new renal parenchymal defects

Correction of VUR

Compared with antibiotics alone, more children in the endoscopic group may have full resolution of VUR after one to two years ([Analysis 4.4.1](#) (2 studies, 164 children): RR 2.69, 95% CI 1.57 to 4.63) and improvement in VUR grade ([Analysis 4.4.2](#) (3 studies, 238 children): RR 2.01 95% CI 1.51 to 2.68; $I^2 = 0\%$; low certainty evidence). Assessment of these studies showed many study design fields with high or unclear levels of bias and the studies were small and imprecise, leading to the conclusion this is low certainty evidence

Microbial resistance to the prophylactic drug

Only [Swedish Reflux Trial 2010](#) reported microbial resistance to the treatment antibiotic and showed little or no difference between the groups in the number of UTIs caused by pathogens resistant

to the antibiotic being used for prophylaxis ([Analysis 4.5](#); RR 0.63 95% CI 0.34 to 1.15; very low certainty evidence), This finding is not surprising given that both groups were taking antibiotics.

Adverse events

None of the children treated with endoscopic injection in [Swedish Reflux Trial 2010](#) experienced vesicoureteric obstruction but three experienced post-operative complications. Three children in the antibiotic group and none in the endoscopic treatment group within the [Nordenstrom 2016](#) study experienced nausea and vomiting while on treatment.

Other outcomes

ESKD and hypertension were not reported.

Endoscopic injection treatment versus no treatment

Urinary tract infection

Symptomatic and febrile UTI

[Swedish Reflux Trial 2010](#) involved three study arms which allows for a comparison between endoscopic injection treatment and no treatment. Compared with no treatment, endoscopic correction of VUR may reduce the risk for symptomatic and febrile UTI ([Analysis 5.1](#), (1 study; 134 participants): RR 0.55 95% CI 0.33 to 0.94 and [Analysis 5.2](#), (1 study; 134 participants): RR 0.52 95% CI 0.30 to 0.89). Since only 134 children were studied this may represent a chance effect.

Renal parenchymal abnormality

[Swedish Reflux Trial 2010](#), compared with no treatment, endoscopic correction of VUR may make little or no difference to the risk for renal parenchymal defects ([Analysis 5.3](#) (1 study; 133 participants): RR 0.70, 95% CI 0.3 to 1.6).

Other outcomes

Resolution of VUR, ESKD, hypertension, and adverse events were not reported.

Different materials for subureteric injection to correct VUR

See [Summary of findings 4](#).

Four studies compared two different products for endoscopic injection to correct VUR, Two studies ([Moore 2014](#); [Oswald 2002](#)) compared endoscopic subureteric injections of Macroplastique with Deflux but these were dissimilar in that [Oswald 2002](#) compared these in children also taking prophylactic antibiotics. [Moore 2014](#) showed no difference in the number of febrile UTIs across the two treatment groups and [Oswald 2002](#) reported no difference in afebrile UTIs between the two groups. For the outcome of persistent VUR both studies favoured Macroplastique with a 50% reduction in risk of persistent VUR at three months ([Analysis 6.1.3](#) (2 studies, 513 participants): RR 0.50, 95% CI 0.33 to 0.78; $I^2 = 0\%$) and a 46% reduction in risk at 12 months ([Analysis 6.1.4](#) (2 studies, 472 participants): RR 0.54 (95% CI 0.35 to 0.83; $I^2 = 0\%$). Assessment of these studies found study design fields of unclear or high risk of bias indicating that subureteric injection with Macroplastique compared with dextranomer hyaluronic acid polymer probably reduces the risk for persistent VUR (moderate certainty evidence). [Oswald 2002](#) reported a moderately higher risk of temporary pelvicaliceal dilatation with Macroplastique ([Analysis](#)

[6.1.5](#) (114 participants): RR 1.85, 95% CI 1.02 to 3.35; low certainty evidence). [Moore 2014](#) reported three cases of ureteral obstruction in the Macroplastique group compared with two cases in the Deflux group (low certainty evidence). There were four cases of flank pain after injection in the Macroplastique compared with three cases in the Deflux group of [Moore 2014](#) (low certainty evidence).

No data on renal parenchymal abnormalities were reported.

One small study of 28 children ([Frey 1997](#)) compared endoscopic subureteric injections of different concentrations of GAX 65 and GAX 35. VUR was five times more likely to persist following GAX 35 than GAX 65 injections ([Analysis 6.2.1](#) (28 children): RR 0.21, 95% CI 0.05 to 0.85). The estimated risk of recurrence of VUR between therapies was imprecise ([Analysis 6.2.2](#) (1 study, 28 children): RR 0.30, 95% CI 0.07 to 1.29). No data on UTIs or renal parenchymal abnormalities were reported. This study represents very low certainty evidence.

[Habeeb Abid 2015](#) compared dextranomer hyaluronic acid polymer with polyacrylate polyalcohol bulking polymer (Vantrix). UTIs were not reported. Different bulking agents may lead to little or no difference in risk of repeat positive urine culture, fever, persistence of VUR or ureteral obstruction ([Analysis 6.3.1](#) to [Analysis 6.3.4](#)) (low certainty evidence).

Probiotic versus antibiotic prophylaxis

See [Summary of findings 5](#).

Two studies ([Lee 2007a](#); [Lee 2015](#)) compared the potential benefits and harms of probiotic versus antibiotic prophylaxis in single centre studies and included 120 and 128 children respectively. Compared with antibiotic prophylaxis, probiotics may make little or no difference in risk of repeat symptomatic UTI ([Analysis 7.1.1](#) (1 study, 24 children): RR 0.82, 95% CI 0.56 to 1.21) and febrile UTI at one year ([Analysis 7.1.2](#) (1 study, 24 children): RR 0.82, 95% CI 0.37 to 1.83; low certainty evidence). Assessment of these two studies showed there were many study design fields with high or unclear risk of bias, the studies were small and imprecise, leading to the conclusion there is low certainty in this evidence. In [Lee 2007a](#) by the end of the one year follow-up period, 2 of 13 patients in the antibiotics group had experienced a repeat UTI and had developed a new renal scar, versus 1 of 11 in the probiotics group. In [Lee 2007a](#) seven symptomatic UTI recurrences (64%) had *Escherichia coli* identified as the causative organism in the probiotics group versus nine (69%) in the antibiotics group. These data were not available for [Lee 2015](#). Across the two studies, six of the 18 repeat UTIs caused by *E.coli* in the probiotic group were resistant to trimethoprim-sulphamethoxazole while 20/20 *E. coli* infections were resistant to the treatment drug in the antibiotic group ([Analysis 7.2](#) (RR 0.38, 95% CI 0.21 to 0.69)(RR 0.38, 95% CI 0.21 to 0.69; $I^2 = 0\%$). Compared with antibiotic prophylaxis, probiotics may make little or no difference to the risk of renal parenchymal abnormality ([Analysis 7.3.1](#) (2 studies, 152 participants): RR 0.69, 95% CI 0.43 to 1.11; $I^2 = 0\%$; low certainty evidence). These studies do not provide evidence of sufficient certainty to support clinical decisions on the efficacy of probiotics to prevent UTI in children.

A single study of 85 children compared probiotics plus antibiotics with antibiotics alone ([Mohseni 2013](#)). No difference between treatment groups was seen for the outcomes repeat febrile UTI, repeat positive urine culture, new DMSA changes, microbial resistance and adverse effects ([Analysis 7.4](#)).

ESKD, hypertension, resolution of VUR, and adverse events were not reported.

Other Interventions

Surgical reimplantation versus endoscopic treatment

A single small scale study ([Garcia-Aparicio 2013](#)) compared surgical reimplantation (Cohens technique) with endoscopic injection of dextranomer hyaluronic acid polymer in 41 children. Compared with endoscopic treatment, surgical implantation may make little or no difference in outcomes of repeat positive urine culture, VUR resolution or complications of the procedures were seen between the surgical and endoscopic treatment groups ([Analysis 8.1](#)) (low certainty evidence).

Cranberry products

A study of 76 children ([Dotis 2014](#)) compared cranberry capsules with no treatment and found repeat positive urine culture was more frequent in the no treatment arm ([Analysis 8.2](#) (1 study, 76 participants): RR 0.29 95% CI 0.14 to 0.59). It is uncertain whether cranberry capsules compared with no treatment prevents UTI since the certainty of the evidence is very low.

Circumcision

[Gucuk 2013](#) compared circumcision plus antibiotics to antibiotics alone in 91 boys with VUR. Repeat positive urine culture was less frequent in the circumcision group ([Analysis 8.3.2](#); (91 participants): RR 0.28 95% CI 0.12 to 0.70) but symptomatic UTI and DMSA changes showed no differences across treatment groups ([Analysis 8.3.1](#); [Analysis 8.3.3](#)). From this single RCT, it remains uncertain whether circumcision plus antibiotics prevents UTI since the certainty of this evidence is very low (very low certainty evidence).

Oxybutynin

[Ahn 2003](#) compared the anticholinergic, Oxybutynin, with placebo for the outcome resolution of VUR at 12 months. 38/56 (67.9%) children treated with oxybutynin had resolution or improvement of their VUR grade while 57/93 (61.3%) children in the placebo group showed an improved or resolved VUR grade after 12 months ([Analysis 8.4](#)). Oxybutynin compared with placebo may make little or no difference to the improvement in VUR (low certainty evidence).

DISCUSSION

Summary of main results

The benefits and harms of interventions for primary VUR were assessed in 34 studies involving 4001 children.

Antibiotic prophylaxis versus surveillance/no treatment

Twelve studies comparing long-term low-dose antibiotic prophylaxis with placebo or no treatment were identified. Although the pooled estimate of risk of repeat symptomatic and febrile UTI favoured antibiotic treatment, precision was poor and included no difference between treatment groups. Overall, long-term low-dose antibiotic prophylaxis may make little or no difference to the number of children who experience repeat symptomatic or febrile UTI (low certainty evidence) or to the number of children, developing new or progressive renal damage compared with placebo or no treatment (low certainty evidence) ([Summary of](#)

[findings for the main comparison](#)). Three studies were at low risk of bias across all study design fields and when only these three studies were considered, antibiotic prophylaxis made little or no difference to the likelihood of further UTIs (moderate certainty evidence) ([Summary of findings for the main comparison](#)). Side effects of the preventive treatment were minor and infrequent but poorly reported (low certainty evidence) ([Summary of findings for the main comparison](#).) Little data were provided as to how side-effects had affected adherence. Reported compliance rates varied between 70% and 100%, but they were inconsistently measured. Since previous studies have shown poor compliance for daily antibiotic regimens for VUR ([Cohen 2005](#); [Greenfield 1997](#)), it raises the question whether antibiotics were inherently ineffective or were not being used as prescribed. Treating VUR patients with long-term low-dose antibiotics probably increases the risk of microbial resistance against the prophylactic drug in breakthrough infections (moderate certainty evidence) ([Summary of findings for the main comparison](#)).

Surgical reimplantation of ureters plus antibiotics (1 to 24 months) versus antibiotics alone

Only the European arm of the IRS study evaluated the differential risk of any symptomatic UTI between medical and surgical treatment and found that by four to five years, there was little or no difference in symptomatic UTI. Both the European and the American arms of the IRS study however, investigated the risk of febrile UTI and found a significant benefit for those children who underwent surgical VUR correction. Assuming a baseline risk of 22% of developing febrile UTI when on antibiotics alone the estimated risk translates into a risk difference of 13% so that eight patients need surgery to prevent one extra febrile UTI over five years. T. Further analysis at 10 years confirmed these results. It supports the idea that although surgery might not provide an added benefit over antibiotic prophylaxis in preventing symptomatic lower UTI, it might keep the infection from spreading to the upper tract, and ultimately prevent subsequent renal damage. No evidence was found to corroborate this theory however, since the risk of developing new or progressive renal damage at five and 10 years was no different between the treatment groups. Although no differences between treatment groups were demonstrated for hypertension or end stage kidney disease, it remains uncertain whether surgical correction of VUR influences the risk of hypertension and CKD because the certainty of this evidence is very low ([Summary of findings 2](#)).

Potential benefits of surgery need to be weighed against its potential adverse effects. Whereas the Birmingham Reflux Study stated that no cases of postoperative obstruction were found after five years, they occurred in 7% of children (10/151) in the European arm of the International Reflux Study, corresponding to one every 14 to 15 patients undergoing the procedure.

Endoscopic injection treatment for VUR correction

If VUR correction were beneficial, endoscopic subureteric injection of various materials could offer an alternative method to open surgery for correcting VUR. Eight studies included in this review used endoscopic injection in at least one study group, and these have demonstrated VUR correction rates of 70% to 90%. However when compared with long-term low-dose antibiotic prophylaxis alone, endoscopic correction with or without antibiotics may make little or no difference to the likelihood of either symptomatic,

febrile UTI or progressive renal damage. Results were derived from three small studies with only 254 children leading to low certainty evidence ([Summary of findings 3](#)). A single small study ([Swedish Reflux Trial 2010](#)) found that fewer children following endoscopic correction of VUR developed UTI compared to no treatment.

Four studies in this review directly compared two different materials with each other, with two suggesting that Macroplastique is superior to dextranomer/hyaluronic acid polymer for correcting reflux. However the quality, size and consistency of findings is inadequate for confident conclusions (low certainty evidence) ([Summary of findings 4](#)).

Other interventions

Probiotic treatment may make little or no difference to the risk of symptomatic UTI or renal damage detected by DMSA scan (low certainty evidence) ([Summary of findings 5](#)). Cranberry capsules, circumcision and oxybutynin were trialled in single, small studies with each study result favouring the intervention but with limited confidence due to imprecision from small sample sizes with infrequent events, and sources of bias from sub-optimal study designs.

Overall completeness and applicability of evidence

Children with grade V reflux are generally viewed as having the highest risk of developing renal damage after pyelonephritis. However in the 12 studies of low-dose antibiotic treatment compared with placebo or no treatment, only three included children with grade V reflux and only [PRIVENT 2009](#) specified the number of children (5) with grade V reflux. Most surgical reimplantation studies included participants with VUR grades II, III and IV and children with grade V VUR were excluded. Hence we cannot extrapolate the results from this review to children with VUR grade V. A randomised comparison between antibiotic treatment and surgical reimplantation alone has not been performed since in all studies, antibiotics were given to children undergoing surgery. Very few studies reported data for analysis of treatment effect within the genders and laterality of VUR, hence no conclusions could be drawn. The few small studies comparing endoscopic injection treatment of VUR with antibiotic prophylaxis are insufficient to determine whether there are real benefits for endoscopic correction over antibiotic prophylaxis or no treatment. Further studies are required to determine if probiotics or other interventions have any role in preventing UTI in children with VUR.

Quality of the evidence

The quality of conduct and reporting of these studies was variable, with many studies omitting crucial methodological information used to assess the risk of bias. Only four of the 34 studies were considered at low risk of bias across all fields of study quality ([Craig 2002](#); [Hari 2015](#); [PRIVENT 2009](#); [RIVUR 2008](#)). The majority of studies had many areas of uncertainty in the risk of bias fields, reflecting missing detail rather than stated poor design.

The overall certainty of the evidence using GRADE was considered to be low in most analyses. Only two analyses were considered to have moderate overall certainty of the evidence.

The effect estimate of incremental benefit of surgical reimplantation was based on only two studies that included 429

patients in the analysis. Given the nature of the intervention, it was not possible to blind participants or health care providers. Unblinded studies tend to report an overestimation of treatment effects and as such bias might have played a part in producing the difference in risk of febrile UTI in favour of surgical management.

Potential biases in the review process

A comprehensive search of the Cochrane Kidney and Transplant's Specialised Register was performed for this review thus reducing the possibility that potential eligible studies were omitted from the review. Eligible studies published after the last search date or published in congress proceedings not routinely searched could have been missed. Authors of this review are also authors of an included study ([PRIVENT 2009](#)) and may have judged that study more favourably. To alleviate this concern, an independent researcher extracted risk of bias data from [PRIVENT 2009](#) and this was incorporated into the review.

Agreements and disagreements with other studies or reviews

For over five decades, scientific published work has suggested the existence of a link between recurrent UTI, VUR, and renal scarring ([Olbing 2003](#); [Smellie 1975](#); [Smellie 1994](#); [Smellie 1998](#)). VUR has been thought to facilitate the involvement of the upper urinary tract during UTI by allowing retrograde passage of infected urine to the ureter. Subsequently all interventions have been targeted at preventing UTI-induced damage to the kidney. Two earlier versions of this review ([Hodson 2007](#); [Wheeler 2004](#)) were not able to provide evidence as to whether the common practice of diagnosing and treating children with VUR conferred important health benefits, since no adequately powered studies had included a no treatment arm. Since then, new studies have compared the administration of antibiotics with placebo or no treatment. Although the addition of these studies produced a risk estimate in favour of long-term low-dose antibiotics for preventing repeat symptomatic UTI, the imprecision of the estimate means there may be no difference in risk of repeat UTI between children taking antibiotics and those not treated.

In this review, we have highlighted symptomatic and febrile UTI as more relevant primary end points than positive urine cultures.

AUTHORS' CONCLUSIONS

Implications for practice

Compared with no treatment, use of long-term low-dose antibiotics may make little or no difference to the number of repeat symptomatic and febrile UTIs in children with VUR. Small studies of variable methodological quality and a large amount of unexplained heterogeneity in the analysis makes drawing firm conclusions challenging. Prophylaxis produced few side effects but was associated with a threefold increase in prophylactic drug resistance in subsequent UTIs.

The added benefit of surgery over long-term low-dose antibiotic use remains uncertain. Although there was a significant reduction in repeat episodes of febrile UTI, there were no differences in either symptomatic UTI or renal damage. Informed decision making should consider the risk of adverse events associated with surgery. Correcting VUR using endoscopic approaches would theoretically reduce these risks. However there were few studies

of low methodological quality so endoscopic correction may make little or no difference to number of symptomatic or febrile UTIs or in new or progressive renal damage.

Implications for research

Identifying which children are at risk of repeat UTI is difficult and relying only on VUR status is not helpful. Correction of severe VUR may alleviate problems in some children and as such the role of surgery and endoscopic treatment for the management of VUR needs further exploration. Of specific interest would be the impact of VUR correction by endoscopic subureteric injection without antibiotics versus no treatment on the incidence of febrile UTI and renal parenchymal injury assessed by DMSA scan. The role of prophylactic treatment using complimentary medicines such as cranberry product and probiotics is uncertain and larger, well designed studies may clarify their efficacy.

ACKNOWLEDGEMENTS

- We are grateful to Danielle Wheeler, Dushyanthi Vimalachandra, L Paul Roy and Grahame Smith who were authors of the original review ([Wheeler 2004](#)), contributing to the design, quality assessment, data collection, entry, analysis and interpretation, and writing.
- We would like to thank Professor Les Irwig for his help with methodology and manuscript preparation, and those authors and experts in the field who replied to our requests for study information.
- This study was initially funded in part by a seeding grant from the Australian Kidney Foundation (Grant no. S2/99).
- We are grateful to Evi Nagler who coordinated, extracted data and drafted text for the 2011 update ([Nagler 2011](#)).

REFERENCES

References to studies included in this review

Ahn 2003 {published data only}

Ahn SH, Choi HR, Cho SJ, Lee SJ. The effect of oxybutynin on spontaneous resolution of primary vesicoureteral reflux in children [abstract no: FC9-02]. *Nephrology* 2003;**8**(Suppl 1):A24.

Ahn SH, Shim SY, Lee JW, Cho SJ, Lee SJ. The effect of an anticholinergic agent (oxybutynin) on spontaneous resolution of primary vesicoureteral reflux. *Journal of the Korean Society of Pediatric Nephrology* 2003;**7**(2):174-80. [CENTRAL: CN-01044096]

BIRSG 1983 {published data only}

A prospective trial of operative versus non-operative treatment of severe vesico-ureteric reflux: 2 years' observation in 96 children. The Birmingham Reflux Study Group. *Contributions to Nephrology* 1984;**39**:169-85. [MEDLINE: 6744870]

Prospective trial of operative versus non-operative treatment of severe vesicoureteric reflux: two years observation in 96 children. *British Medical Journal Clinical Research Ed* 1983;**287**(6386):171-4. [MEDLINE: 6409239]

* Corkery JJ, Gornall P, Shah KJ, Taylor CM, White RHR, Winterborn MH. Prospective trial of operative versus non-operative treatment of severe vesicoureteric reflux in children: five years' observation. Birmingham Reflux Study Group. *British Medical Journal Clinical Research Ed* 1987;**295**(6592):237-41. [MEDLINE: 2888509]

Taylor CM, White R, Winterborn MH, Shah KJ, Corkery JJ, Gornall P. Medical versus surgical management - the Birmingham study five years on. Proceedings of the 1st CJ Hodson Symposium on Reflux Nephropathy; 1987 Mar 5-6; Melbourne, Australia. 1987:79-87.

Capozza 2002 {published data only}

Capozza N, Caione P. Dextranomer/hyaluronic acid copolymer implantation for vesico-ureteral reflux: a randomized comparison with antibiotic prophylaxis. *Journal of Pediatrics* 2002;**140**(2):230-4. [MEDLINE: 11865276]

Craig 2002 {published and unpublished data}

Craig J, Roy L, Burke J, Powell H, Hodson EM, Knight JF, et al. Long-term antibiotics to prevent urinary tract infection in children with isolated vesicoureteric reflux: a placebo-controlled randomized trial [abstract no: 44]. *Nephrology* 2002;**7**(Suppl 3):A55.

Craig J, Roy L, Sureshkumar P, Burke J, Powell H, et al. Long-term antibiotics to prevent urinary tract infection in children with isolated vesicoureteric reflux: a placebo-controlled randomized trial [abstract no: F-FC014]. *Journal of the American Society of Nephrology* 2002;**13**(September, Program & Abstracts):3A.

Dite 2007 {published data only (unpublished sought but not used)}

Dite Z, Kocvara R, Dvoracek J, Langer J, Sedlacek J. Endoscopic and conservative treatment of vesicoureteric reflux [Endoskopicka a konzervativni lecba vezikoureteralniho

refluxu]. *Casopis Lekarů Ceských* 2007;**146**(10):817-21. [MEDLINE: 18020019]

Dotis 2014 {published data only}

Dotis J, Printza N, Stabouli S, Pavlaki A, Samara S, Papachristou F. Efficacy of cranberry capsules to prevent recurrences of urinary tract infections [abstract no: P351]. *Pediatric Nephrology* 2014;**29**(9):1793-4. [EMBASE: 71662748]

Espino Hernandez 2012 {published data only}

Espino Hernandez M, Arcos GB, Melgosa Hijosa M, Orive Olondriz B, Pena A, Rodriguez Do Forno A, et al. Antibiotic prophylaxis in high degree vesicoureteral reflux clinical trial and prospective, observational and multicentric study [abstract no: O80]. *Pediatric Nephrology* 2014;**29**(9):1684. [EMBASE: 71662394]

Espino M, Areses R, Garcia Meseguer C, Pena A, Melgosa M, Ruperez M, et al. Antibiotic prophylaxis in high degree vesicoureteral reflux. Prospective, randomized and multicentric study, preliminary results [abstract no: P10]. *Pediatric Nephrology* 2012;**27**(9):1648.

Frey 1997 {published data only}

Frey P, Gudinchet F, Jenny P. GAX 65: new injectable cross-linked collagen for the endoscopic treatment of vesicoureteral reflux--a double-blind study evaluating its efficiency in children. *Journal of Urology* 1997;**158**(3 Pt 2):1210-2. [MEDLINE: 9258175]

Garcia-Aparicio 2013 {published data only}

Garcia-Aparicio L, Rovira J, Blazquez-Gomez E, Garcia-Garcia L, Gimenez-Llort A, Rodo J, et al. Randomized clinical trial comparing endoscopic treatment with dextranomer hyaluronic acid copolymer and Cohen's ureteral reimplantation for vesicoureteral reflux: long-term results. *Journal of Pediatric Urology* 2013;**9**(4):483-7. [MEDLINE: 23602843]

Garin 2006 {published data only}

Garin E, Olavarria F, Garcia Nieto V, Valenciano Fuente B. Clinical significance of primary vesicoureteral reflux and urinary antibiotic prophylaxis after acute pyelonephritis [abstract no: FP-3-1]. *Pediatric Nephrology* 2007;**22**(7):1070. [CENTRAL: CN-00724967]

* Garin EH, Olavarria F, Garcia Nieto V, Valenciano B, Campos A, Young L. Clinical significance of primary vesicoureteral reflux and urinary antibiotic prophylaxis after acute pyelonephritis: a multicenter, randomized, controlled study. *Pediatrics* 2006;**117**(3):626-32. [MEDLINE: 16510640]

Gucuk 2013 {published data only}

Burgu B, Gokce I, Aydogdu O, Soygun T. Do antibiotic prophylaxis and/or circumcision change periurethral uropathogen colonization and urinary tract infection rates in boys with VUR? [abstract]. *Journal of Urology* 2011;**185**(4 Suppl 1):e231. [EMBASE: 70377838]

* Gućuk A, Burgu B, Gokce I, Mermerkaya M, Soygun T. Do antibiotic prophylaxis and/or circumcision change periurethral uropathogen colonization and urinary tract infection rates

in boys with VUR?. *Journal of Pediatric Urology* 2013;**9**(6 Pt B):1131-6. [MEDLINE: 23721792]

Habeeb Abid 2015 {published data only}

Habeeb Abid M. Comparative study between the use of dextranomer hyaluronic acid copolymer (Dexcel) versus polyacrylate polyalcohol copolymer (Vantris) in the endoscopic treatment of vesicoureteral reflux (VUR) in children [abstract]. *European Urology Supplements* 2015;**14**(2):e504. [EMBASE: 71831430]

Hari 2015 {published data only}

Hari P, Hari S, Sinha A, Kumar R, Kapil A, Bagga A. Antibiotic prophylaxis in management of vesicoureteric reflux: a double-blind placebo controlled trial [abstract no: O-43]. *Pediatric Nephrology* 2013;**28**(8):1363-4. [EMBASE: 71126984]

* Hari P, Hari S, Sinha A, Kumar R, Kapil A, Pandey RM, et al. Antibiotic prophylaxis in the management of vesicoureteric reflux: a randomized double-blind placebo-controlled trial. *Pediatric Nephrology* 2015;**30**(3):479-86. [MEDLINE: 25173357]

Holland 1982 {published data only}

Holland NH, Kazee M, Duff D, McRoberts JW. Antimicrobial prophylaxis in children with urinary tract infection and vesicoureteral reflux. *Reviews of Infectious Diseases* 1982;**4**(2):467-74. [MEDLINE: 6981165]

IRS EUR 1981-2003 {published data only}

Comparison of conservative and surgical treatment of the vesico-uretero-renal reflux. International prospective reflux study in children [Vergleich von medikamentöser und chirurgischer Behandlung beim primären vesiko-uretero-renalen reflux]. *Urologe. Ausgabe A* 1981;**20**(2):107-15. [MEDLINE: 7025416]

Hjalmas K, Lohr G, Tamminen-Mobius T, Seppanen J, Olbing H, Wikstrom S. Surgical results in the International Reflux Study in Children (Europe). *Journal of Urology* 1992;**148**(5 Pt 2):1657-61. [MEDLINE: 1433583]

Jodal U, Koskimies O, Hanson E, Lohr G, Olbing H, Smellie J, et al. Infection pattern in children with vesicoureteral reflux randomly allocated to operation or long-term antibacterial prophylaxis. The International Reflux Study in Children. *Journal of Urology* 1992;**148**(5 Pt 2):1650-2. [MEDLINE: 1433581]

Jodal U, Smellie JM, Lax H, Hoyer PF. Ten-year results of randomized treatment of children with severe vesicoureteral reflux. Final report of the International Reflux Study in Children. *Pediatric Nephrology* 2006;**21**(6):785-92. [MEDLINE: 16565873]

Olbing H. Comparison of the surgical and nonsurgical treatment of primary vesico-ureteral-renal reflux--an international reflux study [Vergleich von chirurgischer und nicht-chirurgischer Behandlung beim primären vesiko-uretero-renalen Reflux--Internationale Refluxstudie]. *Kinderärztliche Praxis* 1986;**54**(9):493-9. [MEDLINE: 3540424]

Olbing H. Risk-oriented recommendations for the management of children with dilating primary vesico-ureteral reflux - report of the International Reflux Study 1 children (IRS) - European branch [abstract]. 9th Congress. International Pediatric

Nephrology Association; 1992 Aug 30 - Sep 4; Jerusalem, Israel. 1992:C37-C8.

Olbing H, Claesson I, Ebel KD, Seppanen U, Smellie JM, Tamminen-Mobius T, et al. Renal scars and parenchymal thinning in children with vesicoureteral reflux: a 5-year report of the International Reflux Study in Children (European branch). *Journal of Urology* 1992;**148**(5 Pt 2):1653-6. [MEDLINE: 1433582]

Olbing H, Hirche H, Koskimies O, Lax H, Seppanen U, Smellie JM, et al. Renal growth in children with severe vesicoureteral reflux: 10-year prospective study of medical and surgical treatment: the International Reflux Study in Children (European branch). *Radiology* 2000;**216**(3):731-7. [MEDLINE: 10966703]

Olbing H, Smellie JM, Jodal U, Lax H. New renal scars in children with severe VUR: a 10-year study of randomized treatment. *Pediatric Nephrology* 2003;**18**(11):1128-31. [MEDLINE: 14523634]

Olbing H, Tamminen-Möbius T, International Reflux Study in Europe. International reflux study patients with stable and unstable vesicoureteral reflux during pre-allocation period [abstract no: W9-2]. *Pediatric Nephrology* 1987;**1**(1):C19. [CENTRAL: CN-00447018]

Piepsz A, Tamminen-Mobius T, Reiners C, Heikkila J, Kivisaari A, Nilsson NJ, et al. Five-year study of medical or surgical treatment in children with severe vesico-ureteral reflux dimercaptosuccinic acid findings. International Reflux Study Group in Europe. *European Journal of Pediatrics* 1998;**157**(9):753-8. [MEDLINE: 9776536]

Smellie JM. The International Reflux Study in Children: an update. Proceedings of the 1st CJ Hodson Symposium on Reflux Nephropathy; 1987 Mar 5-6; Melbourne, Australia. 1987:88-91.

Smellie JM, Jodal U, Lax H, Mobius TT, Hirche H, Olbing H, et al. Outcome at 10 years of severe vesicoureteric reflux managed medically: report of the International Reflux Study in Children. *Journal of Pediatrics* 2001;**139**(5):656-63. [MEDLINE: 11713442]

Smellie JM, Tamminen-Mobius T, Olbing H, Claesson I, Wikstad I, Jodal U, et al. Radiologic findings in the kidney of children with severe reflux. Five-year comparative study of conservative and surgical treatment [Radiologische Nierenbefunde bei Kindern mit schwerem Reflux. Fünfjahresvergleichsstudie konservativer und chirurgischer Behandlung]. *Urologe - Ausgabe A* 1993;**32**(1):22-9. [MEDLINE: 8447041]

Smellie JM, Tamminen-Möbius T, Olbing H, Claesson I, Wikstad I, Jodal U, et al. Five-year study of medical or surgical treatment in children with severe reflux: radiological renal findings. The International Reflux Study in Children. *Pediatric Nephrology* 1992;**6**(3):223-30. [MEDLINE: 1616829]

Tamminen-Mobius T, Brunier E, Ebel KD, Lebowitz R, Olbing H, Seppanen U, et al. Cessation of vesicoureteral reflux for 5 years in infants and children allocated to medical treatment. The International Reflux Study in Children. *Journal of Urology* 1992;**148**(5 Pt 2):1662-6. [MEDLINE: 1433584]

Wingen AM, Koskimies O, Olbing H, Seppanen J, Tamminen-Mobius T. Growth and weight gain in children with

vesicoureteral reflux receiving medical versus surgical treatment: 10-year results of a prospective, randomized study. International Reflux Study in Children (European Branch). *Acta Paediatrica* 1999;**88**(1):56-61. [MEDLINE: 10090549]

IRS USA 1992 {published data only}

Arant BS Jr. Vesicoureteral reflux and evidence-based management. *Journal of Pediatrics* 2001;**139**(5):620-1. [MEDLINE: 11713435]

Duckett JW, Walker RD, Weiss R. Surgical results: International Reflux Study in Children--United States branch. *Journal of Urology* 1992;**148**(5 Pt 2):1674-5. [MEDLINE: 1433586]

Weiss R, Duckett J, Spitzer A. Results of a randomized clinical trial of medical (m) vs surgical (s) management of infants & children with primary grades III & IV vesico-ureteral reflux (VUR) [abstract no: 11P]. *Journal of the American Society of Nephrology* 1992;**3**(3):290. [CENTRAL: CN-00461981]

Weiss R, Duckett J, Spitzer A. Results of a randomized clinical trial of medical versus surgical management of infants and children with grades III and IV primary vesicoureteral reflux (United States). The International Reflux Study in Children. *Journal of Urology* 1992;**148**(5 Pt 2):1667-73. [MEDLINE: 1433585]

Weiss R, Tamminen-Mobius T, Koskimies O, Olbing H, Smellie JM, Hirche H, et al. Characteristics at entry of children with severe primary vesicoureteral reflux recruited for a multicenter, international therapeutic trial comparing medical and surgical management. The International Reflux Study in Children. *Journal of Urology* 1992;**148**(5 Pt 2):1644-9. [MEDLINE: 1433580]

Kojima 2012 {published data only}

Kojima Y, Mizuno K, Umemoto Y, Yasui T, Hayashi Y, Kohri K. Ureteral advancement in patients undergoing laparoscopic extravesical ureteral reimplantation for treatment of vesicoureteral reflux. *Journal of Urology* 2012;**188**(2):582-7. [MEDLINE: 22704095]

Lee 2007a {published and unpublished data}

Lee SJ, Kim HJ, Shim YH, Lee JW. The effect of probiotic prophylaxis for preventing recurrent urinary tract infection in children with persistent primary vesicoureteral reflux [abstract no: COD.PP.81]. *Pediatric Nephrology* 2006;**21**(10):1542. [CENTRAL: CN-00724966]

* Lee SJ, Shim YH, Cho SJ, Lee JW. Probiotics prophylaxis in children with persistent primary vesicoureteral reflux. *Pediatric Nephrology* 2007;**22**(9):1315-20. [MEDLINE: 17530295]

Lee 2015 {published data only}

* Lee SJ, Lee JW. Probiotics prophylaxis in infants with primary vesicoureteral reflux. *Pediatric Nephrology* 2015;**30**(4):609-13. [MEDLINE: 25354903]

Lee SJ, Lee M. Probiotics versus antibiotic prophylaxis in infants with primary vesicoureteral reflux [abstract no: 285]. *Pediatric Nephrology* 2010;**25**(9):1852.

Mohseni 2013 {published data only}

Mohseni MJ, Aryan Z, Emamzadeh-Fard S, Paydary K, Mofid V, Joudaki H, et al. Combination of probiotics and antibiotics in the prevention of recurrent urinary tract infection in children. *Iranian Journal of Pediatrics* 2013;**23**(4):430-8. [MEDLINE: 24427497]

Montini 2008 {published data only (unpublished sought but not used)}

Hewitt IK, Tomasi L, Pavanello L, Maschio F, Molinari PP, Toffolo A, et al. Early treatment of acute pyelonephritis in children fails to reduce renal scarring [abstract no: SA-PO1100]. *Journal of the American Society of Nephrology* 2006;**17**(Abstracts):804A. [CENTRAL: CN-00653791]

Hewitt IK, Zucchetto P, Rigon L, Maschio F, Molinari PP, Tomasi L, et al. Early treatment of acute pyelonephritis in children fails to reduce renal scarring: data from the Italian Renal Infection Study Trials. *Pediatrics* 2008;**122**(3):486-90. [MEDLINE: 18762516]

Montini G. Evaluation of the effectiveness of antibiotic prophylaxis in children with a history of upper urinary tract infections: a multi centre randomised study - Protocol. Personal communication 2004.

Montini G, Rigon L, Gobber D, Zucchetto P, Murer L, Calderan A, et al. A randomised controlled trial of antibiotic prophylaxis in children with a previous documented pyelonephritis. [abstract no: OFC35]. *Pediatric Nephrology* 2004;**19**(9):C70. [CENTRAL: CN-00583479]

* Montini G, Rigon L, Zucchetto P, Fregonese F, Toffolo A, Gobber D, et al. Prophylaxis after first febrile urinary tract infection in children? A multicenter, randomized, controlled, noninferiority trial. *Pediatrics* 2008;**122**(5):1064-71. [MEDLINE: 18977988]

Moore 2014 {published data only}

Moore K, Bolduc S. Prospective study of polydimethylsiloxane vs dextranomer/hyaluronic acid injection for treatment of vesicoureteral reflux. *Journal of Urology* 2014;**192**(6):1794-9. [MEDLINE: 24928269]

Morris 1991 {published data only}

Morris MC, Rothwell DL, Paykel AD. A prospective study of vesico-ureteric reflux and renal function in children. 2nd CJ Hodson Symposium on Reflux Nephropathy; 1991 Feb 26; Christchurch, NZ. 1991:61-2. [CENTRAL: CN-00602103]

Nordenstrom 2016 {published data only}

Nordenstrom J, Holmdahl G, Brandstrom P, Sixt R, Stokland E, Sillen U, et al. The Swedish infant high-grade reflux trial: Study presentation and vesicoureteral reflux outcome. *Journal of Pediatric Urology* 2016;**13**(2):130-8. [MEDLINE: 27889221]

Nordenstrom J, Holmdahl G, Sillen U, Sixt R, Stokland E, Sjostrom S. The Swedish infant high grade reflux trial-VUR grade and bladder function [abstract]. *Pediatric Nephrology* 2015;**30**(9):1565. [EMBASE: 72033229]

Nordenstrom J, Sillen U, Holmdahl G, Linner T, Stokland E, Sjöström S. The Swedish Infant High-grade Reflux Trial - bladder function. *Journal of Pediatric Urology* 2016;**13**(2):139-45. [MEDLINE: 27989639]

* Nordenstrom J, Sjöström S, Sillen U, Sixt R, Brandström P. The Swedish infant high grade reflux trial: UTI and renal damage. *Journal of Pediatric Urology* 2017;**13**(2):146-54. [MEDLINE: 28215835]

Nordenstrom J, Sjöström S, Sixt R, Sillen U, Brandström P. The Swedish infant high grade reflux trial-UTI and renal damage [abstract]. *Pediatric Nephrology* 2015;**30**(9):1565. [EMBASE: 72033228]

Nordenström J, Sillén S, Brandström P, Hansson S, Holmdahl G, Linnér T, et al. The Swedish Infant High Grade Reflux Trial [abstract]. 2nd Gothenburg Conference on Vesicoureteral Reflux; 2015 Jun 11-12; Goteborg, Sweden. 2015:12.

Oswald 2002 {published data only}

Oswald J, Riccabona M, Lusuadi L, Bartsch G, Radmayr C. Prospective comparison and 1-year follow-up of a single endoscopic subureteral polydimethylsiloxane versus dextranomer/hyaluronic acid copolymer injection for treatment of vesicoureteral reflux in children. *Urology* 2002;**60**(5):894-7. [MEDLINE: 12429323]

Pennesi 2008 {published data only (unpublished sought but not used)}

Pennesi M, Peratoner L, Bordugo A, Saccari A, Minisini S, Travan L. Uneffectiveness of antibiotic prophylaxis in children with vesicoureteral reflux. Preliminary data from a controlled study [abstract no: MP239]. *Nephrology Dialysis Transplantation* 2006;**21**(Suppl 4):iv378. [CENTRAL: CN-00602044]

Pennesi M, Peratoner L, Giacomini A, Travan L, Bordugo A. Uneffectiveness of antibiotic prophylaxis in children with VUR. Preliminary data from controlled study [abstract no: COD.OC2]. *Pediatric Nephrology* 2006;**21**(10):1502. [CENTRAL: CN-00602043]

* Pennesi M, Travan L, Peratoner L, Bordugo A, Cattaneo A, Ronfani L, et al. Is antibiotic prophylaxis in children with vesicoureteral reflux effective in preventing pyelonephritis and renal scars? A randomized, controlled trial. *Pediatrics* 2008;**121**(6):e1489-94. [MEDLINE: 18490378]

PRIVENT 2009 {published and unpublished data}

Craig J, Hodson E, Reynolds G, Carapetis J, McTaggart S, Irwig L, et al. A placebo-controlled randomised trial of antibiotics for the prevention of urinary tract infection in children [abstract no: SA-PO3072]. *Journal of the American Society of Nephrology* 2008;**19**(Abstracts Issue):808A.

Craig J, Simpson J, Lowe A, Reynolds G, McTaggart S, Hodson E, et al. A placebo-controlled randomized trial of antibiotics for the prevention of urinary tract infection in children [abstract no: O87]. *Nephrology* 2008;**13**(Suppl 3):A122. [CENTRAL: CN-00689116]

* Craig JC, Simpson JM, Williams GJ, Lowe A, Reynolds GJ, McTaggart SJ, et al. Antibiotic prophylaxis and recurrent urinary

tract infection in children.[Erratum appears in N Engl J Med. 2010 Apr 1;362(13):1250]. *New England Journal of Medicine* 2009;**361**(18):1748-59. [MEDLINE: 19864673]

Reddy 1997 {published data only}

Reddy PP, Evans MT, Hughes PA, Dangman B, Cooper J, Lepow ML, et al. Antimicrobial prophylaxis in children with vesico-ureteral reflux: a randomized prospective study of continuous therapy vs intermittent therapy vs surveillance [abstract]. *Pediatrics* 1997;**100**(3):555-6. [CENTRAL: CN-00427967]

RIVUR 2008 {published data only}

Carpenter MA, Hoberman A, Mattoo TK, Mathews R, Keren R, Chesney RW, et al. The RIVUR trial: profile and baseline clinical associations of children with vesicoureteral reflux. *Pediatrics* 2013;**132**(1):e34-45. [MEDLINE: 23753091]

Chesney RW, Carpenter MA, Moxey-Mims M, Nyberg L, Greenfield SP, Hoberman A, et al. Randomized Intervention for Children with Vesicoureteral Reflux (RIVUR): background commentary of RIVUR investigators. *Pediatrics* 2008;**122** Suppl 5:S233-9. [MEDLINE: 19018047]

Edmonson MB, Eickhoff JC. Weight gain and obesity in infants and young children exposed to prolonged antibiotic prophylaxis. *JAMA Pediatrics* 2017;**171**(2):150-6. [MEDLINE: 28027334]

Greenfield S, Carpenter M, Hoberman A, Mattoo T, Mathews R, Keren R, et al. The RIVUR trial: baseline demographic and clinical characteristics [abstract no: 641]. *Journal of Urology* 2013;**189**(4 Suppl):e262. [EMBASE: 71031583]

Greenfield SP, Carpenter MA, Chesney RW, Zerlin JM, Chow J. The RIVUR voiding cystourethrogram pilot study: experience with radiologic reading concordance. *Journal of Urology* 2012;**188**(4 Suppl):1608-12. [MEDLINE: 22910235]

Keren R. Pediatrics. RIVUR trial. Introduction. *Pediatrics* 2008;**122** Suppl 5:S231-2. [MEDLINE: 19086141]

Keren R, Carpenter M, Greenfield S, Hoberman A, Mathews R, Mattoo T, et al. Is antibiotic prophylaxis in children with vesicoureteral reflux effective in preventing pyelonephritis and renal scars? A randomized, controlled trial. *Pediatrics* 2008;**122**(6):1409-10. [MEDLINE: 19047267]

Keren R, Carpenter MA, Hoberman A, Shaikh N, Mattoo TK, Chesney RW, et al. Rationale and design issues of the Randomized Intervention for Children With Vesicoureteral Reflux (RIVUR) study. *Pediatrics* 2008;**122** Suppl 5:S240-50. [MEDLINE: 19018048]

Keren R, Shaikh N, Pohl H, Gravens-Mueller L, Ivanova A, Zaoutis L, et al. Risk factors for recurrent urinary tract infection and renal scarring. *Pediatrics* 2015;**136**(1):e13-21. [MEDLINE: 26055855]

Mattoo T, Carpenter M, Chesney R, Greenfield S, Hoberman A, Keren R, et al. Screening and enrolling children with vesicoureteral reflux: preliminary data from the RIVUR

trial [abstract no: PS3-SAT-522]. *Pediatric Nephrology* 2011;**26**(9):1729-30. [EMBASE: 70530954]

Mattoo TK. Randomized Intervention for Children with Vesicoureteral Reflux (RIVUR) Trial [abstract]. 2nd Gothenburg Conference on Vesicoureteral Reflux; 2015 Jun 11-12; Goteborg, Sweden. 2015:8.

Mattoo TK, Carpenter MA, Moxey-Mims M, Chesney RW, RIVUR Trial. The RIVUR trial: a factual interpretation of our data. *Pediatric Nephrology* 2015;**30**(5):707-12. [MEDLINE: 25558811]

Mattoo TK, Chesney RW, Greenfield SP, Hoberman A, Keren R, Mathews R, et al. Renal scarring in the Randomized Intervention for Children with Vesicoureteral Reflux (RIVUR) Trial. *Clinical Journal of the American Society of Nephrology: CJASN* 2016;**11**(1):54-61. [MEDLINE: 26555605]

Nelson CP, Hoberman A, Shaikh N, Keren R, Mathews R, Greenfield SP, et al. Antimicrobial resistance and urinary tract infection recurrence. *Pediatrics* 2016;**137**(4):e20152490. [MEDLINE: 26969273]

Perez-Gaxiola G. Antibiotic prophylaxis reduced symptomatic urinary tract infection in children with vesicoureteral reflux, but not scarring. *Archives of Disease in Childhood Education & Practice* 2015;**100**(1):52. [MEDLINE: 25073466]

RIVUR Trial Investigators, Hoberman A, Greenfield SP, Mattoo TK, Keren R, Mathews R, et al. Antimicrobial prophylaxis for children with vesicoureteral reflux. *New England Journal of Medicine* 2014;**370**(25):2367-76. [MEDLINE: 24795142]

Schaeffer AJ, Chow JS, Ivanova A, Cui G, Greenfield SP, Zerlin JM, et al. Variation in the level of detail in pediatric voiding cystourethrogram reports. *Journal of Pediatric Urology* 2017;**13**(3):257-62. [MEDLINE: 28277235]

Schaeffer AJ, Greenfield SP, Ivanova A, Cui G, Zerlin JM, Chow JS, et al. Reliability of grading of vesicoureteral reflux and other findings on voiding cystourethrography. *Journal of Pediatric Urology* 2017;**13**(2):192-8. [MEDLINE: 27666144]

Schwaderer AL, Wang H, Kim S, Kline JM, Liang D, Brophy PD, et al. Polymorphisms in alpha-defensin-Encoding DEFA1A3 associate with urinary tract infection risk in children with vesicoureteral reflux. *Journal of the American Society of Nephrology* 2016;**27**(10):3175-86. [MEDLINE: 26940096]

Seidman C, Lotan Y, Palmer L. Cost effectiveness of antimicrobial prophylaxis for children in the RIVUR trial [abstract]. *Journal of Urology* 2015;**193**(4 Suppl 1):e665. [EMBASE: 71859344]

Shaikh N, Hoberman A, Keren R, Gotman N, Docimo SG, Mathews R, et al. Recurrent urinary tract infections in children with bladder and bowel dysfunction. *Pediatrics* 2016;**137**(1):e20152082. [MEDLINE: 26647376]

Shaikh N, Hoberman A, Keren R, Ivanova A, Gotman N, Chesney RW, et al. Predictors of antimicrobial resistance among pathogens causing urinary tract infection in children. *Journal of Pediatrics* 2016;**171**:116-21. [MEDLINE: 26794472]

Shaikh N, Hoberman A, Keren R, Ivanova A, Ziessman HA, Cui G, et al. Utility of sedation for young children undergoing dimercaptosuccinic acid renal scans. *Pediatric Radiology* 2016;**46**(11):1573-8. [MEDLINE: 27287454]

Shaikh N, Mattoo TK, Keren R, Ivanova A, Cui G, Moxey-Mims M, et al. Early antibiotic treatment for pediatric febrile urinary tract infection and renal scarring. *JAMA Pediatrics* 2016;**170**(9):848-54. [MEDLINE: 27455161]

Williams G, Craig J. RIVUR trial offers confirmatory evidence for a small but real benefit of antibiotics for UTI prevention in children. *Evidence Based Medicine* 2014;**19**(6):229-30. [MEDLINE: 25189165]

Ziessman HA, Majd M. Importance of methodology on (99m)technetium dimercapto-succinic acid scintigraphic image quality: imaging pilot study for RIVUR (Randomized Intervention for Children With Vesicoureteral Reflux) multicenter investigation. *Journal of Urology* 2009;**182**(1):272-9. [MEDLINE: 19450818]

Roussey-Kesler 2008 {published and unpublished data}

De Cunto A, Pennesi M, Salierno P. Re: Antibiotic prophylaxis for the prevention of recurrent urinary tract infection in children with low grade vesicoureteral reflux: results from a prospective randomized study. *Journal of Urology* 2008;**180**(5):2258-9. [MEDLINE: 18804797]

Roussey-Kesler G, Gadjos V, Idres N, Horen B, Guyot C. Antibiotic prophylaxis for the prevention of urinary tract infection in children with low grade vesicoureteral reflux: results from a prospective randomized study [abstract no: 204 (FC)]. *Pediatric Nephrology* 2007;**22**(9):1450.

* Roussey-Kesler G, Gadjos V, Idres N, Horen B, Ichay L, Leclair MD, et al. Antibiotic prophylaxis for the prevention of recurrent urinary tract infection in children with low grade vesicoureteral reflux: results from a prospective randomized study. *Journal of Urology* 2008;**179**(2):674-9. [MEDLINE: 18082208]

Scott 1968 {published data only}

Scott JE, Stansfeld JM. Treatment of vesico-ureteric reflux in children. *Archives of Disease in Childhood* 1968;**43**(229):323-8. [MEDLINE: 5652710]

Smellie 2001 {published and unpublished data}

Barratt TM, Chantler C, Prescod NP, Woolf AS, Smellie JM. Randomised trial comparing medical and surgical treatment in children with bilateral vesicoureteric reflux and nephropathy [abstract]. Renal Association 50th Anniversary Meeting; 2000 Sep 27-29; St John College, Cambridge, UK. 2000.

Roseau E. Vesico-ureteral reflux and nephropathy in the child: medical or surgical treatment? [Reflux vesico-ureteral et nephropathie de l'enfant: traitement medical ou chirurgical?]. *Presse Medicale* 2001;**30**(23):1157-8. [MEDLINE: 11505836]

* Smellie JM, Barratt TM, Chantler C, Gordon I, Prescod NP, Ransley PG, et al. Medical versus surgical treatment in children with severe bilateral vesicoureteric reflux

and bilateral nephropathy: a randomised trial. *Lancet* 2001;**357**(9265):1329-33. [MEDLINE: 11343739]

Swedish Reflux Trial 2010 {published and unpublished data}

Brandstrom P, Esbjorner E, Herthelius M, Holmdahl G, Lackgren G, Neveus T, et al. The Swedish reflux trial in children: I. Study design and study population characteristics. [Erratum appears in *J Urol*. 2011 Jan;185(1):365]. *Journal of Urology* 2010;**184**(1):274-9. [MEDLINE: 20478580]

Brandstrom P, Esbjorner E, Herthelius M, Swerkersson S, Jodal U, Hansson S. The Swedish reflux trial in children: III. Urinary tract infection pattern. *Journal of Urology* 2010;**184**(1):286-91. [MEDLINE: 20488494]

Brandstrom P, Hansson S. Growth in children with dilating VUR - a follow up of the Swedish Reflux Trial [abstract no: P-SAT044]. *Pediatric Nephrology* 2013;**28**(8):1391. [EMBASE: 71127064]

Brandstrom P, Jodal U, Sillen U, Hansson S. The Swedish Reflux Trial: review of a randomized, controlled trial in children with dilating vesicoureteral reflux. *Journal of Pediatric Urology* 2011;**7**(6):594-600. [MEDLINE: 21807562]

Brandstrom P, Neveus T, Sixt R, Stokland E, Jodal U, Hansson S. The Swedish Reflux Trial in children: IV. Renal damage. *Journal of Urology* 2010;**184**(1):292-7. [MEDLINE: 20494369]

Brandström P. Swedish Reflux Trial – a follow-up [abstract]. 2nd Gothenburg Conference on Vesicoureteral Reflux; 2015 Jun 11-12; Goteborg, Sweden. 2015:13.

Holmdahl G, Brandstrom P, Lackgren G, Sillen U, Stokland E, Jodal U, et al. The Swedish Reflux Trial in children: II. Vesicoureteral reflux outcome. *Journal of Urology* 2010;**184**(1):280-5. [MEDLINE: 20488469]

Neveus T, Brandstrom P, Linner T, Jodal U, Hansson S. Parental experiences and preferences regarding the treatment of vesicoureteral reflux. *Scandinavian Journal of Urology & Nephrology* 2012;**46**(1):26-30. [MEDLINE: 22008041]

Sillen U, Brandstrom P, Jodal U, Holmdahl G, Sandin A, Sjoberg I, et al. The Swedish Reflux Trial in children: V. Bladder dysfunction. *Journal of Urology* 2010;**184**(1):298-304. [MEDLINE: 20488486]

Takvani 2015 {published data only}

Takvani A, Gokani C, Malaviya P. Vesicoureteral reflux - a prospective study of 11 years [abstract]. *European Urology Supplements* 2015;**14**(2):e505-e505a. [EMBASE: 71831431]

References to studies excluded from this review

Akin 2014 {published data only}

Akin M, Erginel B, Karadag CA, Yildiz A, Ozelik GS, Sever N, et al. A comparison of the double hydrodistention implantation technique (HIT) and the HIT with a polyacrylate/polyalcohol copolymer (PPC) for the endoscopic treatment of primary vesicoureteral reflux. *International Urology & Nephrology* 2014;**46**(11):2057-61. [MEDLINE: 24966099]

Bailey 1991 {published data only}

Bailey RR, Lynn KL, McRae CU, Smith AH. Prospective randomised study of the effect of nephrectomy on blood pressure of patients with unilateral reflux nephropathy. 2nd CJ Hodson Symposium on Reflux Nephropathy; 1991 Feb 26; Christchurch, NZ. 1991:37-40.

COBSG 1973 {published data only}

Sequelae of covert bacteriuria in schoolgirls. A four-year follow-up study. *Lancet* 1978;**311**(8070):889-93. [MEDLINE: 76841]

Aggarwal VK, Verrier Jones K, Asscher AW, Evans C, Williams LA. Covert bacteriuria: long term follow up. *Archives of Disease in Childhood* 1991;**66**(11):1284-6. [MEDLINE: 1755638]

Asscher AW, McLachlan MS, Jones RV, Meller S, Sussman M, Harrison S, et al. Screening for asymptomatic urinary-tract infection in schoolgirls. A two-centre feasibility study. *Lancet* 1973;**2**(7819):1-4. [MEDLINE: 4123292]

Coupris 1988 {published data only}

Coupris L. Antibiotic prophylaxis for surgery after vesico-ureteral reflux in children. *Drugs* 1988;**35 Suppl 2**:154-7. [MEDLINE: 3135164]

Kajbafzadeh 2010 {published data only}

Kajbafzadeh AM, Baradaran N, Sadeghi Z, Tourchi A, Saeedi P, Madani A, et al. Vesicoureteral reflux and primary bladder neck dysfunction in children: urodynamic evaluation and randomized, double-blind, clinical trial on effect of alpha-blocker therapy. *Journal of Urology* 2010;**184**(5):2128-33. [MEDLINE: 20850812]

Montini 2007 {published data only}

Hewitt IK, Tomasi L, Pavanello L, Maschio F, Molinari PP, Toffolo A, et al. Early treatment of acute pyelonephritis in children fails to reduce renal scarring [abstract no: SA-PO1100]. *Journal of the American Society of Nephrology* 2006;**17**(Abstracts):804A. [CENTRAL: CN-00653791]

Hewitt IK, Zucchetto P, Rigon L, Maschio F, Molinari PP, Tomasi L, et al. Early treatment of acute pyelonephritis in children fails to reduce renal scarring: data from the Italian Renal Infection Study Trials. *Pediatrics* 2008;**122**(3):486-90. [MEDLINE: 18762516]

La Scola C, De Mutiis C, Hewitt IK, Puccio G, Toffolo A, Zucchetto P, et al. Different guidelines for imaging after first UTI in febrile infants: yield, cost, and radiation. *Pediatrics* 2013;**131**(3):e665-71. [MEDLINE: 23439905]

Montini G, Murer L, Gobber D, Comacchio S, Toffolo A, Dall'Amico R, et al. Oral vs initial intravenous antibiotic treatment of urinary tract infections in children: a multicentre study [abstract no: W833]. *Nephrology Dialysis Transplantation* 2003;**18**(Suppl 4):816-7.

Montini G, Toffolo A, Zucchetto P, Dall'Amico R, Gobber D, Calderan A, et al. Antibiotic treatment for pyelonephritis in children: multicentre randomised controlled non-inferiority trial. *BMJ* 2007;**335**(7616):386-8. [MEDLINE: 17611232]

Montini G, Zucchetta P, Gobber D, Murer L, Rigon L, Toffolo A, et al. Usefulness of renal radiology and scintigraphy in children with a first episode of upper urinary tract infection [abstract no: W823]. *Nephrology Dialysis Transplantation* 2003;**18**(Suppl 4):813. [CENTRAL: CN-00446818]

Montini G, Zucchetta P, Tomasi L, Talenti E, Rigamonti W, Picco G, et al. Value of imaging studies after a first febrile urinary tract infection in young children: data from Italian renal infection study 1. *Pediatrics* 2009;**123**(2):e239-46. [MEDLINE: 19139086]

N0012116994 {published data only}

Ransley PG. Clinical trial of deflux for endoscopic correction of severe vesicoureteric reflux and the effects on improving bladder function outcome. National Research Register UK (www.nrr.nhs.uk/) (accessed prior to 10 September 2018).

NCBRG 1981 {published data only}

Covert bacteriuria in schoolgirls in Newcastle upon Tyne: a 5-year follow-up. Newcastle Covert Bacteriuria Research Group. *Archives of Disease in Childhood* 1981;**56**(8):585-92. [MEDLINE: 7271298]

Davison JM, Sprott MS, Selkon JB. The effect of covert bacteriuria in schoolgirls on renal function at 18 years and during pregnancy. *Lancet* 1984;**2**(8404):651-5. [MEDLINE: 6147691]

Scholtmeijer 1988 {published data only}

* Scholtmeijer RJ. Treatment of vesicoureteric reflux. Results of a prospective study. *British Journal of Urology* 1993;**71**(3):346-9. [MEDLINE: 8477320]

Scholtmeijer RJ. Treatment of vesicoureteric reflux: results after 3 years in a prospective study. *Child Nephrology & Urology* 1991;**11**(1):29-32. [MEDLINE: 1868479]

Scholtmeijer RJ, Griffiths DJ. Treatment of vesicoureteric reflux. Preliminary report of a prospective study. *British Journal of Urology* 1988;**61**(3):205-9. [MEDLINE: 3359122]

Uberos 2012 {published data only}

Fernandez-Puentes V, Uberos J, Rodriguez-Belmonte R, Nogueras-Ocana M, Blanca-Jover E, Narbona-Lopez E. Efficacy and safety profile of cranberry in infants and children with recurrent urinary tract infection [Eficacia y perfil de seguridad del arandano americano en lactantes y niños con infección urinaria recurrente]. *Anales de Pediatría* 2015;**82**(6):397-403. [MEDLINE: 25300782]

Uberos J, Nogueras-Ocana M, Fernandez-Puentes V, Rodriguez-Belmonte R, Narbona-Lopez E, Molina-Carballo A, et al. Cranberry syrup vs trimethoprim in the prophylaxis of recurrent urinary tract infections among children: a controlled trial. *Open Access Journal of Clinical Trials* 2012;**4**:31-8. [EMBASE: 2012351759]

Uberos J, Rodriguez-Belmonte R, Fernandez-Puentes V, Narbona-Lopez E, Molina-Carballo A, Munoz-Hoyos A. Cranberry syrup vs. trimethoprim in the prophylaxis of recurrent urinary infection: A double-blind randomized clinical trial [abstract

no: PP003]. *Acta Paediatrica* 2010;**99**(Suppl 462):48. [EMBASE: 70313001]

Additional references

Belman 1995

Belman AB. A perspective on vesicoureteral reflux. *Urologic Clinics of North America* 1995;**22**(1):139-50. [MEDLINE: 7855950]

Cohen 2005

Cohen AL, Rivara FP, Davis R, Christakis DA. Compliance with guidelines for the medical care of first urinary tract infections in infants: a population based study. *Pediatrics* 2005;**115**(6):1474-8. [MEDLINE: 15930206]

Frankenschmidt 1997

Frankenschmidt A, Katzenwadel A, Zimmerhackl LB, Sommerkamp H. Endoscopic treatment of reflux by subureteric collagen injection: critical review of 5 years' experience. *Journal of Endourology* 1997;**11**(5):343-8. [MEDLINE: 9355951]

GRADE 2008

Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;**336**(7650):924-6. [MEDLINE: 18436948]

Greenfield 1997

Greenfield SP, Ng M, Wan J. Experience with vesicoureteral reflux in children: clinical characteristics. *Journal of Urology* 1997;**158**(2):574-7. [MEDLINE: 9224365]

Hellstrom 1991

Hellstrom A, Hanson E, Hansonn S, Hjalmas K, Jodal U. Association between urinary symptoms at 7 years old and previous urinary tract infection. *Archives of Disease in Childhood* 1991;**66**(2):232-4. [MEDLINE: 2001110]

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [MEDLINE: 12958120]

Higgins 2011

Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Lau 1997

Lau J, Ioannidis JP, Schmid CH. Quantitative synthesis in systematic reviews. *Annals of Internal Medicine* 1997;**127**(9):820-6. [MEDLINE: 9382404]

Martinell 1996

Martinell J, Lidin-Janson G, Jagenburg R, Sivertsson R, Claesson I, Jodal U. Girls prone to urinary infections followed into adulthood. Indices of renal disease. *Pediatric Nephrology* 1996;**10**(2):139-42. [MEDLINE: 8703696]

Montini 2007b

Montini G, Toffolo A, Zucchetta P, Dall'Amico R, Gobber D, Calderan A, et al. Antibiotic treatment for pyelonephritis in children: multicentre randomised controlled non-inferiority trial. *BMJ* 2007;**335**(7616):386. [MEDLINE: 17611232]

Olbing 2003

Olbing H, Smellie JM, Jodal U, Lax H. New renal scars in children with severe VUR: a 10-year study of randomized treatment. *Pediatric Nephrology* 2003;**18**(11):1128-31. [MEDLINE: 14523634]

Panaretto 1999

Panaretto KS, Craig JC, Knight JF, Howman-Giles R, Sureshkumar P, Roy LP. Risk factors for recurrent urinary tract infection in preschool children. *Journal of Paediatrics & Child Health* 1999;**35**:454-9. [MEDLINE: 10571758]

Schünemann 2011a

Schünemann HJ, Oxman AD, Higgins JP, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results and 'Summary of findings' tables. In: Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Schünemann 2011b

Schünemann HJ, Oxman AD, Higgins JP, Deeks JJ, Glasziou P, Guyatt GH. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Smellie 1975

Smellie J, Edwards D, Hunter N, Normand IC, Prescod N. Vesico-ureteric reflux and renal scarring. *Kidney International - Supplement* 1975;**4**:S65-72. [MEDLINE: 1104981]

Smellie 1994

Smellie JM, Poulton A, Prescod NP. Retrospective study of children with renal scarring associated with reflux and urinary infection. *BMJ* 1994;**308**(6938):1193-6. [MEDLINE: 8180534]

Smellie 1998

Smellie JM, Prescod NP, Shaw PJ, Risdon RA, Bryant TN. Childhood reflux and urinary infection: a follow-up of 10-41

years in 226 adults. *Pediatric Nephrology* 1998;**12**(9):727-36. [MEDLINE: 9874316]

Wennerstrom 2000a

Wennerstrom M, Hansson S, Jodal U, Sixt R, Stokland E. Renal function 16 to 26 years after the first urinary tract infection in childhood. *Archives of Pediatrics & Adolescent Medicine* 2000;**154**(4):339-45. [MEDLINE: 10768669]

Wennerstrom 2000b

Wennerstrom M, Hansson S, Hedner T, Himmelmann A, Jodal U. Ambulatory blood pressure 16-26 years after the first urinary tract infection in childhood. *Journal of Hypertension* 2000;**18**(4):485-91. [MEDLINE: 10779101]

References to other published versions of this review

Hodson 2007

Hodson EM, Wheeler DM, Smith GH, Craig JC, Vimalachandra D. Interventions for primary vesicoureteric reflux. *Cochrane Database of Systematic Reviews* 2007, Issue 3. [DOI: [10.1002/14651858.CD001532.pub3](https://doi.org/10.1002/14651858.CD001532.pub3)]

Nagler 2011

Nagler EV, Williams G, Hodson EM, Craig JC. Interventions for primary vesicoureteric reflux. *Cochrane Database of Systematic Reviews* 2011, Issue 6. [DOI: [10.1002/14651858.CD001532.pub4](https://doi.org/10.1002/14651858.CD001532.pub4)]

Wheeler 1999

Wheeler D, Vimalachandra D, Roy P, Smith G, Craig JC. Interventions for primary vesicoureteric reflux. *Cochrane Database of Systematic Reviews* 1999, Issue 2. [DOI: [10.1002/14651858.CD001532](https://doi.org/10.1002/14651858.CD001532)]

Wheeler 2003

Wheeler D, Vimalachandra D, Hodson EM, Roy LP, Smith G, Craig JC. Antibiotics and surgery for vesicoureteric reflux: a meta-analysis of randomised controlled trials. *Archives of Disease in Childhood* 2003;**88**(8):688-94. [MEDLINE: 12876164]

Wheeler 2004

Wheeler DM, Vimalachandra D, Hodson EM, Smith GH, Craig JC. Interventions for primary vesicoureteric reflux. *Cochrane Database of Systematic Reviews* 2004, Issue 3. [DOI: [10.1002/14651858.CD001532.pub2](https://doi.org/10.1002/14651858.CD001532.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Ahn 2003

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: October 1996 to April 2002 Duration of follow-up: 12 months
Participants	<ul style="list-style-type: none"> Country: Korea

Interventions for primary vesicoureteric reflux (Review)

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Ahn 2003 (Continued)

- Setting; single centre (Ewha Womens University Mekdong Hospital)
- Children diagnosed with primary VUR which had persisted after 12 months of antibiotic prophylaxis
- Number: treatment group (59); control group (93)
- Mean age $\pm$ SD (years): treatment group (1.8 ± 1.47); control group (1.9 ± 1.62)
- Sex (M/F): not reported
- VUR grades: not reported
- Exclusion criteria: not reported

Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> • Anticholinergic agent oxybutynin: 0.2mg/kg twice daily for 12 months <p>Control group</p> <ul style="list-style-type: none"> • Placebo (no dose details provided) for 12 months
Outcomes	<ul style="list-style-type: none"> • Resolution of VUR at 12 months
Notes	<ul style="list-style-type: none"> • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Selective reporting (reporting bias)	High risk	Only reports VUR resolution, not symptomatic UTI
Other bias	Unclear risk	So few methodological details reported, not possible to exclude other biases

BIRSG 1983

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: July 1975 to December 1986 • Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> • Country: UK

Interventions for primary vesicoureteric reflux (Review)

BIRSG 1983 (Continued)

- Setting: single centre (recruitment from consultant paediatricians, general practitioners and emergency room of investigating centre)
- Children with primary VUR grade II with scarring or grade III, IV, V in absence of UTI within last 12 months
- Number (randomised/analysed): 179/161
 - Two years: group 1 (84); group 2 (77)
 - Five years: group 1 (53); group 2 (51)
- Age: < 15 years
- VUR grades; II (with scarring), III - V
- Exclusion criteria: VUR secondary to obstruction; completely duplicated ureter or neurogenic bladder; urinary calculus

Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> • Trimethoprim or nitrofurantoin <ul style="list-style-type: none"> ◦ Dose: 1 to 2 mg/kg ◦ Duration: antibiotics given for 2 years if resolution of VUR or 5 years <p>Treatment group 2</p> <ul style="list-style-type: none"> • Surgical reimplantation • Antibiotics <ul style="list-style-type: none"> ◦ Trimethoprim or nitrofurantoin (1 to 2 mg/kg) for 2 years
Outcomes	<ul style="list-style-type: none"> • UTI (culture positive) • Renal damage on IVP and DMSA scan • Renal function impairment, measured by GFR • Correction of VUR • Renal length on IVP
Notes	<ul style="list-style-type: none"> • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Central stratified block randomisation, method not clearly stated
Allocation concealment (selection bias)	Unclear risk	Allocation by opening next sealed envelope in batch of 8, so that last ones per batch could theoretically be predicted
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Only radiological outcomes were blinded
Incomplete outcome data (attrition bias) All outcomes	High risk	18/179 (10%) excluded from analysis at 2 years, due to failure to comply with study protocol; lost to follow-up and exclusions not distinguishable
Selective reporting (reporting bias)	High risk	Symptomatic UTI was not primary outcome since assessed by routine screening for bacteriuria or positive culture

Interventions for primary vesicoureteric reflux (Review)

BIRSG 1983 (Continued)

Other bias	Unclear risk	Patient flow from recruitment to randomisation not clearly stated in terms of numbers
------------	--------------	---

Capozza 2002

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: not reported Duration of follow-up: 12 months
Participants	<ul style="list-style-type: none"> Country: Italy Setting: University teaching hospital Children > 1 year with primary VUR grade II to IV for at least 6 months Number: group 1 (21); group 2 (40) Mean age $\pm$ SD (years): group 1 (3.87 ± 2.67); group 2 (4.13 ± 3.02) Sex (%M/F): group 1 (38.1/61.9); group 2 (40/60) VUR grades: II, III and IV Exclusion criteria: recurrent UTI; duplex systems; neurogenic bladder
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic: not reported (drugs and regimen determined on a case-by-case basis) <ul style="list-style-type: none"> Dose: not reported Duration: 1 year <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgery: subureteric implantation of Deflux Antibiotics: not reported (drugs and regimen determined on a case-by-case basis) <ul style="list-style-type: none"> Dose: not reported Duration: 1 month
Outcomes	<ul style="list-style-type: none"> UTI Renal damage on ultrasound at start and after 12 months Kidney function as GFR (change) Resolution of VUR
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	SAS software; 2:1 randomisation
Allocation concealment (selection bias)	Low risk	"blinded allocation"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Open-label study

Capozza 2002 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	High risk	Open-label study
Incomplete outcome data (attrition bias) All outcomes	High risk	One excluded after randomisation but before treatment start not included in analysis Eight exclusions after 6 months due to persistent VUR after endoscopic treatment - included in analysis
Selective reporting (reporting bias)	Unclear risk	UTI definition and method of collection: not reported
Other bias	Low risk	Patient flow from recruitment to randomisation stated

Craig 2002

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: not reported Duration of follow-up: 3 years
Participants	<ul style="list-style-type: none"> Country: Australia Setting: multicentre study (number of sites not reported) Children < 3 months with isolated VUR Number (randomised/analysed): 46/41 Age: < 3 months Sex (M/F): not reported VUR grades: I to V Exclusion criteria: not reported
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX <ul style="list-style-type: none"> Dose: 2/10 mg/kg/d in single daily dose Duration: 3 years <p>Control group</p> <ul style="list-style-type: none"> Matched placebo for 3 years <p>Co-interventions</p> <ul style="list-style-type: none"> Not reported
Outcomes	<ul style="list-style-type: none"> First UTI (definition - not reported) within 3 years New renal parenchymal abnormality (interpreted as new damage on DMSA) at 3 years
Notes	<ul style="list-style-type: none"> Only published as an abstract. Raw data obtained from authors Funding source: no industry funding

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Craig 2002 (Continued)

Random sequence generation (selection bias)	Low risk	Computer generated sequence generation with stratification according to centre, referral source, frequency of previous UTI, VUR status, age, sex, according to method of minimization
Allocation concealment (selection bias)	Low risk	Randomisation centrally by telephone at an independent trials centre
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding of all participants, healthcare providers
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding of outcome collectors and data-analysts
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up accounted for (5/46, 11%) 3 placebo, 2 antibiotic. Bias due to loss-to follow-up unlikely
Selective reporting (reporting bias)	Low risk	Primary outcome appropriate
Other bias	Low risk	Raw data available, no industry funding

Dite 2007

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from 2003 to 2006 Duration of follow-up: 11 to 24 months
Participants	<ul style="list-style-type: none"> Country: Czech Republic Setting: single centre, urology department of a university hospital Children with primary VUR grade III to V Number: treatment group (22); control group (22) Age range: 1 to 32 months VUR grade: III, IV and V Exclusion criteria: posterior urethral valves, decompensated lower UTI, para-urethral diverticulum, or any morphological anomaly of the lower urinary tract
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Surgery: subureteric implantation of Deflux Antibiotics: type not reported <ul style="list-style-type: none"> Dose: not reported Duration: 1 month <p>Control group</p> <ul style="list-style-type: none"> Antibiotics: type not reported <ul style="list-style-type: none"> Dose: not reported Duration: given for entire follow-up period
Outcomes	<ul style="list-style-type: none"> UTI (definition - not reported) Acute pyelonephritis (definition - not reported)

Interventions for primary vesicoureteric reflux (Review)

Dite 2007 (Continued)

- Correction of VUR

Notes

- Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and healthcare providers not blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	No exclusions; loss to follow-up was 0%
Selective reporting (reporting bias)	Unclear risk	Definition of UTI not reported
Other bias	Unclear risk	No information on recruitment, making it impossible to evaluate selection bias. No information on funding.

Dotis 2014

Methods

- Study design: parallel RCT
- Duration of study: 1 year
- Duration of follow-up: 1 year

Participants

- Country: Greece
- Setting: not reported
- Children with a history of recurrent UTI
- Number: treatment group (38); control group (38)
- Age range: 2 to 18 years
- Sex (M/F): 23/56
- VUR grades I and II
- Exclusion criteria; $\geq$ grade III VUR

Interventions

Treatment group

- Cranberry capsules: 2 capsules per day for 12 months

Control group

Dotis 2014 (Continued)

- Placebo capsules: no details reported

Outcomes	<ul style="list-style-type: none"> • UTI • Initiation of antibiotic therapy • Days on antimicrobial therapy • Side effects
Notes	<ul style="list-style-type: none"> • Abstract-only publication • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Selective reporting (reporting bias)	High risk	Issues around symptoms for UTI and renal parenchymal abnormalities were not reported
Other bias	Unclear risk	Too few methodological details reported to exclude other biases

Espino Hernandez 2012

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: 9 months • Duration of follow-up: average was 29 months
Participants	<ul style="list-style-type: none"> • Country: Spain • Setting: multicentre; paediatric renal units (14 sites) • Children with primary VUR • Number; 28 • Age: not reported • Sex (M/F): not reported • VUR grades; III to V • Exclusion criteria: not reported
Interventions	Treatment group

Interventions for primary vesicoureteric reflux (Review)

Espino Hernandez 2012 (Continued)

- Prophylactic treatment: no dose details

Control group

- No treatment

Outcomes	<ul style="list-style-type: none"> • Pyelonephritis (symptomatic UTI) • New scars • Length of scars • Kidney function
Notes	<ul style="list-style-type: none"> • Abstract only, few methodological details • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to permit judgement
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No denominators reported
Selective reporting (reporting bias)	High risk	More outcomes collected than reported
Other bias	Unclear risk	Too few methodological details reported to exclude other biases

Frey 1997

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: recruitment period 1995 to 1997 • Duration of follow-up: 3 months
Participants	<ul style="list-style-type: none"> • Country: Switzerland • Setting: Single centre university hospital study • Children with primary VUR grades I to III • Number: 18 (total analysed unclear due to switch from children to ureters as unit of analysis) • Age: 1.1 to 10.7 years (average 4.6 years) • Sex (M/F): 10/8

Frey 1997 (Continued)

- Exclusion criteria: not reported

Interventions	Treatment group 1 <ul style="list-style-type: none"> • Surgery: Subureteric implantation of GAX 65 Treatment group 2 <ul style="list-style-type: none"> • Surgery: Subureteric implantation of GAX 35
Outcomes	<ul style="list-style-type: none"> • Correction of VUR • Recurrence of VUR
Notes	<ul style="list-style-type: none"> • Data on ureters not patients presented • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Low risk	Centrally prepared injections, coding disclosed post-analysis
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Coded syringes, so blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Coding was only disclosed after analysis of results
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Switch from randomised patients to number of ureters
Selective reporting (reporting bias)	High risk	Primary outcome not appropriate: > surrogate outcome grade of VUR instead of clinical outcome of UTI
Other bias	Unclear risk	Selection bias: patient flow from recruitment to randomisation not reported

Garcia-Aparicio 2013

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: recruitment April 2002 to June 2004 • Duration of follow-up: 5 years
Participants	<ul style="list-style-type: none"> • Country: Spain • Setting: university hospital • Children > 1 year who needed surgery for VUR due to recurrent UTI or impaired function or 2 years antibiotic prophylaxis • Number: 41 • Age: average age 55.1 months in the endoscopic group, 58.9 months in the reimplantation group

Interventions for primary vesicoureteric reflux (Review)

Garcia-Aparicio 2013 (Continued)

- Sex (M/F): 13/28
- VUR grades; II, III and IV
- Exclusion criteria: ESKD; poor kidney function; solitary kidney; complex uropathy; neurogenic bladder

Interventions	Treatment group 1 <ul style="list-style-type: none"> • Reimplantation surgery using Cohens technique Treatment group 2 <ul style="list-style-type: none"> • Endoscopic injection of Deflux • Antibiotics (name of antibiotic and dose not reported)
Outcomes	<ul style="list-style-type: none"> • Resolution of VUR at 6 months • Resolution of VUR after 1 injection • Resolution of VUR at 1 year • Resolution of VUR as 5 years • Complications • Febrile UTI
Notes	<ul style="list-style-type: none"> • Funding source: none

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Used randomiser research tool
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	Units are ureters in some outcomes and impossible to determine number of children
Selective reporting (reporting bias)	High risk	Some outcomes only reported in units of ureter
Other bias	Unclear risk	Poorly reported methods, cannot exclude other biases

Garin 2006

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: recruitment period December 1998 to December 2003 • Follow-up: 1 year follow-up
---------	---

Interventions for primary vesicoureteric reflux (Review)

Garin 2006 (Continued)

Participants	<ul style="list-style-type: none"> Countries: Chile, Spain, USA Setting: international multicentre study Children with acute pyelonephritis (fever 38.5°C, pyuria, > 100,000 colonies/mL) defined as focal/diffuse areas of decreased uptake on DMSA scan performed 2 to 7 days after UTI diagnosis Number: 218; 113 had VUR; 105 had no VUR <ul style="list-style-type: none"> VUR patients: treatment group (55); control group (58) Median age (range): 2 years (3 months to 12 years) Sex (M/F): 22/91 (VUR only) Exclusion criteria: VUR grade IV or V; neurogenic bladder; posterior urethral valves; urinary diversion; bladder diverticulum; ureterocoele; kidney failure; pregnancy
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX (1 to 2 mg/kg/d trimethoprim) or nitrofurantoin (1.5 mg/kg/d) Duration: 12 months <p>Control group</p> <ul style="list-style-type: none"> No treatment <p>Co-interventions</p> <ul style="list-style-type: none"> Not reported
Outcomes	<ul style="list-style-type: none"> Recurrence of UTI Type of recurrence: cystitis (definition not provided) or pyelonephritis Development of renal scars on DMSA scintigraphy
Notes	<ul style="list-style-type: none"> Loss to follow-up and exclusions after randomisation excluded from analysis Study included children with or without VUR stratified before randomisation. Only patients with VUR included Urine screened 3 monthly Funding source: "no financial relationships relevant to this article..."

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomisation stratified for VUR through simple randomisation, but method not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	<p>No clear distinction between loss to follow-up and exclusion</p> <p>Loss to follow-up: 9% of 236 children with or without VUR did not complete follow-up and were excluded from analysis</p>

Garin 2006 (Continued)

Selective reporting (re-reporting bias)	High risk	Positive culture is stated as primary outcome instead of symptomatic UTI
Other bias	Unclear risk	Patent flow from recruitment to randomisation not clearly reported

Gucuk 2013

Methods	<ul style="list-style-type: none"> Study design; parallel RCT Duration of study; not reported Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> Country: Turkey Setting: single centre; Urology Dept of University hospital Boys < 3 years with previous febrile UTI with and without VUR Number (VUR group only): treatment group 1 (45); treatment group 2 (46) Mean age $\pm$ SD (months): treatment group 1 (19.77 $\pm$ 9.12); treatment group 2 (19.10 $\pm$ 9.4) VUR grades: I, II and III Exclusion criteria: recent antibiotics; neurogenic bladder; bladder diverticula; ureterocoele; grade IV or V VUR; allergy to sulphonamide
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Circumcision Cotrimoxazole: 2/10 mg/kg/d for 1 year <p>Treatment group 2</p> <ul style="list-style-type: none"> Cotrimoxazole: 2/10 mg/kg/d for 1 year
Outcomes	<ul style="list-style-type: none"> Periurethral colonisation Repeat positive urine culture Repeat symptomatic UTI Adverse effects DMSA changes
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to permit judgement
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias)	Unclear risk	Insufficient information to permit judgement

Interventions for primary vesicoureteric reflux (Review)

Gucuk 2013 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Cannot determine whether dropouts are included in the analysis
Selective reporting (reporting bias)	High risk	Focus is periurethral colonisation, less complete reporting for other outcomes
Other bias	Unclear risk	Poorly reported methodological details mean other biases cannot be excluded

Habeeb Abid 2015

Methods	<ul style="list-style-type: none"> Study design; parallel RCT Duration of study: not reported Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> Country: Iraq Setting: not reported Children with VUR Number: treatment group 1 (40); treatment group 2 (40) Mean age $\pm$ SD: 4 years $\pm$ 3.4 VUR grades: II, III, IV, V Exclusion criteria: not reported
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Endoscopic injection of Dexcel; 1 year follow-up <p>Treatment group 2</p> <ul style="list-style-type: none"> Endoscopic injection of Vantris; 1 year follow-up
Outcomes	<ul style="list-style-type: none"> Resolution of VUR Fever UTI Ureteral obstruction
Notes	<ul style="list-style-type: none"> Abstract-only publication Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to permit judgement
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias)	Unclear risk	Insufficient information to permit judgement

Interventions for primary vesicoureteric reflux (Review)

Habeeb Abid 2015 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	Clear missing data
Selective reporting (reporting bias)	High risk	Focus on resolution of VUR with few details of other outcomes
Other bias	Unclear risk	Poor reporting of methodology means other biases cannot be excluded

Hari 2015

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: December 2006 to January 2012 Duration of follow-up: 12 months
Participants	<ul style="list-style-type: none"> Country: India Setting: single centre; tertiary care hospital Children < 12 years old with VUR diagnosed after febrile UTI Mean age $\pm$ SD (years): treatment group (5.7 ± 3.2); control group (4.8 ± 3.1) Sex (M/F): not reported VUR grades: I to IV Exclusion criteria; children aged < 1 year; grade V VUR or VUR secondary to urinary tract obstruction; including posterior urethral valves; neurogenic bladder and primary megaureter; history of voiding dysfunction; drug sensitivity to sulphonamides; eGFR < 30 mL/min/1.73 m²
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Cotrimoxazole: TMP/SMX (2/10 mg/kg/d) for 1 year <p>Control group</p> <ul style="list-style-type: none"> Placebo for 1 year
Outcomes	<ul style="list-style-type: none"> Repeat UTI Febrile UTI Repeat positive culture Days to first UTI Hospitalisation for UTI Changes on DMSA Adverse events Microbial resistance
Notes	<ul style="list-style-type: none"> Funding source: Indian Council of Medical Research

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Hari 2015 (Continued)

Random sequence generation (selection bias)	Low risk	Computer generated, permuted blocks
Allocation concealment (selection bias)	Low risk	Opaque sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding specifically stated
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding specified
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants accounted for; 93 started, 85 completed 1 year
Selective reporting (reporting bias)	Low risk	Appropriate outcomes
Other bias	Low risk	Good reporting of good quality methods

Holland 1982

Methods	<ul style="list-style-type: none"> Study design: parallel RCT; allocation to a third arm when patient was deemed ineligible for surgery by urologist or parent Recruitment time-frame: not reported Duration of follow-up 5 months to 3 years with median of 1 year and 5 months
Participants	<ul style="list-style-type: none"> Country: USA Setting: single centre Children with primary VUR grades II to IV Number: treatment group 1 (5); treatment group 2 (5) Mean age (range): 4.75 years (2 months to 10 years) Exclusion criteria: secondary VUR; hypertension; kidney dysfunction
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX (1/5 mg/lb) or nitrofurantoin (1 mg/lb) <ul style="list-style-type: none"> Duration: mean 17 months (range 5 to 36 months) <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgical reimplantation: technique not reported Antibiotic: TMP/SMX (1 / 5 mg/lb) or nitrofurantoin (1 mg/lb) <ul style="list-style-type: none"> Duration: mean 17 months (range 5 to 36 months)
Outcomes	<ul style="list-style-type: none"> UTI: culture positive Renal damage on IVP Adverse effects of antibiotics
Notes	<ul style="list-style-type: none"> Monthly screening by urinalysis the first 6 months, bimonthly screening after that

Holland 1982 (Continued)

- Funding source: "This study was supported by a grant from the Burroughs Wellcome Company, Research Triangle Park, North Carolina. Trimethoprim-sulfamethoxazole and nitrofurantoin were donated by the Burroughs Wellcome Company"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	High risk	Patients and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up; exclusions accounted for; exclusions after randomisation not analysed
Selective reporting (reporting bias)	Unclear risk	Positive urine culture as primary outcome, not clear whether symptoms were required
Other bias	High risk	Concerns about selection bias being introduced by providing alternative non-randomised plan for those after enrolment but excluded by urologist or parents for surgery Patient flow from recruitment to randomisation not clear

IRS EUR 1981-2003

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study recruitment: not reported Duration of follow-up: 10 years
Participants	Inclusion criteria <ul style="list-style-type: none"> Setting: Multicentre international study, university teaching hospitals Countries: Belgium, Finland, Germany, Sweden Children with primary VUR grades III to IV Number: treatment group 1 (155); treatment group 2 (151) Age range: 6 days to 11 years Sex (M/F): treatment group 1 (37/118); treatment group 2 (36/115) Exclusion criteria: primary VUR grades I and II; major urinary tract abnormality; previous urinary tract surgery; kidney dysfunction
Interventions	Treatment group 1

IRS EUR 1981-2003 (Continued)

- Antibiotic: nitrofurantoin or trimethoprim
 - Dose: 1 to 2 mg/kg
 - Duration: continued till resolution of VUR or 5 years

Treatment group 2

- Surgery: PL, Cohen, LG
- Antibiotics: continued for 6 months

Outcomes	<ul style="list-style-type: none"> • UTI: culture positive • Renal damage on IVP and DMSA scan • Obstruction postoperatively • Resolution of VUR • Renal length on IVP
Notes	<ul style="list-style-type: none"> • No intention-to-treat with loss to follow-up and exclusions after randomisation not included in analysis • Funding source: "Supported by VW-Foundation (Grants AZ 35 807, AZ 1/37 504) until 1985 and thereafter by the Bundesministerium für Forschung und Technologie (Grant 07068343)"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random allocation by a computer program
Allocation concealment (selection bias)	Low risk	At the coordinating centre
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Radiologists were blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up at 5 years
Selective reporting (reporting bias)	Low risk	Primary outcome appropriate, secondary outcomes detailed
Other bias	Low risk	Patient flow from recruitment to randomisation clearly stated

IRS USA 1992

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study recruitment: not reported • Duration of follow-up: 5 years
Participants	<ul style="list-style-type: none"> • Country: USA

Interventions for primary vesicoureteric reflux (Review)

IRS USA 1992 (Continued)

- Setting: multicentre; university teaching hospitals
- Children with primary VUR grade III to IV
- Number: 142 (132 analysed)
- Age: < 10 years
- Sex (M/F): 68/64
- Exclusion criteria: major urinary tract abnormality; previous urinary tract surgery; kidney dysfunction

Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> • Antibiotics: nitrofurantoin or trimethoprim • Dose: 1 to 2 mg/kg • Duration: given till resolution of VUR or 5 years <p>Treatment group 2</p> <ul style="list-style-type: none"> • Surgery: PL, Cohen, or other reimplantation • Antibiotics given for 6 months
Outcomes	<ul style="list-style-type: none"> • UTI: culture positive • Renal damage on IVP • Resolution of VUR • Renal area on IVP
Notes	<ul style="list-style-type: none"> • No intention to treat • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation blocking stratified system, stratified for each centre as to sex, age group, and presence or absence of pre-existing renal scar
Allocation concealment (selection bias)	Low risk	Allocation by sealed envelope
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Radiologists were blinded to radiological outcomes
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not clear when participants were excluded from the study, prior or after randomisation and whether they were included in analysis
Selective reporting (reporting bias)	Unclear risk	Positive urine culture as primary outcome, not clear whether symptoms were required
Other bias	Unclear risk	Patent flow from recruitment to randomisation clearly stated but not clear when excluded participants left the study, or how they switched groups, hereby potentially creating selection bias

Kojima 2012

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study recruitment: not reported Duration of follow-up: 3 to 4 months
Participants	<ul style="list-style-type: none"> Country: Japan Setting: single centre, Nagoya City University Hospital Children with VUR and breakthrough UTI; high grade VUR; progressive renal scarring Number: treatment group 1 (15); treatment group 2 (15) Mean age $\pm$ SD (months): treatment group 1 (54.3 ± 26.7); treatment group 2 (66.9 ± 64.1) Sex (M/F): treatment group 1 (9/6); treatment group 2 (8/7) VUR grades; I to V Exclusion criteria: not reported
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Surgery for ureteral reimplantation with advancement <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgery for ureteral reimplantation without advancement
Outcomes	<ul style="list-style-type: none"> VUR resolution at 4 months UTI Perioperative complications
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to permit judgement
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	Outcome data is not comprehensive
Selective reporting (reporting bias)	Unclear risk	Outcome focus is resolution of VUR not a clinical outcome
Other bias	Unclear risk	Unable to exclude other biases, methodology not well reported

Lee 2007a

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from 2002 to 2006 Duration of follow-up: 1year follow-up with intention-to-treat analysis
Participants	<ul style="list-style-type: none"> Country: South Korea Setting: single centre; university teaching hospital Children, VUR status (not clearly stated): persistent primary VUR grade I to IV (method of diagnosis not reported) after 1 year of prophylactic antibiotics with TMP/SMX Number: treatment group 1 (60); treatment group 2 (60) Mean age $\pm$ SD (months): treatment group 1 (21 ± 11.4); treatment group 2 (19 ± 12.1) Sex (M/F): treatment group 1 (45/15); treatment group 2 (44/16) Exclusion criteria: secondary VUR
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX <ul style="list-style-type: none"> Dose: 2/10 mg/kg/d single daily dose Duration: presumed for one year <p>Treatment group 2</p> <ul style="list-style-type: none"> Lactobacillus acidophilus (antibio300 Hanwa Co. Korea) <ul style="list-style-type: none"> Dose: 1.0×10^8 CFU/g ATCC 4356) twice a day Duration: presumed for one year
Outcomes	<ul style="list-style-type: none"> Number of recurrent symptomatic febrile UTI Number of recurrent symptomatic afebrile UTI Number of children with new renal scar Number of recurrent symptomatic UTI caused by <i>E coli</i> resistant to TMP/SMX
Notes	<ul style="list-style-type: none"> No exclusions after randomisation but before intervention UTI defined as 10^5 CFU/L in supra-pubic aspirated urine culture, or $> 10^8$ CFU/L in clean catch urine culture in toilet trained children Renal scar defined as 'Positive' DMSA scan taken at 3 to 6 months after recurrent DMSA + UTI Raw data obtained from authors Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Stratified randomisation according to VUR status with subsequent alternate allocation according to the order of enrolment
Allocation concealment (selection bias)	High risk	Investigators and parents were aware of treatment allocation. The investigator openly communicated there was less evidence for the preventive effect of probiotics versus antibiotics up until the study
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not blinded

Lee 2007a (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Data analyst was blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No exclusions, loss to follow-up: 4%
Selective reporting (reporting bias)	Low risk	Primary outcome appropriate
Other bias	Low risk	Patient flow from recruitment to randomisation clear

Lee 2015

Methods	<ul style="list-style-type: none"> Study design; parallel RCT Duration of study recruitment: not reported Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> Country: South Korea Setting: single centre; university hospital Infants diagnosed with VUR after 1st febrile UTI Number: treatment group 1 (64); treatment group 2 (64) Mean age $\pm$ SD (months): treatment group 1 (5.9 ± 5.5); treatment group 2 (4.7 ± 3.6) Sex (M/F): treatment group 1 (44/20); treatment group 2 (38/26) VUR grades: I to V Exclusion criteria; congenital renal hypodysplasia; secondary VUR
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Cotrimoxazole: TMP/SMX 2/10 mg/kg/d, for 1 year <p>Treatment group 2</p> <ul style="list-style-type: none"> Probiotic Lactobacillus acidophilus: 1×10^8 CFU/g twice daily, for 1 year
Outcomes	<ul style="list-style-type: none"> Repeat UTI Causative organism Microbial resistance Resolution of VUR DMSA abnormalities
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Stratified randomisation suggests good quality methods
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement

Lee 2015 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Open study, no blinding
Blinding of outcome assessment (detection bias) All outcomes	High risk	Open study, no blinding
Incomplete outcome data (attrition bias) All outcomes	Low risk	Data accounts for all children
Selective reporting (reporting bias)	Low risk	Appropriate outcomes
Other bias	Unclear risk	Some methodology details are missing

Mohseni 2013

Methods	<ul style="list-style-type: none"> Study design; parallel RCT Duration of study: March 2007 to April 2011 Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> Country: Iran Setting: single centre; paediatric medical centre Children with VUR and 4 or more UTIs in year prior to study Number (randomised/analysed): treatment group 1 (53/41); treatment group 2 (53/41) Mean age $\pm$ SD (years): treatment group 1 (8.0 ± 3.0); treatment group 2 (8.3 ± 3.1) Sex (M/F): treatment group 1 (14/30); treatment group 2 (15/26) VUR grades: I to V Exclusion criteria: history of secondary VUR; bilateral VUR and urogenital abnormalities; urogenital anomalies; ano-rectal malformations; neuropathic bladder; ureterocoele; concomitant upper real tract anomalies; PUJ obstruction; megaureter
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Nitrofurantoin: 1 mg/kg/d for 2 years <p>Treatment group 2</p> <ul style="list-style-type: none"> Probiotics: Lactobacillus acidophilus and Bifidobacterium lactis <ul style="list-style-type: none"> Dose 0.25 mL/kg of 10^7 CFU/mL Nitrofurantoin: 1 mg/kg/night for 2 years
Outcomes	<ul style="list-style-type: none"> Repeat UTI DMSA anomalies
Notes	<ul style="list-style-type: none"> Funding source: "Yoghurt containing Lactobacillus acidophilus and Bifidobacterium lactis was provided by the Research and Development Department of Iran Dairy Industries Co. (Pegah)."

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Interventions for primary vesicoureteric reflux (Review)

Mohseni 2013 (Continued)

Random sequence generation (selection bias)	Low risk	Random digit tables, computer generated, stratification for sex, age and grade of VUR
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	Appears complete, denominators reported
Selective reporting (reporting bias)	Low risk	Appropriate outcomes
Other bias	High risk	Very select group of patients, > 4 UTIs in past 1 year

Montini 2008

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: May 2000 to August 2006 Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> Country: Italy Setting: multicentre (22 centres) Children aged 2 months to 7 years; with or without primary non-severe VUR (I to III); normal kidney function after first febrile UTI Number: treatment group (211); control group (127) Mean age $\pm$ SD (months): treatment group (14.7 ± 15.5); control group (14.7 ± 15.5) Sex (M/F): treatment group (67/144); control group (37/90) Exclusion criteria: complex urologic malformations; severe kidney damage defined as relative function on 1 kidney of < 30% on DMSA
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Antibiotic: cotrimoxazole (15 mg/kg/d) or amoxicillin clavulanic acid (15 mg/kg/d) Duration: 1 year <p>Control group</p> <ul style="list-style-type: none"> No treatment
Outcomes	<ul style="list-style-type: none"> Recurrent febrile UTI Repeat positive urine cultures Children with new renal scar Repeat positive urine cultures caused by bacteria resistant to TMP/SMX (all patients) Adverse events (all patients)

Interventions for primary vesicoureteric reflux (Review)

Montini 2008 (Continued)

- Notes
- Positive urine culture defined as $>10^5$ CFU/mL of 1 micro-organism in 2 consecutive samples
 - Raw data requested from authors
 - Intention-to-treat analysis
 - Funding source: "Region of Veneto (research project 40/01) and association Il Sogno di Stefano (Stephen's Dream)"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated, stratified block design by centre, gender, and clinical group on the basis of presence or absence of VUR end parenchymal localization of the acute UTI.
Allocation concealment (selection bias)	Low risk	Distribution of codes in opaque envelopes in first part, later centrally
Blinding of participants and personnel (performance bias) All outcomes	High risk	Open-label study
Blinding of outcome assessment (detection bias) All outcomes	High risk	Open-label study
Incomplete outcome data (attrition bias) All outcomes	High risk	Not quite sure of total numbers, and % of VUR with new renal scar that had VUR not clear. Loss-to follow-up: 26/338 (8%) but included in analysis in best and worst case scenario
Selective reporting (reporting bias)	Low risk	Primary outcomes appropriate
Other bias	Unclear risk	Funding not commercial, patient flow from recruitment to randomisation not clearly stated

Moore 2014

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: recruitment April 2005 to June 2011 • Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> • Country: Canada • Setting: single centre • Children with symptomatic VUR • Number: treatment group 1 (122); treatment group 2 (133) • Median age (months): treatment group 1 (58); treatment group 2 (39) • Sex (M/F): treatment group 1 (21/101); treatment group 2 (31/102) • VUR grades: I to V • Exclusion criteria: previous endoscopic treatment of VUR, active infection at injection, untreated elimination dysfunction defined by a 3-day diary and validated questionnaires (Dysfunctional Voiding Symptom Score,11 Rome III12); absence of follow-up imaging

Moore 2014 (Continued)

Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Macroplastique in endoscopic treatment: 1 mL median, 0.5 to 2.6 mL, follow-up at 1 year <p>Treatment group 2</p> <ul style="list-style-type: none"> Deflux as endoscopic agent: 1 mL median 0.5 to 2.8 mL range, follow-up at 1 year
Outcomes	<ul style="list-style-type: none"> Resolution of VUR Adverse events
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Alternate patients, not good quality randomisation technique
Allocation concealment (selection bias)	High risk	Easy to subvert alternate choices
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	Poor reporting
Selective reporting (reporting bias)	High risk	Several appropriate outcomes missing
Other bias	Unclear risk	Other biases cannot be excluded due to poor method reporting

Morris 1991

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment between November 1983 and December 1987 Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> Country: New Zealand Setting: single centre Children with primary VUR grade III to IV Number: 138 (118 analysed) Age: 6 months to 10 years Exclusion criteria: major urological abnormality
Interventions	Treatment group 1

Interventions for primary vesicoureteric reflux (Review)

Morris 1991 (Continued)

- Antibiotic: not reported
 - Dose: not reported
 - Duration: 2 years

Treatment group 2

- Surgery: Cohen reimplantation
- Antibiotics for 3 months

Outcomes	<ul style="list-style-type: none"> • UTI: culture positive • GFR • Resolution of VUR
Notes	<ul style="list-style-type: none"> • Both loss to follow-up and exclusions after randomisation excluded from analysis • Funding source: "This study was supported by three project grants from the Health Research Council of New Zealand"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Stated that patients were stratified (stratified block randomisation) and then randomly assigned to treatment, but method not clear
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	High risk	Patients and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	No intention to treat; loss to follow-up: 20/138 (14%)
Selective reporting (reporting bias)	High risk	Only published in conference proceedings. GFR evolution reported, but renal scarring and UTI not whereas stated in methods
Other bias	Unclear risk	Insufficient detail about patient flow from recruitment to randomisation

Nordenstrom 2016

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT • Duration of study: Recruitment 2004 to 2014 • Follow-up; 1 year
Participants	<ul style="list-style-type: none"> • Country: Sweden • Setting: multicentre (21 sites) • Infants < 8 months with VUR grades 4-5 • Number: treatment group 1 (38); treatment group 2 (39)

Interventions for primary vesicoureteric reflux (Review)

Nordenstrom 2016 (Continued)

- Mean age $\pm$ SD (months): treatment group 1 (1.7 ± 1.8); treatment group 2 (1.6 ± 1.4)
- Sex (M/F): treatment group 1 (26/12); treatment group 2 (29/10)
- Exclusion criteria: kidney function $< 40\%$ of expected; a split function of $< 15\%$ in the refluxing kidney; vesicoureteral obstruction; other urological malformations (except duplication); neurogenic bladder dysfunction; suspected parental non-compliance

Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> • Endoscopic treatment: Deflux • Antibiotics until resolution of VUR <p>Treatment group 2</p> <ul style="list-style-type: none"> • Antibiotics: most cases trimethoprim 0.5 to 1 mg/kg once daily as single drug or combined with sul-famethoxazole. Optional agents were nitrofurantoin 1 mg/kg, cefadroxil 5 mg/kg, and ciprofloxacin 1 mg/kg
Outcomes	<ul style="list-style-type: none"> • Resolution of VUR • Febrile UTI • Adverse events
Notes	<ul style="list-style-type: none"> • Funding source: "This study was funded by regional university grants from Sahlgrenska University Hospital and by grants from The Freemasons' Orphanage Foundation in Göteborg, Sweden"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	All study subjects were matched for sex, presentation, grade of VUR, DMSA, abnormalities, bladder size, duplication, and centre by a computerized minimization procedure
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	High risk	Two patients missing from analysis of febrile UTIs, no explanation
Selective reporting (reporting bias)	Low risk	Included febrile UTI, but didn't report symptomatic UTI, abnormal DMSA, microbial resistance within treatment arms
Other bias	Unclear risk	Patient selection was very unclear

Oswald 2002

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT
---------	--

Interventions for primary vesicoureteric reflux (Review)

Oswald 2002 (Continued)

- Time frame: recruitment from January 2000 to June 2001
- Follow-up: 1 year

Participants	<ul style="list-style-type: none"> • Country: Austria • Setting: single centre, university urological department • Children with primary VUR grades II to IV • Number: treatment group 1 (34); treatment group 2 (38) • Mean age (months): treatment group 1 (33); treatment group 2 (36) • Sex (M/F): treatment group 1 (10/24); treatment group 2 (3/32) • Exclusion criteria: duplex systems; failed surgical reimplantation; neurogenic bladder; voiding dysfunction
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> • Subureteric injection of Macroplastique <p>Treatment group 2</p> <ul style="list-style-type: none"> • Subureteric injection of Deflux
Outcomes	<ul style="list-style-type: none"> • Correction of VUR • UTI • Adverse effects
Notes	<ul style="list-style-type: none"> • Data provided according to ureters for VUR correction. Total analysed unclear due to switch from children to ureters as unit of analysis • Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	High risk	No blinding of the healthcare providers
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Ureters were reported instead of patients, making it difficult to follow; intention-to-treat analysis
Selective reporting (reporting bias)	Unclear risk	Primary outcome not appropriate: grade of VUR as surrogate outcome instead of a clinical outcome UTI; VCUG not available (under progress) for all patients at 12 months
Other bias	Unclear risk	Patent flow from recruitment to randomisation: not reported; representativeness questionable

Pennesi 2008

Methods	<ul style="list-style-type: none"> Study design: parallel group RCT Duration of study: recruitment from November 1999 to March 2003 Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> Country: Italy Setting: multicentre (7 sites) Children with first episode of acute pyelonephritis; VUR grades II to IV Number: treatment group (50); control group (50) Mean age $\pm$ SD (months): treatment group (9 ± 5.9); control group (8.3 ± 5.4) SEX (M/F): treatment group (24/26); control group (24/26) Exclusion criteria: VUR grade I and V; previous pyelonephritis; recurrence pyelonephritis before first DMSA scan if positive for renal scars
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX <ul style="list-style-type: none"> Dose: 1 to 2/5 to 12 mg/kg single daily dose Duration: 1 year <p>Control group</p> <ul style="list-style-type: none"> No treatment
Outcomes	<ul style="list-style-type: none"> Repeat acute pyelonephritis by 2 years Worse renal damage on DMSA after 2 years Development of new renal damage on DMSA after 2 years Persistent VUR Repeat acute pyelonephritis caused by bacteria resistant to TMP/SMX
Notes	<ul style="list-style-type: none"> Urine collected by clean catch unless patient was septic, in which case a bladder tap was performed Standard outcome assessment Funding source: "This study was funded by the Italian Ministry of Health (research grant 35/00). The sponsor had no role in design, data collection, analysis and interpretation, or reporting of this study."

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Centralised randomisation using computerised minimization.
Allocation concealment (selection bias)	Low risk	Centralised, not possible to manipulate by participating centres
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Radiologist and microbiologist were blinded
Incomplete outcome data (attrition bias)	Low risk	No loss to follow-up, intention-to-treat analysis

Interventions for primary vesicoureteric reflux (Review)

Pennesi 2008 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Primary outcome appropriate, Secondary outcome detailed
Other bias	Unclear risk	24 parents refused, not clear why, might introduce selection bias

PRIVENT 2009

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from December 1998 to March 2007 Follow-up: 12 months
Participants	<ul style="list-style-type: none"> Country: Australia Setting: multicentre (4 sites) Children with ≥ 1 symptomatic UTI defined as symptoms consistent with UTI with positive urine culture, defined as any growth of a pathogenic organism from a suprapubic bladder tap, 10^7 CFU/L single organism from catheter or, 10^8 CFU/L midstream voided urine sample; all grades of VUR; recurrent infection; VUR from no VUR separable Number: treatment group (288); control group (288) Median age (months): treatment group (13.1); control group (14.5) Sex (M/F): treatment group (105/183); control group (102/186) Exclusion criteria: known neurologic, skeletal or urologic predisposition; known contra-indication to TMP/SMX therapy
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> Antibiotic: TMP/SMX <ul style="list-style-type: none"> Dose: 2 mg/10 mg/kg single daily dose Duration: 1 year <p>Control group</p> <ul style="list-style-type: none"> Placebo: matched for colour, taste and texture <ul style="list-style-type: none"> Dose: single daily dose Duration: 1 year
Outcomes	<ul style="list-style-type: none"> Recurrent symptomatic UTI (same definition as inclusion criteria) at 1 year Recurrent febrile UTI ($> 38^\circ\text{C}$) at 1 year New renal parenchymal abnormality (interpreted as any change on DMSA) at 1 year Deterioration renal parenchymal abnormality at 1 year Adverse events Recurrent UTI caused by micro-organism resistant to TMP/SMX
Notes	<ul style="list-style-type: none"> Secondary outcomes were extracted from raw data Funding source: "Supported by grants from the National Health and Medical Research Council of Australia (990735, 301999, and 402764) and from the Financial Markets Foundation for Children of Australia (058-2003) and by a private donation by J.T. Honan of the Manildra Group"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated sequence generation with stratification according to centre, referral source, frequency of previous UTI, VUR status, age, sex, according

Interventions for primary vesicoureteric reflux (Review)

PRIVENT 2009 (Continued)

		to method of minimization. Randomisation centrally by telephone by an independent trials centre
Allocation concealment (selection bias)	Low risk	Randomisation centrally by telephone by an independent trials centre
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding of all participants and healthcare providers
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding of outcome collectors and data-analysts
Incomplete outcome data (attrition bias) All outcomes	Low risk	Lost to follow-up small and accounted for (3/243; 1%), intention-to-treat analysis
Selective reporting (reporting bias)	Low risk	Primary outcome appropriate
Other bias	Low risk	Funding not commercial, patient flow from recruitment to randomisation clearly stated

Reddy 1997

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: not reported Duration of follow-up: 1 year
Participants	<ul style="list-style-type: none"> Country: USA Setting: single centre (university teaching hospital) Children with primary VUR (grade not reported); newly diagnosed Number: 43 Age: not reported Sex (M/F): not reported Exclusion criteria: not reported
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic prophylaxis <ul style="list-style-type: none"> Antibiotic: not reported Dose: not reported Duration: 1 year <p>Treatment group 2</p> <ul style="list-style-type: none"> Intermittent antibiotics: given on Monday, Wednesday and Friday <ul style="list-style-type: none"> Daily urine nitrate testing Duration: 1 year <p>Control group</p> <ul style="list-style-type: none"> No antibiotics

Reddy 1997 (Continued)

	<ul style="list-style-type: none"> Surveillance with daily urine nitrate
Outcomes	<ul style="list-style-type: none"> UTI Renal damage on DMSA Resolution of VUR
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Study was described as randomised, method of randomisation was not reported
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	Non-compliant patients were withdrawn from the study but included in analysis; intention-to-treat analysis performed
Selective reporting (reporting bias)	High risk	Primary outcome not appropriate. Both symptomatic and asymptomatic UTI reported together with no distinction between the two
Other bias	Unclear risk	Patent flow from recruitment to randomisation not at all reported

RIVUR 2008

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from June 2007 to May 2011 Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> Country: USA Setting: multicentre (19 sites) Children aged 2 to 72 months, with appropriately treated first or second UTI within 112 days before randomisation and VUR grade I to IV Number: treatment group (302); control group (305) Median age, IQR (months): treatment group (12, 5 to 31); control group (12, 6 to 30) Sex (M/F): treatment group (25/277); control group (24/281) Exclusion criteria: history of other renal injury/disease; congenital or acquired immunodeficiency; underlying anomalies or chronic diseases that could interfere with response to therapy; complex cardiac disease; syndromes associated with VUR or bladder dysfunction
Interventions	Treatment group

Interventions for primary vesicoureteric reflux (Review)

RIVUR 2008 (Continued)

- TMP/SMX: cherry-flavoured liquid suspension in which each 5 mL contains 200 mg sulfamethoxazole and 40 mg trimethoprim. Prophylactic dose is based on trimethoprim component: 3 mg/kg body weight taken once daily for 2 years

Control group

- Placebo: cherry-flavoured liquid suspension matched for taste, colour, odour and consistency to the active comparator. Bottled in identical 500 mL amber container high-density polyethylene containers with child-proof caps, administered for period of 2 years

Outcomes	<ul style="list-style-type: none"> • Recurrent febrile UTI • Recurrent non-febrile, symptomatic UTI • Development of new renal scar • Recurrent febrile or non-febrile symptomatic UTI caused by TMP/SMX resistant bacteria • Stool cultures of <i>E coli</i> resistant to TMP/SMX • Adherence
Notes	<ul style="list-style-type: none"> • Funding source: " Supported by grants from the NIDDK, NIH (U01 DK074059, U01 DK074053, U01 DK074082, U01 DK074064, U01 DK074062, and U01 DK074063); and by Clinical and Translational Science Awards from the National centre for Research Resources, now at the National centre for Advancing Translational Sciences, NIH (UL1RR024153 and UL1TR000005, to the University of Pittsburgh; and UL1TR000003, to the Children's Hospital of Philadelphia)"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Good quality method, Permuted blocks, stratified by site
Allocation concealment (selection bias)	Low risk	Externally managed randomisation and allocation, good quality method
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and clinicians blinded to treatment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Microbiological outcomes external so interpretation was blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	167 stopped treatment but were included in the denominator
Selective reporting (reporting bias)	Low risk	Very inclusive list of outcomes
Other bias	Low risk	Well designed and reported study

Roussey-Kesler 2008

Methods	• Study design: parallel RCT • Duration of study: recruitment from June 2001 to December 2004

Roussey-Kesler 2008 (Continued)

- Duration of follow-up: 1.5 years

Participants	<ul style="list-style-type: none"> • Country: France • Setting: multicentre (17 sites) • Children after a first episode of febrile UTI, with VUR grade I to III • Number: treatment group (103); control group (122) • Mean age $\pm$ SD (months): treatment group (12 ± 8.1); control group (10.6 ± 8.4) • Sex (M/F): treatment group (30/73); control group (39/83) • Exclusion criteria: abnormal renal echography; obstructive uropathy; high grade VUR; allergy to sulfamide
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> • Antibiotic: TMP/SMX <ul style="list-style-type: none"> ◦ Dose: 2/10 mg/kg single daily dose ◦ Duration: 18 months <p>Control group</p> <ul style="list-style-type: none"> • No treatment
Outcomes	<ul style="list-style-type: none"> • Repeat febrile UTI • Repeat symptomatic or asymptomatic afebrile UTI • Repeat UTI caused by <i>E coli</i> resistant to TMP/SMX
Notes	<ul style="list-style-type: none"> • Use of bag-urine in non-toilet trained children • Raw data obtained from authors • Funding source: "Supported by a Grant BRD 00/10-O from the biomedical research delegation from the University Hospital of Nantes"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Stratified block design according to centre and gender
Allocation concealment (selection bias)	Low risk	Centrally prepared sealed opaque envelopes in batches of 20 per gender
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up: 13/225 (6%) included in analysis; intention-to-treat analysis
Selective reporting (reporting bias)	Low risk	Primary outcome: Bag-urine was used however for non-toilet trained kids, hereby increasing risk of contamination in screening-samples
Other bias	Unclear risk	Recruitment process not clearly stated

Interventions for primary vesicoureteric reflux (Review)

Scott 1968

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: not reported Follow-up: 3 years
Participants	<ul style="list-style-type: none"> Country: UK Setting: single centre (university teaching hospital) Children with history of ≥ 1 UTI, with any grade VUR Number (randomised/analysed): treatment group 1 (25/24); treatment group 2 (33/23) Age: not reported Sex (M/F): treatment group 1 (3/21); treatment group 2 (4/19) Exclusion criteria: mechanical obstruction in the urethra, such as posterior valves; neurogenic bladder
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotics: according to microbial sensitivity Treatment after 'several' months of disease free survival. In majority it was given at least 6 months, and in those who developed recurrences of infection it was continued for a year or longer. Two were not infected at moment of enrolment and hence were not given antibiotics <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgery: Bischoff operation (plastic surgery of the ureteral meatus), ureteral reimplantation. not reported how many of each, only that in case when Bischoff had failed, subsequent reimplantation was successful and this case was included in analysis. 2 other cases were excluded after reappearance of VUR and 3 after first failed operation
Outcomes	<ul style="list-style-type: none"> Repeat symptomatic or asymptomatic UTI Pyelonephritic scarring - not outcome only at start of study Correction of VUR
Notes	<ul style="list-style-type: none"> Exclusions: 8/58 (14%), all in surgical group due to persistence of VUR. Loss to follow-up: 3/58 (5%) Some underwent bladder neck plastic operation in both groups: 9 in the antibiotics group, 11 in the operative group for non-obstructive bladder neck obstruction, which was later discarded as an entity altogether. Authors contacted for additional information Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Children were allotted at random to either the operated or control groups according to whether their birthdays fell on even or odd days.
Allocation concealment (selection bias)	High risk	Inadequate see above
Blinding of participants and personnel (performance bias) All outcomes	High risk	No blinding of participants, healthcare providers,
Blinding of outcome assessment (detection bias)	High risk	No blinding of outcome assessors or data-analysts

Scott 1968 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	High risk	Three cases were excluded because they could not be traced at the end of the 3 year follow-up period. Eight in the operation group were also excluded because their VUR had persisted
Selective reporting (reporting bias)	High risk	Symptomatic UTI not reported
Other bias	High risk	Differential chance of being in operation versus antibiotic group; patient flow from enrolment to randomisation not clear

Smellie 2001

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from November 1985 to December 1989 Duration of follow-up: 10 years
Participants	<ul style="list-style-type: none"> Country: UK Setting: single centre (urology and nephrology clinic of a university teaching hospital) Children with primary VUR, grade III to V, renal scarring on IVP Number (randomised/analysed): treatment group 1 (28/26); treatment group 2 (25/24) Mean age, range (years): 1 to 12 years treatment group 1 (47, 1.2 to 12.2); treatment group 2 (4.4, 1.0 to 11.7) Sex (M/F): treatment group 1 (13/14); treatment group 2 (12/13) Exclusion criteria: major urological abnormality
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic: nitrofurantoin, trimethoprim or TMP/SMX <ul style="list-style-type: none"> Dose: 1 to 2 mg/kg Variable duration of antibiotics <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgical: Cohen procedure Antibiotic duration: 6 months
Outcomes	<ul style="list-style-type: none"> UTI: culture positive Renal damage on IVP GFR (change) Renal length (change)
Notes	<ul style="list-style-type: none"> Funding source: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Stratified block randomisation according age and GFR with sealed envelopes containing equal number of cards for each treatment arm, with second investigator picking envelope.

Smellie 2001 (Continued)

Allocation concealment (selection bias)	Unclear risk	Due to small numbers in each stratum, sequence could be anticipated if recruiter knew the stratification method.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Patients and clinicians not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Radiologists blinded to treatment allocation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No loss to follow-up. No intention to treat, but only one exclusion.
Selective reporting (reporting bias)	Unclear risk	Definition of UTI: culture positive; method of collection not reported Secondary outcomes reported
Other bias	Unclear risk	Size of recruitment pool not reported but flow and decision making noted.

Swedish Reflux Trial 2010

Methods	<ul style="list-style-type: none"> Study design: parallel RCT Duration of study: recruitment from 2000 until 2006 Duration of follow-up: 2 years
Participants	<ul style="list-style-type: none"> Country: Sweden Setting: Multicentre (22 sites from county to university hospital covering 80% of Swedish population and 1 paediatric centre in Oslo) Grade III or IV VUR (graded according to International Reflux Study in children standards) at ages 1 to < 2 years; dilating VUR diagnosed before age of 1 year were eligible if repeat VCUG between age 1 and 2 showed grade III or IV Number: treatment group 1 (69); treatment group 2 (66); control group (68) Mean age $\pm$ SD (months; M/F): treatment group 1 ($1.66 \pm 0.27/1.75 \pm 0.28$); treatment group 2 ($1.67 \pm 0.32/1.71 \pm 0.35$); control group ($1.69 \pm 0.31/1.80 \pm 0.29$) Sex (M/F): 75/128 treatment group 1 (26/43); treatment group 2 (23/43); control group (26/42) Exclusion criteria: previous urogenital surgery; malformation except duplication; known urological disease; stone disease; GFR < 70 mL/min/1.73 m²; split kidney function < 15% or suspected non-compliance (inability to understand Swedish or previous non-compliance)
Interventions	<p>Treatment group 1</p> <ul style="list-style-type: none"> Antibiotic (choice at discretion of physician) <ul style="list-style-type: none"> Trimethoprim (primary choice): 0.5 to 1 mg/kg once daily Nitrofurantoin: 1 mg/kg once daily Cefadroxil: 5 mg/kg once daily <p>Treatment group 2</p> <ul style="list-style-type: none"> Surgery: endoscopic injection of Deflux. Each injection was followed by VCUG after three months Antibiotics: Concomitant prophylaxis until new VCUG showed absent VUR or improvement to grade I to II <p>Control group</p>

Interventions for primary vesicoureteric reflux (Review)

Swedish Reflux Trial 2010 (Continued)

- No specific treatment

All groups had run-in period before randomisation

- 2 weeks of TMP/SMX (2/10 mg/kg) single daily dose

Outcomes	<ul style="list-style-type: none"> • Febrile UTI during 2 year time period • Progression of DMSA uptake defects on initial examination or new damage reappearing during 2 year follow-up • VUR status after 2 year follow-up period; correction defined as VUR grade 0 and downgrading as VUR grade I or II • Microbial resistance: rate of UTI caused by resistant bacteria • Symptomatic UTI • Adverse events to endoscopic treatment • Bladder dysfunction
Notes	<ul style="list-style-type: none"> • Funding source: "Supported by Financial by grants from the West Region, Sweden and Futurum, Jönköping County Council, Sweden."

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Quote: "Children were randomly assigned by computer, matching for gender, previous UTI, VUR grade, DMSA uptake defect, bladder size, duplication and centre using minimisation procedures."</p> <p>Judgement: adequate method of sequence generation.</p>
Allocation concealment (selection bias)	Low risk	Patients were randomised immediately after consent form was signed, and allocation status could not be manipulated.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Open-label study, no blinding of participants or clinicians
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All radiological and nuclear medicine investigations were re-evaluated by the same specialist blinded to the treatment group
Incomplete outcome data (attrition bias) All outcomes	Low risk	All data analysed according to intention to treat with censoring of patients with incomplete follow-up in survival analysis
Selective reporting (reporting bias)	Unclear risk	Primary and secondary outcomes appropriate
Other bias	High risk	<p>Sponsor bias: One of the authors states a financial interest with the manufacturer of the Deflux used in the endoscopic treatment</p> <p>Patient flow from enrolment to randomisation not clear</p>

Takvani 2015

Methods	<ul style="list-style-type: none"> • Study design: parallel RCT
---------	--

Interventions for primary vesicoureteric reflux (Review)

Takvani 2015 (Continued)

	<ul style="list-style-type: none"> Duration of study: not reported Duration of follow-up: not reported
Participants	<ul style="list-style-type: none"> Country: India Setting; multicentre (3 sites) Children with antenatal hydronephrosis and low grade VUR Number: unclear Mean age $\pm$ SD: not reported Sex (M/F): not reported VUR grades; I and II Exclusion criteria: high grade VUR and children who'd had febrile UTI and had VUR
Interventions	Treatment group <ul style="list-style-type: none"> Antibiotic prophylaxis (no type, dose or duration reported) Control group <ul style="list-style-type: none"> No treatment
Outcomes	<ul style="list-style-type: none"> Resolution of VUR DMSA changes
Notes	<ul style="list-style-type: none"> Abstract-only publication

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to permit judgement
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Difficult to follow results, poorly reported
Selective reporting (reporting bias)	High risk	Not the most appropriate outcomes
Other bias	Unclear risk	Poorly reported methodological detail

CFU - colony forming units; DMSA scan - 99m-technetium dimercaptosuccinic acid scan; Deflux - dextranomer/hyaluronic acid copolymer; ESKD - end-stage kidney disease; GAX 65 and 35: glutaraldehyde cross-linked collagen with 65 mg/mL and 35 mg/mL of collagen; (e)GFR - (estimated) glomerular filtration rate; IVP - Intravenous pyelogram; IQR - interquartile range; M/F - male/female; PUJ - pelvi-ureteric junction; RCT - randomised controlled trial; Reimplantation - PL (Politano-Leadbetter procedure), Cohen (Cohen procedures), LG (Lich-

Gregoir procedure); SD - standard deviation; TMP/SMX - trimethoprim/sulphamethoxazole; UTI - urinary tract infection; VCUG - voiding cystourethrogram; VUR - vesicoureteric reflux

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Akin 2014	Doubtful that this is a RCT, conflicting details in methods
Bailey 1991	Wrong population: significant abnormality, reflux nephropathy and undergoing nephrectomy
COBSG 1973	Wrong/mixed population: inclusion of children with and without VUR, separate numbers for children with VUR not extractable
Coupris 1988	Wrong intervention: uncertain what comparative treatment was, only states urinary antiseptic
Kajbafzadeh 2010	Wrong population: all patients had primary bladder neck dysfunction, a urological abnormality
Montini 2007	Wrong population: RCT comparing early oral versus intravenous antibiotics for acute treatment of UTI in VUR patients
N0012116994	Study discontinued due to insufficient recruitment numbers and excessive refusal to participate
NCBRG 1981	Wrong/mixed population: inclusion of children with and without VUR, separate numbers for children with VUR not extractable
Scholtmeijer 1988	Unable to differentiate randomised from non-randomised patients
Uberos 2012	Wrong population: majority of participants do not have VUR

RCT - randomised controlled trial; UTI - urinary tract infection; VUR - vesicoureteric reflux

DATA AND ANALYSES

Comparison 1. Antibiotic prophylaxis versus placebo or no treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Urinary tract infection	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Symptomatic UTI by 1-2 years: antibiotics versus no prophylaxis	9	1667	Risk Ratio (M-H, Random, 95% CI)	0.77 [0.54, 1.09]
1.2 Febrile UTI by 1-2 years: antibiotics versus no prophylaxis	9	1667	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.56, 1.21]
1.3 Symptomatic UTI: antibiotics versus placebo in studies with low risk of bias across all domains	3	943	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.40, 1.79]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.4 Repeat positive urine cultures by 1-3 years: continuous antibiotics versus no prophylaxis	7	729	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.63, 1.02]
1.5 Repeat positive urine cultures by 1-3 years: intermittent antibiotics versus no prophylaxis	1	30	Risk Ratio (M-H, Random, 95% CI)	0.46 [0.10, 2.00]
2 All symptomatic UTI by 1-2 years: allocation concealment	8		Risk Ratio (IV, Random, 95% CI)	Subtotals only
2.1 Adequate allocation concealment	7	1533	Risk Ratio (IV, Random, 95% CI)	0.73 [0.50, 1.08]
2.2 Unclear allocation concealment	1	113	Risk Ratio (IV, Random, 95% CI)	7.38 [0.94, 58.07]
3 All symptomatic UTI by 1-2 years: blinding	8		Risk Ratio (IV, Random, 95% CI)	Subtotals only
3.1 Blinded studies	3	943	Risk Ratio (IV, Random, 95% CI)	0.85 [0.40, 1.78]
3.2 Non-blinded studies	5	703	Risk Ratio (IV, Random, 95% CI)	0.81 [0.43, 1.53]
4 Renal parenchymal abnormality on DMSA scan: unit of analysis (children)	10		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
4.1 New renal abnormality by 1-3 years	8	1503	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.33, 1.61]
4.2 Deterioration of existing abnormality by 1-2 years	3	446	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.27, 1.73]
4.3 New and deteriorated abnormalities on DMSA scan by 1-3 years	7	1288	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.42, 1.62]
4.4 All abnormalities detected on DMSA scan by 1-3 years: intermittent antibiotics versus surveillance	1	30	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.02, 8.59]
5 Resolution of VUR	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
5.1 Resolution of VUR after 1-2 years: unit of analysis (children)	3	262	Risk Ratio (M-H, Random, 95% CI)	1.46 [0.71, 2.99]
5.2 Resolution of VUR after 2 years: unit of analysis (ureters)	1	266	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.59, 1.09]
6 Adverse events	4	1056	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.81, 1.08]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7 Microbial resistance to prophylactic drug	6		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
7.1 Microbial resistance to prophylactic drug: unit of analysis (UTI)	6	259	Risk Ratio (M-H, Random, 95% CI)	2.72 [1.85, 4.00]
8 GFR at 3 years	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
9 Renal growth at 3 years	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
10 All symptomatic UTI by 1-2 years: addressing of incomplete outcome data	8	1646	Risk Ratio (IV, Random, 95% CI)	0.81 [0.53, 1.24]
10.1 Adequate addressing of incomplete outcome data	7	1533	Risk Ratio (IV, Random, 95% CI)	0.73 [0.50, 1.08]
10.2 Inadequate addressing of incomplete outcome data	1	113	Risk Ratio (IV, Random, 95% CI)	7.38 [0.94, 58.07]
11 Febrile UTI by 1-2 years: allocation concealment	8	1646	Risk Ratio (IV, Random, 95% CI)	0.86 [0.57, 1.30]
11.1 Adequate allocation concealment	7	1533	Risk Ratio (IV, Random, 95% CI)	0.79 [0.54, 1.15]
11.2 Unclear allocation concealment	1	113	Risk Ratio (IV, Random, 95% CI)	7.38 [0.94, 58.07]
12 Febrile UTI by 1-2 years: blinding	7	1421	Risk Ratio (IV, Random, 95% CI)	0.89 [0.54, 1.46]
12.1 Blinded studies	3	943	Risk Ratio (IV, Random, 95% CI)	0.96 [0.47, 1.96]
12.2 Non-blinded studies	4	478	Risk Ratio (IV, Random, 95% CI)	0.88 [0.39, 1.97]
13 Febrile UTI by 1-2 years: addressing of incomplete outcome data	8	1610	Risk Ratio (IV, Random, 95% CI)	1.04 [0.60, 1.80]
13.1 Adequate addressing of incomplete outcome data	7	1497	Risk Ratio (IV, Random, 95% CI)	0.92 [0.54, 1.58]
13.2 Inadequate addressing of incomplete outcome data	1	113	Risk Ratio (IV, Random, 95% CI)	7.38 [0.94, 58.07]
14 Febrile UTI by 1-2 years according to gender	2	237	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.36, 1.90]
14.1 UTI in girls	2	137	Risk Ratio (M-H, Random, 95% CI)	0.63 [0.17, 2.33]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
14.2 UTI in boys	2	100	Risk Ratio (M-H, Random, 95% CI)	1.26 [0.53, 2.99]

Analysis 1.1. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 1 Urinary tract infection.

Analysis 1.2. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 2 All symptomatic UTI by 1-2 years: allocation concealment.

Analysis 1.3. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 3 All symptomatic UTI by 1-2 years: blinding.

Analysis 1.4. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 4 Renal parenchymal abnormality on DMSA scan: unit of analysis (children).

Analysis 1.5. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 5 Resolution of VUR.

Analysis 1.6. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 6 Adverse events.

Analysis 1.7. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 7 Microbial resistance to prophylactic drug.

Analysis 1.8. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 8 GFR at 3 years.

Analysis 1.9. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 9 Renal growth at 3 years.

Analysis 1.10. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 10 All symptomatic UTI by 1-2 years: addressing of incomplete outcome data. **Analysis 1.11. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 11 Febrile UTI by 1-2 years: allocation concealment.**

Analysis 1.12. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 12 Febrile UTI by 1-2 years: blinding.

Analysis 1.13. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 13 Febrile UTI by 1-2 years: addressing of incomplete outcome data.

Analysis 1.14. Comparison 1 Antibiotic prophylaxis versus placebo or no treatment, Outcome 14 Febrile UTI by 1-2 years according to gender.

Comparison 2. Surgical reimplantation of ureters plus antibiotics versus antibiotics alone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Urinary tract infection	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 All symptomatic UTI by 4-5 years: surgical correction + antibiotics versus antibiotics alone	1	297	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.67, 1.35]
1.2 All symptomatic UTI between 5-10 years: surgical correction + antibiotics versus antibiotics alone	1	252	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.49, 1.26]
1.3 All symptomatic UTI in children followed for 10 years: surgical correction + antibiotics versus antibiotic alone	1	252	Risk Ratio (M-H, Random, 95% CI)	1.06 [0.78, 1.44]
1.4 Febrile UTI by 5 years: surgical correction + antibiotics versus antibiotics alone	2	429	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.27, 0.70]
1.5 Febrile UTI between 5-10 years: surgical correction + antibiotics versus antibiotics alone	1	252	Risk Ratio (M-H, Random, 95% CI)	0.34 [0.14, 0.82]
1.6 Febrile UTI in children followed for 10 years: surgical correction + antibiotics versus antibiotics alone	1	252	Risk Ratio (M-H, Random, 95% CI)	0.54 [0.32, 0.92]
1.7 Repeat positive urine cultures by 4-5 years: surgical correction + antibiotics versus antibiotics alone	3	479	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.79, 1.26]
2 Renal parenchymal defects (scars and thinning) on IVP: unit of analysis (children)	5		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 New defects at 2 years: surgical correction + antibiotics versus antibiotics alone	2	171	Risk Ratio (M-H, Random, 95% CI)	1.06 [0.33, 3.42]
2.2 New defects (scars and thinning of parenchyma) at 4-5 years: surgical correction + antibiotics versus antibiotics alone	4	572	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.79, 1.49]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.3 Progression of existing defects at 2 years: surgical correction + antibiotics versus antibiotics alone	1	10	Risk Ratio (M-H, Random, 95% CI)	7.0 [0.45, 108.26]
2.4 Progression of existing defects (scars and parenchymal thinning) at 4-5 years: surgical correction + antibiotics versus antibiotics alone	3	468	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.69, 1.42]
2.5 Total new and progressive renal parenchymal defects at 2 years: surgical correction + antibiotics versus antibiotics alone	1	10	Risk Ratio (M-H, Random, 95% CI)	9.00 [0.61, 133.08]
2.6 Total new and progressive renal parenchymal defects at 4-5 years: surgical correction + antibiotics versus antibiotics alone	3	468	Risk Ratio (M-H, Random, 95% CI)	1.05 [0.85, 1.29]
3 Renal scars on IVP: unit of analysis (children)	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3.1 New scars at 4-5 years: surgical correction + antibiotics versus antibiotics alone	2	418	Risk Ratio (M-H, Random, 95% CI)	1.28 [0.84, 1.94]
3.2 New scars developing by 5-10 years in children followed for 10 years: surgical correction + antibiotics versus antibiotics alone	1	223	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.07, 16.22]
3.3 New scars at 10 years in children followed for 10 years: surgical correction + antibiotics versus antibiotics alone	1	223	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.53, 2.00]
4 Repeat positive urine culture	4	328	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.55, 1.21]
5 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children)	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
5.1 All abnormalities detected on DMSA at 5 years: surgical correction + antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
5.2 Deterioration of existing abnormalities between 5-10 years: surgical correction + antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
6 Renal parenchymal defects on IVP: unit of analysis (kidneys)	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
6.1 New defects at 2 years: surgical correction + antibiotics versus antibiotics alone	2	235	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.31, 3.37]
6.2 Progression of existing defects at 2 years: surgical correction + antibiotics versus antibiotics alone	2	235	Risk Ratio (M-H, Random, 95% CI)	1.56 [0.24, 10.08]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
6.3 Total new and progressive defects at 2 years: surgical correction + antibiotics versus antibiotics alone	2	235	Risk Ratio (M-H, Random, 95% CI)	1.54 [0.24, 9.95]
6.4 New defects at 4-5 years: surgical correction + antibiotics versus antibiotics alone	2	319	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.24, 3.09]
6.5 Progression of existing defects at 4-5 years: surgical correction + antibiotics versus antibiotics alone	2	319	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.50, 1.41]
6.6 Total new and progressive defects at 4-5 years: surgical correction + antibiotics versus antibiotics alone	2	319	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.53, 1.34]
7 Hypertension and ESKD	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
7.1 ESKD	2	154	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.23, 5.04]
7.2 Hypertension	2	154	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.25, 3.42]
7.3 Hypertension at 10 years	1	252	Risk Ratio (M-H, Random, 95% CI)	0.15 [0.01, 2.78]
8 GFR measured by Schwartz formula	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
8.1 GFR at entry: surgical correction + antibiotics versus antibiotics alone	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
8.2 GFR at 5 years: surgical correction + antibiotics versus antibiotics alone	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
8.3 GFR at 10 years: surgical correction + antibiotics versus antibiotics alone	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
9 Height Standard Deviation Score (SDS)	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
9.1 Height SDS at entry for surgical correction + antibiotics versus antibiotics alone	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
9.2 Height SDS at 10 year follow-up for surgical correction + antibiotics versus antibiotics alone	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
10 Correction of VUR: unit of analysis (children)	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
10.1 Correction of VUR after 3 years: surgical correction + antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
11 Correction of VUR: unit of analysis (ureters)	2		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
11.1 Correction of VUR after 2 years: surgical correction + antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
11.2 Correction of VUR at 5 years: surgical correction + antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Analysis 2.1. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 1 Urinary tract infection.

Analysis 2.2. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 2 Renal parenchymal defects (scars and thinning) on IVP: unit of analysis (children).

Analysis 2.3. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 3 Renal scars on IVP: unit of analysis (children).

Analysis 2.4. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 4 Repeat positive urine culture.

Analysis 2.5. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 5 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children).

Analysis 2.6. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 6 Renal parenchymal defects on IVP: unit of analysis (kidneys).

Analysis 2.7. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 7 Hypertension and ESKD.

Analysis 2.8. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 8 GFR measured by Schwartz formula.

Analysis 2.9. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 9 Height Standard Deviation Score (SDS).

Analysis 2.10. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 10 Correction of VUR: unit of analysis (children).

Analysis 2.11. Comparison 2 Surgical reimplantation of ureters plus antibiotics versus antibiotics alone, Outcome 11 Correction of VUR: unit of analysis (ureters).

Comparison 3. Surgical reimplantation type 1 versus surgical reimplantation type 2

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Repeat positive urine culture	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2 Persistent VUR (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3 Peri-operative complications	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only

Analysis 3.1. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 1 Repeat positive urine culture.

Study or subgroup	Advancement n/N	Std surgery n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
Kojima 2012	0/15	1/15		0%	0.33[0.01,7.58]
Less with advancement Less with Std surgery					

Analysis 3.2. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 2 Persistent VUR (ureters).

Study or subgroup	Advancement n/N	Std surgery n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
Kojima 2012	0/25	4/26		0%	0.12[0.01,2.04]
Less with advancement Less with Std surgery					

Analysis 3.3. Comparison 3 Surgical reimplantation type 1 versus surgical reimplantation type 2, Outcome 3 Peri-operative complications.

Study or subgroup	Advancement n/N	Std surgery n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
Kojima 2012	1/15	0/15		0%	3[0.13,68.26]
Less with advancement Less with Std surgery					

Comparison 4. Endoscopic injection treatment versus antibiotics alone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Urinary tract infection	4		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 All symptomatic UTI by 1-2 year: endoscopic correction + antibiotics versus antibiotics alone	3	254	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.37, 1.75]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 Febrile UTI by 1-2 year: endoscopic correction + antibiotics versus antibiotics alone	3	254	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.31, 1.78]
1.3 Repeat positive urine culture by 1 year	1	60	Risk Ratio (M-H, Random, 95% CI)	7.15 [0.42, 121.04]
1.4 Febrile UTI by 1-2 years in girls	1	86	Risk Ratio (M-H, Random, 95% CI)	1.25 [0.55, 2.86]
1.5 Febrile UTI by 1-2 years in boys	1	49	Risk Ratio (M-H, Random, 95% CI)	2.26 [0.46, 11.22]
2 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children)	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Total new and progressive abnormalities at 1-2 years: endoscopic correction + antibiotics versus antibiotics alone	2	208	Risk Ratio (M-H, Random, 95% CI)	1.20 [0.36, 3.96]
2.2 Deterioration of existing abnormalities at 2 years: endoscopic correction + antibiotics versus antibiotics alone	1	133	Risk Ratio (M-H, Random, 95% CI)	2.09 [0.66, 6.61]
2.3 New abnormalities at 2 years: endoscopic correction + antibiotics versus antibiotics alone	1	133	Risk Ratio (M-H, Random, 95% CI)	11.5 [0.65, 203.91]
3 Renal damage on ultrasound	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
3.1 New renal parenchymal defect at 1 year for endoscopic correction + antibiotics versus antibiotics alone: unit of analysis (children)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.2 New renal scar at 1 year for endoscopic correction + antibiotics versus antibiotics alone: unit of analysis (kidneys)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4 VUR status after treatment: unit of analysis (children)	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
4.1 Resolution of VUR (grade 0) for endoscopic correction + antibiotics versus antibiotics alone	2	164	Risk Ratio (M-H, Random, 95% CI)	2.69 [1.57, 4.63]
4.2 Improvement of VUR grade for endoscopic injection compared to Antibiotics	3	239	Risk Ratio (M-H, Random, 95% CI)	2.01 [1.51, 2.68]
5 Microbial resistance to the prophylactic drug	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
5.1 Unit of analysis (febrile UTI)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Analysis 4.1. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 1 Urinary tract infection.

Analysis 4.2. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 2 Renal parenchymal abnormalities on DMSA scan: unit of analysis (children).

Analysis 4.3. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 3 Renal damage on ultrasound.

Analysis 4.4. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 4 VUR status after treatment: unit of analysis (children).

Analysis 4.5. Comparison 4 Endoscopic injection treatment versus antibiotics alone, Outcome 5 Microbial resistance to the prophylactic drug.

Comparison 5. Endoscopic injection treatment versus no treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Symptomatic urinary tract infection	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
2 Febrile urinary tract infection	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
3 Renal parenchymal abnormality	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected

Analysis 5.1. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 1 Symptomatic urinary tract infection.

Analysis 5.2. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 2 Febrile urinary tract infection.

Analysis 5.3. Comparison 5 Endoscopic injection treatment versus no treatment, Outcome 3 Renal parenchymal abnormality.

Comparison 6. Different materials for endoscopic injection to correct VUR

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Macroplastique versus dextranomer/hyaluronic acid polymer (Deflux)	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Febrile UTI	1	255	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.45, 2.66]
1.2 Number of children with afebrile UTI	1	72	Risk Ratio (M-H, Random, 95% CI)	1.68 [0.52, 5.44]
1.3 Number of ureters with persistent VUR > grade 1 at 3 months	2	513	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.33, 0.78]
1.4 Number of ureters with persistent VUR > grade 1 at 1 year	2	472	Risk Ratio (M-H, Random, 95% CI)	0.54 [0.35, 0.83]
1.5 Number of kidneys with temporary renal dilatation after injection	1	114	Risk Ratio (M-H, Random, 95% CI)	1.85 [1.02, 3.35]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.6 Ureteral obstruction	1	399	Risk Ratio (M-H, Random, 95% CI)	1.46 [0.25, 8.66]
1.7 Post-operative flank pain	1	255	Risk Ratio (M-H, Random, 95% CI)	1.45 [0.33, 6.36]
2 GAX 65 versus GAX 35	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
2.1 Number of ureters with persistent VUR at 3 months	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2.2 Number of ureters with recurrence of VUR	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3 Dexcel versus Vantris	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
3.1 Repeat positive urine culture	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.2 Refluxing ureters after treatment	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.3 Children with fever after treatment	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.4 Ureteral obstruction	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.5 Elimination, or improvement in grade, of VUR (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Analysis 6.1. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 1 Macroplastique versus dextranomer/hyaluronic acid polymer (Deflux).

Analysis 6.2. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 2 GAX 65 versus GAX 35.

Analysis 6.3. Comparison 6 Different materials for endoscopic injection to correct VUR, Outcome 3 Dexcel versus Vantris.

Comparison 7. Probiotics versus antibiotic prophylaxis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Urinary tract infection	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 All symptomatic UTI by 1 year	2	248	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.56, 1.21]
1.2 Febrile UTI by 1 year	1	120	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.37, 1.83]
2 E coli resistance to prophylactic drug	2	38	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.21, 0.69]
3 Renal parenchymal defects on DMSA scan	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3.1 Patient data at 1 year	2	152	Risk Ratio (M-H, Random, 95% CI)	0.69 [0.43, 1.11]
4 Probiotics plus antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
4.1 Repeat febrile UTI	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4.2 Repeat positive urine culture	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.3 New DMSA changes	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.4 Microbial resistance, year 1	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.5 Microbial resistance, year 2	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.6 Microbial resistance, year 3	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.7 Adverse effects	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Analysis 7.1. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 1 Urinary tract infection.

Analysis 7.2. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 2 *E coli* resistance to prophylactic drug.

Analysis 7.3. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 3 Renal parenchymal defects on DMSA scan.

Analysis 7.4. Comparison 7 Probiotics versus antibiotic prophylaxis, Outcome 4 Probiotics plus antibiotics versus antibiotics alone.

Comparison 8. Other interventions

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Reimplantation surgery versus endoscopic treatment	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Repeat febrile UTI	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.2 VUR resolution at 6 months (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.3 VUR resolution at 12 months (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.4 VUR resolution at 5 years (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.5 VUR resolution after 1 procedure (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.6 Complications from the procedure	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2 Cranberry product versus no treatment	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
2.1 Repeat UTI	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3 Circumcision plus antibiotics versus antibiotics alone	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
3.1 Repeat symptomatic UTI	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.2 Repeat positive urine culture	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.3 DMSA changes at 12 months	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4 Oxybutynin versus placebo	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
4.1 Resolution of VUR	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.2 Improved grade of VUR	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.3 Resolved or improved grade of VUR (ureters)	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Analysis 8.1. Comparison 8 Other interventions, Outcome 1 Reimplantation surgery versus endoscopic treatment.

Analysis 8.2. Comparison 8 Other interventions, Outcome 2 Cranberry product versus no treatment.

Analysis 8.3. Comparison 8 Other interventions, Outcome 3 Circumcision plus antibiotics versus antibiotics alone.

Analysis 8.4. Comparison 8 Other interventions, Outcome 4 Oxybutynin versus placebo.

APPENDICES

Appendix 1. Electronic search strategies

Database	Search terms
CENTRAL	<ol style="list-style-type: none"> 1. (Vesico-Ureter* Reflux):ti,ab,kw or (Vesico* Reflux):ti,ab,kw or (ureter* reflux):kw in Clinical Trials 2. (vesic* or ureter*) and (backflow* or reflux):ti,ab,kw in Clinical Trials 3. (vur):ti,ab,kw or (VUR):ti,ab,kw in Clinical Trials 4. MeSH descriptor Vesico-Ureteral Reflux explode all trees 5. (#1 OR #2 OR #3 OR #4)
MEDLINE	<ol style="list-style-type: none"> 1. Vesico-Ureteral Reflux/ 2. (Vesico-Ureter\$ Reflux or Vesico\$ Reflux or ureter\$ reflux).tw. 3. ((vesic\$ or ureter\$) and (backflow\$ or reflux)).tw. 4. vur.tw. 5. or/1-4
EMBASE	<ol style="list-style-type: none"> 1. Vesicoureteral Reflux/ 2. (vesico-ureter\$ reflux or vesico\$ reflux or ureter\$ reflux).tw. 3. ((vesico\$ or ureter\$) and (backflow\$ or reflux)).tw. 4. vur.tw. 5. or/1-4

Appendix 2. Risk of bias assessment tool

Potential source of bias	Assessment criteria
Was there adequate sequence generation?	Yes (<i>low risk of bias</i>): Random number table; computer random number generator; coin tossing; shuffling cards or envelopes; throwing dice; drawing of lots; minimization (minimization may be implemented without a random element, and this is considered to be equivalent to being random).

(Continued)

	<p><i>No (high risk of bias):</i> Sequence generated by odd or even date of birth; date (or day) of admission; sequence generated by hospital or clinic record number; allocation by judgement of the clinician; by preference of the participant; based on the results of a laboratory test or a series of tests; by availability of the intervention.</p> <hr/> <p><i>Unclear:</i> Insufficient information about the sequence generation process to permit judgement.</p>
Was allocation adequately concealed?	<p><i>Yes (low risk of bias):</i> Randomisation method described that would not allow investigator/participant to know or influence intervention group before eligible participant entered in the study (e.g. central allocation, including telephone, web-based, and pharmacy-controlled, randomisation; sequentially numbered drug containers of identical appearance; sequentially numbered, opaque, sealed envelopes).</p> <hr/> <p><i>No (high risk of bias):</i> Using an open random allocation schedule (e.g. a list of random numbers); assignment envelopes were used without appropriate safeguards (e.g. if envelopes were unsealed or non-opaque or not sequentially numbered); alternation or rotation; date of birth; case record number; any other explicitly unconcealed procedure.</p> <hr/> <p><i>Unclear:</i> Randomisation stated but no information on method used is available.</p>
Was knowledge of the allocated interventions adequately prevented during the study?	<p><i>Yes (low risk of bias):</i> No blinding, but the review authors judge that the outcome and the outcome measurement are not likely to be influenced by lack of blinding; blinding of participants and key study personnel ensured, and unlikely that the blinding could have been broken; either participants or some key study personnel were not blinded, but outcome assessment was blinded and the non-blinding of others unlikely to introduce bias.</p> <hr/> <p><i>No (high risk of bias):</i> No blinding or incomplete blinding, and the outcome or outcome measurement is likely to be influenced by lack of blinding; blinding of key study participants and personnel attempted, but likely that the blinding could have been broken; either participants or some key study personnel were not blinded, and the non-blinding of others likely to introduce bias.</p> <hr/> <p><i>Unclear:</i> Insufficient information to permit judgement of 'Yes' or 'No'</p>
Were incomplete outcome data adequately addressed?	<p><i>Yes (low risk of bias):</i> No missing outcome data; reasons for missing outcome data unlikely to be related to true outcome (for survival data, censoring unlikely to be introducing bias); missing outcome data balanced in numbers across intervention groups, with similar reasons for missing data across groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk not enough to have a clinically relevant impact on the intervention effect estimate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes not enough to have a clinically relevant impact on observed effect size; missing data have been imputed using appropriate methods.</p> <hr/> <p><i>No (high risk of bias):</i> Reason for missing outcome data likely to be related to true outcome, with either imbalance in numbers or reasons for missing data across intervention groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk enough to induce clinically relevant bias in intervention effect estimate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes enough to induce clinically relevant bias in observed effect size; 'as-treated' analysis done with substantial departure of the intervention received from that assigned at randomisation; potentially inappropriate application of simple imputation.</p> <hr/> <p><i>Unclear:</i> Insufficient information to permit judgement of 'Yes' or 'No'.</p>
Are reports of the study free of suggestion of selective outcome reporting?	<p><i>Yes (low risk of bias):</i> The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way; the study protocol is not available but it is clear that the published reports include all expected outcomes, including those that were pre-specified (convincing text of this nature may be uncommon).</p>

(Continued)

No (high risk of bias): Not all of the study's pre-specified primary outcomes have been reported; one or more primary outcomes is reported using measurements, analysis methods or subsets of the data (e.g. subscales) that were not pre-specified; one or more reported primary outcomes were not pre-specified (unless clear justification for their reporting is provided, such as an unexpected adverse effect); one or more outcomes of interest in the review are reported incompletely so that they cannot be entered in a meta-analysis; the study report fails to include results for a key outcome that would be expected to have been reported for such a study.

Unclear: Insufficient information to permit judgement of 'Yes' or 'No'.

Was the study apparently free of other problems that could put it at a risk of bias?

Yes (low risk of bias): The study appears to be free of other sources of bias.

No (high risk of bias): Had a potential source of bias related to the specific study design used; stopped early due to some data-dependent process (including a formal-stopping rule); had extreme baseline imbalance; has been claimed to have been fraudulent; had some other problem.

Unclear: Insufficient information to permit judgement of 'Yes' or 'No'.

FEEDBACK

Grade of Reflux

Summary

It is difficult to accept the fact that Surgery does not help. There are specific indications for surgery especially in the realm of paediatric VUR. A more appropriate study would have been to compare the two groups (surgery vs conservative) in a similar grade of reflux and as such the controversy exists only in Gr3 reflux.

Regarding the assessment of children with UTI it is clear cut that the first investigation will be Ultrasound examination Followed by MCU.

Reply

Response to Dr Philipraj

This systematic review of randomised controlled trials has summarised the results of published and unpublished trials identified by a comprehensive search of literature sources. The majority of published trials have compared surgery and antibiotic prophylaxis with antibiotic prophylaxis. The large trials (Birmingham Reflux Study, International Reflux Study) only enrolled children with dilating reflux - equivalent to grade 3 or more on the International Classification. The International Reflux Study only enrolled children with grade 3 and 4 reflux; children with grade 5 reflux were excluded. There were no trials identified that compared surgery and antibiotic prophylaxis only and only enrolled children with grade 3 reflux. When data from these trials are combined in meta-analysis, there were no significant differences in the risk for further urinary tract infection or for renal scarring. In the International Reflux Study, the incidence of febrile urinary tract infections over 5 years of follow up was significantly reduced in children undergoing surgery compared with antibiotic prophylaxis. This was the only benefit of surgery over antibiotic prophylaxis that could be demonstrated. Otherwise the available trials have not demonstrated an additional benefit of surgery over antibiotic prophylaxis.

This systematic review of treatment cannot provide any information on the most appropriate investigations for children following urinary tract infection.

Contributors

Dr S. Joseph Philipraj
Urologist
padisahana@yahoo.com
Dept of Urology, Dr.T.M.A.Pai hospital, Kasturba Medical college, Udupi-576101, Karnataka, India.
Date Received: 27/08/2004

WHAT'S NEW

Date	Event	Description
10 October 2018	New citation required but conclusions have not changed	Summary of findings tables added
10 October 2018	New search has been performed	14 new studies included

HISTORY

Protocol first published: Issue 2, 1999

Review first published: Issue 3, 2004

Date	Event	Description
13 May 2009	Amended	Contact details updated.
9 October 2008	Amended	Converted to new review format.
15 May 2007	New citation required and conclusions have changed	Substantive amendment

CONTRIBUTIONS OF AUTHORS

- DW: Protocol development, study eligibility, data extraction, data analysis, writing review
- DV: Protocol development, study eligibility, data extraction
- EH: Data extraction, data analysis, writing review
- GS: Protocol development, data extraction, writing review
- GW: Data extraction, review updates
- EN: Data extraction, data analysis, writing review
- JC: Protocol development, writing review

DECLARATIONS OF INTEREST

- JC Craig, GJ Williams and E Hodson are authors of the [PRIVENT 2009](#)
- JC Craig is an author of [Craig 2002](#).

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- Australian Kidney Foundation, Seeding Grant number S2/99, Australia.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

The risk of bias assessment tool ([Appendix 2](#)) has replaced the quality checklist.

INDEX TERMS

Medical Subject Headings (MeSH)

Acrylic Resins [therapeutic use]; Anti-Bacterial Agents [therapeutic use]; Antibiotic Prophylaxis; Collagen [therapeutic use]; Dextrans [therapeutic use]; Dimethylpolysiloxanes [therapeutic use]; Fever [etiology] [prevention & control]; Hyaluronic Acid [therapeutic

use]; Kidney [abnormalities]; Probiotics [therapeutic use]; Randomized Controlled Trials as Topic; Replantation; Ureter [surgery]; Urinary Tract Infections [complications] [*prevention & control]; Vesico-Ureteral Reflux [complications] [*therapy]

MeSH check words

Child; Female; Humans; Male